

# P02 Grid Partition V3

Laureano Arcanio (February 2026)

## Problem

Grace and Henry are observing a square grid divided into  $5041 \times 5041$  unit cells arranged in rows and columns. Each cell contains either a *star* or a *moon*, arranged in a checkerboard pattern so that any two cells sharing a side contain different symbols.

## Solution

### Step 1: Balance is automatic

Along any allowed trail, each step moves to a neighboring plot. Because the grid is colored in a checkerboard pattern, each step switches the type of symbol encountered. Thus the symbols along a trail strictly alternate.

If a trail contains an even number of cells, it visits the same number of stars and moons. If it contains an odd number of plots, the difference is exactly one. Hence every allowed trail is automatically balanced, and the balance condition imposes no additional restriction.

Therefore, the problem reduces to determining the minimum number of allowed trails needed to cover all plots.

### Step 2: An explicit construction

Grace assigns one trail to each column of the field. Each trail starts at the top plot of its column and proceeds straight downward through all 5041 plots in that column.

These 5041 trails:

- obey the movement rules,
- are balanced by Step 1, and
- together cover every plot exactly once.

Thus  $M \leq 5041$ .

### Step 3: Why fewer trails are impossible

To show that Bob cannot do better, Henry argues as follows.

Each plot is divided along its diagonal from the upper-left corner to the lower-right corner, producing two triangular half-plots. Think of these diagonals as mirrors.

Light beams are sent into the field through the midpoints of all boundary edges, perpendicular to the boundary. There are exactly  $4 \cdot 5041$  such entry points. Each beam reflects

off the diagonal mirrors and eventually exits the boundary, thereby pairing the boundary edges into  $2 \cdot 5041$  disjoint pairs.

Consider how these beams interact with Bob's trails. Whenever a beam passes through a region corresponding to a segment of a trail, it exits that region on the same type of boundary edge (horizontal or vertical) on which it entered. A beam can change from a horizontal boundary to a vertical boundary, or vice versa, only by passing through a triangular half-plot that is not paired with another half-plot by the trail structure.

Each trail has exactly two endpoints. Each endpoint creates exactly one such unpaired half-plot. Therefore the total number of unpaired half-plots is exactly  $2M$ .

On the other hand, among the  $2 \cdot 5041$  boundary pairs, at least 5041 pairs must connect a horizontal boundary edge with a vertical boundary edge. Each such mixed pair forces a beam to pass through at least one unpaired half-plot.

Hence

$$5041 \leq 2M.$$

Repeating the same argument after shifting all beam entry points by half a unit along the boundary produces an independent pairing and yields the same inequality again. Adding the two inequalities gives

$$2 \cdot 5041 \leq 2M,$$

so

$$M \geq 5041.$$

## Conclusion

Bob has exhibited a covering using 5041 trails, and Marley has shown that no covering can use fewer. Therefore the minimum possible number of trails is

$$M = 5041.$$

## Answer

5041