

Pleasure Rather Than Salience Activates Human Nucleus Accumbens and Medial Prefrontal Cortex

Dean Sabatinelli, Margaret M. Bradley, Peter J. Lang, Vincent D. Costa, and Francesco Versace

National Institute of Mental Health Center for the Study of Emotion and Attention, University of Florida, Gainesville, Florida

Submitted 2 March 2007; accepted in final form 25 June 2007

Sabatinelli D, Bradley MM, Lang PJ, Costa VD, Versace F. Pleasure rather than salience activates human nucleus accumbens and medial prefrontal cortex. *J Neurophysiol* 98: 1374–1379, 2007. First published June 27, 2007; doi:10.1152/jn.00230.2007. Recent human functional imaging studies have linked the processing of pleasant visual stimuli to activity in mesolimbic reward structures. However, whether the activation is driven specifically by the pleasantness of the stimulus, or by its salience, is unresolved. Here we find in two studies that free viewing of pleasant images of erotic and romantic couples prompts clear, reliable increases in nucleus accumbens (NAc) and medial prefrontal cortex (mPFC) activity, whereas equally arousing (salient) unpleasant images, and neutral pictures, do not. These data suggest that in visual perception, the human NAc and mPFC are specifically reactive to pleasant, rewarding stimuli and are not engaged by unpleasant stimuli, despite high stimulus salience.

INTRODUCTION

Recent functional imaging studies have linked activation of human reward structures, including nucleus accumbens (NAc) and medial prefrontal cortex (mPFC), to the perception of pleasant emotional stimuli, through the presentation of attractive faces (Aron et al. 2001; O'Doherty et al. 2003), pictures of loved ones (Aron et al. 2005; Bartels and Zeki 2004; Fisher et al. 2005), aesthetic artwork (Kawabata and Zeki 2004), addictive drug cues (David et al. 2005; Siessmeier et al. 2006), and erotica (Ferreti et al. 2005; Hamann et al. 2004; Karama et al. 2002). Although a commonality of the visual stimuli cited above is pleasant valence, other characteristics shared by these stimuli may be associated with activation of the human NAc and mPFC.

Nonrewarding salient stimuli, operationalized as those that are arousing or elicit attention, have also been shown to prompt ventral striatal activity in humans (Zink et al. 2003, 2006) and in the animal model (Horvitz 2000; Salamone 2005). In these studies, ventral striatal activation is associated with innocuous and infrequent distractor stimuli, irrelevant to a central task. Because these salient stimuli are not pleasant, yet generate measurable responses, it is argued that the ventral striatal system is reward nonspecific. From this perspective, salient pleasant and unpleasant picture stimuli would be expected to prompt equivalent ventral striatal activation.

Past work has determined that picture stimuli rated with high emotional intensity, whether pleasant or unpleasant, are associated not only with greater central and peripheral physiological reactivity (Bradley et al. 2003; Cuthbert et al. 2000; Sabatinelli et al. 2005), but also with extended viewing time,

delayed secondary reaction time, and interest ratings (Bradley et al. 1999; see Lang and Davis 2006). This pattern holds true despite equivalent perceptual characteristics (e.g., color, brightness, spatial frequency; Junghöfer et al. 2001; Sabatinelli et al. 2005). Thus for the purposes of this study, we consider pleasant and unpleasant picture stimuli normatively rated with high emotional arousal to have greater salience than neutral pictures with low ratings of emotional arousal.

The human NAc has been reported to show graded responses to cues of varying predictive value, such that greater signal change is associated with more reliable signals of reward (Abler et al. 2006; Knutson et al. 2005; Preuschoff et al. 2006). The rated intensity of emotional stimuli is similarly associated with the magnitude of physiological response; e.g., given equally appetitive or aversive pictures, those rated as more intense elicit a stronger physiological response (Bradley et al. 2001a; Cuthbert et al. 1996; Lang et al. 1993; Sabatinelli et al. 2005). Thus pictures rated as more emotionally arousing might be expected to elicit greater reactivity in NAc.

In the present research, we study male and female participants free viewing pictures judged to be pleasant, neutral, and unpleasant. Findings from two separate experiments are reported, an initial study, and a subsequent replication and extension with a new sample of participants. Three issues regarding human reward circuit reactivity are addressed. First, to investigate the reward-nonspecific (or saliency-driven) perspective of mesolimbic function, we determine whether unpleasant pictures activate the NAc and mPFC to the same degree as pleasant pictures. We include mPFC in this examination because it is a dense cortical site of NAc interconnectivity (Ferry et al. 2000; Roberts et al. 2007) and is correlated with NAc activity in human reward studies (Knutson et al. 2003; Rogers et al. 2004). We also examine the sensitivity of NAc to pictures of equal valence, but varying emotional arousal, to explore whether the structure's graded responsivity to abstract reward predictors (Abler et al. 2006; Knutson et al. 2005; Preuschoff et al. 2006) extends to inherently emotion-evoking stimuli. Finally, considering that sex differences in arousal ratings (Bradley et al. 2001b; Janssen et al. 2003), skin conductance (Bradley et al. 2001b), and visual cortical blood oxygen level-dependent (BOLD) signal (Sabatinelli et al. 2004) have been reported in response to both pleasant and unpleasant picture stimuli, possible parallel differences in NAc and mPFC reactivity in men and women are assessed.

The costs of publication of this article were defrayed in part by the payment of page charges. The article must therefore be hereby marked "advertisement" in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

Address for reprint requests and other correspondence: D. Sabatinelli, P.O. Box 112766, University of Florida, Gainesville, FL 32611 (E-mail: sabat@ufl.edu).

METHODS

Participants and stimuli

Forty-eight introductory psychology students at the University of Florida participated in the research for course credit or \$20.00 compensation. All volunteers consented to participate after reading a description of the study, approved by the local human subjects review board. Before entering the bore of the Siemens 3T *Allegra* MR scanner, participants were fitted with earplugs and given a patient-alarm squeezeball. A vacuum pillow, padding, and explicit verbal instruction were used to limit head motion. Four participants' data were unusable due to excessive head motion, resulting in 44 participants (21 male) in the final sample. Two experiments were conducted using the same research design: in Study 1 (22, 11 male) participants viewed a set of 24 picture stimuli; in Study 2 (22, 10 male), participants viewed a set of 30 picture stimuli (subsequently described).

Participants were asked to maintain fixation on a dot at the center of a 7-in. LCD screen mounted directly behind the head (25° visual angle), visible by a coil-mounted mirror (IFIS MR-compatible hardware, Intermagnetics, Latham, NY). After three acclimation trials in which checkerboard stimuli were presented, a short series of picture stimuli were presented in an event-related design. In Study 1, the 24 grayscale picture stimuli chosen from the International Affective Picture System depicted eight exemplars of erotic couples, neutral people, and mutilations. The erotic and mutilation pictures were selected to be equivalent in normative ratings of emotional arousal (Lang et al. 2001). In Study 2, 30 picture stimuli depicted five exemplars of erotic couples, romantic couples (clothed), neutral people, dental scenes, snakes, and threatening people. The picture series followed a 28-min image-collection period, in which an unrelated paradigm was conducted. The five erotic and neutral people exemplars were chosen from the stimuli used in Study 1.

The erotic, snake, and threat scenes were chosen to be equivalent in normative ratings of emotional arousal, and erotic and romantic were chosen to be equivalent in normative ratings of valence, but differ in normative ratings of emotional arousal, with erotic greater. All picture categories were matched for luminance and 90% quality JPEG file size using Adobe Photoshop 7. Each picture was presented for 3 s, followed by a 9 s (12 s in group 2) fixation-only period. Picture order was pseudorandomized, allowing no more than two successive presentations of a stimulus category.

Scanning parameters

Once participants were comfortable inside the bore, an 8-min three-dimensional structural volume was collected. The prescription specified 160 sagittal slices, with 1-mm isotropic voxels in a 256-mm field of view. After structural acquisition, a functional prescription

was placed that included 50 coronal slices (2.5 mm thick, 0.5-mm gap), covering the entire cortex in most participants. Image volumes were collected with 3-s temporal resolution and 16-ml voxel size (TR 3 s, TE 30 ms, flip angle 70°, FOV 160 mm, 64 × 64 matrix). Typical functional image acquisition parameters (axial slices ~5 mm thick) are not optimal for investigations of ventral striatal and ventral prefrontal activity because proximity to the nasal sinus cavities leads to excessive susceptibility artifact and thus signal loss (Merboldt et al. 2001).

Data analyses

Each participant's functional time series were mean-intensity adjusted, linearly detrended, high-pass filtered at 0.02 Hz, and smoothed with a 5-mm spatial filter using BrainVoyager QX (www.brainvoyager.com). Trials containing head motion (<2%) were manually excluded from further analyses. Each participant's preprocessed functional activity was coregistered to his or her structural volume, after transformation into standardized Talairach coordinate space (Talairach and Tournoux 1988). Random-effects analyses identified clusters (>100 ml) of voxels showing greater BOLD signal increase during pleasant relative to neutral picture presentations, using a minimum significance threshold of $P < 0.001$ ($t > 3.8$).

Region of interest analyses were conducted using cluster locations revealed in the group analyses. The random-effects contrast maps were relaxed to a threshold of $P < 0.05$, and a 1-cm³ volume was located at the voxel of most reliable signal change in the group average. This cluster location was then sampled in each participant's standardized volume, and the peak of this time course (6 s after stimulus onset) was exported for ANOVA analyses including gender and picture category as factors.

RESULTS

Volume analysis

A random-effects ANOVA of the Study 1 average volume identified significant clusters of voxels with greater BOLD signal change in response to erotica relative to neutral people pictures. As shown in Fig. 1, reliable activation was found in NAc (A; $\pm 10, 9, -2$) and mPFC (C; $-5, 41, 5$). The event-related time course of group-averaged BOLD signal change in these clusters is shown for each picture category in Fig. 1, B and D, and indicates a selective increase in activity elicited by erotic pictures, but no increase over baseline after presentation of neutral people or mutilation pictures. No significant difference in NAc or




FIG. 1. Study 1: functional activity during erotic picture processing, relative to neutral people pictures. Random-effects analyses reveal bilateral nucleus accumbens (A; $\pm 10, 9, -2$) and medial prefrontal cortex (C; $-5, 41, 5$) activation that is greater during erotic relative to neutral picture presentations. Event-related time courses of percentage blood oxygenated level-dependent (BOLD) signal change in nucleus accumbens (NAc, B) and medial prefrontal cortex (mPFC, D) reveal that, in contrast to the response to erotic pictures (yellow), mutilation pictures (red) did not prompt a signal increase and did not differ from activity prompted by neutral people pictures (blue).

mPFC activity was found in contrasts of mutilation and neutral people picture presentations.

Consistent with the Study 1 data, a random-effects ANOVA of the Study 2 average volume identified clusters of voxels in NAc (Fig. 2A: $\pm 4, 2, -3$) and mPFC (Fig. 2C: $-4, 39, 4$) with greater BOLD signal change during erotic and romantic relative to the least-arousing picture contents (i.e., neutral people and dental). The event-related time course of average signal change is shown for each picture category in Fig. 2, B and D, illustrating a selective increase in activity after erotic and romantic pictures, but no increase over baseline after presentation of neutral people, dental, snake, or threatening pictures. No significant activity in NAc or mPFC was found in contrasts of more arousing aversive contents (snakes, human threat) and less-arousing picture presentations.

Region of interest analyses

The modulation of Study 1 NAc activity by picture content was reliable in the individually sampled analysis of peak BOLD signal change [$F(2,40) = 32.48, P < 0.001$] and consistent across men and women [Content \times Gender: $F(2,40) = 1.93$, ns]. Unpleasant and neutral pictures did not elicit differential BOLD signal change in NAc [$F(1,20) = 1.62$, ns]. A similar pattern of signal change was found for peak mPFC activity, where signal change was reliably modulated by picture content [$F(2,40) = 28.47, P < 0.001$] and did not interact with gender [$F(2,40) < 1$, ns]. As in NAc, unpleasant and neutral pictures elicited equivalent reactivity in mPFC [$F(1,20) = 1.28$, ns] Table 1.

The modulation of Study 2 NAc activity by picture category was reliable in the individually sampled analysis of peak BOLD signal change [$F(5,100) = 6.32, P < 0.001$] and consistent across men and women [Category \times Gender: $F(5,100) < 1$, ns]. Unpleasant (snake and threat) and neutral (people and dental) pictures did not elicit differential BOLD signal change in NAc [$F(1,20) < 1$, ns]. A similar pattern of signal change was found for peak mPFC activity, in which signal change was reliably modulated by picture category [$F(5,100) = 8.62, P < 0.001$] and did not interact with gender [$F(5,100) = 1.09$, ns]. As in NAc, unpleasant and neutral pictures elicited equivalent reactivity in mPFC [$F(1,20) < 1$, ns] Table 2.

DISCUSSION

Pleasant picture perception was associated with highly reliable increases in NAc and mPFC BOLD signal. This activation is unlikely to be a result of picture salience because images of horrific injuries, threatening snakes, or aggressive people did not elicit signal increases, nor did activity in primary visual cortex differ across picture categories. The clear distinction between NAc and mPFC activity elicited by pleasant pictures and all other picture categories was evident despite a brief presentation series of 24 to 30 total trials. Previous studies with virtually identical stimuli have found strong indications of orienting and novelty responses to aversive pictures, including self-determined viewing time (Lang et al. 1993), skin conductance (Bradley et al. 2001), and the electrocortical event-related potential (Cuthbert et al. 2000; Sabatinelli et al. 2007). Therefore these data do not support the perspective that human ventral striatal reactivity is driven by salience, rather than reward. Selective activation of the NAc and mPFC to pleasant visual stimuli (a valence effect) contrasts with reliable BOLD signal increases elicited by both pleasant and unpleasant visual stimuli (an arousal effect) in amygdala (Breiter et al. 1996; Garavan et al. 2001; Sabatinelli et al. 2005), anterior cingulate (Britton et al. 2006; Bush et al. 2000), and inferior temporal visual cortex (Bradley et al. 2003; Sabatinelli et al. 2005; Vuilleumier et al. 2001) and suggests a dedicated role of NAc and mPFC in reward processing.

The magnitude of NAc and mPFC signal did not differ as a function of whether pictures depicted erotic or romantic content, despite clear differences in normative ratings of emotional arousal. These data suggest that during picture perception, the human NAc and mPFC react specifically to stimulus pleasantness, rather than to emotional arousal. This relative insensitivity to the rated arousal of pictures is unlike the patterns of reactivity seen in a variety of peripheral and central physiological measures, including skin conductance and facial electromyography (Bernat et al. 2006; Lang et al. 1993), the modulation of the startle blink reflex (Bradley et al. 2001; Cuthbert et al. 1996), the amplitude of the electrocortical late positive potential (Cuthbert et al. 2000; Schupp et al. 2003), and in BOLD signal change in ventral visual cortex and amygdala (Bradley et al. 2003; Phan et al. 2004; Sabatinelli et al. 2005; Winston et al. 2005). It should be noted that the pleasant stimuli used here—erotic and romantic couples—were variants on a theme of sexual opportunity, one explicit




FIG. 2. Study 2: functional activity during erotic and romantic picture processing, relative to neutral people and dental scenes. Random-effects analyses reveal bilateral nucleus accumbens (A; $\pm 4, 2, -3$) and medial prefrontal cortex (C; $-4, 39, 4$) activation that is greater during erotic and romantic relative to neutral people and dental scene presentations. Event-related time courses of percentage BOLD signal change in NAc (B) and mPFC (D) reveal that, in contrast to the response to erotic (yellow) and romantic (white) pictures, snake (magenta) and human threat (red) contents did not prompt a signal increase and did not differ from activity prompted by neutral people (blue) and dental scene (light blue) pictures.

TABLE 1. *Region of interest analysis after transformation into standardized Talairach coordinate space for Study 1*

Region	Pleasant > Neutral				Region	Unpleasant > Neutral			
	x	y	z	t(21)		x	y	z	t(21)
Medial prefrontal cortex	-5	41	5	5.87	Amygdala	±21	-6	-11	4.62
Nucleus accumbens	±10	9	-2	7.85	Fusiform gyrus	±34	-69	-10	6.69
Amygdala	±23	-6	-10	6.03	Lateral occipital cortex	±39	-78	-6	4.97
Fusiform gyrus	±39	-60	-7	7.17	Inferior frontal gyrus	±47	15	21	4.37
Lateral occipital cortex	±47	-73	-10	7.11	R intraparietal cortex	28	-72	31	4.92
Inferior frontal gyrus	±44	0	29	5.55	R lingual gyrus	17	-86	-6	4.48
Intraparietal sulcus	±32	-61	54	6.23	Cingulate	-4	-7	33	3.94
Inferior parietal cortex	±57	-29	27	5.38	Postcentral gyrus	±53	-23	37	4.92
Cingulate	8	17	32	5.58	R orbitofrontal cortex	23	33	-7	4.23
Precuneus	-1	-61	40	6.46					
R medial frontal gyrus	26	-6	56	6.65					
L anterior insula	-31	23	4	5.29					

and one implicit. Thus it will be important to reexamine the issue when other pleasant (but nonsexual) stimuli are processed.

No differences between men and women were apparent in subcortical NAc or cortical mPFC, in either sample. It appears that at the level of the mesolimbic reward circuit, both men and women are receptive to this type of pleasant cue. Sex differences seen previously in sensory cortical, peripheral physiological, and self-report measures (Bradley et al. 2001; Costa et al. 2003; Janssen et al. 2003; Karama et al. 2002; Lang et al. 1998; Sabatinelli et al. 2004) and associated sensitivity to variations in emotional arousal may occur in downstream or parallel circuits.

Consistent with previous studies using pleasant visual stimuli (Kawabata and Zeki 2004; O'Doherty et al. 2003), the time course of mPFC activity after pleasant picture onset indicated a clear increase in BOLD signal, in Studies I and II. However, neutral and unpleasant picture stimuli apparently evoke a decrease in mPFC BOLD signal, relative to prepicture baseline. In fact, tests of mPFC signal change reveal reliable decreases in signal intensity after neutral and unpleasant pictures in Study 1 [$F(1,21) = 29.30$, $P < 0.001$] and Study 2 [$F(1,21) = 8.81$, $P < 0.01$]. Although the nature of BOLD signal decrease is less well understood than BOLD increase, and may reflect an interruption of ongoing "default" activity (Gusnard et al. 2001), this pattern of reactivity in mPFC may be a result of an unmet expectation of a pleasant picture (reward) presentation (Knutson et al. 2003; Rogers et al. 2004;

Schultz 1998), similar to responses seen in conditioning studies in which predicted rewarding unconditioned stimuli fail to appear (Abler et al. 2006; Pessiglione et al. 2006). Clearly the possibility of decreasing BOLD signal as a function of unmet reward delivery will require more focused research designs.

In summary, the human NAc and mPFC are selectively reactive to pleasant, relative to neutral or unpleasant, picture contents. Furthermore, the magnitude of BOLD signal was determined specifically by the emotional valence of the pictures and not by arousal. That is, highly arousing erotic pictures could not be distinguished from moderately arousing romantic pictures, nor were differences found among the unpleasant picture contents, suggesting that unlike many other neural structures studied in imaging research, the NAc and mPFC are sensitive particularly to picture pleasantness and not to emotional intensity. Finally, these effects appear to be gender general because no difference between men and women was found in the substantial samples of participants studied in two experiments. Overall, these data suggest that activation of the human mesolimbic reward system (i.e., NAc and mPFC) is not enhanced by the salience (arousal, attention capture) of visual scene contents during free viewing, but is specifically a reaction to image pleasantness.

GRANTS

This research was supported by National Institute of Mental Health Grant P50-MH-072850.

TABLE 2. *Region of interest analysis after transformation into standardized Talairach coordinate space for Study 2*

Region	Pleasant > Neutral				Region	Unpleasant > Neutral			
	x	y	z	t(21)		x	y	z	t(21)
Medial prefrontal cortex	-4	39	4	7.99	L amygdala	-22	-5	-18	4.41
Nucleus accumbens	±4	2	-3	6.35	Fusiform gyrus	±37	-69	-5	4.91
L amygdala	-20	-8	-8	4.17	Lateral occipital cortex	36	-77	-11	4.31
Fusiform gyrus	±42	-56	-20	4.77	Inferior frontal gyrus	-29	12	-13	5.34
Lateral occipital cortex	±46	-68	-9	5.21	Precuneus	8	-61	10	4.09
Inferior frontal gyrus	±23	15	-9	6.17	Cingulate	9	-13	38	4.15
Precuneus	5	-61	27	5.32	Cuneus	4	-83	19	5.07
Cingulate	1	-14	37	5.08	Anterior cingulate	1	23	22	4.02
Intraparietal sulcus	30	-53	55	3.96					
Precentral gyrus	±28	-10	49	5.27					

REFERENCES

- Abler B, Walter H, Erk S, Kammerer H, Spitzer M. Prediction error as a linear function of reward probability is coded in human nucleus accumbens. *Neuroimage* 31: 790–795, 2006.
- Aharon I, Etcoff N, Ariely D, Chabris CF, O'Connor E, Breiter HC. Beautiful faces have variable reward value: fMRI and behavioral evidence. *Neuron* 32: 537–551, 2001.
- Aron A, Fisher H, Mashek DJ, Strong G, Li H, Brown LL. Reward, motivation, and emotion systems associated with early-stage intense romantic love. *J Neurophysiol* 94: 327–337, 2005.
- Bartels A, Zeki S. The neural correlates of maternal and romantic love. *Neuroimage* 21: 1155–1166, 2004.
- Bernat E, Patrick CJ, Benning SD, Tellegen A. Effects of picture content and intensity on affective physiological response. *Psychophysiology* 43: 93–103, 2006.
- Bradley MM, Codispoti M, Cuthbert BN, Lang PJ. Emotion and motivation I: defensive and appetitive reactions in picture processing. *Emotion* 1: 276–298, 2001a.
- Bradley MM, Codispoti M, Sabatinelli D, Lang PJ. Emotion and motivation II: sex differences in picture processing. *Emotion* 1: 300–319, 2001b.
- Bradley MM, Cuthbert BN, Lang PJ. Affect and the startle reflex. In: *Startle Modification: Implications for Neuroscience, Cognitive Science, and Clinical Science*, edited by Dawson ME, Schell A, Boehmelt A. New York: Cambridge Univ. Press, 1999.
- Bradley MM, Sabatinelli D, Lang PJ, Fitzsimmons JR, King W, Desai P. Activation of the visual cortex in motivated attention. *Behav Neurosci* 117: 369–380, 2003.
- Breiter HC, Etcoff NL, Whalen PJ, Kennedy WA, Rauch SL, Buckner RL, Strauss MM, Hyman SE, Rosen BR. Response and habituation of the human amygdala during visual processing of facial expression. *Neuron* 17: 875–887, 1996.
- Britton JC, Phan KL, Taylor SF, Welsh RC, Berridge KC, Liberzon I. Neural correlates of social and nonsocial emotions: an fMRI study. *Neuroimage* 31: 397–409, 2006.
- Bush G, Luu P, Posner MI. Cognitive and emotional influences in anterior cingulate cortex. *Trends Cogn Sci* 4: 215–222, 2000.
- Costa M, Braun C, Birbaumer N. Gender differences in response to pictures of nudes: a magnetoencephalographic study. *Biol Psychol* 63: 129–147, 2003.
- Cuthbert BN, Bradley MM, Lang PJ. Probing picture perception: activation and emotion. *Psychophysiology* 33: 103–111, 1996.
- Cuthbert BN, Schupp HT, Bradley MM, Birbaumer N, Lang PJ. Brain potentials in affective picture processing: covariation with autonomic arousal and affective report. *Biol Psychol* 52: 95–111, 2000.
- David SP, Munafo MR, Johansen-Berg H, Smith SM, Rogers RD, Matthews PM, Walton RT. Ventral striatum/nucleus accumbens activation to smoking-related pictorial cues in smokers and nonsmokers: a functional magnetic resonance imaging study. *Biol Psychiatry* 58: 488–494, 2005.
- Ferretti A, Caulo M, Del Gratta C, Matteo RD, Merla A, Montorsi F, Pizzella V, Pompa P, Rigatti P, Rossini PM, Salonia A, Tartaro A, Romani GL. Dynamics of male sexual arousal: distinct components of brain activation revealed by fMRI. *Neuroimage* 26: 1086–1096, 2005.
- Ferry AT, Öngür D, An X, Price JL. Prefrontal cortical projections to the striatum in macaque monkeys: evidence for an organization related to prefrontal networks. *J Comp Neurol* 425: 447–470, 2000.
- Fisher H, Aron A, Brown LL. Romantic love: an fMRI study of a neural mechanism for mate choice. *J Comp Neurol* 493: 58–62, 2005.
- Garavan H, Pendergrass JC, Ross TJ, Stein EA, Risinger RC. Amygdala responses to both positively and negatively valenced stimuli. *Neuroreport* 12: 2779–2783, 2001.
- Gusnard DA, Akbudak E, Shulman GL, Raichle ME. Medial prefrontal cortex and self-referential mental activity: relation to a default mode of brain function. *Proc Natl Acad Sci USA* 98: 4259–4264, 2001.
- Hamann S, Herman RA, Nolan CL, Wallen K. Men and women differ in amygdala response to sexual stimuli. *Nat Neurosci* 7: 411–416, 2004.
- Horvitz JC. Mesolimbic and nigrostriatal dopamine responses to salient non-reward events. *Neuroscience* 96: 651–656, 2000.
- Janssen E, Carpenter D, Graham CA. Selecting films for sex research: gender differences in erotic film preference. *Arch Sex Behav* 32: 243–251, 2003.
- Junghöfer M, Bradley MM, Elbert TR, Lang PJ. Fleeting images: a new look at early emotion discrimination. *Psychophysiology* 38: 175–178, 2001.
- Karama S, Lecours AR, Leroux JM, Bourgouin P, Beaudoin G, Joubert S, Beauregard M. Areas of brain activation in males and females during viewing of erotic film excerpts. *Hum Brain Mapp* 16: 1–13, 2002.
- Kawabata H, Zeki S. Neural correlates of beauty. *J Neurophysiol* 91: 1699–1705, 2004.
- Knutson B, Fong GW, Bennett SM, Adams CM, Hommer D. A region of mesial prefrontal cortex tracks monetarily rewarding outcomes: characterization with rapid event-related fMRI. *Neuroimage* 18: 263–272, 2003.
- Knutson B, Taylor J, Kaufman M, Peterson R, Glover G. Distributed neural representation of expected value. *J Neurosci* 25: 4806–4812, 2005.
- Lang PJ, Bradley MM, Cuthbert BN. *International Affective Picture System (IAPS): Instruction Manual and Affective Ratings*. Technical Report A-4. Gainesville, FL: The Center for Research in Psychophysiology, Univ. of Florida, 2001.
- Lang PJ, Bradley MM, Fitzsimmons JR, Cuthbert BN, Scott JD, Moulder B, Nangia V. Emotional arousal and activation of the visual cortex: an fMRI analysis. *Psychophysiology* 35: 199–210, 1998.
- Lang PJ, Davis M. Emotion, motivation, and the brain: reflex foundations in animal and human research. *Prog Brain Res* 156: 3–29, 2006.
- Lang PJ, Greenwald MK, Bradley MM, Hamm AO. Looking at pictures: affective, facial, visceral, and behavioral reactions. *Psychophysiology* 30: 261–273, 1993.
- Merboldt KD, Franssen P, Bruhn H, Frahm J. Functional MRI of the human amygdala? *Neuroimage* 14: 253–257, 2001.
- O'Doherty J, Winston J, Critchley H, Perrett D, Burt DM, Dolan RJ. Beauty in a smile: the role of medial orbitofrontal cortex in facial attractiveness. *Neuropsychologia* 41: 147–155, 2003.
- Pessiglione M, Seymour B, Flandin G, Dolan RJ, Frith CD. Dopamine-dependent prediction errors underpin reward-seeking behavior in humans. *Nature* 422: 1042–1045, 2006.
- Phan KL, Taylor SF, Welsh RC, Ho SH, Britton JC, Liberzon I. Neural correlates of individual ratings of emotional salience: a trial-related fMRI study. *Neuroimage* 21: 768–778, 2004.
- Preusschoff K, Bossaerts P, Quartz SR. Neural differentiation of expected reward and risk in human subcortical structures. *Neuron* 51: 381–390, 2006.
- Roberts AC, Tomic DL, Parkinson CH, Roeling TA, Cutter DJ, Robbins TW, Everitt BJ. Forebrain connectivity of the prefrontal cortex in the marmoset monkey (*Callithrix jacchus*): an anterograde and retrograde tract-tracing study. *J Comp Neurol* 502: 86–112, 2007.
- Rogers RD, Ramnani N, Mackay C, Wilson JL, Jezard P, Carter CS, Smith SM. Distinct portions of anterior cingulate and medial prefrontal cortex are activated by reward processing in separable phases of decision-making cognition. *Biol Psychiatry* 55: 594–602, 2004.
- Sabatinelli D, Bradley MM, Fitzsimmons JR, Lang PJ. Parallel amygdala and inferotemporal activation reflect emotional intensity and fear relevance. *Neuroimage* 24: 1265–1270, 2005.
- Sabatinelli D, Flaisch T, Bradley MM, Fitzsimmons JR, Lang PJ. Affective picture perception: gender differences in visual cortex? *Neuroreport* 15: 1109–1112, 2004.
- Sabatinelli D, Lang PJ, Keil A, Bradley MM. Emotional perception: correlation of functional MRI and event-related potentials. *Cereb Cortex* 17: 1085–1091, 2007.
- Salamone JD, Correa M, Mingote SM, Weber SM. Beyond the reward hypothesis: alternative functions of nucleus accumbens dopamine. *Curr Opin Pharmacol* 5: 34–41, 2005.
- Schultz W. Predictive reward signal of dopamine neurons. *J Neurophysiol* 80: 1–27, 1998.
- Schupp HT, Junghöfer M, Weike AI, Hamm AO. Emotional facilitation of sensory processing in the visual cortex. *Psychol Sci* 14: 7–13, 2003.
- Siessmeier T, Kienast T, Wrase J, Larsen JL, Braus DF, Smolka MN, Buchholz HG, Schreckenberger M, Rösch F, Cumming P, Mann K, Bartenstein P, Heinz A. Net influx of plasma 6-[18F]fluoro-L-DOPA (FDOPA) to the ventral striatum correlates with prefrontal processing of affective stimuli. *Eur J Neurosci* 24: 305–313, 2006.
- Talairach J, Tournoux P. *Co-Planar Stereotaxic Atlas of the Human Brain*. New York: Thieme, 1988.

- Vuilleumier P, Armony JL, Driver J, Dolan RJ.** Effects of attention and emotion on face processing in the human brain: an event related fMRI study. *Neuron* 30: 829–841, 2001.
- Winston JS, Gottfried JA, Kilner JM, Dolan RJ.** Integrated neural representations of odor intensity and affective valence in human amygdala. *J Neurosci* 25: 8903–8907, 2005.
- Zink CF, Pagnoni G, Chappelow JC, Martin-Skurski ME, Berns GS.** Human striatal activation reflects degree of stimulus saliency. *Neuroimage* 29: 977–983, 2006.
- Zink CF, Pagnoni G, Martin ME, Dhamala M, Berns GS.** Human striatal response to salient nonrewarding stimuli. *J Neurosci* 23: 8092–8097, 2003.