

Coronal Seismology

ASTR 598


Laurel Farris

Spring 2016



Magnetohydrodynamics (MHD)

Theory



Model

- Straight cylindrical flux tube in uniform magnetic field.
- Equations of “ideal” MHD
- Characteristic wave speeds are determined by ρ , T , P , and \vec{B}

Sound speed

- $$C_s \propto \sqrt{\frac{P}{\rho}}$$
$$\propto \sqrt{T}$$

Alfvén speed

- $$V_A \propto \frac{B}{\sqrt{\rho}}$$


MHD modes in the solar corona

Main categories

- Magnetoacoustic
 - Fast
 - Slow
- Alfvén

Research Topics


1. Kink oscillations
2. Sausage oscillations
3. Acoustic oscillations
4. Propagating acoustic waves
5. Propagating fast waves
6. Torsional (Alfvén) modes



- $C_k = V_A \sqrt{\frac{2}{1 + \rho_e / \rho_o}}$
- $\xi(x) = \xi(r) e^{i(kz + m\phi)}$

Coronal seismology

Technique and motivation



Elusive coronal properties

- magnetic field strength, \vec{B}
- density, ρ
- Alfvén velocity, V_A

Motivation


- Coronal heating
- Space weather prediction

Coronal seismology

1. Observe disturbances
2. Measure physical parameters
3. Identify wave properties
4. Extract physical parameters

Fast standing oscillations

Kinks vs. Sausages



Period

- $P = \frac{2\ell}{V_{ph}} \quad (\lambda = 2\ell)$

Sausage


- No loop spatial displacement
- Symmetric
- Intensity change
→ density change

Kink

- loop spatial displacement
- Asymmetric
- No intensity change

“Observations of sausage modes in magnetic pores”


Morton et al. 2011



- Periods ~ 30 -450 seconds (0.5-7.5 minutes)
- Possibly driven by 5-minute acoustic oscillations

Alfvén waves

Verwichte et al.



Objective


- Determine Alfvén speed in two ways:
 1. Coronal seismology
 2. Magnetic extrapolation and spectral methods

Observed

- Two transversely oscillating loops triggered by flare
- AIA/SDO 171 Å

Alfvén waves

Verwichte et al.



“The determination of the Alfvén speed from the observed phase speed lies at the heart of the seismological method. . .”

Research

AIA/SDO


Fe XII, XXIV

193 Å



Research


Bright point (BP)



- Image:
~ 45000 km across
- Bright point:
~ 9000 km across




Research

Plots



Research


Cross-correlations




Research

Cross-correlation & timelag images

Cross-correlation



Timelag



2 pixels \sim 1 arcsec \sim 700 km

Other questions and future work

Other questions

- What is the excitation mechanism for the observed disturbances?
- How are they damped, and what determines the timescales?

My future work

- Download data in other wavelengths (i.e. coronal heights).
- Download data from other instruments, e.g. the Extreme Ultraviolet Variability Experiment (EVE) on SDO.
- Characterize other bright points in coronal hole, quiet sun, and active regions.



Acknowledgements

Advisor: James McAteer

Extra slides here