

## Part I: Gaussian Elimination

1. Solve the following system using Gaussian elimination and then use the Grapher utility on a Mac to visualize the solution point:

$$3x + y + z = 3$$

$$12x - 2y + 3z = 3$$

$$-6x - 2y - 4z = 12$$

subtract ~~four~~ <sup>two</sup> times eqn 1 from eqn 2:

$$3x + y + z = 3$$

$$-6y - z = -9$$

$$-6x - 2y - 4z = 12$$

Add 2 times eqn 1 to eqn 3:

$$3x + y + z = 3$$

$$-6y - z = -9$$

$$-2z = 18$$

Back substitute:

$$z = -9$$

$$y = 3$$

$$x = 3$$

$$\begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 3 \\ 3 \\ -9 \end{bmatrix} \text{ point of intersection}$$

2. Solve the following system using Gaussian elimination and then use the Grapher utility on a Mac to visualize the solution line:

$$\begin{cases} 2x + 3y + 4z = 12 \\ x + y + z = 6 \\ 4x + 6y + 8z = 24 \end{cases} \text{ same plane}$$

subtract  $\frac{1}{2}$  of eqn #1 from eqn #3:

$$2x + 3y + 4z = 12$$

$$-\frac{1}{2}y - z = 0$$

$$4x + 6y + 8z = 24$$

subtract 2 times eqn #1 from eqn #3

$$2x + 3y + 4z = 12$$

$$-\frac{1}{2}y - z = 0$$

$$0 = 0$$

Back substitute:  $z = -\frac{1}{2}y$

$$2x + 3y + 4(-\frac{1}{2}y) = 12$$

$$2x + y = 12$$

$$y = 12 - 2x$$

$$z = -\frac{1}{2}(12 - 2x) = x - 6$$

$$\begin{cases} x \text{ can be anything} \\ y = -2x + 12 \\ z = x - 6 \end{cases}$$

$$\rightarrow \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 0 \\ 12 \\ -6 \end{bmatrix} + t \begin{bmatrix} 1 \\ -2 \\ 1 \end{bmatrix} \text{ where } t \text{ is any real \#}$$

→ a line in 3D

3. Solve the following system using Gaussian elimination and then use the Grapher utility on a Mac to visualize the solution plane:

$$\begin{cases} 3x + 2y + 4z = 12 \\ 6x + 4y + 8z = 24 \\ 12x + 8y + 16z = 48 \end{cases} \quad \left. \begin{array}{l} \\ \\ \end{array} \right\} \text{3 of the same plane}$$

$$3x + 2y + 4z = 12$$

$$0 = 0$$

$$0 = 0$$

Everything on the  $3x + 2y + 4z = 12$  plane is a solution.

Since  $x$  and  $y$  can be anything and

$$z = \frac{12 - 3x - 2y}{4} = -\frac{3}{4}x - \frac{1}{2}y + 3, \text{ we}$$

can write this plane as

$$\begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 3 \end{bmatrix} + s \begin{bmatrix} 1 \\ 0 \\ -3/4 \end{bmatrix} + t \begin{bmatrix} 0 \\ 1 \\ -1/2 \end{bmatrix}$$

where  $s$  and  $t$  are any real numbers.

4. (Try) to solve the following system using Gaussian elimination and then use the Grapher utility on a Mac to visualize why there is no solution:

$$x + y + z = 2$$

$$2x + 2y + 2z = 6$$

$$4x + 6y + 8z = 10$$

subtract 2 times eqn #1 from eqn 2:

$$x + y + z = 2$$

$$0 = 2$$

Impossible  $\rightarrow$  no solution.

# Homework:

1. Solve:

$$x + y + z = 2$$

$$x + 3y + 3z = 0$$

$$x + 3y + 5z = 2$$

$$x + y + z = 2$$

$$2y + 2z = -2$$

$$2y + 4z = 0$$

$$x + y + z = 2$$

$$2y + 2z = -2$$

$$2z = 2$$

$$z = 1$$

$$\rightarrow y = -2$$

$$\rightarrow x = 3$$

$$\begin{bmatrix} 3 \\ -2 \\ 1 \end{bmatrix}$$

2. Solve:

$$x + 2y + z = 6$$

$$3x + 6y + 3z = 18$$

$$x + y + z = 12$$

$$x + 2y + z = 6$$

$$0 = 0$$

$$-y = 6$$

$$y = -6$$

$$x + 2(-6) + z = 6$$

$$x + z = 18$$

$$\begin{cases} z = 18 - x \\ y = -6 \end{cases}$$

translate into :

$$\begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 0 \\ -6 \\ 18 \end{bmatrix} + t \begin{bmatrix} 1 \\ 0 \\ -1 \end{bmatrix} \quad \begin{matrix} \uparrow \\ \text{intercept} \end{matrix} \quad \begin{matrix} \uparrow \\ \text{slope} \end{matrix}$$

$$x = 1t$$

for any  $t \in \mathbb{R}$

## Part II: Matrix Multiplication & Elementary Matrices

1. Find the **inner product** (aka dot product) of the row vector  $\begin{bmatrix} 1 & 2 & 3 \end{bmatrix}$  and the column

vector  $\begin{bmatrix} 4 \\ 5 \\ 6 \end{bmatrix}$ .

$$\begin{bmatrix} 1 & 2 & 3 \end{bmatrix} \cdot \begin{bmatrix} 4 \\ 5 \\ 6 \end{bmatrix} = 1 \cdot 4 + 2 \cdot 5 + 3 \cdot 6 \\ = 4 + 10 + 18 \\ = 14 + 18 \\ = 32$$

2. Consider the system of equations:

$$2x + y + z = 5$$

$$4x - 6y = -2$$

$$-2x + 7y + 2z = 9$$

We can think of the left hand side as a matrix multiplication in two different ways.

- A. Write the left hand side as three rows where each row includes an inner product.

$$\begin{bmatrix} 2 \cdot x + 1 \cdot y + 1 \cdot z \\ 4 \cdot x - 6 \cdot y + 0 \cdot z \\ -2 \cdot x + 7 \cdot y + 2 \cdot z \end{bmatrix} = \begin{bmatrix} 5 \\ -2 \\ 9 \end{bmatrix}$$

- B. Write the left hand side as a linear combination of three column vectors.

$$x \begin{bmatrix} 2 \\ 4 \\ -2 \end{bmatrix} + y \begin{bmatrix} 1 \\ -6 \\ 7 \end{bmatrix} + z \begin{bmatrix} 1 \\ 0 \\ 2 \end{bmatrix} = \begin{bmatrix} 5 \\ -2 \\ 9 \end{bmatrix}$$

3. Write out what the general  $i^{\text{th}}$  row of the product  $Ax$  is using sigma notation where:

$$\begin{bmatrix} a_{i1} & \dots & \dots & a_{in} \\ \vdots & & & \vdots \\ a_{m1} & \dots & \dots & a_{mn} \end{bmatrix} \begin{bmatrix} x_1 \\ \vdots \\ x_n \end{bmatrix} = a_{i1} \cdot x_1 + a_{i2} \cdot x_2 + \dots + a_{in} \cdot x_n \\ = \sum_{j=1}^n a_{ij} x_j$$

4. What are the requirements on the size of matrices A and B in order to multiply  $A \cdot B$ ? What is the size of the resulting matrix?

$$(m \times n)(n \times p) = m \times p$$

5. Find the following products:

a.  $\begin{bmatrix} 1 & 2 & 3 \\ 0 & 1 & -1 \\ 4 & 0 & 2 \end{bmatrix} \begin{bmatrix} 1 & 0 \\ -2 & -2 \\ -0 & -1 \end{bmatrix}$

$$= \begin{bmatrix} -3 & -7 \\ -2 & -1 \\ 4 & -2 \end{bmatrix}$$

b.  $\begin{bmatrix} 1 & 2 & 0 & 1 \end{bmatrix} \begin{bmatrix} -1 \\ 1 \\ 0 \\ 3 \end{bmatrix}$

$$(1)(-1) + 2(1) + 0(0) + (1)(3) = 4$$

c.  $\begin{bmatrix} -1 \\ 1 \\ 0 \\ 3 \end{bmatrix} \begin{bmatrix} 1 & 2 & 0 & 1 \end{bmatrix} \cdot \begin{bmatrix} -1 & -2 & 0 & -1 \\ 1 & 2 & 0 & 1 \\ 0 & 0 & 0 & 0 \\ 3 & 6 & 0 & 3 \end{bmatrix}$

6. More generally, write the entry in the  $i$ th row and  $j$ th column, denoted  $(AB)_{ij}$ , of this product:

$$\begin{bmatrix} a_{i1} & \dots & a_{in} \\ \vdots & & \vdots \\ a_{m1} & \dots & a_{mn} \end{bmatrix} \begin{bmatrix} b_{11} & b_{1j} & \dots & b_{1p} \\ \vdots & \vdots & & \vdots \\ b_{n1} & b_{nj} & \dots & b_{np} \end{bmatrix} = a_{i1}b_{1j} + a_{i2}b_{2j} + \dots + a_{in}b_{nj}$$

$$= \sum_{k=1}^n a_{ik}b_{kj}$$

7. Matrix multiplication is associative, meaning that  $(AB)C = A(BC)$ . Show that this is true for:

$$A = \begin{bmatrix} 1 & 1 \\ 0 & 2 \end{bmatrix}, B = \begin{bmatrix} 2 & 1 \\ 3 & 0 \end{bmatrix}, C = \begin{bmatrix} -1 & -1 \\ 2 & 0 \end{bmatrix} \quad (AB)C = \begin{bmatrix} 5 & 1 \\ 6 & 0 \end{bmatrix} \begin{bmatrix} -1 & -1 \\ 2 & 0 \end{bmatrix} = \begin{bmatrix} -3 & -5 \\ -6 & -6 \end{bmatrix}$$

$$A(BC) = \begin{bmatrix} 1 & 1 \\ 0 & 2 \end{bmatrix} \begin{bmatrix} 0 & -2 \\ -3 & -3 \end{bmatrix} = \begin{bmatrix} -3 & -5 \\ -6 & -6 \end{bmatrix}$$

8. In general, matrix multiplication is NOT commutative, meaning that  $AB$  does not usually

$$A = \begin{bmatrix} 1 & 1 \\ 0 & 2 \end{bmatrix}, B = \begin{bmatrix} 2 & 1 \\ 3 & 0 \end{bmatrix}$$

equal  $BA$ . Show that  $AB$  and  $BA$  are not equal for

$$AB = \begin{bmatrix} 5 & 1 \\ 6 & 0 \end{bmatrix}$$

$$BA = \begin{bmatrix} 2 & 4 \\ 3 & 3 \end{bmatrix}$$

Some definitions:

The  $I_{n \times n}$  **identity matrix** has 1's along its diagonal and zero's everywhere else. For any matrix A,  $A = AI = IA$  when the dimensions match up appropriately.

Ex:  $\begin{matrix} A & I \\ \begin{bmatrix} 1 & 1 \\ 0 & 2 \end{bmatrix} & \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \end{matrix} = \begin{bmatrix} 1 & 1 \\ 0 & 2 \end{bmatrix}$

The elementary matrices  $E_{ij}$  are the identity matrices with an extra  $-L$  term in the  $ij$ th entry. Multiplying an elementary matrix on the left has the effect of subtracting  $L$  times row  $j$  from row  $i$ . For example:

$$E_{31} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ -l & 0 & 1 \end{bmatrix} \text{ has the effect of subtracting } L \text{ times row 1 from row 3.}$$

Gaussian elimination is done by multiplying by elementary matrices.

9. Consider the system:

$$\begin{bmatrix} 2 & 1 & 1 \\ 4 & -6 & 0 \\ -2 & 7 & 2 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 5 \\ -2 \\ 9 \end{bmatrix}$$

a. Use an elementary matrix in which you multiply row 1 by 2 and subtract it from row 2.

$$\begin{bmatrix} 1 & 0 & 0 \\ -2 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} 2 & 1 & 1 \\ 4 & -6 & 0 \\ -2 & 7 & 2 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \\ -2 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} 5 \\ -2 \\ 9 \end{bmatrix} \rightarrow \begin{bmatrix} 2 & 1 & 1 \\ 0 & -8 & -2 \\ -2 & 7 & 2 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 5 \\ -12 \\ 9 \end{bmatrix}$$

b. From there, use an elementary matrix in which you add row 1 to row 3.

$$\begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 1 & 0 & 1 \end{bmatrix} \begin{bmatrix} 2 & 1 & 1 \\ 0 & -8 & -2 \\ -2 & 7 & 2 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 1 & 0 & 1 \end{bmatrix} \begin{bmatrix} 5 \\ -12 \\ 9 \end{bmatrix} \rightarrow \begin{bmatrix} 2 & 1 & 1 \\ 0 & -8 & -2 \\ 0 & 8 & 3 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 5 \\ -12 \\ 14 \end{bmatrix}$$

c. From there, use an elementary matrix in which you add row 2 to row 3.

$$\begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 1 & 1 \end{bmatrix} \begin{bmatrix} 2 & 1 & 1 \\ 0 & -8 & -2 \\ 0 & 8 & 3 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 1 & 1 \end{bmatrix} \begin{bmatrix} 5 \\ -12 \\ 14 \end{bmatrix} \rightarrow \begin{bmatrix} 2 & 1 & 1 \\ 0 & -8 & -2 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 5 \\ -12 \\ 2 \end{bmatrix}$$

d. You should now have an upper triangular matrix. Solve for  $x, y, z$ .

$$\begin{aligned} z &= 2 \\ y &= 1 \\ x &= 1 \end{aligned}$$

Homework:

Use elementary matrices to solve:

a.

$$3x + y + z = 3$$

$$12x - 2y + 3z = 3$$

$$-6x - 2y - 4z = 12$$

Answer:  
(3, 3, -9)

$$\begin{bmatrix} 1 & 0 & 0 \\ -4 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} 3 & 1 & 1 \\ 12 & -2 & 3 \\ -6 & -2 & -4 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \\ -4 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} 3 \\ 3 \\ 12 \end{bmatrix} \rightarrow \begin{bmatrix} 3 & 1 & 1 \\ 0 & -6 & -1 \\ -6 & -2 & -4 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 3 \\ -9 \\ 12 \end{bmatrix}$$

$$\begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ +2 & 0 & 0 \end{bmatrix} \begin{bmatrix} 3 & 1 & 1 \\ 0 & -6 & -1 \\ -6 & -2 & -4 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 2 & 0 & 0 \end{bmatrix} \begin{bmatrix} 3 \\ -9 \\ 12 \end{bmatrix} \rightarrow \begin{bmatrix} 3 & 1 & 1 \\ 0 & -6 & -1 \\ 0 & 0 & -2 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 3 \\ -9 \\ 18 \end{bmatrix}$$

$$-2z = 18$$

$$z = -9$$

$$-6y - (-9) = -9$$

$$-6y = -18$$

$$y = 3$$

$$x = 3$$

b.

$$x + y + z = 2$$

$$x + 3y + 3z = 0$$

$$x + 3y + 5z = 2$$

Answer:  
(3, -2, 1)

$$\begin{bmatrix} 1 & 0 & 0 \\ -1 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & 1 & 1 \\ 1 & 3 & 3 \\ 1 & 3 & 5 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \\ -1 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} 2 \\ 0 \\ 2 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & 1 & 1 \\ 0 & 2 & 2 \\ 1 & 3 & 5 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 2 \\ -2 \\ 2 \end{bmatrix}$$

$$\begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ -1 & 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & 1 & 1 \\ 0 & 2 & 2 \\ 1 & 3 & 5 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ -1 & 0 & 1 \end{bmatrix} \begin{bmatrix} 2 \\ -2 \\ 2 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & 1 & 1 \\ 0 & 2 & 2 \\ 0 & 2 & 4 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 2 \\ -2 \\ 0 \end{bmatrix}$$

$$\begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & -1 & 1 \end{bmatrix} \begin{bmatrix} 1 & 1 & 1 \\ 0 & 2 & 2 \\ 0 & 2 & 4 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & -1 & 1 \end{bmatrix} \begin{bmatrix} 2 \\ -2 \\ 0 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & 1 & 1 \\ 0 & 2 & 2 \\ 0 & 0 & 2 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 2 \\ -2 \\ 2 \end{bmatrix}$$

$$2z = 2 \rightarrow z = 1$$

$$2y + 2(1) = -2$$

$$y = -2$$

$$x = 3$$



### Part III: Inverses

Def: A matrix  $A$  is **invertible** if there exists a matrix  $A^{-1}$  such that  $AA^{-1} = A^{-1}A = I$ .

1. Prove this theorem: If a matrix does have an inverse, then it is unique.

Suppose there be two inverses,  $B$  and  $C$ .  
Then  $BA = I$  and  $CA = I$

$$\text{Then } B = B \cdot I = B \cdot (CA) = (BA)C = I \cdot C = C$$

2. The inverse of a  $2 \times 2$  matrix, if it exists, is given by the formula:  $\Rightarrow B = C$ .

$$\begin{bmatrix} a & b \\ c & d \end{bmatrix}^{-1} = \frac{1}{ad-bc} \begin{bmatrix} d & -b \\ -c & a \end{bmatrix}$$

We'll soon get to that the determinant of matrix  $A$  is  $ad-bc$ .

$\uparrow$   
 $\det A$

3. What is the inverse of a diagonal matrix  $A = \begin{bmatrix} d_1 & \dots & \dots & 0 \\ \vdots & d_2 & & \vdots \\ \vdots & & \ddots & \vdots \\ 0 & \dots & \dots & d_n \end{bmatrix}$ ?

$$A^{-1} = \begin{bmatrix} 1/d_1 & & & \\ & 1/d_2 & & \\ & & \ddots & \\ & & & 1/d_n \end{bmatrix} = I$$

4. Prove this theorem:  $(AB)^{-1} = B^{-1}A^{-1}$ .

Show that  $(AB)(B^{-1}A^{-1}) = I$

$$\begin{aligned} (AB)(B^{-1}A^{-1}) &= ABB^{-1}A^{-1} \\ &= A \cdot I \cdot A^{-1} \\ &= AA^{-1} = I \end{aligned}$$

Showing that  $(B^{-1}A^{-1})(AB) = I$  is analogous.

Gauss-Jordan Substitution: To solve for the inverse of a matrix, we can start out with this equation  $AX=I$  below:

$$\begin{bmatrix} a_{11} & \dots & \dots & a_{1n} \\ \vdots & & & \vdots \\ \vdots & & & \vdots \\ a_{n1} & \dots & \dots & a_{nn} \end{bmatrix} \begin{bmatrix} x_{11} & \dots & \dots & x_{1n} \\ \vdots & & & \vdots \\ \vdots & & & \vdots \\ x_{n1} & \dots & \dots & x_{nn} \end{bmatrix} = \begin{bmatrix} 1 & \dots & \dots & 0 \\ \vdots & & & \vdots \\ \vdots & & & \vdots \\ 0 & \dots & \dots & 1 \end{bmatrix}$$

And then perform Gaussian elimination until the A matrix is transformed into the identity matrix. Whatever is left in the X position will be in the inverse matrix.

5. Explain why this process works.

$$\begin{aligned} AX &= I \\ A^{-1}AX &= A^{-1}I \\ IX &= A^{-1} \end{aligned}$$

6. Use Gauss-Jordan elimination to find the inverse of:

$$\begin{bmatrix} 2 & 1 & 1 \\ 4 & -6 & 0 \\ -2 & 7 & 2 \end{bmatrix} \rightarrow \left[ \begin{array}{ccc|ccc} 2 & 1 & 1 & 1 & 0 & 0 \\ 4 & -6 & 0 & 0 & 1 & 0 \\ -2 & 7 & 2 & 0 & 0 & 1 \end{array} \right] \rightarrow \left[ \begin{array}{ccc|ccc} 2 & 0 & 3/4 & 3/4 & 1/8 & 1 \\ 0 & -8 & -2 & -2 & 1 & 0 \\ 0 & 0 & 1 & -1 & 1 & 1 \end{array} \right]$$

$$\rightarrow \left[ \begin{array}{ccc|ccc} 2 & 0 & 0 & 3/2 & -5/8 & -3 \\ 0 & -8 & -2 & -2 & 1 & 0 \\ 0 & 0 & 1 & -1 & 1 & 1 \end{array} \right] \rightarrow \left[ \begin{array}{ccc|ccc} 2 & 0 & 0 & 3/2 & -5/8 & -3 \\ 0 & -8 & 0 & -4 & 3 & 2 \\ 0 & 0 & 1 & -1 & 1 & 1 \end{array} \right] \rightarrow \left[ \begin{array}{ccc|ccc} 1 & 0 & 0 & 3/4 & -5/16 & -3/8 \\ 0 & 1 & 0 & 1/2 & -3/8 & -1/4 \\ 0 & 0 & 1 & -1 & 1 & 1 \end{array} \right]$$

$A^{-1}$

Homework:

1. Find the inverse of:


a.  $\begin{bmatrix} 5 & 6 \\ -1 & 2 \end{bmatrix}$

$$\begin{aligned} A^{-1} &= \frac{1}{5 \cdot 2 - 6 \cdot -1} \begin{bmatrix} 2 & -6 \\ 1 & 5 \end{bmatrix} \\ &= \frac{1}{16} \begin{bmatrix} 2 & -6 \\ 1 & 5 \end{bmatrix} \end{aligned}$$

b.

$$\begin{bmatrix} -1 & 2 & -1 \\ 2 & -1 & 0 \\ 0 & -1 & 2 \end{bmatrix}$$


$$A^{-1} = \begin{bmatrix} 1/2 & 3/4 & 1/4 \\ 1 & 1/2 & 1/2 \\ 1/2 & 1/4 & 3/4 \end{bmatrix}$$



c. Explain why  $\begin{bmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{bmatrix}$  is a rotation matrix in 2D using 90 degrees as an example.

$$\begin{bmatrix} \cos 90 & -\sin 90 \\ \sin 90 & \cos 90 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} -y \\ x \end{bmatrix}$$

Explain why  $\begin{bmatrix} \cos \theta & \sin \theta \\ -\sin \theta & \cos \theta \end{bmatrix}$  is its inverse using:



a.) Your 2x2 formula.

$$\frac{1}{\cos^2 \theta + \sin^2 \theta} \begin{bmatrix} \cos \theta & \sin \theta \\ -\sin \theta & \cos \theta \end{bmatrix} = \begin{bmatrix} \cos \theta & \sin \theta \\ -\sin \theta & \cos \theta \end{bmatrix}$$

b.) Geometric intuition of what would "undo" a rotation by an angle. Hint: You'll need to use even and odd properties of sine and cosine.

$$\begin{bmatrix} \cos(-\theta) & -\sin(-\theta) \\ \sin(-\theta) & \cos(-\theta) \end{bmatrix} = \begin{bmatrix} \cos \theta & \sin \theta \\ -\sin \theta & \cos \theta \end{bmatrix}$$

( $\cos \theta$  is even)  
( $\sin \theta$  is odd)

## Part IV: Determinants

Recall our 2x2 formula for the determinant:

$$\det \begin{bmatrix} a & b \\ c & d \end{bmatrix} = ad - bc$$

The following properties will extend to higher order matrices, but let's gain some intuition with the 2D case.

1. Show that when rows are exchanged, the determinant changes sign.

$$\det \begin{bmatrix} c & d \\ a & b \end{bmatrix} = bc - ad = -(ad - bc)$$

2. Show that multiplying a row by a scalar multiple of the original row multiplies the determinant by that amount.

$$\det \begin{bmatrix} ta & tb \\ c & d \end{bmatrix} = atd - btc = t(ad - bc)$$

3. Show that if two rows are equal, then the determinant is zero.

$$\det \begin{bmatrix} a & b \\ a & b \end{bmatrix} = ab - ba = 0$$

4. Show that if the matrix has a row (or column) of zeros, then the determinant is zero.

$$\det \begin{bmatrix} a & b \\ 0 & 0 \end{bmatrix} = 0$$

5. Show that subtracting a multiple of one row from another row leaves the same determinant (so that Gaussian elimination does not affect the determinant).

$$\det \begin{bmatrix} a & b \\ c - ta & d - tb \end{bmatrix} = (ad - atb) - (bc - tab) = ad - bc$$

6. Show that if A is diagonal, then the determinant is the product of its diagonal elements.

$$\det \begin{bmatrix} a & 0 \\ 0 & d \end{bmatrix} = ad$$

7. Show that if A is triangular, then the determinant is the product of its diagonal elements.

$$\det \begin{bmatrix} a & b \\ 0 & d \end{bmatrix} = ad$$

8. Show that A is invertible if and only if the determinant does not equal zero.

$$\frac{1}{ad-bc} \begin{bmatrix} d & -b \\ -c & a \end{bmatrix} \text{ only exists if } ad-bc \neq 0$$

9. Show that  $\det(AB) = \det(A)\det(B)$

$$\det \begin{pmatrix} a_1 & a_2 \\ a_3 & a_4 \end{pmatrix} \cdot \begin{pmatrix} b_1 & b_2 \\ b_3 & b_4 \end{pmatrix} = \det \begin{pmatrix} a_1 b_1 + a_2 b_3 & a_1 b_2 + a_2 b_4 \\ a_3 b_1 + a_4 b_3 & a_3 b_2 + a_4 b_4 \end{pmatrix}$$

$$= (a_1 b_1 + a_2 b_3)(a_3 b_2 + a_4 b_4) - (a_1 b_2 + a_2 b_4)(a_3 b_1 + a_4 b_3)$$

$$= a_1 b_1 a_3 b_2 + a_1 b_1 a_4 b_4 + a_2 b_3 a_3 b_2 + a_2 b_3 a_4 b_4$$

10. The transpose of a matrix A, denoted  $A^T$ , is obtained by putting the  $ij^{th}$  elements of A in the

$$ji^{th} \text{ position in } A^T. \text{ For example, the transpose of}$$

$$= (a_1 a_4 - a_2 a_3) \cdot (b_1 b_4 - b_2 b_3)$$

$$A = \begin{bmatrix} 1 & 2 \\ 3 & 4 \\ 5 & 6 \end{bmatrix} \text{ is } A^T = \begin{bmatrix} 1 & 3 & 5 \\ 2 & 4 & 6 \end{bmatrix}$$

Show that  $\det A = \det A^T$ .

$$\det \begin{bmatrix} a & b \\ c & d \end{bmatrix} = ad - bc$$

$$\det A^{-1} = \frac{1}{\det A}$$

11. Show that in the general case (non 2x2). To do this, use property #9 and the definition of inverse.

$$1 = \det I = \det (A \cdot A^{-1}) = \det A \cdot \det A^{-1}$$

### Higher Order Determinant Formula:

$\det(A)$  is equal to a linear combination of any row  $i$  (or column  $j$ ) times its cofactors:

$$\det A = a_{i1}C_{i1} + a_{i2}C_{i2} + \dots + a_{in}C_{in},$$

where the cofactor,  $C_{ij}$ , is the determinant of  $M_{ij}$  with the correct sign:

$$C_{ij} = (-1)^{i+j} \det M_{ij},$$

where  $M_{ij}$  is the minor matrix formed by deleting the  $i$ th row and  $j$ th column of  $A$ .

Here's a 3x3 example:

$$\begin{vmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{vmatrix} = a_{11}(-1)^{1+1} \begin{vmatrix} a_{22} & a_{23} \\ a_{32} & a_{33} \end{vmatrix} + a_{12}(-1)^{1+2} \begin{vmatrix} a_{21} & a_{23} \\ a_{31} & a_{33} \end{vmatrix} + a_{13}(-1)^{1+3} \begin{vmatrix} a_{21} & a_{22} \\ a_{31} & a_{32} \end{vmatrix}$$

Note: Because you'll multiply each determinant by a coefficient  $a_{ij}$ , you'll want to choose the row or column with the most zeros in order to simplify your calculations.

Example:

$$\begin{vmatrix} 2 & -1 & 0 & 0 \\ -1 & 2 & -1 & 0 \\ 1 & 0 & 1 & 3 \\ 2 & 4 & 1 & 2 \end{vmatrix} = 2 \begin{vmatrix} 2 & -1 & 0 \\ 0 & 1 & 3 \\ 4 & 1 & 2 \end{vmatrix} + (-1)(-1) \begin{vmatrix} -1 & -1 & 0 \\ 1 & 1 & 3 \\ 2 & 1 & 2 \end{vmatrix} \\ = 2 \left( 2 \begin{vmatrix} 1 & 3 \\ 1 & 2 \end{vmatrix} + (-1)(-1) \begin{vmatrix} 0 & 3 \\ 4 & 2 \end{vmatrix} \right) + 1 \left( (-1) \begin{vmatrix} 1 & 3 \\ 1 & 2 \end{vmatrix} + (-1)(-1) \begin{vmatrix} 1 & 3 \\ 2 & 2 \end{vmatrix} \right) \\ = 2(2(-1) + (1)(-12)) + ((-1)(-1) + (1)(-4)) \\ = -28 - 3 = -31$$

### Cramer's Rule:

The solution of  $Ax = b$  can be found by computing  $x_j = \frac{\det B_j}{\det A}$ , where  $B_j$  is the  $A$  matrix with its  $j$ th column replaced by the  $b$  vector.

Example:

The solution to  $x_1 + 3x_2 = 0$   
 $2x_1 + 4x_2 = 6$  is:

$$x_1 = \frac{\begin{vmatrix} 0 & 3 \\ 6 & 4 \end{vmatrix}}{\begin{vmatrix} 1 & 3 \\ 2 & 4 \end{vmatrix}} = \frac{-9}{-2} = 4.5$$

and

$$x_2 = \frac{\begin{vmatrix} 1 & 0 \\ 2 & 6 \end{vmatrix}}{\begin{vmatrix} 1 & 3 \\ 2 & 4 \end{vmatrix}} = \frac{-6}{-2} = 3$$

$$A \cdot x = b$$

$$\begin{bmatrix} 1 & 3 \\ 2 & 4 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 6 \end{bmatrix}$$

$$\begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} 4.5 \\ 3 \end{bmatrix}$$

### Homework:

Calculate the determinant of the following matrices. Try using determinant properties to simplify your calculations on a few of them.

a.  $\begin{bmatrix} 2 & 1 & 1 \\ 3 & 2 & -1 \\ 0 & 3 & 4 \end{bmatrix}$  19

$$(2)(+1) \begin{vmatrix} 2 & -1 \\ 3 & 4 \end{vmatrix} + (3)(-1) \begin{vmatrix} 1 & 1 \\ 3 & 4 \end{vmatrix}$$

$$= 2(8+3) - 3(4-3)$$

$$= 22 - 3 = 19$$

b.  $\begin{bmatrix} 2 & 1 & 5 & 1 \\ 3 & 1 & 1 & 0 \\ 0 & 0 & -2 & 1 \\ 0 & 3 & 0 & 1 \end{bmatrix}$  -55

$$(2)(+1) \begin{vmatrix} 1 & 0 \\ 0 & -2 \end{vmatrix} + 3(-1) \begin{vmatrix} 1 & 5 \\ 0 & -2 \end{vmatrix}$$

$$= 2[(-1)(+1) - 2(0)] - 3[(-1)(+1) - 2(5)]$$

$$= 2[-1 - 0] - 3[-1 - 10]$$

$$= 2(-1) - 3(-11) = -2 + 33 = 31$$

$$c. \begin{bmatrix} 2 & 2 & 2 & 2 \\ 2 & 2 & 2 & 2 \\ 0 & 0 & -2 & 1 \\ 0 & 3 & 0 & 1 \end{bmatrix}$$

0 since 2 rows same

$$d. \begin{bmatrix} 0 & 2 & 2 & 2 \\ 0 & 2 & 2 & 2 \\ 0 & 0 & -2 & 1 \\ 0 & 3 & 0 & 1 \end{bmatrix}$$

0 since column of 0's

$$3x_1 + x_2 = 2$$

2. Solve  $x_1 - x_2 = 6$  using Cramer's rule.

$$x_1 = \frac{\begin{vmatrix} 2 & 1 \\ 6 & -1 \end{vmatrix}}{\begin{vmatrix} 3 & 1 \\ 1 & -1 \end{vmatrix}} = 2$$

$$x_2 = \frac{\begin{vmatrix} 3 & 2 \\ 1 & 6 \end{vmatrix}}{\begin{vmatrix} 3 & 1 \\ 1 & -1 \end{vmatrix}} = -4$$

Inverse

$$\begin{bmatrix} 1 & -2 & 1 \\ -1 & 2 & 2 \\ 1 & 0 & 1 \end{bmatrix}^{-1}$$

$$\begin{bmatrix} -1/3 & -1/3 & 1 \\ -1/2 & 0 & 1/2 \\ 1/3 & 1/3 & 0 \end{bmatrix}$$



# Extra Inverse Problem

$$\left[ \begin{array}{ccc|ccc} 1 & -2 & 1 & 1 & 0 & 0 \\ -1 & 2 & 2 & 0 & 1 & 0 \\ 1 & 0 & 1 & 0 & 0 & 1 \end{array} \right]$$

New Row 2 = Row 1 + Row 2 :

$$\left[ \begin{array}{ccc|ccc} 1 & -2 & 1 & 1 & 0 & 0 \\ 0 & 0 & 3 & 1 & 1 & 0 \\ 1 & 0 & 1 & 0 & 0 & 1 \end{array} \right]$$

Switch Row 2 and Row 3 :

$$\left[ \begin{array}{ccc|ccc} 1 & -2 & 1 & 1 & 0 & 0 \\ 1 & 0 & 1 & 0 & 0 & 1 \\ 0 & 0 & 3 & 1 & 1 & 0 \end{array} \right]$$

New Row 2 = Row 1 - Row 2

$$\left[ \begin{array}{ccc|ccc} 1 & -2 & 1 & 1 & 0 & 0 \\ 0 & -2 & 0 & 1 & 0 & -1 \\ 0 & 0 & 3 & 1 & 1 & 0 \end{array} \right]$$

New Row 1 = Row 1 - Row 2

$$\left[ \begin{array}{ccc|ccc} 1 & 0 & 1 & 0 & 0 & 1 \\ 0 & -2 & 0 & 1 & 0 & -1 \\ 0 & 0 & 3 & 1 & 1 & 0 \end{array} \right]$$

continued

<?xml version="1.0" encoding="UTF-8"?>

<!DOCTYPE xpif SYSTEM "xpif-v02025.dtd">

<x

$$\text{New Row 1} = 3(\text{Row 1}) - \text{Row 3}$$

$$\left[ \begin{array}{ccc|ccc} 3 & 0 & 0 & -1 & -1 & 3 \\ 0 & -2 & 0 & 1 & 0 & -1 \\ 0 & 0 & 3 & 1 & 1 & 0 \end{array} \right]$$

$$\text{New Row 1} = \frac{1}{3} \text{Row 1}$$

$$\text{New Row 2} = -\frac{1}{2} \text{Row 2}$$

$$\text{New Row 3} = \frac{1}{3} \text{Row 3}$$

$$\left[ \begin{array}{ccc|ccc} 1 & 0 & 0 & -\frac{1}{3} & -\frac{1}{3} & 1 \\ 0 & 1 & 0 & -\frac{1}{2} & 0 & \frac{1}{2} \\ 0 & 0 & 1 & \frac{1}{3} & \frac{1}{3} & 0 \end{array} \right]$$

## Part V: Projections and Least Squares

Consider systems with more equations than unknowns:

$$2x = b_1$$

$$3x = b_2$$

$$4x = b_3$$

1. For what values of  $b$ 's will this system be solvable? for  $b$ 's in ratio 2:3:4

$$\begin{bmatrix} 2b_1 \\ 3b_1 \\ 4b_1 \end{bmatrix}$$

Suppose the  $b$ 's are not in that special form. Then this system is inconsistent, which arises all the time in real life, and still must be solved. We can determine  $x$  by solving part of the system and ignoring the rest of the equations, but this is hard to justify. Instead, let's choose  $x$  that minimizes the average error  $E$  from all  $n$  equations. Meaning, let's minimize:

$$E = (2x - b_1)^2 + (3x - b_2)^2 + (4x - b_3)^2$$

2. Why do we square each term above instead of just adding up the differences?

So errors don't cancel

3. Take the derivative of  $E(x)$  with respect to  $x$  and set it equal to zero in order to solve for your approximation,  $\hat{x}$ .

$$\begin{aligned} \frac{dE}{dx} &= 2(2x - b_1)(2) + 2(3x - b_2)(3) + 2(4x - b_3)(4) \\ &= 2[2(2x - b_1) + 3(3x - b_2) + 4(4x - b_3)] = 0 \\ &\Rightarrow \hat{x} = \frac{4x - 2b_1 + 9x - 3b_2 + 16x - 4b_3}{4 + 9 + 16} = \frac{2b_1 + 3b_2 + 4b_3}{2^2 + 3^2 + 4^2} \end{aligned}$$

4. Show that  $\hat{x} = \frac{a^T b}{a^T a}$ . This is called our **least squares solution**.

if  $a = \begin{bmatrix} 2 \\ 3 \\ 4 \end{bmatrix}$  and  $b = \begin{bmatrix} b_1 \\ b_2 \\ b_3 \end{bmatrix}$

then  $\hat{x} = \frac{a^T \cdot b}{a^T \cdot a}$

$$2x = 1$$

$$3x = 1$$

In our case, if we were trying to solve  $4x = 1$ , then our least squares solution would be

$\hat{x} = \frac{2+3+4}{2^2+3^2+4^2} = \frac{9}{29}$ , which is in between  $\frac{1}{4}$  and  $\frac{1}{2}$  (the solutions to the first and last equations individually).

How close does this approximation get to the  $b$  vector  $(1,1,1)$ ? The **projection** is given by

$$A\hat{x} = \begin{bmatrix} 2 \\ 3 \\ 4 \end{bmatrix} \frac{9}{29} = \begin{bmatrix} 18/29 \\ 27/29 \\ 36/29 \end{bmatrix}$$

More generally, let's try to solve  $A_{m \times n} x_{n \times 1} = b_{m \times 1}$ . We'll need find an approximation  $\hat{x}$  that minimizes the error of  $E = \|Ax - b\|$ . The proof for the formula for  $\hat{x}$  is a bit beyond the scope of this class, as we haven't talked about column and null space, but the general idea is that you need to solve  $A^T(b - A\hat{x}) = 0$ .

5. Show that  $A^T(b - A\hat{x}) = 0$  can be transformed to  $\hat{x} = (A^T A)^{-1} A^T b$ , our **least squares solution** formula.  $A^T b - A^T A \hat{x} = 0 \rightarrow A^T A \hat{x} = A^T b$

$$(A^T A)^{-1} A^T A \hat{x} = (A^T A)^{-1} A^T b$$

$$\hat{x} = (A^T A)^{-1} A^T b$$

How well does our solution do at approximating  $b$ ? The closest we get is the **projection**

$$A\hat{x} = A(A^T A)^{-1} A^T b$$

6. Show that if  $A$  is invertible then the above formula  $\hat{x} = (A^T A)^{-1} A^T b$  simplifies to the exact solution  $\hat{x} = A^{-1} b$ .

$$\begin{aligned} \hat{x} &= A^{-1} (A^T)^{-1} A^T b \\ &= A^{-1} \cdot I \cdot b \\ &= A^{-1} b \end{aligned}$$

## Part VI: Eigenvalues and Eigenvectors

Consider the following system of differential equations:

$$\frac{dx_1}{dt} = 4x_1 - 5x_2$$

$$\frac{dx_2}{dt} = 2x_1 - 3x_2$$

$$x_1(0) = 8, x_2(0) = 5$$

$$\begin{bmatrix} \frac{dx_1}{dt} \\ \frac{dx_2}{dt} \end{bmatrix} = \begin{bmatrix} 4 & -5 \\ 2 & -3 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix}$$

also written as  $\frac{dx}{dt} = Ax$

$$\vec{x} = \begin{bmatrix} x_1 \\ x_2 \end{bmatrix}$$

$$\frac{d\vec{x}}{dt} = \begin{bmatrix} 4 & -5 \\ 2 & -3 \end{bmatrix} \vec{x}$$

Note that if this was just a one dimensional system, then by separation of variables we would

find that  $\frac{dx}{dt} = ax$  has the general exponential solution  $x(t) = ce^{at}$

We shall take a direct approach and look for solutions with the same exponential dependence on  $t$  as in the one dimensional case.

1. Assume that  $x_1(t) = c_1 e^{\lambda t}$  and  $x_2(t) = c_2 e^{\lambda t}$  and then plug these formulas and their

derivatives into the original system above to obtain

$$\lambda c_1 = 4c_1 - 5c_2$$

$$\lambda c_2 = 2c_1 - 3c_2$$

$$\begin{aligned} \lambda c_1 e^{\lambda t} &= 4c_1 e^{\lambda t} - 5c_2 e^{\lambda t} \\ \lambda c_2 e^{\lambda t} &= 2c_1 e^{\lambda t} - 3c_2 e^{\lambda t} \end{aligned}$$

Note that

$$\lambda c_1 = 4c_1 - 5c_2$$

$$\lambda c_2 = 2c_1 - 3c_2$$

can be written in matrix form as  $Av = \lambda v$ , where

$$v = \begin{bmatrix} c_1 \\ c_2 \end{bmatrix}$$

$$\lambda \vec{v} = \begin{bmatrix} 4 & -5 \\ 2 & -3 \end{bmatrix} \vec{v}$$

2. Show that solving  $Av = \lambda v$  is equivalent to solving  $(A - \lambda I)v = 0$

$$Av - \lambda v = 0$$

$$Av - \lambda I v = 0$$

$$(A - \lambda I)v = 0$$

$$\det(A - \lambda I) = 0$$

$$\text{or } v = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

$$A \cdot B = 0$$

$$\begin{aligned} A &= 0 \\ B &= 0 \end{aligned}$$

For example, to solve:

$$x = 0$$

$$x + y = 8$$

$$x + 3y = 8$$

$$x + 4y = 20$$

$$A = \begin{bmatrix} 1 & 0 \\ 1 & 1 \\ 1 & 3 \\ 1 & 4 \end{bmatrix}$$

, we could calculate  $\hat{x} = (A^T A)^{-1} A^T b$ , where

$$\hat{x} = \begin{bmatrix} 1 \\ 4 \end{bmatrix} \quad A\hat{x} = \begin{bmatrix} 1 \\ 5 \\ 13 \\ 17 \end{bmatrix} \quad A^T = \begin{bmatrix} 1 & 1 & 1 & 1 \\ 0 & 1 & 3 & 4 \end{bmatrix}$$

$$A^T A = \begin{bmatrix} 4 & 8 \\ 8 & 26 \end{bmatrix} \quad (A^T A)^{-1} = \frac{1}{20} \begin{bmatrix} 13 & -4 \\ -4 & 2 \end{bmatrix}$$

Homework:  $\hat{x} = \frac{1}{20} \begin{bmatrix} 13 & -4 \\ -4 & 2 \end{bmatrix} \begin{bmatrix} 1 & 1 & 1 & 1 \\ 0 & 1 & 3 & 4 \end{bmatrix} \begin{bmatrix} 0 \\ 8 \\ 8 \\ 20 \end{bmatrix} = \frac{1}{20} \begin{bmatrix} 13 & -4 \\ -4 & 2 \end{bmatrix} \begin{bmatrix} 36 \\ 112 \end{bmatrix}$

Find the a.) least squares solution and b.) projection of the following system:

$$\begin{bmatrix} 1 & 2 \\ 1 & 3 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 4 \\ 5 \\ 6 \end{bmatrix}$$

$$\begin{bmatrix} 2 \\ 1 \end{bmatrix} \quad \begin{bmatrix} 4 \\ 5 \\ 0 \end{bmatrix}$$

$$A^T A = \begin{bmatrix} 2 & 5 \\ 5 & 13 \end{bmatrix}$$

$$(A^T A)^{-1} = \begin{bmatrix} 13 & -5 \\ -5 & 2 \end{bmatrix}$$

$$\begin{bmatrix} 13 & -5 \\ -5 & 2 \end{bmatrix} \begin{bmatrix} 1 & 1 & 0 \\ 2 & 3 & 0 \end{bmatrix} \begin{bmatrix} 4 \\ 5 \\ 6 \end{bmatrix} = \begin{bmatrix} 2 \\ 1 \end{bmatrix}$$

2. The goal of regression is to fit a mathematical model to a set of observed points. Say we're collecting data on the number of machine failures per day in some factory. Imagine we've got three data points: (day, number of failures) (1,1) (2,2) (3,2).

$$(x, b)$$

The goal is to find a linear equation that fits these points. We believe there's an underlying mathematical relationship that maps "days" uniquely to "number of machine failures" in the form  $b = C + Dx$ .

$$b = \frac{2}{3} + \frac{1}{2}x$$

Put this data into a matrix system and then use least squares to solve it.

$$C + D = 1$$

$$C + 2D = 2$$

$$C + 3D = 2$$

$$\begin{bmatrix} 1 & 1 \\ 1 & 2 \\ 1 & 3 \end{bmatrix} \begin{bmatrix} C \\ D \end{bmatrix} = \begin{bmatrix} 1 \\ 2 \\ 2 \end{bmatrix}$$

$$\hat{x} = \begin{bmatrix} 2/3 \\ 1/2 \end{bmatrix}$$

$$A\hat{x} = \begin{bmatrix} 1.1\bar{6} \\ 1.6 \\ 2.1\bar{6} \end{bmatrix}$$

If we are assuming that  $v$  is a nonzero eigenvector, then we must have the **characteristic equation**  $\det(A - \lambda I) = 0$ .

In the 2x2 case,

$$\det \begin{bmatrix} a_{11} - \lambda & a_{12} \\ a_{21} & a_{22} - \lambda \end{bmatrix} = (a_{11} - \lambda)(a_{22} - \lambda) - a_{12}a_{21} = 0$$

Once we solve this quadratic equation for  $\lambda$ , we will have our **eigenvalues**  $\lambda_1$  and  $\lambda_2$ , and then we can plug each value into  $(A - \lambda I)v = 0$  to solve for each **eigenvector**  $v_1$  and  $v_2$ . Note

that  $x(t) = e^{\lambda_1 t} v_1$  and  $x(t) = e^{\lambda_2 t} v_2$  are the pure exponential solutions to  $\frac{dx}{dt} = Ax$  and the superposition of both of them gives us the most general purely exponential solution:

$$x(t) = c_1 e^{\lambda_1 t} v_1 + c_2 e^{\lambda_2 t} v_2$$

Note: As this is not a full linear algebra course, we will only deal with real, unique roots (not repeated roots or imaginary roots). Their most general solution formulas are a bit different. A few things to know about eigenvalues:

a. Their form greatly affect the type of solution behavior. Purely imaginary eigenvalues lead to strictly periodic (sine and cosine) behavior. Whether or not the real part of the root is positive or negative affects whether there is exponential growth or decay.

b. Eigenvalues are related to the frequency of oscillations of the solutions. As a historical note, soldiers do not march in step as they go across a bridge because if they happen to march at the same frequency as one of the eigenvalues of the bridge, then the bridge begins to oscillate. (Just as a child's swing, you soon notice the natural frequency of the swing, and by matching it you go higher). An engineer tries to keep natural frequencies of his bridge away from those of wind. The Tacoma Narrows Bridge actually crashed in 1940 due to wind and the Broughton Bridge collapsed in 1831 due to soldiers marching.

Let's return to solving this example:

$$\frac{dx_1}{dt} = 4x_1 - 5x_2$$

$$\frac{dx_2}{dt} = 2x_1 - 3x_2$$

$$x_1(0) = 8, x_2(0) = 5$$

3. Find the eigenvalues.

$$\begin{vmatrix} 4-\lambda & -5 \\ 2 & -3-\lambda \end{vmatrix} = (4-\lambda)(-3-\lambda) + 10 = 0$$

$$-12 - \lambda + \lambda^2 + 10 = 0$$

$$\lambda^2 - \lambda - 2 = 0$$

$$(\lambda - 2)(\lambda + 1) = 0$$

$$\lambda = 2, -1$$

$$\lambda = 2 \quad \lambda = -1$$

4. Find each eigenvector.

$$\lambda = 2:$$

$$\begin{bmatrix} 4-2 & -5 \\ 2 & -3-2 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix} \rightarrow \begin{bmatrix} 2 & -5 \\ 2 & -5 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

$$2v_1 - 5v_2 = 0 \quad v = \begin{bmatrix} 5 \\ 2 \end{bmatrix}$$

$$\begin{bmatrix} 4+1 & -5 \\ 2 & -3+1 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix} \rightarrow \begin{bmatrix} 5 & -5 \\ 2 & -2 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

$$\begin{bmatrix} 5 & -5 \\ 2 & -2 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

$$\begin{bmatrix} 1 \\ 1 \end{bmatrix}$$

$$x(t) = c_1 e^{-t} \begin{bmatrix} 1 \\ 1 \end{bmatrix} + c_2 e^{2t} \begin{bmatrix} 5 \\ 2 \end{bmatrix}$$

5. Thus, the general solution is

$$x_1 = 3e^{-t} + 5e^{2t}$$

Use the initial conditions to

show that the particular solution is  $x_2 = 3e^{-t} + 2e^{2t}$ .

$$8 = c_1 + 5c_2$$

$$5 = c_1 + 2c_2$$

$$3 = 3c_2$$

$$c_2 = 1$$

$$c_1 = 3$$

$$x = 3e^{-t} \begin{bmatrix} 1 \\ 1 \end{bmatrix} + e^{2t} \begin{bmatrix} 5 \\ 2 \end{bmatrix}$$

Homework: Find the particular solution to:



Homework: Find the particular solution to:

$$x' = \begin{bmatrix} 1 & 2 \\ 3 & 2 \end{bmatrix} x,$$

$$x(0) = \begin{bmatrix} 0 \\ -5 \end{bmatrix}$$

a.

$$\det \begin{vmatrix} 1-\lambda & 2 \\ 3 & 2-\lambda \end{vmatrix} = (1-\lambda)(2-\lambda) - 6 = 0$$

$$2 - \lambda - 2\lambda + \lambda^2 - 6 = 0$$

$$\lambda^2 - 3\lambda - 4 = 0$$

$$(\lambda - 4)(\lambda + 1) = 0$$

$$\lambda = 4, \lambda = -1$$

$$\begin{bmatrix} 1-4 & 2 \\ 3 & 2-4 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

$$\begin{bmatrix} -3 & 2 \\ 3 & -2 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

$$-3v_1 + 2v_2 = 0$$

$$v_1 = 2, v_2 = 3$$

$$\begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 2 \\ 3 \end{bmatrix}$$

$$x' = \begin{bmatrix} -5 & 1 \\ 4 & -2 \end{bmatrix} x,$$

$$x(0) = \begin{bmatrix} 5 \\ 5 \end{bmatrix}$$

b.

$$x(t) = (-1)e^{4t} \begin{bmatrix} 2 \\ 3 \end{bmatrix} + (+2)e^{-1t} \begin{bmatrix} 1 \\ -1 \end{bmatrix}$$

$$x_1 = -2e^{4t} + 2e^{-1t}$$

$$x_2 = -3e^{4t} + 2e^{-1t}$$

$$\begin{bmatrix} 1-(-1) & 2 \\ 3 & 2-(-1) \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

$$\begin{bmatrix} 2 & 2 \\ 3 & 3 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

$$2v_1 + 2v_2 = 0$$

$$v_1 = -v_2$$

$$\begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 1 \\ -1 \end{bmatrix}$$

$$\begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = c_1 e^{4t} \begin{bmatrix} 2 \\ 3 \end{bmatrix} + c_2 e^{-1t} \begin{bmatrix} 1 \\ -1 \end{bmatrix}$$

$$0 = c_1 e^0 \cdot 2 + c_2 e^0 \cdot 1$$

$$-5 = c_1 e^0 \cdot 3 + c_2 e^0 \cdot (-1)$$

$$0 = 2c_1 + c_2$$

$$-5 = 3c_1 - c_2$$

$$-5 = 5c_1$$

$$-1 = c_1$$

$$c_2 = 2$$

$$A = S \cdot \Delta \cdot S^{-1}$$

$\uparrow$  eigenvectors of  $A$        $\uparrow$  eigenvalues of  $A$

## Part VII: Singular Value Decomposition

Suppose the  $n \times n$  matrix  $A$  has  $n$  distinct, real eigenvalues. If their corresponding eigenvectors are the columns of the matrix  $S$ , then  $S^{-1}AS$  is a diagonal matrix, with eigenvalues along the diagonal:

**Diagonalization**       $S^{-1}AS = \Lambda = \begin{bmatrix} \lambda_1 & & \\ & \lambda_2 & \\ & & \ddots \\ & & & \lambda_n \end{bmatrix}$

also written as  $S^{-1}AS = \Delta$  (1)

We call  $S$  the "eigenvector matrix" and  $\Lambda$  the "eigenvalue matrix"—using a capital lambda because of the small lambdas for the eigenvalues on its diagonal.

**Proof** Put the eigenvectors  $x_i$  in the columns of  $S$ , and compute  $AS$  by columns:

$$AS = A \begin{bmatrix} | & | & & | \\ x_1 & x_2 & \cdots & x_n \\ | & | & & | \end{bmatrix} = \begin{bmatrix} | & | & & | \\ \lambda_1 x_1 & \lambda_2 x_2 & \cdots & \lambda_n x_n \\ | & | & & | \end{bmatrix}$$

Then the trick is to split this last matrix into a quite different product  $S\Lambda$ :

$$\begin{bmatrix} \lambda_1 x_1 & \lambda_2 x_2 & \cdots & \lambda_n x_n \end{bmatrix} = \begin{bmatrix} | & | & & | \\ x_1 & x_2 & \cdots & x_n \\ | & | & & | \end{bmatrix} \begin{bmatrix} \lambda_1 & & & \\ & \lambda_2 & & \\ & & \ddots & \\ & & & \lambda_n \end{bmatrix}$$

$S$        $\Delta$

1. Show that  $AS = S\Lambda$  implies  $S^{-1}AS = \Lambda$  and  $A = S\Lambda S^{-1}$

$$A = S \cdot \Lambda \cdot S^{-1}$$

$$S^{-1}A = S^{-1}S \Lambda S^{-1}$$

$$S^{-1}A = \Lambda S^{-1}$$

$$S^{-1}AS = \Lambda S^{-1}S$$

$$S^{-1}AS = \Lambda$$

$$\frac{dx_1}{dt} = 4x_1 - 5x_2$$

$$\frac{dx_2}{dt} = 2x_1 - 3x_2$$

2. Recall our example from before  $x_1(0) = 8, x_2(0) = 5$ , where  $A = \begin{bmatrix} 4 & -5 \\ 2 & -3 \end{bmatrix}$  with

eigenvalues -1 and 2 and corresponding eigenvectors  $\begin{bmatrix} 1 \\ 1 \end{bmatrix}$  and  $\begin{bmatrix} 5 \\ 2 \end{bmatrix}$ . Show that  $A = S\Delta S^{-1}$ .

$$\begin{aligned} & \begin{bmatrix} 1 & 5 \\ 1 & 2 \end{bmatrix} \begin{bmatrix} -1 & 0 \\ 0 & 2 \end{bmatrix} \begin{bmatrix} 1 & 5 \\ 1 & 2 \end{bmatrix}^{-1} \\ & \begin{bmatrix} 1 & 5 \\ 1 & 2 \end{bmatrix} \begin{bmatrix} -1 & 0 \\ 0 & 2 \end{bmatrix} \frac{1}{-3} \begin{bmatrix} 2 & -5 \\ -1 & 1 \end{bmatrix} \\ & = \begin{bmatrix} 4 & -5 \\ 2 & -3 \end{bmatrix} \end{aligned}$$

The above formula works if  $A$  is a square matrix. What if  $A$  is not a square matrix? Well,  $AA^T$  and  $A^T A$  are always square. The diagonal matrix  $\Sigma$  has eigenvalues from  $A^T A$ . Those positive entries are called the singular values and will be placed along the diagonal.

**Singular Value Decomposition:** Any  $m$  by  $n$  matrix  $A$  can be factored into

$$A = U\Sigma V^T = (\text{orthogonal})(\text{diagonal})(\text{orthogonal}).$$

The columns of  $U$  ( $m$  by  $m$ ) are eigenvectors of  $AA^T$ , and the columns of  $V$  ( $n$  by  $n$ ) are eigenvectors of  $A^T A$ . The  $r$  singular values on the diagonal of  $\Sigma$  ( $m$  by  $n$ ) are the square roots of the nonzero eigenvalues of both  $AA^T$  and  $A^T A$ .

We won't prove this theorem but we will use it.

3. We'll decompose  $A = \begin{bmatrix} 3 & 2 & 2 \\ 2 & 3 & -2 \end{bmatrix}$  into this form using the following steps:

a. Find the eigenvalues of  $AA^T$ .  $AA^T = \begin{bmatrix} 17 & 8 \\ 8 & 17 \end{bmatrix}$   
 $\downarrow \lambda^2 - 34\lambda + 25 = 0 \rightarrow \lambda = 25, 9$   
 $0 = 5, 3$

b. Find the eigenvalues of  $A^T A$ .  $\begin{bmatrix} 13 & 12 & 2 \\ 12 & 13 & -2 \\ 2 & -2 & 8 \end{bmatrix}$   $-\lambda^3 + 34\lambda^2 - 25\lambda = 0$   
 $\lambda = 0, 25, 9$   
 positive

c. The only positive eigenvalues are 25 and 9. Stack the square roots of these eigenvalues

together to get  $\Sigma = \begin{bmatrix} 5 & 0 & 0 \\ 0 & 3 & 0 \end{bmatrix}$ . Note that the size of this matrix will always have the number of rows as  $AA^T$  and the number of columns of  $A^T A$ .

d. Find the normalized eigenvectors of  $AA^T$ . Stack these to form U:

$$U = \begin{bmatrix} 1/\sqrt{2} & 1/\sqrt{2} \\ 1/\sqrt{2} & -1/\sqrt{2} \end{bmatrix}$$

$\begin{bmatrix} 17-25 & 8 \\ 8 & 17-25 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix} \rightarrow \begin{bmatrix} -8 & 8 \\ 8 & -8 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix} \rightarrow x = y$   
 $\frac{1}{\sqrt{2}}(1, 1)$

$\begin{bmatrix} 17-9 & 8 \\ 8 & 17-9 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix} \rightarrow x = y$   
 $\frac{1}{\sqrt{2}}(1, -1)$

$$\begin{bmatrix} -12 & 12 & 2 \\ 12 & -12 & -2 \\ 2 & -2 & -17 \end{bmatrix} \rightarrow \begin{bmatrix} -12 & 12 & 2 \\ 0 & 0 & 0 \\ 0 & 0 & -17 \end{bmatrix}$$

e. Find the normalized eigenvectors of  $A^T A$ . Stack these to form V:

$$V = \begin{bmatrix} 1/\sqrt{2} & 1/\sqrt{18} & 2/3 \\ 1/\sqrt{2} & -1/\sqrt{18} & -2/3 \\ 0 & 4/\sqrt{18} & -1/3 \end{bmatrix}$$

$$\begin{bmatrix} -12 & 12 & 2 \\ 0 & 0 & 0 \\ 0 & 0 & -10 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

$$z=0, x=y \quad \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix} \rightarrow \begin{bmatrix} 1/\sqrt{2} \\ 1/\sqrt{2} \\ 0 \end{bmatrix}$$

$$\lambda = 9$$

$$\begin{bmatrix} 4 & 0 & -1 & | & 0 \\ 0 & 4 & 1 & | & 0 \\ 0 & 0 & 0 & | & 0 \end{bmatrix} \rightarrow$$

$$x = 1/4 z$$

$$y = -1/4 z$$

$$\begin{bmatrix} 1 \\ -1 \\ 4 \end{bmatrix} \rightarrow$$

$$\begin{bmatrix} 1/\sqrt{18} \\ -1/\sqrt{18} \\ 4/\sqrt{18} \end{bmatrix}$$

$$\lambda = 0$$

$$\begin{bmatrix} 0 & 0 & 0 & | & 0 \\ 0 & 0 & 0 & | & 0 \\ 0 & 0 & -2 & | & 0 \end{bmatrix} \rightarrow$$

$$x = -y, z = 1/2 y$$

$$\begin{bmatrix} 2 \\ -2 \\ -1 \end{bmatrix} \rightarrow \begin{bmatrix} 2/3 \\ -2/3 \\ -1/3 \end{bmatrix}$$

f. Transpose V and we now have our decomposition of A:

$$A = \begin{bmatrix} 1/\sqrt{2} & 1/\sqrt{2} \\ 1/\sqrt{2} & -1/\sqrt{2} \end{bmatrix} \begin{bmatrix} 5 & 0 & 0 \\ 0 & 3 & 0 \end{bmatrix} \begin{bmatrix} 1/\sqrt{2} & 1/\sqrt{2} & 0 \\ 1/\sqrt{18} & -1/\sqrt{18} & 4/\sqrt{18} \\ 2/3 & -2/3 & -1/3 \end{bmatrix}$$

Q1 Q2 Q3 Q4 Q5 Q6 Q7 Q8 Q9 Q10 Q11 Q12 Q13 Q14 Q15 Q16 Q17 Q18 Q19 Q20 Q21 Q22 Q23 Q24 Q25 Q26 Q27 Q28 Q29 Q30 Q31 Q32 Q33 Q34 Q35 Q36 Q37 Q38 Q39 Q40 Q41 Q42 Q43 Q44 Q45 Q46 Q47 Q48 Q49 Q50 Q51 Q52 Q53 Q54 Q55 Q56 Q57 Q58 Q59 Q60 Q61 Q62 Q63 Q64 Q65 Q66 Q67 Q68 Q69 Q70 Q71 Q72 Q73 Q74 Q75 Q76 Q77 Q78 Q79 Q80 Q81 Q82 Q83 Q84 Q85 Q86 Q87 Q88 Q89 Q90 Q91 Q92 Q93 Q94 Q95 Q96 Q97 Q98 Q99 Q100

# Application of SVD: Image compression

Suppose a satellite takes a picture and wants to send it back to earth. The picture may contain 1000x1000 pixels – a million little squares, each with color. We can code the colors and send back 1,000,000 numbers. It is better to find the *essential* information inside the 1000x1000 matrix and send only that.

Suppose we know the SVD. The key is in the singular values (in  $\Sigma$ ). Typically, some  $\sigma$ 's are significant and others are extremely small. If we keep 20 and throw away 980, then we send only the corresponding 20 columns of  $U$  and  $V$ . The other 980 columns are multiplied in  $U\Sigma V^T$  by the small  $\sigma$ 's that are being ignored. We can do the matrix multiplication as columns times rows:

$$A = U\Sigma V^T = \underbrace{u_1\sigma_1 v_1^T + u_2\sigma_2 v_2^T + \dots + u_{20}\sigma_{20} v_{20}^T}_{20} \quad (3)$$

Any matrix is the sum of  $r$  matrices of rank 1. If only 20 terms are kept, we send 20 times 2000 numbers instead of a million (25 to 1 compression).

The pictures are really striking, as more and more singular values are included. At first you see nothing, and suddenly you recognize everything. The cost is in computing the SVD—this has become much more efficient, but it is expensive for a big matrix.

## Homework:

1. Explain the previous claim that only 20 times 2000 numbers will need to be sent.

Handwritten work for homework:

$$\begin{aligned}
 & \text{Matrix } U \text{ (size } 1000 \times 1000 \text{)} \text{ is partitioned into } U_{20} \text{ (size } 1000 \times 20 \text{)} \text{ and } U_{980} \text{ (size } 1000 \times 980 \text{)}. \\
 & \text{Matrix } \Sigma \text{ (size } 1000 \times 1000 \text{)} \text{ is partitioned into } \Sigma_{20} \text{ (size } 20 \times 20 \text{)} \text{ and } \Sigma_{980} \text{ (size } 980 \times 980 \text{)}. \\
 & \text{Matrix } V^T \text{ (size } 1000 \times 1000 \text{)} \text{ is partitioned into } V_{20}^T \text{ (size } 20 \times 1000 \text{)} \text{ and } V_{980}^T \text{ (size } 980 \times 1000 \text{)}. \\
 & \text{The product } U\Sigma V^T \text{ is approximated by } U_{20}\Sigma_{20}V_{20}^T. \\
 & \text{Dimensions: } (1000 \times 20) \times (20 \times 20) \times (20 \times 1000) = 1000 \times 20 \times 1000. \\
 & \text{Total numbers sent: } 20 \times 1000 + 20 + 20 \times 1000 = 40,020.
 \end{aligned}$$

$$A = \begin{bmatrix} -1 \\ 2 \\ 2 \end{bmatrix}$$

2. Show that SVD decomposition of  $A$  is given by:

$$A = \begin{bmatrix} -1/3 & 2/\sqrt{5} & 2/\sqrt{5} \\ 2/3 & 0 & 1/\sqrt{5} \\ 2/3 & 1/\sqrt{5} & 0 \end{bmatrix} \begin{bmatrix} 3 \\ 0 \\ 0 \end{bmatrix} [1]$$

3. Show that the SVD decomposition of  $A = \begin{bmatrix} 1 & 2 \\ 2 & 1 \end{bmatrix}$  is given by:

$$A = \begin{bmatrix} 1/\sqrt{2} & 1/\sqrt{2} \\ 1/\sqrt{2} & -1/\sqrt{2} \end{bmatrix} \begin{bmatrix} 3 & 0 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} 1/\sqrt{2} & 1/\sqrt{2} \\ -1/\sqrt{2} & 1/\sqrt{2} \end{bmatrix}$$



$$A = \begin{bmatrix} -1 \\ 2 \\ 2 \end{bmatrix} \quad A^T = \begin{bmatrix} -1 & 2 & 2 \end{bmatrix}$$

$$1.) \quad AA^T = \begin{bmatrix} 1 & -2 & -2 \\ -2 & 4 & 4 \\ -2 & 4 & 4 \end{bmatrix}$$

Find eigenvalues:

$$\det \begin{bmatrix} 1-\lambda & -2 & -2 \\ -2 & 4-\lambda & 4 \\ -2 & 4 & 4-\lambda \end{bmatrix} = (1-\lambda) \begin{vmatrix} 4-\lambda & 4 \\ 4 & 4-\lambda \end{vmatrix} + 2 \begin{vmatrix} -2 & 4 \\ -2 & 4-\lambda \end{vmatrix} - 2 \begin{vmatrix} -2 & 4-\lambda \\ -2 & 4 \end{vmatrix}$$

$$= (1-\lambda)[(4-\lambda)^2 - 16] + 2[-2(4-\lambda) + 8] - 2[-8 + 2(4-\lambda)]$$

$$= (1-\lambda)[16 - 8\lambda + \lambda^2 - 16] + 2[-8 + 2\lambda + 8] - 2[-8 + 8 - 2\lambda]$$

$$= (1-\lambda)[-8\lambda + \lambda^2] + 2[2\lambda] - 2[-2\lambda]$$

$$= -8\lambda + \lambda^2 + 8\lambda^2 - \lambda^3 + 4\lambda + 4\lambda$$

$$= -\lambda^3 + 9\lambda^2 = -\lambda^2(\lambda - 9) = 0 \Rightarrow \lambda = 0, \lambda = 9$$

$$2.) \quad A^T A = \begin{bmatrix} -1 & 2 & 2 \end{bmatrix}_{1 \times 3} \begin{bmatrix} -1 \\ 2 \\ 2 \end{bmatrix}_{3 \times 1} = \begin{bmatrix} 9 \end{bmatrix}_{1 \times 1}$$

Find eigenvalues:

$$\det[9 - \lambda] = 0 \Rightarrow \lambda = 9$$

3.)  $\Sigma$  must have same # of rows as  $AA^T$  and same # of columns as  $A^T A$ , so  $3 \times 1$

$$\Sigma = \begin{bmatrix} 3 \\ 0 \\ 0 \end{bmatrix} \quad \leftarrow \text{take square root}$$

4.) To find  $V$ , we'll need to find the eigenvectors of  $A^T A$ .

$$\lambda = 9:$$

$$\text{Solve } [9-9][v_1] = [0]$$

$\Rightarrow v_1$  can be anything so  
by convention, choose  $v_1 = [1]$ .

$$\Rightarrow \boxed{V^T = [1]}$$

5.) To find  $U$ , we'll need to find the eigenvectors of  $A A^T$ .

$$\lambda = 9:$$

$$\begin{bmatrix} 1-9 & -2 & -2 \\ -2 & 4-9 & 4 \\ -2 & 4 & 4-9 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \\ v_3 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix} \rightarrow \begin{bmatrix} -8 & -2 & -2 \\ -2 & -5 & 4 \\ -2 & 4 & -5 \end{bmatrix} \left| \begin{array}{l} 0 \\ 9 \\ 0 \end{array} \right.$$

$$\rightarrow \begin{bmatrix} -8 & -2 & -2 & | & 0 \\ -2 & -5 & 4 & | & 0 \\ 0 & 9 & -9 & | & 0 \end{bmatrix} \rightarrow \begin{bmatrix} -8 & -2 & -2 & | & 0 \\ -2 & -5 & 4 & | & 0 \\ 0 & 1 & -1 & | & 0 \end{bmatrix} \rightarrow$$

$$\rightarrow \begin{bmatrix} -8 & -2 & -2 & | & 0 \\ 0 & -18 & 18 & | & 0 \\ 0 & 1 & -1 & | & 0 \end{bmatrix} \rightarrow \begin{bmatrix} -8 & -2 & -2 & | & 0 \\ 0 & -1 & 1 & | & 0 \\ 0 & 1 & -1 & | & 0 \end{bmatrix} \rightarrow \begin{bmatrix} -4 & -1 & -1 & | & 0 \\ 0 & -1 & 1 & | & 0 \\ 0 & 0 & 0 & | & 0 \end{bmatrix}$$

$$\rightarrow \begin{bmatrix} -4 & 0 & 0 & | & 0 \\ 0 & -1 & 1 & | & 0 \\ 0 & 0 & 0 & | & 0 \end{bmatrix} \Rightarrow \begin{bmatrix} 2 & 0 & 1 & | & 0 \\ 0 & -1 & 1 & | & 0 \\ 0 & 0 & 0 & | & 0 \end{bmatrix}$$

$$-v_2 + v_3 = 0$$

$$\Rightarrow v_2 = v_3 \rightarrow v_2 = v_3 = 1$$

$$2v_1 + v_3 = 0 \Rightarrow v_1 = -\frac{1}{2}$$

$$\begin{bmatrix} v_1 \\ v_2 \\ v_3 \end{bmatrix} = \begin{bmatrix} -1 \\ 2 \\ 2 \end{bmatrix}$$

$$\text{Normalize: } \begin{bmatrix} v_1 \\ v_2 \\ v_3 \end{bmatrix} = \frac{1}{\sqrt{2^2 + 2^2 + 1^2}} = \frac{1}{3} \begin{bmatrix} -1 \\ 2 \\ 2 \end{bmatrix} =$$

$$\begin{bmatrix} -1/3 \\ 2/3 \\ 2/3 \end{bmatrix}$$

(2)

$$\lambda = 0:$$

$$\left[ \begin{array}{ccc|c} 1-0 & -2 & -2 & 0 \\ -2 & 4-0 & 4 & 0 \\ -2 & 4 & 4-0 & 0 \end{array} \right] \rightarrow \left[ \begin{array}{ccc|c} 1 & -2 & -2 & 0 \\ -2 & 4 & 4 & 0 \\ -2 & 4 & 4 & 0 \end{array} \right]$$

$$\rightarrow \left[ \begin{array}{ccc|c} 1 & -2 & -2 & 0 \\ -2 & 4 & 4 & 0 \\ 0 & 0 & 0 & 0 \end{array} \right] \Rightarrow \left[ \begin{array}{ccc|c} 1 & -2 & -2 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{array} \right]$$

$$\rightarrow v_1 - 2v_2 - 2v_3 = 0 \rightarrow v_1 = 2v_2 + 2v_3$$

There are actually two different types of solutions to this system.

First: let  $v_2 = 0, v_3 = 1$ . Then  $\begin{bmatrix} v_1 \\ v_2 \\ v_3 \end{bmatrix} = \begin{bmatrix} 2 \\ 0 \\ 1 \end{bmatrix}$

Normalized, gives

$$\begin{bmatrix} v_1 \\ v_2 \\ v_3 \end{bmatrix} = \begin{bmatrix} 2/\sqrt{5} \\ 0 \\ 1/\sqrt{5} \end{bmatrix}$$

Second, let  $v_2 = 1, v_3 = 0$ . Then  $\begin{bmatrix} v_1 \\ v_2 \\ v_3 \end{bmatrix} = \begin{bmatrix} 2 \\ 1 \\ 0 \end{bmatrix}$

$$\text{So... } U = \begin{bmatrix} -1/3 & 2/\sqrt{5} & 2/\sqrt{5} \\ 2/3 & 0 & 1/\sqrt{5} \\ 2/3 & 1/\sqrt{5} & 0 \end{bmatrix}$$

In summary,

$$A = U \Sigma V^T = \begin{bmatrix} -1/3 & 2/\sqrt{5} & 2/\sqrt{5} \\ 2/3 & 0 & 1/\sqrt{5} \\ 2/3 & 1/\sqrt{5} & 0 \end{bmatrix} \begin{bmatrix} 3 \\ 0 \\ 0 \end{bmatrix} \begin{bmatrix} 1 \end{bmatrix}$$





EX 4)

Ex 3:  $A = \begin{bmatrix} -1 \\ 2 \\ 2 \end{bmatrix}$   $A^T = [-1 \ 2 \ 2]$

1.)  $AA^T = \begin{bmatrix} 1 & -2 & -2 \\ -2 & 4 & 4 \\ -2 & 4 & 4 \end{bmatrix}$

$\begin{vmatrix} 1-\lambda & -2 & -2 \\ -2 & 4-\lambda & 4 \\ -2 & 4 & 4-\lambda \end{vmatrix} = -\lambda^3 + 9\lambda^2 = -\lambda^2(\lambda - 9) \Rightarrow \lambda = 9, \lambda_2 = 0, \lambda_3 = 0$   
 $\sigma_1 = \sqrt{9} = 3$   
 3 rows of  $\Sigma$

2.)  $ATA = [9]$   $|9 - \lambda| = 0 \rightarrow \lambda = 9$   $\sigma_1 = 3$   
 the col of  $\Sigma$

$\Sigma = \begin{bmatrix} 3 \\ 0 \\ 0 \end{bmatrix}$

3.) e vectors of  $ATA$ :

$[9 - 9][x] = [0] \rightarrow x = [1] \rightarrow v = [1]$

4.)  $v^T = [1]$

5.)  $u_1 = \frac{Av_1}{\sigma_1} = \frac{\begin{bmatrix} -1 \\ 2 \\ 2 \end{bmatrix} [1]}{3} = \begin{bmatrix} -1/3 \\ 2/3 \\ 2/3 \end{bmatrix}$

6.) Need two more e vectors for  $U$ :

$\lambda_2 = 0$ :  ~~$[9 - 0][x] = [0] \rightarrow x =$~~

$\begin{bmatrix} 1 & -2 & -2 \\ -2 & 4 & 4 \\ -2 & 4 & 4 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & -2 & -2 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$   
 $x = 2y + 2z$   
 let  $\begin{cases} y=0 \\ z=1 \end{cases} \rightarrow v_2 = \begin{bmatrix} 2 \\ 0 \\ 1 \end{bmatrix}$

let  $\begin{cases} z=0 \\ y=1 \end{cases} \rightarrow \begin{bmatrix} 2 \\ 1 \\ 0 \end{bmatrix}$  but normalize  $v_2 = \begin{bmatrix} 2/\sqrt{5} \\ 0 \\ 1/\sqrt{5} \end{bmatrix}$   $v_3 = \begin{bmatrix} 2/\sqrt{5} \\ 1/\sqrt{5} \\ 0 \end{bmatrix}$   
 $U = \begin{bmatrix} -1/3 & 2/\sqrt{5} & 2/\sqrt{5} \\ 2/3 & 0 & 1/\sqrt{5} \\ 2/3 & 1/\sqrt{5} & 0 \end{bmatrix}$

$A = \begin{bmatrix} -1 \\ 2 \\ 2 \end{bmatrix} = \begin{bmatrix} -1/3 & 2/\sqrt{5} & 2/\sqrt{5} \\ 2/3 & 0 & 1/\sqrt{5} \\ 2/3 & 1/\sqrt{5} & 0 \end{bmatrix} \begin{bmatrix} 3 \\ 0 \\ 0 \end{bmatrix} [1]$

$$A = \begin{bmatrix} -1 & 1 & 0 \\ 0 & -1 & 1 \end{bmatrix} \quad A^T = \begin{bmatrix} -1 & 0 \\ 1 & -1 \\ 0 & 1 \end{bmatrix}$$

HW don't do

$$AA^T = \begin{bmatrix} 2 & -1 \\ -1 & 2 \end{bmatrix}$$

$$\begin{vmatrix} 2-\lambda & -1 \\ -1 & 2-\lambda \end{vmatrix} = \lambda^2 - 4\lambda + 3 = (\lambda-1)(\lambda-3) = 0 \quad \sigma_1 = \sqrt{3}, \sigma_2 = 1$$

$\lambda = \frac{1 \pm \sqrt{3}}{2}$  rows of  $\Sigma$

$$A^T A = \begin{bmatrix} 1 & -1 & 0 \\ -1 & 2 & -1 \\ 0 & -1 & 1 \end{bmatrix}$$

$$\begin{vmatrix} 1-\lambda & -1 & 0 \\ -1 & 2-\lambda & -1 \\ 0 & -1 & 1-\lambda \end{vmatrix} = -\lambda^3 + 4\lambda^2 - 3\lambda = -\lambda(\lambda-3)(\lambda-1)$$

3 cols of  $\Sigma \rightarrow \lambda_1 = 3, \lambda_2 = 1, \lambda_3 = 0$   
 $\sigma_1 = \sqrt{3}, \sigma_2 = 1$

$$\Sigma = \begin{bmatrix} \sqrt{3} & 0 & 0 \\ 0 & 1 & 0 \end{bmatrix}$$

3) e. vector of  $A^T A$

$$\lambda_1 = 3: \begin{bmatrix} -2 & -1 & 0 \\ -1 & -1 & -1 \\ 0 & -1 & -2 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix} \rightarrow \begin{bmatrix} -2 & -1 & 0 \\ 0 & 1 & 2 \\ 0 & 0 & 0 \end{bmatrix} \rightarrow \begin{matrix} x = -\frac{1}{2}y \\ z = -\frac{1}{2}y \end{matrix}$$

$$\rightarrow v_1 = \begin{bmatrix} 1/\sqrt{6} \\ -2/\sqrt{6} \\ 1/\sqrt{6} \end{bmatrix}$$

$$\lambda_2 = 1: \begin{bmatrix} 0 & -1 & 0 \\ -1 & 1 & -1 \\ 0 & -1 & 0 \end{bmatrix} \rightarrow y=0 \rightarrow x=-z \quad \begin{bmatrix} -1 \\ 0 \\ 1 \end{bmatrix} \rightarrow v_2 = \begin{bmatrix} -1/\sqrt{2} \\ 0 \\ 1/\sqrt{2} \end{bmatrix}$$

$$\lambda_3 = 0: \begin{bmatrix} 1 & -1 & 0 \\ -1 & 2 & -1 \\ 0 & -1 & 1 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & -1 & 0 \\ 0 & 1 & -1 \\ 0 & 0 & 0 \end{bmatrix} \rightarrow x=y=z \quad v_3 = \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} \rightarrow v_3 = \begin{bmatrix} 1/\sqrt{3} \\ 1/\sqrt{3} \\ 1/\sqrt{3} \end{bmatrix}$$

$$V = \begin{bmatrix} 1/\sqrt{6} & -1/\sqrt{2} & 1/\sqrt{3} \\ -2/\sqrt{6} & 0 & 1/\sqrt{3} \\ 1/\sqrt{6} & 1/\sqrt{2} & 1/\sqrt{3} \end{bmatrix} \quad V^T = \begin{bmatrix} 1/\sqrt{6} & -2/\sqrt{6} & 1/\sqrt{6} \\ -1/\sqrt{2} & 0 & 1/\sqrt{2} \\ 1/\sqrt{3} & 1/\sqrt{3} & 1/\sqrt{3} \end{bmatrix}$$

$$u_1 = \frac{1}{\sigma_1} A v_1 = \frac{1}{\sqrt{3}} \begin{bmatrix} -1 & 1 & 0 \\ 0 & -1 & 1 \end{bmatrix} \begin{bmatrix} 1/\sqrt{6} \\ -2/\sqrt{6} \\ 1/\sqrt{6} \end{bmatrix} = \frac{1}{\sqrt{3}} \begin{bmatrix} -3/\sqrt{6} \\ 3/\sqrt{6} \end{bmatrix} = \begin{bmatrix} -1/\sqrt{2} \\ 1/\sqrt{2} \end{bmatrix}$$

$$u_2 = \frac{1}{\sigma_2} \begin{bmatrix} -1 & 1 & 0 \\ 0 & -1 & 1 \end{bmatrix} \begin{bmatrix} -1/\sqrt{2} \\ 0 \\ 1/\sqrt{2} \end{bmatrix} = \begin{bmatrix} 1/\sqrt{2} \\ 1/\sqrt{2} \end{bmatrix}$$

$$U = \begin{bmatrix} -1/\sqrt{2} & 1/\sqrt{2} \\ 1/\sqrt{2} & 1/\sqrt{2} \end{bmatrix}$$

$$A = U \Sigma V^T = \begin{bmatrix} -1/\sqrt{2} & 1/\sqrt{2} \\ 1/\sqrt{2} & 1/\sqrt{2} \end{bmatrix} \begin{bmatrix} \sqrt{3} & 0 & 0 \\ 0 & 1 & 0 \end{bmatrix} \begin{bmatrix} 1/\sqrt{6} & -2/\sqrt{6} & 1/\sqrt{6} \\ -1/\sqrt{2} & 0 & 1/\sqrt{2} \\ 1/\sqrt{3} & 1/\sqrt{3} & 1/\sqrt{3} \end{bmatrix}$$



Ex 3)  $A = \begin{bmatrix} 1 & 2 \\ 2 & 1 \end{bmatrix}$

1.)  $AA^T = \begin{bmatrix} 5 & 4 \\ 4 & 5 \end{bmatrix}$   $\begin{vmatrix} 5-\lambda & 4 \\ 4 & 5-\lambda \end{vmatrix} = \lambda^2 - 10\lambda + 9 = (\lambda-1)(\lambda-9) = 0$

2 rows of  $\Sigma \rightarrow \lambda_1 = 9 \quad \lambda_2 = 1$   
 $\sigma_1 = 3 \quad \sigma_2 = 1$

2.)  $A^T A = \begin{bmatrix} 5 & 4 \\ 4 & 5 \end{bmatrix} \rightarrow$  same  
 $\Sigma = \begin{bmatrix} 3 & 0 \\ 0 & 1 \end{bmatrix}$

3.)  $\lambda_1 = 9$   $\begin{bmatrix} 5-9 & 4 \\ 4 & 5-9 \end{bmatrix} = \begin{bmatrix} -4 & 4 & | & 0 \\ 4 & -4 & | & 0 \end{bmatrix} \rightarrow x = y \rightarrow v_1 = \begin{bmatrix} 1/\sqrt{2} \\ 1/\sqrt{2} \end{bmatrix}$   
 $\lambda_2 = 1$   $\begin{bmatrix} 4 & 4 & | & 0 \\ 4 & 4 & | & 0 \end{bmatrix} \rightarrow x = -y \rightarrow v_2 = \begin{bmatrix} -1/\sqrt{2} \\ 1/\sqrt{2} \end{bmatrix}$

$V = \begin{bmatrix} 1/\sqrt{2} & -1/\sqrt{2} \\ 1/\sqrt{2} & 1/\sqrt{2} \end{bmatrix}$

4.)  $V^T = \begin{bmatrix} 1/\sqrt{2} & 1/\sqrt{2} \\ -1/\sqrt{2} & 1/\sqrt{2} \end{bmatrix}$

4.)  $u_1 = \frac{1}{3} \begin{bmatrix} 1 & 2 \\ 2 & 1 \end{bmatrix} \begin{bmatrix} 1/\sqrt{2} \\ 1/\sqrt{2} \end{bmatrix} = \begin{bmatrix} 1/\sqrt{2} \\ 1/\sqrt{2} \end{bmatrix}$

$u_2 = \frac{1}{1} \begin{bmatrix} 1 & 2 \\ 2 & 1 \end{bmatrix} \begin{bmatrix} -1/\sqrt{2} \\ 1/\sqrt{2} \end{bmatrix} = \begin{bmatrix} 1/\sqrt{2} \\ -1/\sqrt{2} \end{bmatrix}$

$U = \begin{bmatrix} 1/\sqrt{2} & 1/\sqrt{2} \\ 1/\sqrt{2} & -1/\sqrt{2} \end{bmatrix}$

$A = \begin{bmatrix} 1/\sqrt{2} & 1/\sqrt{2} \\ 1/\sqrt{2} & -1/\sqrt{2} \end{bmatrix} \begin{bmatrix} 3 & 0 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} 1/\sqrt{2} & 1/\sqrt{2} \\ -1/\sqrt{2} & 1/\sqrt{2} \end{bmatrix}$

(Exs)  $A = \begin{bmatrix} 0 & 1 & 1 \\ \sqrt{2} & 2 & 0 \\ 0 & 1 & 1 \end{bmatrix}$   $A^T = \begin{bmatrix} 0 & \sqrt{2} & 0 \\ 1 & 2 & 1 \\ 1 & 0 & 1 \end{bmatrix}$

1)  $AA^T = \begin{bmatrix} 2 & 2 & 2 \\ 2 & 6 & 2 \\ 2 & 2 & 2 \end{bmatrix}$   $\det: -\lambda^3 + 10\lambda^2 - 16\lambda$   
 $= -\lambda(\lambda-8)(\lambda-2) = 0$   
 3 rows of  $\Sigma \rightarrow \lambda_1 = 8 \quad \lambda_2 = 2 \quad \lambda_3 = 0$   
 $\sigma_1 = 2\sqrt{2} \quad \sigma_2 = \sqrt{2}$

2)  $A^T A = \begin{bmatrix} 2 & 2\sqrt{2} & 0 \\ 2\sqrt{2} & 6 & 2 \\ 0 & 2 & 2 \end{bmatrix} \rightarrow \lambda = 8, 2, 0$   
 $\Sigma = \begin{bmatrix} 2\sqrt{2} & 0 & 0 \\ 0 & \sqrt{2} & 0 \\ 0 & 0 & 0 \end{bmatrix}$

3) e vects of  $A^T A$ .  $\lambda_1 = 8$   $\begin{bmatrix} -6 & 2\sqrt{2} & 0 & | & 0 \\ 2\sqrt{2} & -2 & 2 & | & 0 \\ 0 & 2 & -6 & | & 0 \end{bmatrix} \rightarrow \begin{matrix} x = \frac{\sqrt{2}}{3}y \\ z = \frac{1}{3}y \end{matrix} \begin{bmatrix} \sqrt{2} \\ 3 \\ 1 \end{bmatrix}$   
 $\rightarrow v_1 = \frac{1}{\sqrt{2}} \begin{bmatrix} \sqrt{2} \\ 3 \\ 1 \end{bmatrix} = \begin{bmatrix} 1/\sqrt{2} \\ 3/\sqrt{2} \\ 1/\sqrt{2} \end{bmatrix}$

$\lambda_2 = 2$ :  $\begin{bmatrix} 0 & 2\sqrt{2} & 0 \\ 2\sqrt{2} & 4 & 2 \\ 0 & 2 & 0 \end{bmatrix} \rightarrow y=0 \rightarrow x = -\frac{1}{\sqrt{2}}z$   $\begin{bmatrix} 1 \\ 0 \\ -\sqrt{2} \end{bmatrix} \rightarrow v_2 = \begin{bmatrix} 1/\sqrt{2} \\ 0 \\ -\sqrt{2}/\sqrt{2} \end{bmatrix}$

$\lambda_3 = 0$ :  $\begin{bmatrix} 2 & 2\sqrt{2} & 0 \\ 2\sqrt{2} & 4 & 2 \\ 0 & 2 & 2 \end{bmatrix} \rightarrow \begin{matrix} x = -\sqrt{2}y \\ z = -y \end{matrix}$   $\begin{bmatrix} \sqrt{2} \\ -1 \\ 1 \end{bmatrix} \rightarrow v_3 = \begin{bmatrix} 1/\sqrt{2} \\ 1/2 \\ 1/2 \end{bmatrix}$

4)  $V^T = \begin{bmatrix} 1/\sqrt{2} & 3/\sqrt{2} & 1/\sqrt{2} \\ 1/\sqrt{2} & 0 & -\sqrt{2}/\sqrt{2} \\ 1/\sqrt{2} & -1/2 & 1/2 \end{bmatrix}$

5.  $u_1 = \frac{1}{\sqrt{8}} \begin{bmatrix} 0 & 1 & 1 \\ \sqrt{2} & 2 & 0 \\ 0 & 1 & 1 \end{bmatrix} \begin{bmatrix} 1/\sqrt{2} \\ 3/\sqrt{2} \\ 1/\sqrt{2} \end{bmatrix} = \begin{bmatrix} 1/\sqrt{6} \\ 2/\sqrt{6} \\ 1/\sqrt{6} \end{bmatrix}$

$u_2 = \dots \begin{bmatrix} -1/\sqrt{3} \\ 1/\sqrt{3} \\ -1/\sqrt{3} \end{bmatrix}$

$u_3 = \dots$  need to get

6.) manually  $\begin{bmatrix} 2 & 0 & 2 & 2 & | & 0 \\ 2 & 6 & 0 & 2 & | & 0 \\ 2 & 2 & 2 & 0 & | & 0 \end{bmatrix} \rightarrow \begin{bmatrix} -1 & 0 & -1 \\ 0 & 1 & 0 \\ 0 & 0 & 0 \end{bmatrix} \rightarrow y = -x, x = -z$   $\begin{bmatrix} 1/\sqrt{2} \\ 0 \\ -1/\sqrt{2} \end{bmatrix}$

$\begin{bmatrix} 1/\sqrt{6} & -1/\sqrt{3} & 1/\sqrt{2} \\ 2/\sqrt{6} & 1/\sqrt{3} & 0 \\ 1/\sqrt{6} & -1/\sqrt{3} & -1/\sqrt{2} \end{bmatrix} \begin{bmatrix} 2\sqrt{2} & 0 & 0 \\ 0 & \sqrt{2} & 0 \\ 0 & 0 & 0 \end{bmatrix} \begin{bmatrix} 1/\sqrt{2} & 3/\sqrt{2} & 1/\sqrt{2} \\ 1/\sqrt{2} & 0 & -\sqrt{2}/\sqrt{2} \\ 1/\sqrt{2} & -1/2 & 1/2 \end{bmatrix}$