

# Laplace Transform of an IVP

Given

$$ay'' + by' + cy = f(t) \quad y(0) = y_0, y'(0) = v_0$$

we want to find  $Y(s)$ .

$$\begin{aligned} \mathcal{L}\{ay'' + by' + cy\} &= \mathcal{L}\{f(t)\} \\ a\mathcal{L}\{y''\} + b\mathcal{L}\{y'\} + cY(s) &= F(s) \end{aligned}$$

In order to isolate  $Y(s)$ , we need to be able to compute  $\mathcal{L}\{y'\}$  and  $\mathcal{L}\{y''\}$ .

# Laplace Transform of Derivatives

Suppose  $\mathcal{L}\{y(t)\} = Y(s)$ . What is  $\mathcal{L}\{y'(t)\}$ ?

$$\begin{aligned}\mathcal{L}\{y'(t)\} &= \int_0^{\infty} \underbrace{e^{-st}}_u \underbrace{y'(t)dt}_{dv} \\ &= e^{-st}y(t)\Big|_0^{\infty} - (-s) \underbrace{\int_0^{\infty} e^{-st}y(t)dt}_{\mathcal{L}\{y(t)\}}\end{aligned}$$

$$\stackrel{s \geq 0}{=} -y(0) + s\mathcal{L}\{y(t)\}$$

$$= \boxed{sY(s) - y_0}$$

$$\begin{aligned}v &= y(t) \\ du &= -se^{-st}dt\end{aligned}$$

# Laplace Transform of Derivatives

Given that  $\mathcal{L}\{y'(t)\} = sY(s) - y_0$ . What is  $\mathcal{L}\{y''(t)\}$ ?

$$\mathcal{L}\{y''(t)\} = s\mathcal{L}\{y'(t)\} - y'(0) = s[sY(s) - y_0] - \underbrace{y'(0)}_{v_0}$$

$$= \boxed{s^2 Y(s) - sy_0 - v_0}$$

# Laplace Transform of an IVP

Given

$$ay'' + by' + cy = f(t) \quad y(0) = y_0, y'(0) = v_0$$

we want to find  $Y(s)$ .

$$a\mathcal{L}\{y''\} + b\mathcal{L}\{y'\} + cY(s) = F(s)$$

$$a(s^2 Y(s) - sy_0 - v_0) + b(sY(s) - y_0) + cY(s) = F(s)$$

$$\underbrace{(as^2 + bs + c)}_{\text{char. poly.}} Y(s) = F(s) + (as + b)y_0 + av_0$$

$$Y(s) = \frac{F(s) + (as + b)y_0 + av_0}{as^2 + bs + c}$$

ex: Use Laplace Transforms to solve  $y' + 6y = 3$  with  $y(0) = 2$ .

$$(\mathcal{L}\{y'\} + 6\mathcal{L}\{y\} = \mathcal{L}\{3\}$$

$$\mathcal{L}\{y'\} + 6\mathcal{L}\{y\} = \frac{3}{s}$$

$$sY(s) - \underbrace{y(0)}_2 + 6Y(s) = \frac{3}{s}$$

$$(s + 6)Y(s) = \frac{3}{s} + 2$$

$$Y(s) = \frac{3}{s(s+6)} + \frac{2}{s+6}$$

This is the example from pg. 8 of lecture 14.

$$y(t) = \frac{1}{2} + \frac{3}{2}e^{-6t}$$

ex: Solve  $y'' + 6y' + 25y = 0$  with  $y(0) = 0$   
 $y'(0) = 4$  using Laplace Trans.

$$s^2 Y(s) - s\cancel{y(0)} - \cancel{y'(0)} + 6(sY(s) - \cancel{y(0)}) + 25Y(s) = 0$$

$$(s^2 + 6s + 25)Y(s) = \underbrace{y'(0)}_4$$

$$Y(s) = \frac{4}{s^2 + 6s + 25}$$

This is the example from pg. 5 of lecture 14.

$$y(t) = e^{-3t} \sin(4t)$$

ex: Solve  $y'' + 6y' + 25y = -8e^{-4t}$  with  $y(0) = 0$   
 $y'(0) = 1$  using Lap. Trans.

$$\begin{aligned} s^2 Y(s) - \cancel{sy(0)} - y'(0) + 6(sY(s) - \cancel{y(0)}) + 25Y(s) &= \frac{-8}{s+4} \\ \underbrace{(s^2 + 6s + 25)}_{(s+3)^2 + 16} Y(s) &= \frac{-8}{s+4} + 1 = \frac{s-4}{s+4} \\ Y(s) &= \frac{s-4}{((s+3)^2 + 16)(s+4)} \end{aligned}$$

This is the example from pg. 12 of lecture 14.

$$y(t) = e^{-3t} \left( \frac{8}{17} \cos(4t) + \frac{33}{68} \sin(4t) \right) + \frac{8}{17} e^{-4t}$$

ex: Solve  $y'' - 3y' - 4y = 8e^{4t}$  with  $y(0) = 0$   
 $y'(0) = 1$  using Laplace Trans.

$$s^2 Y(s) - \cancel{sy(0)} - y'(0) - 3(sY(s) - \cancel{y(0)}) - 4Y(s) = \frac{-8}{s+4}$$

$$\underbrace{(s^2 - 3s - 4)}_{(s-4)(s+1)} Y(s) = \frac{8}{s-4} + 1 = \frac{s+4}{s-4}$$

$$Y(s) = \frac{s+4}{(s-4)^2(s+1)}$$

This is the example from pg. 10 of lecture 14.

$$y(t) = \underbrace{\frac{40}{25}te^{4t}}_{y_p} - \underbrace{\frac{3}{25}e^{4t} + \frac{3}{25}e^{-t}}_{y_h}$$

This is a case with mathematical resonance, where we multiply our guess for  $y_p$  by  $t$ .



# Multiplication by $t \Leftrightarrow$ Differentiation in $s$ -domain

$$\mathcal{L}\{t^k f(t)\} = \int_0^\infty e^{-st} t^k f(t) dt = \int_0^\infty \underbrace{e^{-st} t}_{\frac{d}{ds} e^{-st}} t^{k-1} f(t) dt$$

$$= -\frac{d}{ds} \int_0^\infty e^{-st} t^{k-1} f(t) dt$$

repeat same thing

$$\dots = (-1)^k \frac{d^k}{ds^k} F(s)$$

k-1 more times

with  $k=1$

$$\mathcal{L}\{t f(t)\} = -\frac{d}{ds} F(s)$$

$$\text{ex : } \mathcal{L}\{t \sin(\omega t)\} = \frac{2\omega s}{(s^2 + \omega^2)^2}$$

$$\mathcal{L}\{t \cos(\omega t)\} = \frac{s^2 - \omega^2}{(s^2 + \omega^2)^2}$$

ex: Solve  $y'' + 4y = 4 \cos(2t)$  with  $y(0) = 0$   
 $y'(0) = 0$  using Laplace Transforms

$$s^2 Y(s) + 4Y(s) = \frac{4s}{s^2 + 4}$$

$$Y(s) = \frac{4s}{(s^2 + 4)^2} = \underbrace{\frac{2\omega s}{(s^2 + \omega^2)^2}}_{\mathcal{L}\{t \sin(\omega t)\}} \quad \text{with } \omega = 2$$

$$y(t) = t \sin(2t)$$

ex: Find the Laplace Transform of  $f(t) = t^2 e^{3t}$

$$\begin{aligned} F(s) &= (-1)^2 \frac{d^2}{ds^2} \mathcal{L}\{e^{3t}\} \\ &= \frac{d^2}{ds^2} \frac{1}{s-3} \\ &= \frac{d}{ds} \frac{-1}{(s-3)^2} \\ &= \frac{2}{(s-3)^3} \end{aligned}$$

ex: Solve  $y'' - 6y' + 9y = e^{3t}$  with  $y(0) = 0$   
 $y'(0) = 0$  using Laplace Transforms

$$\underbrace{(s^2 - 6s + 9)}_{(s-3)^2} Y(s) = \frac{1}{s-3}$$

$$Y(s) = \frac{1}{(s-3)^2}$$

$$y(t) = \frac{1}{2} t^2 e^{3t}$$