

Recall: Solutions flows and fixed points (critical points)

The origin $\vec{0}$ is the unique fixed point for the linear system $\frac{d}{dt}\vec{x} = \mathbf{A}(t)\vec{x}$.

How about the (constant) non-homogeneous linear system $\frac{d}{dt}\vec{x} = \mathbf{A}\vec{x} + \vec{f}$?

$$\vec{0} = \mathbf{A}\vec{x}^* + \vec{f} \quad \Rightarrow \quad \vec{x}^* = -\mathbf{A}^{-1}\vec{f} \quad \text{is the fixed point or critical point}$$

A constant inhomogeneity just shifts everything from the origin.

The global behaviour of solutions is fully determined by the eigenvalues and eigenvectors of \mathbf{A} :

1. Eigendirections with real eigenvalues and:

- $\lambda < 0$ attract solutions towards the fixed point.
- $\lambda > 0$ repel solutions away from the fixed point.

2. Complex conjugate eigenvalues produce oscillations.

2-Dimensional Autonomous Nonlinear Systems

Consider the autonomous nonlinear system

$$\frac{d}{dt}x = f(x, y), \quad \frac{d}{dt}y = g(x, y)$$

where f and g are nonlinear function with no explicit time-dependence.


This can be written as

$$\frac{d}{dt} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} f(x, y) \\ g(x, y) \end{bmatrix} \quad \text{or} \quad \frac{d}{dt} \vec{x} = \vec{f}(\vec{x})$$

We can plot the vector field $\begin{bmatrix} x' \\ y' \end{bmatrix}$ over the (x, y) -plane.

This vector field gives us qualitative information about solution flows.

ex: The vector field for the the system $x' = y$ and $y' = -x + x^2$



This system has two critical points:

$$\begin{array}{ll} x = y = 0 & x = 1, \quad y = 0 \\ \text{(center)} & \text{(saddle)} \end{array}$$

Linearization & Local Solution Flow

To classify critical points of a nonlinear system, we linearize it around its critical points \vec{x}^* (e.g., for a 2D system $\vec{x}^* = \begin{bmatrix} x^* \\ y^* \end{bmatrix}$).

Any solution that starts at \vec{x}^* stays there forever.

So we perturb the fixed point by setting $\vec{x}(t) = \vec{x}^* + \vec{u}(t)$ where $\vec{u}(0)$ is infinitesimally small.

Thus we are interested in

$$\frac{d}{dt}\vec{x} = \vec{f}(\vec{x}), \quad \text{with } \vec{x}(0) = \vec{x}^* + \vec{u}(0) \quad \text{where} \quad \vec{u}(0) \approx \vec{0}$$

expanding $\vec{x}(t)$ we get

$$= \vec{f}(\vec{x}^* + \vec{u})$$

Linearization & Local Solution Flow

To classify critical points of a nonlinear system, we linearize it around its critical points \vec{x}^* (e.g., for a 2D system $\vec{x}^* = \begin{bmatrix} x^* \\ y^* \end{bmatrix}$).

$$\frac{d}{dt}\vec{u} = \vec{f}(\vec{x}^* + \vec{u}) \quad \text{where} \quad \vec{u}(0) \approx \vec{0}$$

make a first-order Taylor expansion in \vec{u} around $\vec{u} = \vec{0}$

$$\frac{d}{dt}\vec{x} \approx \vec{f}(\vec{x}^*) + \frac{\partial \vec{f}(\vec{x})}{\partial \vec{x}} \bigg|_{\vec{x}=\vec{x}^*} \vec{u} + \dots + h.o.t.$$

where is $\frac{\partial \vec{f}}{\partial \vec{x}}$ called the Jacobian matrix. For our 2D system, it is given by

$$\mathbf{J} = \frac{\partial \vec{f}}{\partial \vec{x}} = \begin{bmatrix} \frac{\partial}{\partial x} f(x, y) & \frac{\partial}{\partial y} f(x, y) \\ \frac{\partial}{\partial x} g(x, y) & \frac{\partial}{\partial y} g(x, y) \end{bmatrix} = \begin{bmatrix} f_x & f_y \\ g_x & g_y \end{bmatrix}$$

Linearization & Local Solution Flow

To classify critical points of a nonlinear system, we linearize it around its critical points \vec{x}^* (e.g., for a 2D system $\vec{x}^* = \begin{bmatrix} x^* \\ y^* \end{bmatrix}$).

After the substitution $\vec{u} = \vec{x} - \vec{x}^*$, near the critical point \vec{x}^*

$$\frac{d}{dt} \begin{bmatrix} x \\ y \end{bmatrix} \approx \mathbf{J}^* \left(\begin{bmatrix} x \\ y \end{bmatrix} - \begin{bmatrix} x^* \\ y^* \end{bmatrix} \right) \quad \text{where } \mathbf{J}^* = \begin{bmatrix} f_x(x^*, y^*) & f_y(x^*, y^*) \\ g_x(x^*, y^*) & g_y(x^*, y^*) \end{bmatrix}$$

$$\frac{d}{dt} \vec{x} \approx \mathbf{J}^* \vec{x} - \underbrace{\mathbf{J}^* \vec{x}^*}_{\text{constant}} = \text{a (constant) non-homogeneous linear system}$$

This linear approximation holds locally.

Thus the local behaviour of solutions near critical points is determined by the eigenvalues and eigenvectors of \mathbf{J} .

ex: Find the Jaobian matrix for the system

$$x' = f(x, y), \quad y' = g(x, y) \quad \text{with} \quad f(x, y) = y, \quad g(x, y) = -x + x^2$$

$$\mathbf{J} = \begin{bmatrix} f_x & f_y \\ g_x & g_y \end{bmatrix} = \begin{bmatrix} 0 & 1 \\ 2x - 1 & 0 \end{bmatrix}$$

ex: Find a linear approximation to the above system at the two following critical points

$$\underline{x = y = 0}$$

$$\underline{x = 1, \quad y = 0}$$

$$\frac{d}{dt} \begin{bmatrix} x \\ y \end{bmatrix} \approx \begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix}$$

$$\frac{d}{dt} \begin{bmatrix} x \\ y \end{bmatrix} \approx \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} \begin{bmatrix} x - 1 \\ y \end{bmatrix}$$

Approximate Local Solution Flows

$$\underline{x = y = 0}$$

$$\underline{x = 1, \quad y = 0}$$

$$\frac{d}{dt} \begin{bmatrix} x \\ y \end{bmatrix} \approx \begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix}$$

$$\frac{d}{dt} \begin{bmatrix} x \\ y \end{bmatrix} \approx \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} \begin{bmatrix} x - 1 \\ y \end{bmatrix}$$

Eigenvalues: $\pm i$ (center)

Eigenvalues: ± 1 (saddle)

