

Differential Equations

1. What are they and why do we solve them?
2. Terminology
3. Graphical intuition and the direction field

1. What are they and why do we solve them?
2. Terminology
3. Graphical intuition and the direction field

What is a Differential Equation?

A differential equation (DE) is an equation involving an unknown function y and atleast one derivative of y w.r.t. an independent variable.

Given: A DE with an unknown function $y(t)$. e.x., $\frac{dy}{dt} = -3y(t)$ or

$$y' = -3y$$

Task: Find the function(s) $y(t)$.

Solution: $y(t) = C_1 e^{-3t}$

- Tools:
- Calculus (i.e., integration/differentiation)
 - Guess and check (does some function $f(t)$ make LHS=RHS?)
 - Specialized procedures (informed by experience geussing)
 - Geometry/Linear Algebra (useful for systems of DEs)

A differential equation (DE) is an equation involving an unknown function y and atleast one derivative of y w.r.t. an independent variable.

Given: A DE with an unknown function $y(t)$. e.x., $\frac{dy}{dt} = -3y(t)$ or


$$y' = -3y$$

Task: Find the function(s) $y(t)$.

Solution: $y(t) = C_1 e^{-3t}$

- Tools:
- Calculus (i.e., integration/differentiation)
 - Guess and check (does some function $f(t)$ make LHS=RHS?)
 - Specialized procedures (informed by experience geussing)
 - Geometry/Linear Algebra (useful for systems of DEs)

Example: Skydiving




Newton's Second Law:

$$F = ma$$

$$ma = \underbrace{-mg}_{\text{gravitational force}} \quad \underbrace{-\mu v}_{\text{drag force}}$$

$$a = v'$$

$$\boxed{mv' = -mg - \mu v} \quad \text{DE for } v(t)$$



Newton's Second Law:

$$F = ma$$

$$ma = \underbrace{-mg}_{\text{gravitational force}} \quad \underbrace{-\mu v}_{\text{drag force}}$$

$$a = v'$$

$$\boxed{mv' = -mg - \mu v} \quad \text{DE for } v(t)$$

Example: Epidemiology

Kermack & McKendrick's SIR model

- Susceptible \rightarrow Infectious \rightarrow Recovered

System of 3 ordinary differential equations:

$$\begin{aligned}\frac{dS}{dt} &= \mu(I + R) - \beta SI \\ \frac{dI}{dt} &= \beta SI - \gamma I - \mu I \\ \frac{dR}{dt} &= \gamma I - \mu R\end{aligned}$$

t = time

β = infection rate

γ = recovery rate

μ = birth/death rate

Reproduction number: $R_0 = \frac{\beta}{\mu + \gamma}$

1. $R_0 > 1$: endemic equilibrium

2. $R_0 < 1$: disease dies out - basic idea behind "flatten the curve"

Kermack & McKendrick's SIR model

- Susceptible \rightarrow Infectious \rightarrow Recovered

System of 3 ordinary differential equations:

$$\frac{dS}{dt} = \mu(I + R) - \beta SI$$

$$\frac{dI}{dt} = \beta SI - \gamma I - \mu I$$

$$\frac{dR}{dt} = \gamma I - \mu R$$

t = time

β = infection rate

γ = recovery rate

μ = birth/death rate

Reproduction number: $R_0 = \frac{\beta}{\mu + \gamma}$

1. $R_0 > 1$: endemic equilibrium

2. $R_0 < 1$: disease dies out - basic idea behind "flatten the curve"

Terminology: ODEs vs PDEs

- Ordinary differential equation (ODE) (covered in this course)
 - A DE with derivatives w.r.t. only one independent variable.
 - $\frac{dy}{dt} = y(t) + 3$ or $\frac{dy}{dt} = \sin(y) + \cos(t)$
- Partial differential equation (PDE) (not covered in this course)
 - A DE with derivatives w.r.t multiple independent variable.
 - Heat/Diffusion eq: $\frac{\partial u}{\partial t} = D \frac{\partial^2 u}{\partial x^2}$
 - Wave eq: $\frac{\partial^2 u}{\partial t^2} = c \frac{\partial^2 u}{\partial x^2}$
 - Partial derivatives are necessary for solutions to agree when changing coordinate systems (e.g., switch from cartesian to polar coordinates)

- Ordinary differential equation (ODE) (covered in this course)
 - A DE with derivatives w.r.t. only one independent variable.
 - $\frac{dy}{dt} = y(t) + 3$ or $\frac{dy}{dt} = \sin(y) + \cos(t)$
- Partial differential equation (PDE) (not covered in this course)
 - A DE with derivatives w.r.t multiple independent variable.
 - Heat/Diffusion eq: $\frac{\partial u}{\partial t} = D \frac{\partial^2 u}{\partial x^2}$
 - Wave eq: $\frac{\partial^2 u}{\partial t^2} = c \frac{\partial^2 u}{\partial x^2}$
 - Partial derivatives are necessary for solutions to agree when changing coordinate systems (e.g., switch from cartesian to polar coordinates)

Terminology: Order of a DE

The highest derivative that appears in the DE.

- $y' = y + 3$ first order
- $y' = y^2 + 9$ first order
- $\left(\frac{dy}{dt}\right)^2 = \tan(t)$ first order
- $y'' = -y$ second order
- $\frac{d^4y}{dx^4} = ky$ fourth order

The highest derivative that appears in the DE.

- $y' = y + 3$ first order
- $y' = y^2 + 9$ first order
- $\left(\frac{dy}{dt}\right)^2 = \tan(t)$ first order
- $y'' = -y$ second order
- $\frac{d^4y}{dx^4} = ky$ fourth order

Terminology: Operator Form $\Rightarrow \mathcal{L}[y(t)] = f(t)$

Everything that depends on the unknown function goes on one side of the equal sign and everything else on the other.

- $\frac{dy}{dt} = y(t) + 3 \quad \rightarrow \quad \frac{dy}{dt} - y(t) = 3$
 - $\mathcal{L}[y] = f' - y, \quad f(t) = 3$
- $\frac{dy}{dt} = \sin(y) + \cos(t) \quad \rightarrow \quad \frac{dy}{dt} - \sin(y) = \cos(t)$
 - $\mathcal{L}[y] = f' - \sin(y), \quad f(t) = \cos(t)$

The operator $\mathcal{L}[\cdot]$, encodes the "intrinsic" dynamics that the ODE is modelling.

- Force-displacement relationship of a spring.
- Velocity-drag relationship of a viscous fluid.

Everything that depends on the unknown function goes on one side of the equal sign and everything else on the other.

- $\frac{dy}{dt} = y(t) + 3 \quad \rightarrow \quad \frac{dy}{dt} - y(t) = 3$
 - $\mathcal{L}[y] = f' - y, \quad f(t) = 3$
- $\frac{dy}{dt} = \sin(y) + \cos(t) \quad \rightarrow \quad \frac{dy}{dt} - \sin(y) = \cos(t)$
 - $\mathcal{L}[y] = f' - \sin(y), \quad f(t) = \cos(t)$

The operator $\mathcal{L}[\cdot]$, encodes the "intrinsic" dynamics that the ODE is modelling.

- Force-displacement relationship of a spring.
- Velocity-drag relationship of a viscous fluid.

Terminology: Linearity of DEs - $L[y(t)] = f(t)$

If the operator $L[\cdot]$ is linear, then the DE is linear.

Conditions for linearity:

Given any two functions f and g and a constant c , a linear operator satisfies

$$1. L[f + g] = L[f] + L[g]$$

$$2. L[cf] = cL[f]$$

In practice: does the operator have any nonlinear functions?

$$\underline{\text{ex:}} L[y] = y'' + y$$

Linear

$$\underline{\text{ex:}} L[y] = y' + \sin(y)$$

Nonlinear

If the operator $L[\cdot]$ is linear, then the DE is linear.

Conditions for linearity:

Given any two functions f and g and a constant c , a linear operator satisfies

$$1. L[f + g] = L[f] + L[g]$$

$$2. L[cf] = cL[f]$$

In practice: does the operator have any nonlinear functions?

$$\underline{\text{ex:}} L[y] = y'' + y$$

Linear

$$\underline{\text{ex:}} L[y] = y' + \sin(y)$$

Nonlinear

Terminology: Autonomous DEs - $L[y(t)] = f(t)$

If both $L[\cdot]$ and $f(t)$ do not explicitly depend on the independent variable, then the DE is autonomous.

- $y' = y \rightarrow y' - y = 0$ Autonomous
- $y' = y^2 + 3 \rightarrow \frac{dy}{dt} - y^2 = 3$ Autonomous
- $\frac{dy}{dt} = y + \tan(t) \rightarrow \frac{dy}{dt} - y = \tan(t)$ Non-autonomous
- $\frac{dy}{dt} = -3ty \rightarrow \frac{dy}{dt} + 3ty = 0$ Non-autonomous

$f(t)$ is often called the (external) forcing term.

constant or zero-forcing \Rightarrow Autonomous DE

If both $L[\cdot]$ and $f(t)$ do not explicitly depend on the independent variable, then the DE is autonomous.

- $y' = y \rightarrow y' - y = 0$ Autonomous
- $y' = y^2 + 3 \rightarrow \frac{dy}{dt} - y^2 = 3$ Autonomous
- $\frac{dy}{dt} = y + \tan(t) \rightarrow \frac{dy}{dt} - y = \tan(t)$ Non-autonomous
- $\frac{dy}{dt} = -3ty \rightarrow \frac{dy}{dt} + 3ty = 0$ Non-autonomous

$f(t)$ is often called the (external) forcing term.

constant or zero-forcing \Rightarrow Autonomous DE

Classifying ODEs

- $x'' + x^2 = t$
 - Order: 2
 - Linear: No
 - Autonomous: No

- $\frac{d^4x}{dt^4} = 0$
 - Order: 4
 - Linear: Yes
 - Autonomous: Yes

- $x'' + x^2 = t$
 - Order: 2
 - Linear: No
 - Autonomous: No

- $\frac{d^4x}{dt^4} = 0$
 - Order: 4
 - Linear: Yes
 - Autonomous: Yes

Terminology: Solution to an ODE

A solution of an ODE is a function that satisfies the ODE.

ex: Is $y = Ce^{-t} + t - 1$ a solution to $y' + y = t$?

$$\begin{aligned} y' &= -Ce^{-t} + 1 \\ y' + y &= \cancel{-Ce^{-t} + 1} + \cancel{Ce^{-t}} + t - \cancel{1} \\ &= t \quad \checkmark \end{aligned}$$

Here C is an arbitrary constant that can have any value.

Any solution with an arbitrary constant is called a general solution

We can obtain a unique solution by imposing some constraint, a solution with no arbitrary constants is called a particular solution

A solution of an ODE is a function that satisfies the ODE.

ex: Is $y = Ce^{-t} + t - 1$ a solution to $y' + y = t$?

$$\begin{aligned} y' &= -Ce^{-t} + 1 \\ y' + y &= \cancel{-Ce^{-t} + 1} + \cancel{Ce^{-t}} + t - \cancel{1} \\ &= t \quad \checkmark \end{aligned}$$

Here C is an arbitrary constant that can have any value.

Any solution with an arbitrary constant is called a general solution

We can obtain a unique solution by imposing some constraint, a solution with no arbitrary constants is called a particular solution

Initial Value Problems

ODEs of the form

$$L[y] = f(t), \text{ with } y(t_0) = y_0,$$

where t_0 and y_0 are numerical values (usually real-valued).

ex: Find the particular solution to $y' + y = t$ with $y(0) = 0$?

Start with the general solution

$$y(t) = Ce^{-t} + t - 1$$

evaluate at $t = t_0 = 0$, make that equal to $y_0 = 0$

$$y(0) = C - 1 = 0 \Rightarrow C = 1$$

$$\boxed{y(t) = e^{-t} + t - 1}$$

ODEs of the form

$$L[y] = f(t), \text{ with } y(t_0) = y_0,$$

where t_0 and y_0 are numerical values (usually real-valued).

ex: Find the particular solution to $y' + y = t$ with $y(0) = 0$?

Start with the general solution

$$y(t) = Ce^{-t} + t - 1$$

evaluate at $t = t_0 = 0$, make that equal to $y_0 = 0$

$$y(0) = C - 1 = 0 \Rightarrow C = 1$$

$$\boxed{y(t) = e^{-t} + t - 1}$$

Summary

1. What are DEs?

- Equations involving unknown function(s) and function derivatives.
- Specify rates of change of certain quantities.
- Useful for modelling many natural phenomena.

2. Terminology

- ODEs (& PDEs).
- Order of DEs, Linear DEs, Autonomous DEs, Solutions to DEs

3. Initial Value Problems

- The most "standard" way to obtain a unique solution
- Specify solution value at some initial time

1. What are DEs?

- Equations involving unknown function(s) and function derivatives.
- Specify rates of change of certain quantities.
- Useful for modelling many natural phenomena.

2. Terminology

- ODEs (& PDEs).
- Order of DEs, Linear DEs, Autonomous DEs, Solutions to DEs

3. Initial Value Problems

- The most "standard" way to obtain a unique solution
- Specify solution value at some initial time