

# Dataset for use in data poor methods

*Laurence Kell and Alex Tidd*

*25 December, 2018*

## MYDAS

Task 5 of the MYDAS project <https://github.com/laurieKell/mydas> requires a review of data available on length composition, mean size, catch and indices of abundance (commercial or survey).

Data are available from the MI or from collective sources such as DATRAS <http://www.ices.dk/marine-data/data-portals/Pages/DATRAS.aspx> for survey data and the STECF <https://stecf.jrc.ec.europa.eu/data-dissemination> for commercial data.

## Work plan

- Condition the operating model using life histories (from table above) from brill, turbot, skate and Pollack as per the Mydas project (see the quick start). Will have to use MI estimates for sprat. Use estimates of length (brill, sprat, skate). Obtain length frequencies for turbot and pollack from MI.

The life histories were reviewed by Hans, using data from the MI databases.

- With further investigation look at predicted abundance estimates for brill, sprat (although on high side) and skate (abundance was for all species of skates and rays, will have to make consistent with commercial species... maybe use thornback only). Pollack and turbot catches are greater than abundance so cannot use these.

If the absolute abundance estimates are wrong relative indices can be used, as normal practice.

- With such good time series of survey length data for brill, sprat and skate it would be good to simulate the mean size by using the mlz package <https://cran.r-project.org/web/packages/MLZ/vignettes/MLZ.html#introduction>. This would give an estimate of Z (assuming constant M) and thus look at changes in F.

MLZ uses mean size, it is unlikely to provide good estimates of Z for sprat due to high M and high recruitment variability, the mean size is therefore determined by recruitment not F.

- It may also be worth pursuing the LBSR for length-based composition for the above (and compare) with life-history parameters M/K ratio and Linf to estimate F/M and F/FMSY. With SPR being the biological reference point. <https://cran.r-project.org/web/packages/LBSPR/vignettes/LBSPR.html>

LBSPR is better at estimating F.

- For turbot and pollack it may be beneficial to use an empirical approach (as survey info pretty poor) based on the commercial CPUE (biomass index from MI lengths or estimated by weights) such as the ICES 2/3 rule (as per Simon Fischer).

The surveys could be used in the

For the observation model commercial CPUEs could be used.


Further: Need to discuss with MI what data we can get for Lobsters and Razors in terms of lengths. We have 1 set of priors from Iyves.

## Survey data extraction


Extract via sql the case study stocks from Mydas survey database and get numbers at length

For more in depth detail of the data see <https://data.marine.gov.scot/sites/default/files//SMFS%200816.pdf>


Calculate weighted mean length for case study species




**Figure 1** Plot of numbers at length for pollock with weighted mean length represented by dashed red line.




**Figure 2** Plot of numbers at length for turbot with weighted mean length represented by dashed red line.




**Figure 3** Plot of numbers at length for brill with weighted mean length represented by dashed red line.




**Figure 4** Plot of numbers at length for brill with weighted mean length represented by dashed red line.



**Figure 5** Plot of numbers at length for skate with weighted mean length represented by dashed red line.



**Figure 6** Plot of numbers at length for sprat with weighted mean length represented by dashed red line.




**Figure 7** Time series of weighted mean length.

Get all survey data from sql query


Estimate fish weight in kgs





**Figure 8** Catch per unit effort in kgs (CPUE) time series with red line depicting mean cpue.

Fish numbers




**Figure 9** Catch per unit effort in numbers (CPUE) time series with red line depicting mean cpue.

## Commercial data


Part of the issues with the main catch data in the stecf database is that the effort does not match up by gear and area with the effort estimations. Here the STECF Annual Economic Report database is used.

Extract the data via sql

Estimate catch in kgs per day



**Figure 10** Time series of commercial cpue.



**Figure 11** Time series of total catch in tonnes.


## Estimate abundance from survey data

See <http://www.fao.org/docrep/w5449e/w5449e0f.htm> section 13.7

Get survey stations

Calculate total area/total number of stations the survey covered by division and merge with catch tonnes perkm2

Biomass = totalsurveyarea x 1/total number of stations within the survey area x sum of catch



## Life history parameters from the literature

Table 1: life history paramters

id	species	speciesgp	t.0	linf	k	lmat	tm	sex	area	source
1	Amblyraja radiata	SKA		66.00	0.2330	46.0	4.0	U	North Sea	[12]
7	Leucoraja circularis	SKA		98.80	0.1210	51.2	5.3	U	North Sea	[12]
8	Leucoraja fullonica	SKA		98.80	0.1210	51.2	5.3	U	North Sea	[12]
17	Pollachius pollachius	POL		85.60	0.1900	NA	NA	U	North Sea	[18]
18	Pollachius pollachius	POL		85.60	0.1860	44.8	3.7	U	North Sea	[12]
23	Psetta maxima	TUR	-0.05	64.80	0.2600	46.0	NA	F	North Sea	[15]
24	Psetta maxima	TUR	-0.51	49.20	0.3700	NA	NA	M	North Sea	[15]

id	species	speciesgp	t.0	linf	k	lmat	tm	sex	area	source
25	Psetta maxima	TUR		65.20	0.3240	49.0	NA	M	Bay of Biscay	[9], [7]
26	Psetta maxima	TUR		73.60	0.2770	49.0	NA	F	Bay of Biscay	[9], [7]
27	Psetta maxima	TUR	-1.79	69.60	0.2497	49.0	4.5	U	North Sea	[19], [8], [16], [17]
28	Psetta maxima	TUR		57.00	0.3200	46.0	4.5	U	North Sea	[12]
29	Psetta maxima	TUR		52.50	0.3200	NA	NA	M	North Sea	[17]
30	Psetta maxima	TUR	-1.79	70.00	0.1480	NA	NA	F	North Sea	[17]
31	Raja batis	SKA	-1.63	254.00	0.0600	130.0	11.0	U	North Sea	[11]
32	Raja batis	SKA		253.70	0.0570	155.0	15.0	U	North Sea	[12]
33	Raja brachyura	SKA		139.00	0.1200	100.0	9.3	U	North Sea	[12]
34	Raja brachyura	SKA	-0.8	118.00	0.1900	NA	NA	M	Irish Sea	[13]
35	Raja brachyura	SKA	-1.52	139.00	0.1200	NA	NA	F	Irish Sea	[13]
36	Raja clavata	SKA	-1.32	128.00	0.0900	NA	NA	F	Irish Sea	[13]
37	Raja clavata	SKA	-0.6	85.60	0.2100	NA	NA	M	Irish Sea	[13]
38	Raja clavata	SKA		105.00	0.2200	65.0	5.0	U	North Sea	[12]
39	Raja montagui	SKA	-0.37	72.80	0.1800	NA	NA	F	Irish Sea	[13]
40	Raja montagui	SKA	-0.56	68.70	0.1900	NA	NA	M	Irish Sea	[13]
41	Raja montagui	SKA		97.80	0.1480	67.0	6.0	U	North Sea	[12]
42	Raja naevus	SKA	-0.465	92.00	0.1100	59.0	9.0	U	North Sea	[11]
43	Raja naevus	SKA		91.60	0.1090	59.0	9.0	U	North Sea	[12]
44	Raja radiata	SKA		66.00	0.2300	46.0	4.0	U	North Sea	[22]
45	Raja undulata	SKA		112.00	0.1000	57.5	6.3	U	North Sea	[12]
46	Scophthalmus rhombus	BLL		74.88	0.1400	37.0	NA	M	Bay of Biscay	[9], [7]
47	Scophthalmus rhombus	BLL		85.23	0.1470	37.0	NA	F	Bay of Biscay	[9], [7]
48	Scophthalmus rhombus	BLL		50.00	0.2700	37.0	4.5	U	North Sea	[12]
49	Sprattus sprattus	SPR	*	16.00	0.6500	13.0	NA	U	North Sea	[20]

## Conclusion from the above analysis for 5 of the 7 case studies

- Pollack and turbot have poor coverage in terms of numbers at length from the survey data and hence potentially poor abundance estimates.
- Brill, sprat and skates have long time series of numbers at length. Brill has adequate numbers from 1993 to 2016 as has skate, while sprat contains a longer time series from 1985-to 2016.
- Observations from the commercial time-series show that for all species there are 9 years of data.
- There are enough life history parameters from the literature to condition an operating model for all case-study species apart from sprat.

## Software Versions

- R version 3.5.1 (2018-07-02)
- plyr: 1.8.4
- dplyr: 0.7.8
- ggplot2: 3.1.0
- DBI: 1.0.0
- RPostgreSQL: 0.6.2
- reshape 0.8.8
- **Compiled:** Tue Dec 25 20:37:56 2018

## Author information

Alex Tidd. emperorfish@gmail.com

## Acknowledgements

This vignette and many of the methods documented in it were developed under the MyDas project funded by the Irish exchequer and EMFF 2014-2020. The overall aim of MyDas is to develop and test a range of assessment models and methods to establish Maximum Sustainable Yield (MSY) reference points (or proxy MSY reference points) across the spectrum of data-limited stocks.

## References

7. Deniel, C. (1981). Les poissons plats (TelCostCens, Pleuronecti- formes) en baie de Douarnenez. Reproduction croissance et migration des Bothidae, Scophthalmidae, Pleuronectidae et Soleidae. These, Universite Bretagne Occidentale, Brest. 476 PP.
8. Deniel, C. (1990). Comparative study of growth of flatfishes on the west coast of Brittany. *Journal of Fish Biology*, 37: 149–166.
9. Dorel, D. (1986). Poissons de l’Atlantique Nord-Est Relations tailles-poids.
10. Greenstreet, S.P.R., Rossberg, A.G., Fox, C.J., Le Quesne, W.J.F., Blasdale, T., Boulcott, P., Mitchell, I., Millar, C., Moffat, C.F. (2012). Demersal fish biodiversity: species- level indicators and trends-based targets for the Marine Strategy Framework Directive. *ICES J. Mar. Sci.* 69, 1789–1801.
11. Holden, M. J. (1972). The growth rates of *R. brachyura*, *R. clavata* and *R. montagui* as determined by tagging data. *Journal du Conseil International pour l’Exploration de la Mer* 34:161–168.
12. Jones, A. (1974). Sexual maturity, fecundity and growth of the turbot, *Scophthalmus maximus* L. *Journal of the Marine Biological Association of the UK*, 54: 109–125.
13. Knijn, R.J., Boon, T.W., Heessen, H.J.L., Hislop, J.F.G. (1993). “Atlas of North Sea Fishes” (ICES Coop. Res. Rep. 194, ICES, Copenhagen, 1993).
14. Mengi, T. (1963). *Ber. Deut. Wiss. Komm.* 7, 119.
15. Moreau, J. (1964). Contribution a’ l’e ´tude du lieu jaune (*Gadus pollachius* L.). *Revue de Travaux Institut de Pe ´ches Maritimes*, 28:237–255.
16. Muus, B.J., Nielson, J. G. (1999). *Sea Fish (Scandinavian Fishing Year Book, Hedeusene, Denmark.*
17. Pauly, D. (1978). A preliminary compilation of fish length growth parameters. *Berichte des Institut fu ¨r Meereskunde an der Universita ¨t Kiel*, 55. 200 pp.

18. Vinther, M. (1989). Some notes on the biology of the starry ray, *Raja radiata*, in the North Sea. Working document: ICES Study Group on Elasmobranch Fisheries, 4/1989, 1–20. mimeo.



## Session Info

R version 3.5.1 (2018-07-02)  
Platform: x86\_64-pc-linux-gnu (64-bit)  
Running under: Ubuntu 16.04.2 LTS

Matrix products: default  
BLAS: /usr/lib/libblas/libblas.so.3.6.0  
LAPACK: /usr/lib/lapack/liblapack.so.3.6.0

locale:  
[1] LC\_CTYPE=en\_US.UTF-8 LC\_NUMERIC=C  
[3] LC\_TIME=en\_GB.UTF-8 LC\_COLLATE=en\_US.UTF-8  
[5] LC\_MONETARY=en\_GB.UTF-8 LC\_MESSAGES=en\_US.UTF-8  
[7] LC\_PAPER=en\_GB.UTF-8 LC\_NAME=C  
[9] LC\_ADDRESS=C LC\_TELEPHONE=C  
[11] LC\_MEASUREMENT=en\_GB.UTF-8 LC\_IDENTIFICATION=C

attached base packages:  
[1] splines stats graphics grDevices utils datasets methods  
[8] base

other attached packages:  
[1] gam\_1.16 foreach\_1.4.4 bindrcpp\_0.2.2  
[4] reshape\_0.8.8 RPostgreSQL\_0.6-2 DBI\_1.0.0  
[7] ggplot2\_3.1.0 dplyr\_0.7.8 plyr\_1.8.4  
[10] knitr\_1.20

loaded via a namespace (and not attached):  
[1] Rcpp\_1.0.0 highr\_0.7 pillar\_1.3.1 compiler\_3.5.1  
[5] bindr\_0.1.1 iterators\_1.0.10 tools\_3.5.1 digest\_0.6.18  
[9] evaluate\_0.12 tibble\_1.4.2 gtable\_0.2.0 pkgconfig\_2.0.2  
[13] rlang\_0.3.0.1 rstudioapi\_0.8 yaml\_2.2.0 withr\_2.1.2  
[17] stringr\_1.3.1 rprojroot\_1.3-2 grid\_3.5.1 tidyselect\_0.2.5  
[21] glue\_1.3.0 R6\_2.3.0 rmarkdown\_1.10 reshape2\_1.4.3  
[25] purrr\_0.2.5 magrittr\_1.5 backports\_1.1.2 scales\_1.0.0  
[29] codetools\_0.2-15 htmltools\_0.3.6 assertthat\_0.2.0 colorspace\_1.3-2  
[33] labeling\_0.3 stringi\_1.2.4 lazyeval\_0.2.1 munsell\_0.5.0  
[37] crayon\_1.3.4