



**Predicting recruitment density dependence and intrinsic growth rate for all fishes worldwide using a data-integrated life-history model**

Journal:	<i>Fish and Fisheries</i>
Manuscript ID	Draft
Wiley - Manuscript type:	Original Article
Date Submitted by the Author:	n/a
Complete List of Authors:	Thorson, James; Alaska Fisheries Science Center, Habitat and Ecosystem Process Research Program
Key terms:	steepness, recruitment, meta-analysis, life-history theory, taxonomy, stock assessment
Abstract:	<p>Fisheries managers and scientists use biological models to determine sustainable fishing rates, understand environmental responses, and forecast future dynamics. These models typically require both life-history parameters (mortality, maturity, growth, and size) and stock-recruit parameters (production of juveniles). However, there has been surprisingly little research to simultaneously predict life-history and stock-recruit parameters. I therefore develop the first "data-integrated life-history model," which modifies the recent FishLife model to fit a simple model of evolutionary dynamics to field-measurements of life-history parameters as well as historical estimates of spawning output and subsequent recruitment. This evolutionary model predicts recruitment productivity ("steepness") and variability (variance and autocorrelation in recruitment deviations) as well as mortality, maturity, growth, and size, and uses these to predict intrinsic growth rate (<math>r</math>) for all described fishes. The model confirms previous analysis showing little correlation between recruitment productivity ("steepness") and either natural mortality or asymptotic maximum size (<math>W_{\infty}</math>). However, it does reveal strong taxonomic patterns, where family Sebastidae has lower steepness (mean=0.63, SD=0.18) and Lutjanidae has elevated steepness (mean=0.77, SD=0.16) relative to the prediction for bony fishes (class Actinopterygii, mean=0.69, SD=0.22). Similarly, genus <i>Sebastes</i> has growth rate <math>r</math> (0.07) similar to several shark families (Lamniformes: 0.07; Carcharhiniformes: 0.10). Predictions are tighter for species than for higher-level organizations (e.g., families), and predictions (including multivariate confidence intervals) are available for &gt;32,000 described fishes worldwide in R package FishLife. I conclude by outlining how multivariate predictions of life-history and stock-recruit parameters could be useful for stock-assessment, decision theory, ensemble modelling, and strategic management.</p>



**Predicting recruitment density dependence and intrinsic growth rate for all  
fishes worldwide using a data-integrated life-history model**

Running title: Predicting productivity for all fishes

James Thorson

Habitat and Ecosystem Process Research program, Alaska Fisheries Science Center, National  
Marine Fisheries Service, NOAA, Seattle, WA, USA

Previously at: Fisheries Resource Assessment and Monitoring Division, Northwest Fisheries  
Science Center, National Marine Fisheries Service, NOAA, Seattle, WA, USA

[James.Thorson@noaa.gov](mailto:James.Thorson@noaa.gov)

Keywords: steepness; recruitment; meta-analysis; life-history theory; taxonomy; stock  
assessment

19	<b>Outline</b>	
20	1.	Abstract
21	2.	Introduction
22	3.	Methods
23	3.1	Evolutionary model for life-history parameters
24	3.2	Hierarchical model for stock-recruit parameters
25	3.3	Parameter estimation
26	4.	Results
27	5.	Discussion
28	6.	Acknowledgements
29		
30		

## 1. Abstract

Fisheries managers and scientists use biological models to determine sustainable fishing rates, understand environmental responses, and forecast future dynamics. These models typically require both life-history parameters (mortality, maturity, growth, and size) and stock-recruit parameters (production of juveniles). However, there has been surprisingly little research to simultaneously predict life-history and stock-recruit parameters. I therefore develop the first “data-integrated life-history model,” which modifies the recent *FishLife* model to fit a simple model of evolutionary dynamics to field-measurements of life-history parameters as well as historical estimates of spawning output and subsequent recruitment. This evolutionary model predicts recruitment productivity (“steepness”) and variability (variance and autocorrelation in recruitment deviations) as well as mortality, maturity, growth, and size, and uses these to predict intrinsic growth rate ( $r$ ) for all described fishes. The model confirms previous analysis showing little correlation between recruitment productivity (“steepness”) and either natural mortality or asymptotic maximum size ( $W_{\infty}$ ). However, it does reveal strong taxonomic patterns, where family Sebastidae has lower steepness (mean = 0.63,  $SD$  = 0.18) and Lutjanidae has elevated steepness (mean = 0.77,  $SD$  = 0.16) relative to the prediction for bony fishes (class Actinopterygii, mean = 0.69,  $SD$  = 0.22). Similarly, genus *Sebastes* has growth rate  $r$  (0.07) similar to several shark families (Lamniformes: 0.07; Carcharhiniformes: 0.10). Predictions are tighter for species than for higher-level organizations (e.g., families), and predictions (including multivariate confidence intervals) are available for >32,000 described fishes worldwide in R package *FishLife*. I conclude by outlining how multivariate predictions of life-history and stock-recruit parameters could be useful for stock-assessment, decision theory, ensemble modelling, and strategic management.

## 2. Introduction

Fisheries managers worldwide seek to regulate fishing industries based on biological determination of maximum sustainable yield. Maximum sustainable yield is generally identified by fitting a population-dynamics model to data obtained from a given population. Population-dynamics models are a simplified representation of basic biological processes including natural mortality, individual growth, sexual maturation, and production of juveniles (termed “recruitment”). These processes are typically represented using a small set of population-dynamics parameters, and identifying plausible values for these parameters is a central task for theoretical and applied ecologists for the management of both poor- and well-studied species.

Unfortunately, available data are often insufficient to determine the rate of important biological processes for a given stock. Parameters representing size and growth are often unknown for low-value species, where resources have not been dedicated to collect the age and size of representative individuals. Similarly, sexual maturity requires field sampling of ovaries, and natural mortality rates are estimated from some combination of tag-capture data and by treating the history of a fishery as a depletion experiment. In the following, I refer to adult mortality, growth, and sexual maturity rates as “life-history parameters,” and there is an extensive literature regarding the expected relationship among life-history parameters in fishes (Roff 1984; Winemiller 1989; Charnov and Berrigan 1990; Beverton 1992). Recent research has sought to integrate life-history and evolutionary theory to predict life-history parameters for all fishes worldwide while distributing results in a public, reproducible, and well-documented manner (e.g., Froese *et al.* 2014). For example, *FishLife* is an R package that distributes predictions of adult life-history parameters for all fishes while combining information from taxonomic relatedness and fish records worldwide, and while estimating the degree of

uncertainty from the quantity and quality of records for a given taxon and its relatives (Thorson *et al.* 2017). Improved documentation and access to these meta-analytic results can presumably improve fisheries management for species that currently have little direct biological information.

Despite progress in predicting adult life-history parameters for fishes, these parameters are insufficient to predict population dynamics. In fact, fisheries scientists have recognized for nearly 100 years that fish populations are often composed of a single or few small cohorts (Hjort 1926), and that fluctuations in population size are largely driven by the production of juveniles. Recruitment is typically predicted as a function of spawning output, and parameters governing the expected relationship between spawning output and recruitment are rarely included in classical life-history theory. Instead, stock-recruit parameters have historically been predicted using meta-analyses (e.g., Myers 2001) applied to a database of spawning biomass and subsequent recruitment (termed “stock-recruit data”) or via ecological theories of marine community structure (Andersen and Beyer 2006). The stock-recruit relationship has received a great deal of research attention over the past thirty years both using theoretical arguments (Charnov *et al.* 1991; Neill *et al.* 1994; Iles and Beverton 1998; Mangel *et al.* 2010) and empirical methods (Rose *et al.* 2001; Minto *et al.* 2008; Thorson *et al.* 2014; Foss-Grant *et al.* 2016; Britten *et al.* 2016) to predict variation in stock-recruit parameters. However, there have been surprisingly few empirical studies that simultaneously analyze variation in stock-recruit and adult life-history parameters.

Ecologists increasingly use models combining data from multiple collection protocols (termed “integrated models”) to estimate ecological parameters. These integrated models are useful to estimate processes that are not identifiable using one data set in isolation and are common, e.g., when estimating population-dynamics for assessing stock status (Maunder and

Punt 2013), when estimating growth from tagging and length sampling data (Eveson *et al.* 2004), or when estimating population density using encounter and count data (Zipkin *et al.* 2017). However, data-integrated modelling has not to my knowledge been used in life-history analysis (which typically analyzes databases of field-measurements of population rates) or stock-recruit analysis (which typically analyzes stock-recruit data while assuming that population-dynamics parameters are known and fixed exogenously). Combining these two analyses is sufficient to describe adult fish dynamics (e.g., spawning biomass per recruit without fishing) and recruit dynamics (e.g., recruits per spawning biomass) and therefore is sufficient to describe the entire life-cycle under alternative scenarios regarding fishing effort and selectivity (e.g., Thorson *et al.* 2015b).

In this study, I develop the first data-integrated model for life-history analysis and demonstrate this approach by combining data from life-history and stock-recruit databases. By combining life-history and stock-recruit information, I am able to generate predictions of life-cycle dynamics for all described fishes. I specifically modify the evolutionary model used by *FishLife* to fit both adult life-history parameters as well as stock-recruit measurements for over 150 fish populations worldwide, and use this integrated model for population-dynamics parameters to estimate the strength of recruitment compensation for all fishes. I then use these predictions to generate a full life-cycle model and estimate the intrinsic growth rate  $r$  for all fishes. This parameter is widely used in theoretical and applied studies, e.g., as one of two key parameters in the surplus production models that are taught in introductory fisheries science classes (Haddon 2010). In the discussion, I compare resulting estimates with previous studies to evaluate plausibility of model results. Finally, I distribute predictions of recruitment density dependence and intrinsic growth rate (and associated uncertainty) as a fully documented,



publicly available R package *FishLife* release 2.0.0, so that future users can generate multivariate predictions of life-history and stock-recruit parameters for any described fish species.

### 3. Methods

I seek to predict stock-recruit parameters (the strength of recruitment compensation and the variance and autocorrelation in recruitment variation) and use results to describe life-cycle dynamics (i.e., intrinsic growth rate) for all fishes worldwide. To do so, I integrate two previous meta-analytic models:

1. An evolutionary model of life-history parameters (Thorson *et al.* 2017) fitted to field-measurements of size, growth, mortality, and maturity for thousands of species worldwide as compiled by FishBase (Froese 1990);
2. A hierarchical model for stock-recruit parameters (Thorson *et al.* 2014) fitted to stock and recruitment measurements from the original RAM database (Myers *et al.* 1995).

By fitting an evolutionary model to both data sets simultaneously, I am able to characterize uncertainty arising from estimates of individual life-history parameters as well as stock-recruit processes. I then distribute results using an R package *FishLife* release 2.0.0.

For this analysis, I draw heavily upon previous advice regarding good-practices in fisheries meta-analysis (Thorson *et al.* 2015a). Specifically, I address both experimental variability, e.g., errors in field-measurements of life-history parameters, and parametric variability, e.g., differences in larval and adult survival rates (defined by Osenberg *et al.* 1999). To account for both types of variability, I apply a hierarchical model that explicitly estimates the variance of measurement vs. process errors (Thorson and Minto 2015). Hierarchical models automatically implement “shrinkage,” i.e., the tendency for parameter estimates for each species to be shrunk towards both (1) the average of available data and (2) the prediction given other

model components (Gelman and Hill 2007). Shrinkage permits parameter estimates even for species with no field-records. Shrinkage will also bias estimates for a given taxon towards estimates for related species, but theory suggests that it will minimize expected errors even in cases when some species are mis-categorized or otherwise greatly different from other taxa (Efron and Morris 1977).

### 3.1 Evolutionary model for life-history parameters

I first summarize *FishLife* (Thorson *et al.* 2017), an evolutionary model for size, growth, mortality, and maturity parameters for all fishes worldwide. This model adapts the coalescent model for molecular evolution (Felsenstein 1973) to predict a vector  $\mathbf{x}_g$  of life-history values  $x_{g,j}$  for taxon  $g$  and parameter  $j$  (of  $G$  and  $J$  modeled taxa and parameters).

Life-history parameters  $\mathbf{x}_g$  evolve along a phylogenetic lineage following a multivariate random walk:

$$\mathbf{x}_g \sim MVN(\mathbf{x}_{p(g)}, \mathbf{\Sigma}_{l(g)}) \quad (1)$$

where  $p(g)$  is the taxonomic “parent” of taxon  $g$  (e.g., if  $\mathbf{x}_g$  is the average life-history values for genus *Sebastes*, then  $\mathbf{x}_{p(g)}$  is the average values for family Sebastidae).  $\mathbf{\Sigma}_{l(g)}$  is the evolutionary covariance among life-history parameters:

$$\mathbf{\Sigma}_{l(g)} = \lambda_{l(g)}(\mathbf{L}_{\Sigma}\mathbf{L}_{\Sigma}^T + \mathbf{D}) \quad (2)$$

where  $\mathbf{L}_{\Sigma}$  is a lower-diagonal matrix that approximates covariance among parameters,  $\mathbf{D}$  is a diagonal matrix representing additional variance for the evolution of each trait, and  $l(g)$  represents the taxonomic level of taxon  $g$  (e.g., Class, Order, Family, Genus, or Species) such that  $\lambda_{l(g)}$  represents the proportion of evolutionary covariance occurring at taxonomic level  $l(g)$ . This specification is completed by specifying the mean traits for the top of the fish taxonomy (phylum Chordata) as fixed effects with expected value  $\mu_x$ . The evolutionary covariance from

phylum Chordata to any fish species can be calculated as  $\Sigma = \sum_{l=1}^5 \Sigma_l$ . This model approximates many different evolutionary mechanisms, e.g., by assuming that evolution rates of life-history and stock-recruit parameters are identical for all evolutionary lineages despite likely differences in evolutionary mechanisms and effective population size, and future research could expand the model by incorporating additional information regarding evolutionary rates and divergence times within the fish taxonomic tree.

The association of one trait (e.g., natural mortality  $M$ ) and another trait (e.g., asymptotic mass  $W_\infty$ ) can be interpreted using major axis regression (Warton *et al.* 2006). Major axis regression (MAR) involves calculating the eigen-decomposition of evolutionary covariance  $\Sigma$ , and the ratio of elements in a given eigen-vector is interpreted as the regression “slope” between two variables. MAR is useful given that the slope does not require defining which variable is “dependent” or “independent”, and an estimated slope of  $\beta$  can be interpreted as “a 1% change in variable  $X_1$  is associated with a  $\beta\%$  change in variable  $X_2$ ” whenever both variables have been log-transformed prior to analysis (see Thorson *et al.* 2017 for an example). By contrast, conventional linear modelling (e.g., a regression of natural mortality  $M$  on growth coefficient  $K$ ) requires specifying a dependent and independent variable, and estimates of the slope will depend upon that choice.

The evolutionary model in *FishLife* is completed by specifying a distribution for life-history measurements  $\tilde{\mathbf{y}}_i$  for study  $i$ :

$$\tilde{\mathbf{y}}_i \sim MVN(\mathbf{x}_{g(i)}, \mathbf{V}) \quad (3)$$

where  $g(i)$  is the taxon for study  $i$ , such that  $\mathbf{x}_{g(i)}$  is the expected value for each field-measurement for that study, and  $\mathbf{V}$  represents covariance among studies for a given taxon (resulting from a combination of field-measurement errors and biological variation among stocks

for a given species, years for a given stock, etc). However, most studies do not measure all life-history parameters simultaneously, so I use a “missing-data” model under the assumption that data are missing at random:

$$\tilde{y}_i = \begin{cases} \varepsilon_{i,j} & \text{if } y_{i,j} \text{ was not measured by study } i \\ y_{i,j} & \text{if } y_{i,j} \text{ was measured by study } i \end{cases} \quad (4)$$

where  $\varepsilon_{i,j}$  is assigned a uniform prior distribution,  $\varepsilon_{i,j} \sim Uniform(LB, UB)$  (where the lower bound  $LB$  and upper bound  $UB$  are chosen such that further changes in their values have no effect on model results).

This model was previously fitted to seven life history parameters and average environmental temperature (which I collectively call “life-history parameters”; Table 1), where all were log-transformed prior to analysis (except for average temperature  $Temp$ ). Results from Thorson et al. (2017) showed:

1. Well-known life-history patterns, e.g., a positive association between observed values for growth coefficient ( $K$ ) and natural mortality rate ( $M$ ) and the close-to-isometric relationship between asymptotic length ( $L_\infty$ ) and asymptotic mass ( $W_\infty$ ).
2. Major axes of evolution in life-history parameters, wherein the first axis is associated with temperature, the second with body size, and the third represents a nonlinear relationship between mortality rate and growth coefficient ( $M/K$ ) and the timing of maturation ( $L_{mat}/L_\infty$ ).
3. Useful predictions of uncertainty for both well- and poorly-studied species, where predicted life-history parameters for individual species are more precise (smaller predictive intervals) than predictions for higher-level taxa.

However, that analysis could not predict population dynamics because it did not include parameters governing recruitment.

### 3.2 Hierarchical model for stock-recruit parameters

I next extend the existing FishLife model to fit to estimates of spawning stock size  $S_{i,t}$  and subsequent recruitment  $R_{i,t}$  for stock-recruit study  $i$  and spawning biomass in year  $t$ . In the following, I use notation from Thorson et al. (2014), but modify this model by (1) embedding it within the evolutionary model for life-history parameters, and (2) defining a hyper-distribution for maximum recruits per spawning biomass for each stock. I note that the original RAM database records the age-at-recruitment for each stock  $i$ , and I align data such that  $\hat{R}_{i,t}$  is the record of recruitment arising from spawning that occurs given spawning biomass  $S_{i,t}$ .

I specify a Beverton-Holt model for expected recruitment  $\hat{R}_{i,t}$  for study  $i$  in year  $t$ :

$$\hat{R}_{i,t} = \frac{\alpha_i S_{i,t}}{1 + S_{i,t}/\beta_i} \quad (5)$$

where  $\alpha_i$  is maximum recruits per spawning biomass for stock  $i$  and  $\beta_i$  represents the density-dependent decrease in recruits per spawning biomass with increasing  $S_{i,t}$ . I then specify a distribution for process errors around this assumed relationship:

$$\log(R_{i,t}) = \begin{cases} \log(\hat{R}_{i,t}) + \rho_i(\log(R_{i,t-1}) - \log(\hat{R}_{i,t-1})) + \delta_{i,t} & \text{if } t > 1 \\ \log(\hat{R}_{i,t}) + \delta_{i,t} & \text{if } t = 1 \end{cases} \quad (6)$$

where  $\rho_i$  is the first-order autocorrelation coefficient for study  $i$  and residual errors are normally distributed,  $\delta_{i,t} \sim N(0, \tau_i^2)$ , where  $\tau_i^2$  is the conditional variance for recruitment deviations. This model involves estimating four parameters for every study  $i$ :  $\alpha_i$ ,  $\beta_i$ ,  $\tau_i^2$ , and  $\rho_i$ .

I apply a hierarchical model that shrinks  $\alpha_i$ ,  $\tau_i^2$ , and  $\rho_i$  towards expected values for a given taxon. I assume that carrying capacity varies greatly among species (e.g., because spawning biomass and recruits are expressed in different units among different studies, or mechanisms differ among species), so I treat  $\beta_i$  as a fixed effect for every study  $i$  such that it is

not shrunk towards a common value. By contrast, I specify a hierarchical model for maximum recruits per spawning biomass  $\alpha_i$ , which varies among taxa but is shrunk towards a predicted value in the following way. First, I use the observation that maximum annual spawners per spawner is less variable than other measures of density dependence (Myers *et al.* 1999). I therefore define maximum annual spawners per spawner in excess of replacement  $r_i$  for each study  $i$ :

$$p_i = 1 + \frac{r_i}{1 - \exp(-M_i^*)} \quad (7)$$

where  $p_i$  is maximum lifetime spawners per spawner and  $\exp(-M_i^*)$  is the annual survival fraction given adult natural mortality rate  $M_i^*$ . The  $p_i = 1 + \dots$  in the right-hand-side of Eq. 7 ensures that  $p_i > 1$  for all values of  $\log(r_i)$ , which is a necessary condition for persistence of the population in the absence of fishing, such that  $r_i$  is defined as maximum annual spawners per spawner *in excess of replacement*. Converting between  $r_i$  and  $\alpha_i$  is made possible by the observation that maximum annual recruits per spawning biomass ( $\alpha_i$ ) times lifetime spawning biomass per recruit in the absence of fishing ( $SPR_{F=0}$ ) must equal maximum lifetime spawners per spawner ( $p_i$ ) (Myers and Mertz 1998), such that  $\alpha_i = p_i / SPR_{F=0}^*$ . In summary, I predict  $r_i$ , convert  $r_i$  to  $p_i$  (Eq. 7) and then convert  $p_i$  to  $\alpha_i$ , where  $\alpha_i$  and  $\beta_i$  are used to predict recruitment  $\hat{R}_{i,t}$  for the levels of spawning biomass  $S_{i,t}$  that have been observed for each population. In this analysis, I use values  $SPR_{F=0}^*$  and  $M_i^*$  compiled by Ram Myers in the original stock-recruit database.

Given this model, I then define the variables used in the evolutionary model as the set of life-history parameters that are comparable across different fish species worldwide (see Table 1 for list).

$$\mathbf{x}_g \equiv \{\log(M_g), \log(A_{max,g}), \log(K_g), \log(L_{\infty,g}), \log(W_{\infty,g}), \log(A_{mat,g}), \log(L_{mat,g}), T_i, \log(\tau_g), \log(\rho_{R,g}), \log(r_g)\} \quad (8)$$

where the vector of life-history parameters  $\mathbf{x}_g$  includes  $r_g$  such that maximum annual spawners per spawner in excess of replacement is shrunk towards a value that is correlated both taxonomically and with other life-history parameters.

Given predicted values for life-history parameters  $\mathbf{x}_g$  for each taxon  $g$ , I calculate derived quantities that are useful for other models or management purposes:

1. *Intrinsic growth rate*: Many theoretical and applied studies in ecology have used the intrinsic growth rate ( $r$ ) as an integrated measure of population resilience (Levins 1969). This growth rate is also one of two parameters in the Schaeffer surplus production model, which is the first model that many students learn in introductory fisheries science (Haddon 2010). I calculate  $r$  as the dominant eigenvalue for a Leslie matrix representing population growth in numbers and reproductive output, and it represents the instantaneous rate of population growth at asymptotically low population sizes; it is identical if expressed as growth rate in numbers or biomass (McAllister *et al.* 2001). The calculation requires the parameters predicted by FishLife (Table 1), as well as several that are not available including: the allometric scaling of asymptotic biomass and length (fixed at 3.04 based on the average from Froese *et al.* (2014)), the extrapolated age at zero length in the von Bertalanffy growth curve (fixed at -0.1), and the slope of the maturity ogive at 50% maturity (fixed at 25% of age-at-maturity). I recommend future research to incorporate these parameters into FishLife, although sensitivity analysis suggested that results are relatively insensitive to small changes in these values.

2. *Steepness*: Many stock assessment models use the steepness parameterization of the Beverton-Holt stock-recruit model (Mace 1994)

$$h_g = \frac{p_g}{4 + p_g} \tag{9}$$

where this calculation is derived e.g. by Myers and Mertz (1998) and has been re-derived with small differences elsewhere (Mangel *et al.* 2010 Eq. 26).

3. *Standard deviation for recruitment*: the marginal (pointwise) standard deviation of recruitment is calculated as:

$$\sigma_g = \sqrt{\tau_g^2 / (1 - \rho_g^2)} \tag{10}$$

where this standard deviation (and its standard error) can be used, e.g., when fitting an age-structured surplus production model in data-poor assessments (Thorson *et al.* In press).

I calculate these derived quantities for each taxon  $g$ , append them to the vector of estimated parameters,  $\tilde{\mathbf{x}}_g = (\mathbf{x}_g, \sigma_g^2, h_g, \phi_g)$ , and calculate the expectation and covariance of  $\tilde{\mathbf{x}}_g$  using Monte Carlo sampling.

**3.3 Parameter estimation**

I estimate parameters for this evolutionary model of life-history and stock-recruit parameters given life-history measurements  $y_{i,j}$  in FishBase (Froese 1990) as downloaded using *rfishbase* (Boettiger *et al.* 2015) on Aug. 25, 2016, and stock-recruit data  $R_{i,t}$  and  $S_{i,t}$  from the Myers stock-recruit database (Myers and Mertz 1998). Parameters are estimated using maximum marginal likelihood as implemented using Template Model Builder (Kristensen *et al.* 2016), while using the Laplace approximation (Skaug and Fournier 2006) to integrate the joint likelihood of data and random effects with respect to random effects  $\mathbf{x}_g$  for all taxa. Further details are available in Appendix S1: Computational details. I then distribute results as a



publicly available R package *FishLife* version 2.0.0 (URL will be provided upon acceptance). I specifically provide predicted values  $\hat{\mathbf{x}}_g$  for every described fish species, as well as the predictive covariance  $\widehat{\text{Cov}}(\hat{\mathbf{x}}_g)$  as calculated using a generalization of the delta-method (Kass and Steffey 1989).

#### 4. Results

I first illustrate the predictive intervals for life-history and recruitment parameters for three commercially important species, *Sebastes pinniger*, *Thunnus albacares*, and *Salmo trutta* and the taxonomic family for each species (Fig. 1). As expected, predictions for each individual species are much more precise (have smaller confidence interval width) than predictions for a given family, and predictions for species generally fall within the 95% confidence interval for a given family.

I next present the relationship between parameters using major axis regression (Table 2). This show that the model fitted to life-history and stock-recruit records preserves the life-history relationships documented by Thorson et al. (2017), e.g., the nonlinear relationship between the ratio of mortality and growth rates ( $M/K$ ) and the timing of maturation ( $L_{mat}/L_{inf}$ ). However, FishLife also now fits stock-recruit records, and there is no evidence that the model fits these data poorly (Appendix 2, Suppl. Fig. S1). By fitting to both data sets simultaneously, FishLife is able to estimate the relationship between stock-recruit and life-history parameters. The 1<sup>st</sup> axis (temperature) indicates that a 1°C increase in temperature is associated with a 0.01 increase in steepness on average and a 1% decrease in the standard deviation of recruitment. Similarly, the 2<sup>nd</sup> axis (body size) indicates that a 10% increase in asymptotic body size ( $W_{\infty}$ ) is associated with a 0.4% ( $\frac{0.039}{0.833} = 0.04$ ) increase in recruitment autocorrelation ( $\rho_R$ ), but has an essentially negligible effect on steepness. Finally, the 3<sup>rd</sup> axis (timing of maturation) shows that a 10%

increase in the ratio of mortality and growth (and resulting decrease in  $\frac{L_{mat}}{L_{\infty}}$ ) is associated with a 6% decrease in recruitment variation ( $\frac{-0.076}{0.472 - 0.346} = -0.60$ ). In summary, steepness has a weak-to-nonexistent relationship with body size and timing of maturity, and recruitment autocorrelation ( $\rho_R$ ) has a positive association with age-at-maturity ( $A_{mat}$ ) via axes representing increased body size and the timing of maturation.

I next visualize the predictive relationships between steepness ( $h$ ) or intrinsic growth rate ( $r$ ) and both natural mortality and asymptotic mass (Fig. 2). This shows that steepness has a weakly positive relationship with natural mortality (Fig. 2, 1<sup>st</sup> column), and a weakly negative relationship with asymptotic mass (Fig. 2, 2<sup>nd</sup> column), resulting from the net effect of temperature, body size, and timing of maturation patterns. Although steepness does not have a strong association with either mortality or body size, there does appear to be some taxonomic clustering in both natural mortality and steepness values. For example, Sebastidae generally has lower values for natural mortality and steepness while Scombridae has elevated values of natural mortality and steepness. By contrast, intrinsic growth rate has a strong positive relationship with natural mortality (Fig. 2, 3<sup>rd</sup> column) and a strong negative relationship with asymptotic body size (Fig. 2, 4<sup>th</sup> column). Again, there is strong taxonomic clustering along this axis of variation, where Sebastidae has relatively low natural mortality and intrinsic growth rate, and is somewhat of an outlier in having a asymptotic body size in the middle range of predicted values. Finally, there is also some taxonomic signal in recruitment variance and autocorrelation (Appendix 2, Suppl. Fig. S2), where Sebastidae has somewhat higher recruitment variation than other taxa.

Exploring these taxonomic patterns for steepness in detail (Fig. 3, left column), I confirm that the mean of the predictive distribution for Sebastidae is 0.63 while the mean for Lutjanidae is 0.77, and Salmonidae (0.74) and Scombridae (0.69) are intermediate. By contrast, intrinsic

growth rate for Sebastidae is on the low-end (0.11) and genus *Sebastes* is particularly low (0.07); these values are near those for several Chondrythian families (Carcharhiniformes: 0.10; Lamniformes: 0.07). Meanwhile, Scombridae has a relatively high intrinsic growth rate among the fish families explored here (0.45) and Lutjanidae (0.40) and Salmonidae (0.36) are intermediate.

## 5. Discussion

In this study, I have developed the first integrated life-history model for population-dynamics parameters in fishes, which simultaneously fits to public life-history and stock-recruit databases. The analysis specifically builds upon a simple model for evolution along phylogenetic lineages to reconcile both data sets, and results can be used to predict parameters for species that otherwise have no known information (Thorson *et al.* 2017). Unlike past life-history studies, this model estimates the uncertainty for each parameter (including the correlation between parameters) in a logical way, e.g., where species with many life-history records are more precise than those with few or no records, where species are shrunk towards related species that are well-studied, and where uncertainty is greater for poorly studied species that have few well-studied relatives than those with many well-studied relatives.

This analysis shows that the strength of recruitment compensation (“steepness”) is not strongly correlated with the major axes of life-history variation in fishes (temperature, body size, or timing of maturation), but does cluster taxonomically (e.g., with lower steepness for family Sebastidae and higher for Scombridae). By contrast, intrinsic growth rate has strong correlations with body size, as expected given the dependence of intrinsic growth rate upon natural mortality rates. By using a simple evolutionary model, I am able to predict a unique value for life-history parameters for every species in either data set, as well as all higher-level taxonomic groups of

those species. This contrasts with previous attempts to estimate steepness and intrinsic growth rate, which have typically estimated a single, average value for these quantities for all species (e.g., Jensen *et al.* 2012) or within a given taxonomic grouping (Shertzer and Conn 2012).

I evaluate results by comparing predictions of steepness for higher-level taxa (e.g., for a given genus, family, or order) with those from previous studies to look for differences with previously-published studies (see Table 3). For example, the mode of the predictive distribution of steepness for bony fishes (Actinopterygii) is 0.69 in this study, compared with the predictive mean of 0.75 for all data sets analyzed by Shertzer and Conn (2012). The current analysis did not observe the strong, positive association between steepness and asymptotic size that is predicted from size-structured community theory (Andersen and Beyer 2015). Instead, results were consistent with Shertzer and Conn (2012), who found no strong life-history predictor for steepness. Exploring this in more detail, FishLife predictions of the slope-at-the-origin of the stock-recruit relationship ( $\alpha$ ) shows a strong negative correlation with asymptotic biomass (Appendix S2 and Suppl. Fig. S3) as previously shown by Hall *et al.* (2006 Fig. 1). However, this negative relationship is counter-balanced by a strong positive correlation with unfished spawning biomass per recruit ( $SPR_{F=0}^*$ ), as previously shown by Goodwin *et al.* (2006), such that *FishLife* identifies little correlation of body size with either maximum lifetime spawners per spawner ( $p = \alpha \times SPR_{F=0}^*$ ) or steepness ( $h = \frac{p}{4+p}$ ).

The predictive mode for steepness of genus *Sebastes* (Pacific rockfishes) is 0.54 in this study, compared with 0.58-0.78 obtained from six biannual iterations of a regional meta-analysis for these species on the US West Coast (Thorson *et al.* In press). Steepness predictions from that biannual regional meta-analysis rose sharply between 2007-2011, likely reflecting improved environmental conditions for West Coast rockfishes in the mid-2000s. Predictions in this paper,

by contrast, are based on data from the Myers and Mertz (1998) database, prior to these years of good recruitment. Different steepness predictions from these two meta-analyses emphasizes that meta-analytic results are reflective of stock-recruitment parameters during an historical period that might differ from current environmental conditions. Finally, the current analysis shows that family Sebastidae has higher recruitment variance and autocorrelation than Salmonidae (Supporting Information S1), and this is consistent with results from Thorson et al. (2014) for orders Scorpaeniformes and Salmoniformes. The low value for steepness for genus *Sebastes*, combined with unusually low mortality for their body size, causes *Sebastes* species to have an intrinsic growth rate (0.07) that is similar to several shark families (Lamniformes: 0.07; Carcharhiniformes: 0.10). Large differences in productivity among different fish genera suggest the importance of predicting life-history parameters using taxonomic information in addition to life-history theory.

I note that any meta-analysis is confronted with a trade-off between using aggregated data (e.g., output from an assessment model) for many stocks, or using lower-level data (e.g., output from an index standardization model) for a small number of stocks (Thorson *et al.* 2015a). In this study, I have fitted to the largest data sets available for both life-history parameters (FishBase) and stock-recruit estimates (the RAM database), and results are therefore sensitive to the biases that arise from using model output as “data” (Dickey-Collas *et al.* 2014; Brooks and Deroba 2015). Although it is possible to conduct a meta-analysis for steepness using lower-level data, this has been done previously for only a small number of species (Thorson *et al.* In press). I therefore recommend ongoing research to use lower-level data (e.g., individual records of size and age to estimate growth parameters) within future data-integrated life-history models to eliminate the potential bias arising from analyzing model output as data in a meta-analysis.

I believe that multivariate predictions for multiple population-dynamics parameters will be useful for stock-assessment, decision theory, ensemble modelling, and strategic decision-making, and discuss each of these briefly:

1. *Stock assessment*: These multivariate predictions can be used to define a joint prior distribution (or penalty) for parameters used in a Bayesian (or maximum likelihood) stock assessment model. For example, a recent data-poor assessment model using length-composition data to estimate stock status requires values for maturity, mortality, size, steepness, and recruitment variability (Rudd and Thorson 2017), and a joint prior for all of these parameters can be generated using the results. Using a joint prior distribution prevents assessments from estimating a combination of life-history parameters that are highly unlikely (Brandon *et al.* 2007), e.g., by estimating high mortality  $M$  in combination with large maximum body size  $W_{\infty}$ . The concern with biologically implausible life-history values has motivated development of data-poor assessment models that employ life-history invariants to restrict parameter inputs to plausible values (Hordyk *et al.* 2015; Kokkalis *et al.* 2017). Similarly, using multivariate predictions as a joint prior distribution would restrict parameter estimates to plausible life-history strategies for any assessment model that can use multivariate prior distributions.
2. *Decision theory*: Similarly, a multivariate distribution for life-history parameters can guide the design of simulation experiments that are widely used to evaluate likely performance for ecological models or management procedures (Sainsbury *et al.* 2000). In particular, analysts could use predictions to generate plausible population-dynamics parameters for different life-history types, e.g., by using the predictive distribution for all parameters for genus *Sebastes* to evaluate harvest strategies given simulated dynamics for Pacific rockfishes (Punt 2008).

This would allow researchers to link management decision-theory to the likely population-dynamics for all species worldwide.

3. *Ensemble modelling*: Stock assessment will generally be more robust if management advice is based on an ensemble of alternative life-history assumptions (Stewart and Martell 2015). Results from a model ensemble can be presented using a decision-table (Hilborn *et al.* 1993) or via ensemble-weighting of model results (Anderson *et al.* 2017), but either presentation requires some objective method for determining the weight of different unknown “states of nature”. I therefore recommend that stock assessments increasingly present results using an ensemble of life-history values, where model weights can be obtained from the multivariate distribution for these parameters.
4. *Strategic decisions*: Finally, strategic decision-making in ocean management is increasingly informed by models that include dynamics for multiple species and physical drivers (Fulton *et al.* 2011). For example, ecosystem models have been used to forecast likely impacts of ocean acidification, temperature changes, or invasive species on fisheries potential (Cheung *et al.* 2008; Morello *et al.* 2014; Marshall *et al.* 2017). However, these models often require values for population-dynamics parameters for several to hundreds of species simultaneously, and there is increased interest in characterizing uncertainty in ecosystem model predictions given uncertainty about population-dynamics parameters (Link *et al.* 2012; Collie *et al.* 2016). One source of uncertainty is fixing biological parameters (and their associated standard errors), and this uncertainty could be dealt with uniformly across different species or taxonomic groups using results from this analysis.

For these reasons and more, I encourage continued research regarding data-integrated life-history modelling. These models lie at the intersection of evolutionary, life-history, and stock-recruit

theory, and can translate insights from these theories into tools for applied ecologists and fisheries managers. For this purpose, I distribute the predictions of life-history and stock-recruit parameters (and associated confidence intervals) within R package *FishLife* version 2.0.0 and interested users can read the vignette<sup>1</sup> to learn more.

## 6. Acknowledgements

As always, I thank K. Kristensen, H. Skaug, and the developers of Template Model Builder, without which this analysis would not be feasible. I also thank R. Myers, D. Hively, and C. Minto for assembling and preserving the original RAM stock-recruit database, and R. Froese, D. Pauly, and others for assembling and maintaining the FishBase life-history archive. These two databases are a tremendous asset for fisheries scientists worldwide. Finally, I thank K. Shertzer, J. Hastie, M. McClure, J. Munyandorero, and two anonymous reviewers for comments on an earlier draft.

---

<sup>1</sup> The vignette can be accessed using the following commands within an R terminal:

```
``R
install.packages("devtools")
devtools::install_packages("james-thorson/FishLife")
vignette(topic="tutorial", package="FishLife")`
``
```



## References

- Andersen, K.H. and Beyer, J.E. (2006) Asymptotic size determines species abundance in the marine size spectrum. *The American Naturalist* **168**, 54–61.
- Andersen, K.H. and Beyer, J.E. (2015) Size structure, not metabolic scaling rules, determines fisheries reference points. *Fish and Fisheries* **16**, 1–22.
- Anderson, S.C., Cooper, A.B., Jensen, O.P., et al. (2017) Improving estimates of population status and trend with superensemble models. *Fish and Fisheries*.
- Beverton, R.J.H. (1992) Patterns of reproductive strategy parameters in some marine teleost fishes. *Journal of Fish Biology* **41**, 137–160.
- Boettiger, C., Chamberlain, S., Lang, D.T. and Wainwright, P. (2015) *rfishbase: R Interface to "FishBase."*
- Brandon, J.R., Breiwick, J.M., Punt, A.E. and Wade, P.R. (2007) Constructing a coherent joint prior while respecting biological realism: application to marine mammal stock assessments. *ICES Journal of Marine Science: Journal du Conseil* **64**, 1085–1100.
- Britten, G.L., Dowd, M. and Worm, B. (2016) Changing recruitment capacity in global fish stocks. *Proceedings of the National Academy of Sciences* **113**, 134–139.
- Brooks, E.N. and Deroba, J.J. (2015) When "data" are not data: the pitfalls of post hoc analyses that use stock assessment model output. *Canadian Journal of Fisheries and Aquatic Sciences* **72**, 634–641.
- Charnov, E.L. and Berrigan, D. (1990) Dimensionless numbers and life history evolution: age of maturity versus the adult lifespan. *Evolutionary Ecology* **4**, 273–275.
- Charnov, E.L., Berrigan, D. and Beverton, R.J.H. (1991) Dimensionless numbers and the assembly rules for life histories [and discussion]. *Philosophical Transactions of the Royal Society of London. Series B: Biological Sciences* **332**, 41–48.
- Cheung, W.W.L., Lam, V.W.Y. and Pauly, D. (2008) Modelling present and climate-shifted distribution of marine fishes and invertebrates.
- Collie, J.S., Botsford, L.W., Hastings, A., et al. (2016) Ecosystem models for fisheries management: finding the sweet spot. *Fish and Fisheries* **17**, 101–125.
- Dickey-Collas, M., Payne, M.R., Trenkel, V.M. and Nash, R.D.M. (2014) Hazard warning: model misuse ahead. *ICES Journal of Marine Science* **71**, 2300–2306.
- Efron, B. and Morris, C.N. (1977) *Stein's paradox in statistics*. WH Freeman.
- Eveson, J.P., Laslett, G.M. and Polacheck, T. (2004) An integrated model for growth incorporating tag recapture, length frequency, and direct aging data. *Canadian Journal of Fisheries and Aquatic Sciences* **61**, 292–306.
- Felsenstein, J. (1973) Maximum-likelihood estimation of evolutionary trees from continuous characters. *American Journal of Human Genetics* **25**, 471.
- Forrest, R.E., McAllister, M.K., Dorn, M.W., Martell, S.J.D. and Stanley, R.D. (2010) Hierarchical Bayesian estimation of recruitment parameters and reference points for Pacific rockfishes (*Sebastes* spp.) under alternative assumptions about the stock-recruit function. *Canadian Journal of Fisheries and Aquatic Sciences* **67**, 1611–1634.
- Foss-Grant, A.P., Zipkin, E.F., Thorson, J.T., Jensen, O.P. and Fagan, W.F. (2016) Hierarchical analysis of taxonomic variation in intraspecific competition across fish species. *Ecology* **97**, 1724–1734.
- Froese, R. (1990) FishBase: An information system to support fisheries and aquaculture research. *ICLARM Fishbyte* **8**, 21–24.
- Froese, R., Thorson, J.T. and Reyes, R.B. (2014) A Bayesian approach for estimating length-weight relationships in fishes. *Journal of Applied Ichthyology* **30**, 78–85.
- Fulton, E.A., Link, J.S., Kaplan, I.C., et al. (2011) Lessons in modelling and management of marine ecosystems: the Atlantis experience. *Fish and Fisheries*.

- Gelman, A. and Hill, J. (2007) *Data analysis using regression and multilevel/hierarchical models*. Cambridge University Press, Cambridge, UK.
- Goodwin, N.B., Grant, A., Perry, A.L., Dulvy, N.K. and Reynolds, J.D. (2006) Life history correlates of density-dependent recruitment in marine fishes. *Canadian Journal of Fisheries and Aquatic Sciences* **63**, 494–509.
- Haddon, M. (2010) *Modelling and Quantitative Methods in Fisheries, Second Edition*. CRC Press.
- Hall, S.J., Collie, J.S., Duplisea, D.E., Jennings, S., Bravington, M. and Link, J. (2006) A length-based multispecies model for evaluating community responses to fishing. *Canadian Journal of Fisheries and Aquatic Sciences* **63**, 1344–1359.
- Hilborn, R., Pikitch, E.K. and Francis, R.C. (1993) Current Trends in Including Risk and Uncertainty in Stock Assessment and Harvest Decisions. *Canadian Journal of Fisheries and Aquatic Sciences* **50**, 874–880.
- Hjort, J. (1926) Fluctuations in the year classes of important food fishes. *ICES Journal of Marine Science* **1**, 5.
- Hordyk, A., Ono, K., Sainsbury, K., Loneragan, N. and Prince, J. (2015) Some explorations of the life history ratios to describe length composition, spawning-per-recruit, and the spawning potential ratio. *ICES Journal of Marine Science: Journal du Conseil* **72**, 204–216.
- Iles, T.C. and Beverton, R.J.H. (1998) Stock, recruitment and moderating processes in flatfish. *Journal of Sea Research* **39**, 41–55.
- Jensen, O.P., Branch, T.A. and Hilborn, R. (2012) Marine fisheries as ecological experiments. *Theoretical Ecology* **5**, 3–22.
- Kass, R.E. and Steffey, D. (1989) Approximate bayesian inference in conditionally independent hierarchical models (parametric empirical bayes models). *Journal of the American Statistical Association* **84**, 717–726.
- Kokkalis, A., Eikeset, A.M., Thygesen, U.H., Steingrund, P. and Andersen, K.H. (2017) Estimating uncertainty of data limited stock assessments. *ICES Journal of Marine Science* **74**, 69–77.
- Kristensen, K., Nielsen, A., Berg, C.W., Skaug, H. and Bell, B.M. (2016) TMB: Automatic Differentiation and Laplace Approximation. *Journal of Statistical Software* **70**, 1–21.
- Levins, R. (1969) Some Demographic and Genetic Consequences of Environmental Heterogeneity for Biological Control. *Bulletin of the ESA* **15**, 237–240.
- Link, J.S., Ihde, T.F., Harvey, C.J., et al. (2012) Dealing with uncertainty in ecosystem models: The paradox of use for living marine resource management. *Progress in Oceanography* **102**, 102–114.
- Mace, P.M. (1994) Relationships between Common Biological Reference Points Used as Thresholds and Targets of Fisheries Management Strategies. *Canadian Journal of Fisheries and Aquatic Sciences* **51**, 110–122.
- Mangel, M., Brodziak, J. and DiNardo, G. (2010) Reproductive ecology and scientific inference of steepness: a fundamental metric of population dynamics and strategic fisheries management. *Fish and Fisheries* **11**, 89–104.
- Marshall, K.N., Kaplan, I.C., Hodgson, E.E., et al. (2017) Risks of ocean acidification in the California Current food web and fisheries: ecosystem model projections. *Global Change Biology* **23**, 1525–1539.
- Maunder, M.N. and Punt, A.E. (2013) A review of integrated analysis in fisheries stock assessment. *Fisheries Research* **142**, 61–74.
- McAllister, M.K., Pikitch, E.K. and Babcock, E.A. (2001) Using demographic methods to construct Bayesian priors for the intrinsic rate of increase in the Schaefer model and implications for stock rebuilding. *Canadian Journal of Fisheries and Aquatic Sciences* **58**, 1871–1890.

- 559 Minto, C., Myers, R.A. and Blanchard, W. (2008) Survival variability and population density in fish  
560 populations. *Nature* **452**, 344–347.
- 561 Morello, E.B., Plagányi, É.E., Babcock, R.C., Sweatman, H., Hillary, R. and Punt, A.E. (2014) Model to  
562 manage and reduce crown-of-thorns starfish outbreaks. *Marine Ecology Progress Series* **512**,  
563 167–183.
- 564 Myers, R. and Mertz, G. (1998) Reducing uncertainty in the biological basis of fisheries management by  
565 meta-analysis of data from many populations: a synthesis. *Fisheries Research* **37**, 51–60.
- 566 Myers, R.A. (2001) Stock and recruitment: generalizations about maximum reproductive rate, density  
567 dependence, and variability using meta-analytic approaches. *ICES Journal of Marine Science: Journal du Conseil* **58**, 937–951.
- 568 Myers, R.A., Bowen, K.G. and Barrowman, N.J. (1999) Maximum reproductive rate of fish at low  
569 population sizes. *Canadian Journal of Fisheries and Aquatic Sciences* **56**, 2404–2419.
- 570 Myers, R.A., Bridson, J. and Barrowman, N.J. (1995) Summary of worldwide spawner and recruitment  
571 data. 2024. Department of Fisheries and Oceans Canada, Northwest Atlantic Fisheries Centre, St.  
572 Johns, Newfoundland.
- 573 Neill, W.H., Miller, J.M., Van Der Veer, H.W. and Winemiller, K.O. (1994) Ecophysiology of marine fish  
574 recruitment: A conceptual framework for understanding interannual variability. *Netherlands Journal of Sea Research* **32**, 135–152.
- 575 Osenberg, C.W., Sarnelle, O., Cooper, S.D. and Holt, R.D. (1999) Resolving ecological questions through  
576 meta-analysis: goals, metrics, and models. *Ecology* **80**, 1105–1117.
- 577 Punt, A.E. (2008) Refocusing stock assessment in support of policy evaluation. In: *Fisheries for Global*  
578 *Welfare and Environment*. (eds K. Tsukamoto, T. Kawamura, T. Takeuchi, T.D. Beard and M.J.  
579 Kaiser). TerraPub, Tokyo, pp 139–152.
- 580 Roff, D.A. (1984) The evolution of life history parameters in teleosts. *Canadian Journal of Fisheries and*  
581 *Aquatic Sciences* **41**, 989–1000.
- 582 Rose, K.A., Cowan, J.H., Winemiller, K.O., Myers, R.A. and Hilborn, R. (2001) Compensatory density  
583 dependence in fish populations: importance, controversy, understanding and prognosis. *Fish*  
584 *and Fisheries* **2**, 293–327.
- 585 Rudd, M.B. and Thorson, J.T. (2017) Accounting for variable recruitment and fishing mortality in length-  
586 based stock assessments for data-limited fisheries. *Canadian Journal of Fisheries and Aquatic*  
587 *Sciences*.
- 588 Sainsbury, K.J., Punt, A.E. and Smith, A.D.. (2000) Design of operational management strategies for  
589 achieving fishery ecosystem objectives. *ICES Journal of Marine Science: Journal du Conseil* **57**,  
590 731–741.
- 591 Shertzer, K.W. and Conn, P.B. (2012) Spawner-Recruit Relationships of Demersal Marine Fishes: Prior  
592 Distribution of Steepness. *Bulletin of Marine Science* **88**, 39–50.
- 593 Skaug, H. and Fournier, D. (2006) Automatic approximation of the marginal likelihood in non-Gaussian  
594 hierarchical models. *Computational Statistics & Data Analysis* **51**, 699–709.
- 595 Stewart, I.J. and Martell, S.J.D. (2015) Reconciling stock assessment paradigms to better inform fisheries  
596 management. *ICES Journal of Marine Science* **72**, 2187–2196.
- 597 Thorson, Cope, J.M., Kleisner, K.M., Samhouri, J.F., Shelton, A.O. and Ward, E.J. (2015a) Giants’  
598 shoulders 15 years later: lessons, challenges and guidelines in fisheries meta-analysis.
- 599 Thorson, Dorn, M. and Hamel, O.S. (In press) Steepness for West Coast rockfishes: Results from a  
600 twelve-year experiment in iterative regional meta-analysis.
- 601 Thorson, Jensen, O.P. and Hilborn, R. (2015b) Probability of stochastic depletion: an easily interpreted  
602 diagnostic for stock assessment modelling and fisheries management. *ICES Journal of Marine*  
603 *Science: Journal du Conseil* **72**, 428–435.

Thorson, Jensen, O.P. and Zipkin, E.F. (2014) How variable is recruitment for exploited marine fishes? A hierarchical model for testing life history theory. *Canadian Journal of Fisheries and Aquatic Sciences* **71**, 973–983.

Thorson, J.T. and Minto, C. (2015) Mixed effects: a unifying framework for statistical modelling in fisheries biology. *ICES Journal of Marine Science: Journal du Conseil* **72**, 1245–1256.

Thorson, J.T., Munch, S.B., Cope, J.M. and Gao, J. (2017) Predicting life history parameters for all fishes worldwide. *Ecological Applications* **27**, 2262–2276.

Thorson, J.T., Rudd, M. and Winker, H. (In review) The case for estimating recruitment variation in data-moderate and data-poor age-structured models.

Warton, D.I., Wright, I.J., Falster, D.S. and Westoby, M. (2006) Bivariate line-fitting methods for allometry. *Biological Reviews* **81**, 259–291.

Winemiller, K.O. (1989) Patterns of variation in life history among South American fishes in seasonal environments. *Oecologia* **81**, 225–241.

Zipkin, E.F., Rossman, S., Yackulic, C.B., Wiens, J.D., Thorson, J.T., Davis, R.J. and Grant, E.H.C. (2017) Integrating count and detection–nondetection data to model population dynamics. *Ecology* **98**, 1640–1650.

625 Table 1 – List of life-history parameters (including their name, symbol, the database used to  
 626 estimate them, and the type of parameter) that are predicted (along with estimated covariance)  
 627 for all fishes worldwide

Name	Symbol	Database	Type
Mortality rate	$M$	FishBase	Mortality
Maximum age	$A_{max}$	FishBase	Mortality
Growth coefficient	$K$	FishBase	Size
Asymptotic length	$L_{\infty}$	FishBase	Size
Asymptotic mass	$W_{\infty}$	FishBase	Size
Age at maturity	$A_{mat}$	FishBase	Maturity
Length at maturity	$L_{mat}$	FishBase	Maturity
Average temperature	$T$	FishBase	Environmental conditions
Marginal standard deviation of recruitment variability	$\sigma_R$	RAM	Recruitment
Autocorrelation of recruitment variability	$\rho_R$	RAM	Recruitment
Steepness	$h$	RAM	Recruitment
Fishing mortality rate at maximum sustainable yield	$F_{msy}$	FishBase, RAM	Management

628

Table 2 – Summary of the dominant axes of covariation for evolutionary covariance  $\Sigma$ . The ratio of two elements of a given eigenvector (columns) is interpreted as the “slope” between these two variables (see text below Eq. 2 in main text for details), while the proportion of variance explained by a given eigenvector (row “Proportion”) measures what proportion of total variance is explained by a given eigenvector.

	Eigenvectors		
	1	2	3
Variance decomposition			
Eigenvalues	59.918	11.959	1.298
Proportion	0.808	0.161	0.018
Cumulative proportion	0.808	0.97	0.987
Loadings			
$\log(M)$	0.055	-0.209	0.472
$\log(A_{max})$	-0.042	0.202	-0.505
$\log(K)$	0.046	-0.203	0.346
$\log(L_{\infty})$	-0.020	0.256	0.133
$\log(W_{\infty})$	-0.043	0.833	0.387
$\log(A_{mat})$	-0.037	0.219	-0.459
$\log(L_{mat})$	-0.025	0.237	0.087
$T$	0.994	0.084	-0.059
$\log(\sigma_R)$	-0.011	-0.039	-0.076
$\rho_R$	0.006	0.039	-0.085
$h$	0.010	-0.001	0.005

Table 3 – Mode of the predictive distribution for steepness ( $h$ ) and intrinsic growth rate ( $r$ ) for bony fishes (Actinopterygii) and common taxonomic orders and families (with predictive standard deviation in parentheses). Also shown is the range of predictive mean steepness (and range of predictive standard deviations) for West Coast rockfishes from six biannual meta-analyses compiled by Thorson et al. (In press), the sample mean (and sample standard deviation) for steepness from all compiled data sets from Shertzer and Conn (2012).

	This study	Steepness ( $h$ ) Thorson et al. (In press)	Shertzer and Conn (2012)	Intrinsic growth rate ( $r$ ) This study
Class				
Actinopterygii	0.69 (0.22)	-	0.75 (0.15)	1.91 (1.37)
Order				
Carcharhiniformes	0.50 (0.22)	-		0.10 (0.19)
Lamniformes	0.51 (0.22)	-		0.07 (0.12)
Clupeiformes	0.72 (0.21)	-		0.55 (0.66)
Gadiformes	0.63 (0.22)	-		0.23 (0.32)
Perciformes	0.71 (0.21)	-		0.41 (0.48)
Pleuronectiformes	0.76 (0.19)	-		0.50 (0.57)
Scorpaeniformes	0.62 (0.22)	-		0.26 (0.38)
Family				
Sebastidae	0.63 (0.18)	-		0.11 (0.12)
Salmonidae	0.74 (0.17)	-		0.36 (0.30)
Scombridae	0.69 (0.18)	-		0.45 (0.39)
Lutjanidae	0.77 (0.16)	-		0.40 (0.31)
Genus				
Sebastes	0.54 (0.13)	0.58-0.78 (0.15-0.20)		0.07 (0.05)


Fig. 1 – Predictive distribution for selected life-history and stock-recruit parameters (see Table 1) for three commercially important families (Sebastidae, Scombridae, and Salmonidae) and a selected species from each family (*Sebastes pinniger*, *Thunnus albacares*, and *Salmo trutta*).

Fig. 2 – Visualization of the natural mortality rate ( $M$ , 1<sup>st</sup> and 3<sup>rd</sup> columns and plotted on x-axis on log-scale) or asymptotic mass ( $W_{\infty}$ , 2<sup>nd</sup> and 4<sup>th</sup> columns and plotted on x-axis on log-scale) and steepness ( $h$ , 1<sup>st</sup> and 2<sup>nd</sup> columns and plotted on y-axis using logit-scale from 0.2 to 1.0) or intrinsic growth rate ( $r$ , 2<sup>nd</sup> and 3<sup>rd</sup> columns and plotted on y-axis on log-scale) for every Class, Order, Family, Genus, and Species in the database, while highlighting the predictive distribution for commercially important families (Sebastidae, Salmonidae, Scombridae, and Lutjanidae). For Class, Order and Family I show the 95% predictive distribution (an ellipse, given that the model uses a multivariate normal distribution), while for Genus and Species I show the for each modeled stock in that taxon.

Fig. 3 – Predictive probability (y-axis) for steepness  $h$  (left column) or intrinsic growth rate  $r$  (right column) for every described class, order, family, genus, and species in FishBase (panels, ordered top to bottom), with families Sebastidae, Salmonidae, Scombridae, and Lutjanidae highlighted to emphasize these particular, commercially-important species




662 Fig. 1




663

664

665 Fig. 2



667 Fig. 3



668

669

## Appendix 1: Computational details

I estimate parameters using Template Model Builder (Kristensen *et al.* 2016), which applies automatic differentiation to calculate gradients of the joint likelihood with respect to fixed and random effects, and uses an inner-optimize to optimize the joint likelihood with respect to random effects. It then applies the Laplace approximation, and provides the gradient of the marginal likelihood with respect to fixed effects. I use these gradients in a gradient-based Nelder-Mead optimizer within the R statistical environment (R Core Team 2017) to identify fixed effects that optimize the marginal likelihood, followed by two iterations of a Newton optimizer step to ensure that remaining gradients are very low. I check convergence by ensuring that the hessian matrix is positive definite, and that the gradient of the marginal log-likelihood is within  $\pm 10^{-6}$  for all fixed effects.

To improve convergence, I use the following numerical and analytical techniques:

1. I use a nonlinear generalization of restricted maximum likelihood (REML), which involves treating non-variance fixed effects (e.g., density dependence  $\beta_i$  and expected values for shared ancestor  $\mu_x$ ) as random effects with a “flat” hyperdistribution. This improves calculation of gradients for combinations of fixed and random effects that are highly correlated (e.g., density dependence  $\beta_i$  and recruitment compensation  $r_i$  for each stock-recruit data set).
2. I add a small penalty to the objective function which favors plausible values for stock-recruit parameters ( $\alpha_i$  and  $\beta_i$  for each stock  $i$ ). Specifically, I regularize density dependence towards a shared mean,  $\beta_i \sim \text{Norm}(0,10)$  for each stock  $i$ . I also add a penalty towards stock-recruit fits where predicted recruitment is constant for all observations of a given stock, where

penalty  $p_i = 0.5 \log(\text{Var}_t(\hat{R}_{i,t}))$  is added to the log-likelihood function. This regularization penalty on the variance of predicted recruitment was unnecessary in previous Bayesian hierarchical models for stock-recruit meta-analysis (Thorson *et al.* 2014; Foss-Grant *et al.* 2016), but appears to be necessary when using the Laplace approximation for Empirical Bayes estimation.


3. I have removed two stocks where the standard error for density dependence  $\beta_i$  approached the regularization penalty (i.e.,  $\hat{SE}(\beta_i) \rightarrow 10$ ). Exploration showed that eliminating these stocks improved model stability without noticeably impacting results. This decision resulted in 151 stocks available for analysis.

These three techniques allow the model to converge with random starting values for different complexities of evolutionary and measurement covariance (i.e., number of columns in  $\mathbf{L}_\Sigma$  and  $\mathbf{L}_\mathbf{V}$ ).

**Appendix 2: Visualizing other model results**

Suppl. Fig. S1 – Fits to stock-recruit records for 151 stocks in the original RAM database. For each stock (panel), I show recruitment (y-axis) for a given record of spawning biomass (x-axis), visualizing the recruitment record (black point), predicted recruitment from the stock-recruit relationship (blue point), or predicted recruitment from the stock-recruit relationship and the impact of autocorrelated errors (red point). I also list the marginal standard deviation of recruitment, the magnitude of autocorrelation, and the steepness parameter.

Suppl. Fig. S2 – Visualization of the conditional variance of recruitment ( $\tau_t^2$ , x-axis left column) or marginal standard deviation of recruitment ( $\sigma_R$ , x-axis right column) and the first-order autocorrelation in recruitment ( $\rho_R$ , y-axis), see Fig. 2 caption for details.




49


50

51    Suppl. Fig. S3 – Visualization of stock-recruit parameters (y-axis) against maximum biomass (x-  
52    axis on log-scale) for the 150 stocks that had available records of spawning biomass and  
53    subsequent recruitment in the original RAM database, showing slope-at-the-origin of the stock-  
54    recruit relationship (top-left panel), maximum annual spawners per spawner (top-right panel),  
55    and steepness (bottom-left panel). Each panel also shows fits to a linear regression, and the p-  
56    value for a two-sided Wald test in the bottom-right corner.


For Review Only








CHUMNBC




CHUMQCI




Fish and Fisheries

CHUMWVAN




CMACKCAL




Page 42 of 49


COD2J3KL




COD3M3




COD3NO




COD3Ps




COD4TVn




COD4VsW




COD4X




COD5Z




CODBA2224




CODBA2532




CODCS




CODFAPL







HADNS2



HADVIa




Fish and Fisheries




HADVIa2

HAKENOR




Page 44 of 49


HERR25-27




HERR25-2932GOR2




HERR302




HERR4-5




HERRBOTB2




HERRCC




HERRCS




HERRDOWN




HERRGM




HERRGOR



















HERRIspr




HERRIsun






PERCHEBS




PERCHGA




Fish and Fisheries


PERCHGOOSE




PHAKEVANCON




PIKEWINN




PIKEWINS




PLAIC7d




PLAIC7e




PLAICCS




PLAICIIIa




PLAICIS




PLAICKAT




PLAICNS




POLLFA




POLLICE




POLLNEAR








SOCKCHIK




SOCKEG3




Fish and Fisheries

SOCKFRAZ




SOCKHELY




Page 48 of 49


SOCKIGU2




SOCKKAR




SOCKKAS




SOCKKEN




SOCKKVIC




SOCKLSTU




SOCKNAK3




SOCKNUSH




SOCKNUY




SOCKREDR




SOCKRINL




SOCKSK2






**WHITVla**




**WHITVla2**




Fish and Fisheries


**WPOLLEBS**




**WPOLLGA**



**YELL5Z**



**YELLSNE**



**YFINEPAC**

