

Unnecessarily Complicated Research Title

John Smith

California, United States

Abstract

Keywords: Science, Publication, Complicated

1. Introduction

- Risk and uncertainty
- One rule for all?
- Impact of life histories
- Comparison of constant catch v changing catch based on trends in an empirical index.
- catch only - need good catch data if you havent got this how can you set catch limits

Sustainability and risks to non target exploited marine fish stock populations requires both estimates of current stock status, the effects of fishing pressure (catchability and fishing effort) and the effects of management measures on target populations, however these data are often lacking. Subsequently there is increasing concern and a growing need for the development of innovative approaches so that management of all marine stocks not just those of high commercial value can be included into the Common Fisheries Policy (CFP [1]) framework. Under the CFP management objectives are to recover stocks and to maintain stocks within safe biological limits to levels that can produce Maximum Sustainable Yield (MSY), including by-catch species by 2015 (Implementation Plan adopted at the World Summit on Sustainable Development, Johannesburg in 2002) and no later than 2020. These conservation objectives are currently being achieved by introducing

biological target (can fluctuate around targets) and limit (i.e must not be exceeded) reference points e.g. population size (stock biomass) and/or yields (catches) and/or long-term yields and fishing mortality against which the preservation of stocks within such limits are assessed. These targets or limit reference points are often referred to as harvesting strategies which include an operational component called a harvest control rule (HCR) that are based on indicators (e.g. monitoring data or models) of stock status and to prevent overfishing.

The International Council for the Exploration of the Sea (ICES) categorises stocks in to classes *data-rich*, (categories 1 and 2) i.e those that have a quantitative assessment based on conventional methods that require large amounts of data that include a long historical time series of catches and sound biological information [2]; or *data-limited* [3](categories 3 and 4) (often called data poor) those without assessment, forecasts and have limited funding for research. For data-rich stocks ICES uses two types of reference points for providing fisheries advice;

1. Precautionary Approach (PA) reference points (those relating to stock status and exploitation relative to precautionary objectives) and
2. MSY reference points (those relating to achieving MSY)

In contrast for data limited stocks MSY *proxy* reference points are used to estimate stock status and exploitation. Often many of the methods used to estimate MSY proxy reference points require length based inputs as they are cheap, easy to collect [4] and are related to life history parameters such as fish size, mortality and fecundity as well as fishery selectivity. For example many methods are being developed to estimate MSY, but currently only 4 are approved by ICES, these include, Surplus Production model in Continuous Time (catch based) (SPiCT; [5], Mean Length Z (MLZ; [6]), Length Based Spawner Per Recruit (LBSPR; [7]) and Length Based Indicators (LBI; e.g. [8]). The aforementioned data limited procedures have differing data requirements, intended uses and obviously have their own strengths and weaknesses.

To test the performance of candidate management procedures often requires evaluation of alternative hypothesis about the dynamics of the system e.g. population dynamics (life history dynamics such as growth parameters which are an indication of fishery exploitation levels and management) and the behaviour of the fishery (e.g range contraction and density dependence) etc.. Due to the nature of conflicting objectives, stakeholder interests and the uncertainty in the dynamics of the resource and/ or the plausibility of alter-

native hypotheses can lead to poor decision making and can be problematic when defining management policy.

An intense area of work being researched over the last 2 decades is Management Strategy Evaluation (MSE), which focuses on the broader aspects of fishing (the Ecosystem) whereby different management options are tested against a range of objectives (see [9]). The approach is not to come up with a definitive answer, but to lay-bare the trade offs associated with each management objective, along with identifying and incorporating uncertainties in the evaluation and communicating the results effectively to client groups and decision-makers. MSE is not intended to be complex but to provide a robust framework that account for conflicting poorly defined objectives and uncertainties that have been absent in conventional management [9].

To assess case specific harvest strategies (via simulation) within the MSE, we will implement a management procedure based on a empirical HCR that adjusts yield depending on stock status for a given set of tunable parameters for each of the harvest strategies and to test their robustness to uncertainty. This approach could also help identify similar conditions across species where particular advice rules are likely to work well, and where they perform poorly for a given a set of parameters.

Often empirical harvest control rules require extensive exhaustive parameter searches to tune or optimise 'hyper-parameters' (external parameters to a model) that aren't directly learnt from estimators. This requires a technique known as a grid search that extensively searches for all combinations of all parameters. In contrast, and some what less time consuming alternative and efficient parameter search strategies can be considered for a given range of parameter space and a known distribution. As such a random sample can be obtained and used to perform the different experiments for parameter optimisation. This approach differs from the simplest constant catch HCR strategy whereby catches are kept constant and low to ensure no lasting damage is done in periods of low stock productivity or whereby the stock is highly variable year on year, therefore the empirical approach can help optimise catch by setting a precautionary Total Allowable Catch (TAC).

Here we describe methods to assess the performance of each empirical HCR via a set of utilities and show the benefits compared to a constant catch strategy: safety ($B/B_{MSY} > 1$), yield ($yield/MSY$), kobe proportion (proportion of years that stay in the green zone of kobe plot ($B/B_{MSY} > 1$), and Yield Annual Variation (yield changes by 10% year on year).

2. Material and Methods

2.1. Materials

Life history parameters for growth, natural mortality and maturity were used to develop an age-based Operating Model. To do this the parameters were first used to parameterise functional forms for mass (W), proportion mature (Q), natural mortality (M) and fishing mortality (F) at age. These were then used to calculate the spawner (S/R) and yield-per-recruit (Y/R) which were then combined with a stock recruitment relationship [10] to calculate the equilibrium stock size as a function of fishing mortality (F).

This analysis allows a variety of reference points such as those based on Maximum Sustainable Yield (MSY), i.e. B_{MSY} the spawning stock biomass (S) and F_{MSY} the fishing mortality that produces MSY at equilibrium to be estimated. Other reference points are $F_{0.1}$ the fishing mortality on the yield per recruit curve where the slope is 10% of that at the origin, a conservative proxy for F_{MSY} ; and F_{Crash} which is the fishing mortality that will drive the stock to extinction since it is equivalent to a R/S greater than the slope at the origin of the stock recruitment relationship, i.e. recruitment can not replace removals for a fishing mortality equal to F_{Crash} .

The equilibrium relationships can then be turned into a forward dynamic model and projected forward.

A variety of functional forms can be assumed for all of the various process, i.e. growth, mortality, maturity, the selection pattern of the fisheries and the stock recruitment relationship. Commonly processes such as growth and maturity-at-age are well known while those for natural mortality and the stock recruitment relationship are poorly known [11]. In the later case assumptions have to be made and to evaluate the sensitivity of any analysis to those assumptions a variety of scenarios are considered. Below a base case is defined that can then be modified to create a variety of scenarios.

2.2. Methods

Individual Growth

Growth in length is modelled by the Von Bertalanffy growth equation @vonbert1957quantitative

$$L = L_{\infty}(1 - \exp(-k(t - t_0))) \quad (1)$$

128 where k is the rate at which the rate of growth in length declines as
 129 length approaches the asymptotic length L_∞ and t_0 is the hypothetical time
 130 at which an individual is of zero length.

131 Length is converted to mass using the length-weight relationship

$$W = aL_t^b \quad (2)$$

132 where a is the condition factor and b is the allometric growth coefficient.

133 Maturity-at-age

134 Proportion mature-at-age is modelled by the logistic equation with 2 pa-
 135 rameters: age at 50% (a_{50}) and 95% (a_{95}) mature.

$$f(x) = \begin{cases} 0 & \text{if } (a_{50} - x)/a_{95} > 5 \\ a_\infty & \text{if } (a_{50} - x)/a_{95} < -5 \\ \frac{m_\infty}{1.0 + 19.0^{(a_{50} - x)/a_{95}}} & \text{otherwise} \end{cases} \quad (3)$$

136 2.2.1. Operating Model

137 Age based.

138 2.2.2. Management Procedure

139 The management procedure was based on an empirical MP, where an
 140 increase in an index of abundance resulted in an increase in the TAC, while
 141 a decrease in the index results in an decrease in the TAC.

142 2.2.3. Random Search

143 When running an MSE commonly a set of MP scenarios are run to tune
 144 the MP, this requires running the MSE for each OM scenario for a range of
 145 fixed values in the HCR and then choosing the rule that best meets manage-
 146 ment objectives. If there are a lot of parameters to tune then a grid search
 147 may become unfeasible. An alternative is random search [12] as randomly
 148 chosen trials are more efficient for parameter optimisation than trials based
 149 on a grid.

150 3. Results

151 Estimates of the simulated life history parameters obtained from Fishbase
 152 (<http://www.fishbase.org>) are presented in Fig.1. These show that for fast
 153 growing species which are small in size l_∞ (asymptotic length parameter)

species such as sprat, the growth parameter k is high. The sprats age-at-50%-maturity a_{50} are low, in contrast to a slower growing larger longer lived species l_{∞} such as rays or pollack.

Observations in Fig.2 shows that the assumed maturity in the OM is related to selectivity and that the faster growing species are more susceptible to fishing, although the slower growing larger (by mass and length) species (e.g. pollack has a higher natural mortality rate at lower ages) with the most significant rate increases associated with turbot. A levelling off in the mortality rate is evident for ray just prior to age 4.5. In contrast, there have been less steep declines in natural mortality estimates for brill, but most notably for sprat.

Fig.3 displays the equilibrium relationships of the operating model. Comparisons of reference points estimates can be made across species. The m/k plot shows interesting trends with lower values for sprat where the growth rate k is considerably higher than the natural mortality rate with little uncertainty around the estimate. In contrast to a slower growing species such as pollack where natural mortality is higher, as is the uncertainty around the estimate. The relationships when compared the proxy for fishing pressure f/m show that the estimate is considerably higher in sprat than pollack, however the intrinsic population growth rate r shows that sprats reproductive capacity is higher and thus its surplus production.

[EXAMPLES TO BE UPDATED]

- Figure 3 shows the life history parameters
- Figure 2 shows the vectors
- Figure 4 shows the time series relative to reference points
- Figure ?? shows the performance statistics; points are
 - 1.
 - 2.
 - 3.
 - 4.
 5. Figure 5 shows the utility functions for the seven study stocks points area
- 1.
- 2.

- 188 3.
189 4.

190 **4. Discussion**

191 Fisheries management is often faced with multiple conflicting objectives,
192 and it is widely recognised that there is a need to incorporate these objectives
193 into management plans although some what impossible to perform such an
194 experiment on large scale fish stocks. Therefore performing computer sim-
195 ulations to develop robust management procedures is particularly valuable
196 in data poor situations where knowledge and data are limited, but also in
197 data rich situations as simulation testing an assessment procedure using a
198 model conditioned on the same assumptions is not necessarily a true test of
199 robustness. This paper has shown that the desired performance measure can
200 be met via tweaking of the management procedure by adjusting a particular
201 HCR a specific management objective can be achieved.

- 202 • Bullet point one
203 • Bullet point two

204 **5. Conclusions**

- 205 • Bullet point one
206 • Bullet point two

207 **6. References**

208 **References**

- 209 [1] E. Commission, Regulation (eu) no 1380/2013 of the european par-
210 liament and of the council of 11 december 2013 on the common fish-
211 eries policy, amending council regulations (ec) no 1954/2003 and (ec)
212 no 1224/2009, and repealing council regulations (ec) no 2371/2002 and
213 (ec) no 639/2004, and council decision 2004/585/ec (2013).
- 214 [2] N. Bentley, Data and time poverty in fisheries estimation: potential
215 approaches and solutions, ICES J. Mar. Sci. 72 (2015) 186–193.

- 216 [3] C. Costello, D. Ovando, R. Hilborn, S. D. Gaines, O. Deschenes, S. E.
217 Lester, Status and solutions for the world’s unassessed fisheries, *Science*
218 (2012) 1224768.
- 219 [4] T. J. Quinn, R. B. Deriso, Quantitative fish dynamics, Oxford University
220 Press, 1999.
- 221 [5] M. W. Pedersen, C. W. Berg, A stochastic surplus production model in
222 continuous time, *Fish and Fisheries* 18 (2017) 226–243.
- 223 [6] T. Gedamke, J. M. Hoenig, Estimating mortality from mean length
224 data in nonequilibrium situations, with application to the assessment
225 of goosfish, *Transactions of the American Fisheries Society* 135 (2006)
226 476–487.
- 227 [7] A. Hordyk, K. Ono, K. Sainsbury, N. Loneragan, J. Prince, Some
228 explorations of the life history ratios to describe length composition,
229 spawning-per-recruit, and the spawning potential ratio, *ICES Journal*
230 *of Marine Science* 72 (2014) 204–216.
- 231 [8] W. N. Probst, M. Kloppmann, G. Kraus, Indicator-based status assess-
232 ment of commercial fish species in the north sea according to the eu
233 marine strategy framework directive (msfd), *ICES Journal of Marine*
234 *Science* 70 (2013) 694–706.
- 235 [9] L. Kell, I. Mosqueira, P. Grosjean, J. Fromentin, D. Garcia, R. Hillary,
236 E. Jardim, S. Mardle, M. Pastoors, J. Poos, et al., FLR: an open-
237 source framework for the evaluation and development of management
238 strategies, *ICES J. Mar. Sci.* 64 (2007) 640.
- 239 [10] M. Sissenwine, J. Shepherd, An alternative perspective on recruitment
240 overfishing and biological reference points, *Can. J. Fish. Aquat. Sci.* 44
241 (1987) 913–918.
- 242 [11] C. G. Michielsens, M. K. McAllister, A bayesian hierarchical analysis of
243 stock recruit data: quantifying structural and parameter uncertainties,
244 *Can. J. Fish. Aquat. Sci.* 61 (2004) 1032–1047.
- 245 [12] J. Bergstra, Y. Bengio, Random search for hyper-parameter optimiza-
246 tion, *Journal of Machine Learning Research* 13 (2012) 281–305.




Figure 1:





Figure 2:



spp

Figure 3:




Figure 4:




Figure 5: