

Multiple solutions for 1-D quasilinear indefinite Schrödinger equations

Shibo Liu

Florida Institute of Technology

<http://lausb.github.io>

sliu@fit.edu

Contents

1	Introduction	3
2	Proof of Thm 1	8
3	Proof of Thm 2	14

1. Introduction

To find standing waves $\psi(t, x) = e^{-i\omega t} u(x)$ of QL Schrödinger equation

$$i\partial_t \psi = -\Delta \psi + U(x) - \psi \Delta(|\psi|^2) - \bar{g}(|\psi|^2) \psi \quad (t, x) \in \mathbb{R}^+ \times \mathbb{R}^N,$$

1. Introduction

To find standing waves $\psi(t, x) = e^{-i\omega t}u(x)$ of QL Schrödinger equation

$$i\partial_t\psi = -\Delta\psi + U(x) - \psi\Delta(|\psi|^2) - \bar{g}(|\psi|^2)\psi \quad (t, x) \in \mathbb{R}^+ \times \mathbb{R}^N,$$

we need to solve ($V = U - \omega$)

$$-\Delta u + V(x)u - u\Delta(u^2) = g(u), \quad u \in H^1(\mathbb{R}^N), \quad (1)$$

initiated by [Poppenberg et al. \(2002\)](#) for $N = 1$.

1. Introduction

To find standing waves $\psi(t, x) = e^{-i\omega t} u(x)$ of QL Schrödinger equation

$$i\partial_t \psi = -\Delta \psi + U(x) - \psi \Delta(|\psi|^2) - \bar{g}(|\psi|^2) \psi \quad (t, x) \in \mathbb{R}^+ \times \mathbb{R}^N,$$

we need to solve ($V = U - \omega$)

$$-\Delta u + V(x)u - u\Delta(u^2) = g(u), \quad u \in H^1(\mathbb{R}^N), \quad (1)$$

initiated by Poppenberg et al. (2002) for $N = 1$. Solutions are critical pts of

$$J(u) = \frac{1}{2} \int (1 + 2u^2) |\nabla u|^2 + \frac{1}{2} \int V(x)u^2 - \int G(u),$$

which is not well-defined ($J(u) = \infty$ for some $u \in H^1(\mathbb{R}^N)$) unless $N = 1$.

1. Introduction

To find standing waves $\psi(t, x) = e^{-i\omega t} u(x)$ of QL Schrödinger equation

$$i\partial_t \psi = -\Delta \psi + U(x) - \psi \Delta(|\psi|^2) - \bar{g}(|\psi|^2) \psi \quad (t, x) \in \mathbb{R}^+ \times \mathbb{R}^N,$$

we need to solve ($V = U - \omega$)

$$-\Delta u + V(x)u - u\Delta(u^2) = g(u), \quad u \in H^1(\mathbb{R}^N), \quad (1)$$

initiated by Poppenberg et al. (2002) for $N = 1$. Solutions are critical pts of

$$J(u) = \frac{1}{2} \int (1 + 2u^2) |\nabla u|^2 + \frac{1}{2} \int V(x)u^2 - \int G(u),$$

which is not well-defined ($J(u) = \infty$ for some $u \in H^1(\mathbb{R}^N)$) unless $N = 1$.

Liu et al. (2003); Colin & Jeanjean (2004) introduced a nonlinear transform $u = f(v)$ so that $\Phi : H^1(\mathbb{R}^N) \rightarrow \mathbb{R}$,

$$\Phi(v) = \frac{1}{2} \int (|\nabla v|^2 + V(x)f^2(v)) - \int G(f(v))$$

is well defined,

1. Introduction

To find standing waves $\psi(t, x) = e^{-i\omega t}u(x)$ of QL Schrödinger equation

$$i\partial_t \psi = -\Delta \psi + U(x) - \psi \Delta(|\psi|^2) - \bar{g}(|\psi|^2)\psi \quad (t, x) \in \mathbb{R}^+ \times \mathbb{R}^N,$$

we need to solve ($V = U - \omega$)

$$-\Delta u + V(x)u - u\Delta(u^2) = g(u), \quad u \in H^1(\mathbb{R}^N), \quad (1)$$

initiated by [Poppenberg et al. \(2002\)](#) for $N = 1$. Solutions are critical pts of

$$J(u) = \frac{1}{2} \int (1 + 2u^2) |\nabla u|^2 + \frac{1}{2} \int V(x)u^2 - \int G(u),$$

which is not well-defined ($J(u) = \infty$ for some $u \in H^1(\mathbb{R}^N)$) unless $N = 1$.

[Liu et al. \(2003\)](#); [Colin & Jeanjean \(2004\)](#) introduced a nonlinear transform $u = f(v)$ so that $\Phi : H^1(\mathbb{R}^N) \rightarrow \mathbb{R}$,

$$\Phi(v) = \frac{1}{2} \int (|\nabla v|^2 + V(x)f^2(v)) - \int G(f(v))$$

is well defined, if $\Phi'(v) = 0$ then $u = f(v)$ solves (1).

Then many results for $\inf V > 0$ appear.

1. Introduction

To find standing waves $\psi(t, x) = e^{-i\omega t} u(x)$ of QL Schrödinger equation

$$i\partial_t \psi = -\Delta \psi + U(x) - \psi \Delta(|\psi|^2) - \bar{g}(|\psi|^2) \psi \quad (t, x) \in \mathbb{R}^+ \times \mathbb{R}^N,$$

we need to solve ($V = U - \omega$)

$$-\Delta u + V(x)u - u\Delta(u^2) = g(u), \quad u \in H^1(\mathbb{R}^N), \quad (1)$$

initiated by [Poppenberg et al. \(2002\)](#) for $N = 1$. Solutions are critical pts of

$$J(u) = \frac{1}{2} \int (1 + 2u^2) |\nabla u|^2 + \frac{1}{2} \int V(x)u^2 - \int G(u),$$

which is not well-defined ($J(u) = \infty$ for some $u \in H^1(\mathbb{R}^N)$) unless $N = 1$.

[Liu et al. \(2003\)](#); [Colin & Jeanjean \(2004\)](#) introduced a nonlinear transform $u = f(v)$ so that $\Phi : H^1(\mathbb{R}^N) \rightarrow \mathbb{R}$,

$$\Phi(v) = \frac{1}{2} \int (|\nabla v|^2 + V(x)f^2(v)) - \int G(f(v))$$

is well defined, if $\Phi'(v) = 0$ then $u = f(v)$ solves (1).

Then many results for $\inf V > 0$ appear. Until [Shen & Han \(2015\)](#) for $N = 1$ and [Liu & Zhou \(2018\)](#) for $N \geq 1$, no results if $-\Delta + V$ is indefinite.

In Liu & Zhou (2018): $g(t) \approx |t|^{p-2} t$ for some $p \in (4, 2 \cdot 2^*)$.

In Liu & Zhou (2018): $g(t) \approx |t|^{p-2} t$ for some $p \in (4, 2 \cdot 2^*)$. It is crucial that

$$\lim_{|t| \rightarrow \infty} V(x) = +\infty, \quad (2)$$

so that the working space

$$E = \left\{ u \in H^1(\mathbb{R}^N) \left| \|u\| = \left(\int (|\nabla u|^2 + V u^2) \right)^{1/2} < \infty \right\} \hookrightarrow L^2(\mathbb{R}^N).$$

See also Silva & Silva (2019).

In Liu & Zhou (2018): $g(t) \approx |t|^{p-2} t$ for some $p \in (4, 2 \cdot 2^*)$. It is crucial that

$$\lim_{|t| \rightarrow \infty} V(x) = +\infty, \quad (2)$$

so that the working space

$$E = \left\{ u \in H^1(\mathbb{R}^N) \left| \|u\| = \left(\int (|\nabla u|^2 + V u^2) \right)^{1/2} < \infty \right\} \hookrightarrow L^2(\mathbb{R}^N).$$

See also Silva & Silva (2019).

If $N = 1$ we can study the case that $|V|_\infty < \infty$.

In Liu & Zhou (2018): $g(t) \approx |t|^{p-2} t$ for some $p \in (4, 2 \cdot 2^*)$. It is crucial that

$$\lim_{|t| \rightarrow \infty} V(x) = +\infty, \quad (2)$$

so that the working space

$$E = \left\{ u \in H^1(\mathbb{R}^N) \mid \|u\| = \left(\int (|\nabla u|^2 + V u^2) \right)^{1/2} < \infty \right\} \hookrightarrow L^2(\mathbb{R}^N).$$

See also Silva & Silva (2019).

If $N = 1$ we can study the case that $|V|_\infty < \infty$. We consider

$$-u'' + V(x)u - (u^2)''u = f(x, u), \quad u \in H^1(\mathbb{R}).$$

In Liu & Zhou (2018): $g(t) \approx |t|^{p-2} t$ for some $p \in (4, 2 \cdot 2^*)$. It is crucial that

$$\lim_{|t| \rightarrow \infty} V(x) = +\infty, \quad (2)$$

so that the working space

$$E = \left\{ u \in H^1(\mathbb{R}^N) \mid \|u\| = \left(\int (|\nabla u|^2 + V u^2) \right)^{1/2} < \infty \right\} \hookrightarrow L^2(\mathbb{R}^N).$$

See also Silva & Silva (2019).

If $N = 1$ we can study the case that $|V|_\infty < \infty$. We consider

$$-u'' + V(x)u - (u^2)''u = f(x, u), \quad u \in H^1(\mathbb{R}). \quad (3)$$

Assume $V \in C(\mathbb{R})$ is bounded from below. Let

$$\lambda_n = \inf_{X \in \mathcal{X}_n} \sup_{u \in X \setminus \{0\}} \frac{\int (\dot{u}^2 + V(x)u^2)}{\int u^2},$$

where \mathcal{X}_n is the collection of all n -dimensional subspaces of $C_0^\infty(\mathbb{R})$.

In Liu & Zhou (2018): $g(t) \approx |t|^{p-2} t$ for some $p \in (4, 2 \cdot 2^*)$. It is crucial that

$$\lim_{|t| \rightarrow \infty} V(x) = +\infty, \quad (2)$$

so that the working space

$$E = \left\{ u \in H^1(\mathbb{R}^N) \left| \|u\| = \left(\int (|\nabla u|^2 + V u^2) \right)^{1/2} < \infty \right\} \hookrightarrow L^2(\mathbb{R}^N).$$

See also Silva & Silva (2019).

If $N = 1$ we can study the case that $|V|_\infty < \infty$. We consider

$$-u'' + V(x)u - (u^2)''u = f(x, u), \quad u \in H^1(\mathbb{R}). \quad (3)$$

Assume $V \in C(\mathbb{R})$ is bounded from below. Let

$$\lambda_n = \inf_{X \in \mathcal{X}_n} \sup_{u \in X \setminus \{0\}} \frac{\int (\dot{u}^2 + V(x)u^2)}{\int u^2},$$

where \mathcal{X}_n is the collection of all n -dimensional subspaces of $C_0^\infty(\mathbb{R})$. Assume

$$\lambda_n \rightarrow \lambda_\infty < \infty,$$

then λ_∞ is the bottom of the essential spectrum of $S = -\frac{d^2}{dx^2} + V$

In [Liu & Zhou \(2018\)](#): $g(t) \approx |t|^{p-2} t$ for some $p \in (4, 2 \cdot 2^*)$. It is crucial that

$$\lim_{|t| \rightarrow \infty} V(x) = +\infty, \quad (2)$$

so that the working space

$$E = \left\{ u \in H^1(\mathbb{R}^N) \mid \|u\| = \left(\int (|\nabla u|^2 + V u^2) \right)^{1/2} < \infty \right\} \hookrightarrow L^2(\mathbb{R}^N).$$

See also [Silva & Silva \(2019\)](#).

If $N = 1$ we can study the case that $|V|_\infty < \infty$. We consider

$$-u'' + V(x)u - (u^2)''u = f(x, u), \quad u \in H^1(\mathbb{R}). \quad (3)$$

Assume $V \in C(\mathbb{R})$ is bounded from below. Let

$$\lambda_n = \inf_{X \in \mathcal{X}_n} \sup_{u \in X \setminus \{0\}} \frac{\int (\dot{u}^2 + V(x)u^2)}{\int u^2},$$

where \mathcal{X}_n is the collection of all n -dimensional subspaces of $C_0^\infty(\mathbb{R})$. Assume

$$\lambda_n \rightarrow \lambda_\infty < \infty,$$

then λ_∞ is the bottom of the essential spectrum of $S = -\frac{d^2}{dx^2} + V$ and $\lambda_n < \lambda_\infty$ implies that λ_n is an eigenvalue of S of finite multiplicity.

We assume

(V₁) $V \in C(\mathbb{R})$ is such that $0 \in (\lambda_k, \lambda_{k+1})$ for some $k \in \mathbb{N}$.

We assume

(V₁) $V \in C(\mathbb{R})$ is such that $0 \in (\lambda_k, \lambda_{k+1})$ for some $k \in \mathbb{N}$.

(f₁) $f \in C(\mathbb{R} \times \mathbb{R})$ and there are constants $p > 2$ and $c > 0$ such that

$$|f(x, t)| \leq c (1 + |t|^{p-1}) \quad \text{for } (x, t) \in \mathbb{R} \times \mathbb{R}.$$

We assume

(V₁) $V \in C(\mathbb{R})$ is such that $0 \in (\lambda_k, \lambda_{k+1})$ for some $k \in \mathbb{N}$.

(f₁) $f \in C(\mathbb{R} \times \mathbb{R})$ and there are constants $p > 2$ and $c > 0$ such that

$$|f(x, t)| \leq c (1 + |t|^{p-1}) \quad \text{for } (x, t) \in \mathbb{R} \times \mathbb{R}.$$

(f₂) $f(x, t) = o(t)$ as $t \rightarrow 0$ uniformly in $x \in \mathbb{R}$.

We assume

(V₁) $V \in C(\mathbb{R})$ is such that $0 \in (\lambda_k, \lambda_{k+1})$ for some $k \in \mathbb{N}$.

(f₁) $f \in C(\mathbb{R} \times \mathbb{R})$ and there are constants $p > 2$ and $c > 0$ such that

$$|f(x, t)| \leq c (1 + |t|^{p-1}) \quad \text{for } (x, t) \in \mathbb{R} \times \mathbb{R}.$$

(f₂) $f(x, t) = o(t)$ as $t \rightarrow 0$ uniformly in $x \in \mathbb{R}$.

(f₃^{*}) There exists $h \in (0, \lambda_\infty)$ such that $tf(x, t) \leq ht^2$ for all $(x, t) \in \mathbb{R} \times \mathbb{R}$.

We assume

(V₁) $V \in C(\mathbb{R})$ is such that $0 \in (\lambda_k, \lambda_{k+1})$ for some $k \in \mathbb{N}$.

(f₁) $f \in C(\mathbb{R} \times \mathbb{R})$ and there are constants $p > 2$ and $c > 0$ such that

$$|f(x, t)| \leq c(1 + |t|^{p-1}) \quad \text{for } (x, t) \in \mathbb{R} \times \mathbb{R}.$$

(f₂) $f(x, t) = o(t)$ as $t \rightarrow 0$ uniformly in $x \in \mathbb{R}$.

(f₃^{*}) There exists $h \in (0, \lambda_\infty)$ such that $tf(x, t) \leq ht^2$ for all $(x, t) \in \mathbb{R} \times \mathbb{R}$.

Increasing h a little bit we may assume that h in (f₃^{*}) is not an eigenvalue.

Thm 1. Suppose (V₁), (f₁), (f₂) and (f₃^{*}) are satisfied, then (3) has at least two nontrivial solutions.

We assume

(V₁) $V \in C(\mathbb{R})$ is such that $0 \in (\lambda_k, \lambda_{k+1})$ for some $k \in \mathbb{N}$.

(f₁) $f \in C(\mathbb{R} \times \mathbb{R})$ and there are constants $p > 2$ and $c > 0$ such that

$$|f(x, t)| \leq c(1 + |t|^{p-1}) \quad \text{for } (x, t) \in \mathbb{R} \times \mathbb{R}.$$

(f₂) $f(x, t) = o(t)$ as $t \rightarrow 0$ uniformly in $x \in \mathbb{R}$.

(f₃^{*}) There exists $h \in (0, \lambda_\infty)$ such that $tf(x, t) \leq ht^2$ for all $(x, t) \in \mathbb{R} \times \mathbb{R}$.

Increasing h a little bit we may assume that h in (f₃^{*}) is not an eigenvalue.

Thm 1. Suppose (V₁), (f₁), (f₂) and (f₃^{*}) are satisfied, then (3) has at least two nontrivial solutions. If in addition $f(x, \cdot)$ is odd for all $x \in \mathbb{R}$, then (3) has k pairs of nontrivial solutions.

We assume

(V₁) $V \in C(\mathbb{R})$ is such that $0 \in (\lambda_k, \lambda_{k+1})$ for some $k \in \mathbb{N}$.

(f₁) $f \in C(\mathbb{R} \times \mathbb{R})$ and there are constants $p > 2$ and $c > 0$ such that

$$|f(x, t)| \leq c(1 + |t|^{p-1}) \quad \text{for } (x, t) \in \mathbb{R} \times \mathbb{R}.$$

(f₂) $f(x, t) = o(t)$ as $t \rightarrow 0$ uniformly in $x \in \mathbb{R}$.

(f₃^{*}) There exists $h \in (0, \lambda_\infty)$ such that $tf(x, t) \leq ht^2$ for all $(x, t) \in \mathbb{R} \times \mathbb{R}$.

Increasing h a little bit we may assume that h in (f₃^{*}) is not an eigenvalue.

Thm 1. Suppose (V₁), (f₁), (f₂) and (f₃^{*}) are satisfied, then (3) has at least two nontrivial solutions. If in addition $f(x, \cdot)$ is odd for all $x \in \mathbb{R}$, then (3) has k pairs of nontrivial solutions.

Rem 1. Under (V₁), (f₁), (f₂) and the following condition weaker than (f₃^{*}):

(f₃) There exists $h \in (0, \lambda_\infty)$ such that $F(x, t) \leq \frac{1}{2}ht^2$ for all $(x, t) \in \mathbb{R} \times \mathbb{R}$,

a nontrivial solution is obtained by Wang & Yang (2015).

Thm1 is motivated by [Chen & Wang \(2014\)](#) who obtained similar results for

$$\begin{cases} -\Delta u + V(x)u + \phi u = f(x, u) & \text{in } \mathbb{R}^3, \\ -\Delta \phi = u^2 & \text{in } \mathbb{R}^3. \end{cases}$$

Thm1 is motivated by Chen & Wang (2014) who obtained similar results for

$$\begin{cases} -\Delta u + V(x)u + \phi u = f(x, u) & \text{in } \mathbb{R}^3, \\ -\Delta \phi = u^2 & \text{in } \mathbb{R}^3. \end{cases} \quad (4)$$

Unlike our case, weak limits of (PS) sequences of

$$\Phi(u) = \frac{1}{2} \int (|\nabla u|^2 + V(x)u^2) + \frac{1}{4} \int \phi_u u^2 - \int F(x, u), \quad \phi_u(x) = \frac{1}{4\pi} \int \frac{u^2(y)}{|x-y|} dy$$

are critical points of Φ .

Thm1 is motivated by Chen & Wang (2014) who obtained similar results for

$$\begin{cases} -\Delta u + V(x)u + \phi u = f(x, u) & \text{in } \mathbb{R}^3, \\ -\Delta \phi = u^2 & \text{in } \mathbb{R}^3. \end{cases} \quad (4)$$

Unlike our case, weak limits of (PS) sequences of

$$\Phi(u) = \frac{1}{2} \int (|\nabla u|^2 + V(x)u^2) + \frac{1}{4} \int \phi_u u^2 - \int F(x, u), \quad \phi_u(x) = \frac{1}{4\pi} \int \frac{u^2(y)}{|x-y|} dy$$

are critical points of Φ .

Lacking this property is the reason that Wang & Yang (2015) could not get multiplicity results for (3) as in Thm1.

Thm1 is motivated by Chen & Wang (2014) who obtained similar results for

$$\begin{cases} -\Delta u + V(x)u + \phi u = f(x, u) & \text{in } \mathbb{R}^3, \\ -\Delta \phi = u^2 & \text{in } \mathbb{R}^3. \end{cases} \quad (4)$$

Unlike our case, weak limits of (PS) sequences of

$$\Phi(u) = \frac{1}{2} \int (|\nabla u|^2 + V(x)u^2) + \frac{1}{4} \int \phi_u u^2 - \int F(x, u), \quad \phi_u(x) = \frac{1}{4\pi} \int \frac{u^2(y)}{|x-y|} dy$$

are critical points of Φ .

Lacking this property is the reason that Wang & Yang (2015) could not get multiplicity results for (3) as in Thm1.

Rem 2. Initiated by Benci & Fortunato (1998), (4) attracted great interest for definite case in which $u = 0$ is loc min of Φ .

Thm1 is motivated by Chen & Wang (2014) who obtained similar results for

$$\begin{cases} -\Delta u + V(x)u + \phi u = f(x, u) & \text{in } \mathbb{R}^3, \\ -\Delta \phi = u^2 & \text{in } \mathbb{R}^3. \end{cases} \quad (4)$$

Unlike our case, weak limits of (PS) sequences of

$$\Phi(u) = \frac{1}{2} \int (|\nabla u|^2 + V(x)u^2) + \frac{1}{4} \int \phi_u u^2 - \int F(x, u), \quad \phi_u(x) = \frac{1}{4\pi} \int \frac{u^2(y)}{|x-y|} dy$$

are critical points of Φ .

Lacking this property is the reason that Wang & Yang (2015) could not get multiplicity results for (3) as in Thm1.

Rem 2. Initiated by Benci & Fortunato (1998), (4) attracted great interest for definite case in which $u = 0$ is loc min of Φ .

The first work on the indefinite case that $u = 0$ is saddle pt is due to Chen & Liu (2015).

Thm1 is motivated by Chen & Wang (2014) who obtained similar results for

$$\begin{cases} -\Delta u + V(x)u + \phi u = f(x, u) & \text{in } \mathbb{R}^3, \\ -\Delta \phi = u^2 & \text{in } \mathbb{R}^3. \end{cases} \quad (4)$$

Unlike our case, weak limits of (PS) sequences of

$$\Phi(u) = \frac{1}{2} \int (|\nabla u|^2 + V(x)u^2) + \frac{1}{4} \int \phi_u u^2 - \int F(x, u), \quad \phi_u(x) = \frac{1}{4\pi} \int \frac{u^2(y)}{|x-y|} dy$$

are critical points of Φ .

Lacking this property is the reason that Wang & Yang (2015) could not get multiplicity results for (3) as in Thm1.

Rem 2. Initiated by Benci & Fortunato (1998), (4) attracted great interest for definite case in which $u = 0$ is loc min of Φ .

The first work on the indefinite case that $u = 0$ is saddle pt is due to Chen & Liu (2015).

Motivated by Liu & Wu (2017) on indefinite problem (4), we consider (3) when $f(x, \cdot)$ is 4-superlinear:

$$\lim_{|t| \rightarrow \infty} \frac{F(x, t)}{t^4} = +\infty \quad \text{a.e. } x \in \mathbb{R}, \quad \text{where } F(x, t) = \int_0^t f(x, \cdot).$$

(f₄) $0 < 4F(x, t) \leq tf(x, t)$ for all $(x, t) \in \mathbb{R} \times \mathbb{R}$,

$$\lim_{|t| \rightarrow \infty} \frac{F(x, t)}{t^4} = +\infty \quad \text{a.e. } x \in \mathbb{R}.$$

(f₄) $0 < 4F(x, t) \leq tf(x, t)$ for all $(x, t) \in \mathbb{R} \times \mathbb{R}$,

$$\lim_{|t| \rightarrow \infty} \frac{F(x, t)}{t^4} = +\infty \quad \text{a.e. } x \in \mathbb{R}. \quad (5)$$

(f₅) if $u_n \rightarrow u$ in $H^1(\mathbb{R})$, then $\overline{\lim}_{n \rightarrow \infty} \int f(x, u_n)(u_n - u) \leq 0$.

(f₄) $0 < 4F(x, t) \leq tf(x, t)$ for all $(x, t) \in \mathbb{R} \times \mathbb{R}$,

$$\lim_{|t| \rightarrow \infty} \frac{F(x, t)}{t^4} = +\infty \quad \text{a.e. } x \in \mathbb{R}. \quad (5)$$

(f₅) if $u_n \rightarrow u$ in $H^1(\mathbb{R})$, then $\overline{\lim}_{n \rightarrow \infty} \int f(x, u_n)(u_n - u) \leq 0$.

Thm 2. Suppose (V_1) , (f_1) , (f_2) , (f_4) and (f_5) are satisfied, then (3) has at least one nontrivial solutions.

(f₄) $0 < 4F(x, t) \leq tf(x, t)$ for all $(x, t) \in \mathbb{R} \times \mathbb{R}$,

$$\lim_{|t| \rightarrow \infty} \frac{F(x, t)}{t^4} = +\infty \quad \text{a.e. } x \in \mathbb{R}. \quad (5)$$

(f₅) if $u_n \rightarrow u$ in $H^1(\mathbb{R})$, then $\overline{\lim}_{n \rightarrow \infty} \int f(x, u_n)(u_n - u) \leq 0$.

Thm 2. Suppose (V_1) , (f_1) , (f_2) , (f_4) and (f_5) are satisfied, then (3) has at least one nontrivial solutions. If in addition $f(x, \cdot)$ is odd for $x \in \mathbb{R}$, then (3) has a sequence of solutions $\{u_n\}$ such that $\Phi(u_n) \rightarrow +\infty$.

(f₄) $0 < 4F(x, t) \leq tf(x, t)$ for all $(x, t) \in \mathbb{R} \times \mathbb{R}$,

$$\lim_{|t| \rightarrow \infty} \frac{F(x, t)}{t^4} = +\infty \quad \text{a.e. } x \in \mathbb{R}. \quad (5)$$

(f₅) if $u_n \rightarrow u$ in $H^1(\mathbb{R})$, then $\overline{\lim}_{n \rightarrow \infty} \int f(x, u_n)(u_n - u) \leq 0$.

Thm 2. Suppose (V_1) , (f_1) , (f_2) , (f_4) and (f_5) are satisfied, then (3) has at least one nontrivial solutions. If in addition $f(x, \cdot)$ is odd for $x \in \mathbb{R}$, then (3) has a sequence of solutions $\{u_n\}$ such that $\Phi(u_n) \rightarrow +\infty$.

Rem 3. Condition (f_5) holds in e.g. one of the following:

(1) for $\forall r > 0$, $\lim_{|x| \rightarrow \infty} \sup_{0 < |t| \leq r} \left| \frac{f(x, t)}{t} \right| = 0$, Bartsch et al. (2004)

(f_4) $0 < 4F(x, t) \leq tf(x, t)$ for all $(x, t) \in \mathbb{R} \times \mathbb{R}$,

$$\lim_{|t| \rightarrow \infty} \frac{F(x, t)}{t^4} = +\infty \quad \text{a.e. } x \in \mathbb{R}. \quad (5)$$

(f_5) if $u_n \rightarrow u$ in $H^1(\mathbb{R})$, then $\overline{\lim}_{n \rightarrow \infty} \int f(x, u_n)(u_n - u) \leq 0$.

Thm 2. Suppose (V_1) , (f_1) , (f_2) , (f_4) and (f_5) are satisfied, then (3) has at least one nontrivial solutions. If in addition $f(x, \cdot)$ is odd for $x \in \mathbb{R}$, then (3) has a sequence of solutions $\{u_n\}$ such that $\Phi(u_n) \rightarrow +\infty$.

Rem 3. Condition (f_5) holds in e.g. one of the following:

(1) for $\forall r > 0$, $\lim_{|x| \rightarrow \infty} \sup_{0 < |t| \leq r} \left| \frac{f(x, t)}{t} \right| = 0$, Bartsch et al. (2004)

Example: $f(x, t) = a(x)|t|^{p-2}t$, $\lim_{|x| \rightarrow \infty} a(x) = 0$.

(f₄) $0 < 4F(x, t) \leq tf(x, t)$ for all $(x, t) \in \mathbb{R} \times \mathbb{R}$,

$$\lim_{|t| \rightarrow \infty} \frac{F(x, t)}{t^4} = +\infty \quad \text{a.e. } x \in \mathbb{R}. \quad (5)$$

(f₅) if $u_n \rightarrow u$ in $H^1(\mathbb{R})$, then $\overline{\lim}_{n \rightarrow \infty} \int f(x, u_n)(u_n - u) \leq 0$.

Thm 2. Suppose (V_1) , (f_1) , (f_2) , (f_4) and (f_5) are satisfied, then (3) has at least one nontrivial solutions. If in addition $f(x, \cdot)$ is odd for $x \in \mathbb{R}$, then (3) has a sequence of solutions $\{u_n\}$ such that $\Phi(u_n) \rightarrow +\infty$.

Rem 3. Condition (f_5) holds in e.g. one of the following:

(1) for $\forall r > 0$, $\lim_{|x| \rightarrow \infty} \sup_{0 < |t| \leq r} \left| \frac{f(x, t)}{t} \right| = 0$, Bartsch et al. (2004)

Example: $f(x, t) = a(x)|t|^{p-2}t$, $\lim_{|x| \rightarrow \infty} a(x) = 0$.

(2) $|f(x, t)| \leq \alpha_+(x)|t|^{p_+-1} + \alpha_-(x)|t|^{p_--1}$, $\alpha_{\pm} \in L^{q_{\pm}}(\mathbb{R})$ for some $q_{\pm} > 1$.

2. Proof of Thm 1

We denote $X = H^1(\mathbb{R})$. By [Poppenberg et al. \(2002\)](#), $N : X \rightarrow \mathbb{R}$ given by

$$N(u) = \int \dot{u}^2 u^2$$

is of class C^1 ,

2. Proof of Thm 1

We denote $X = H^1(\mathbb{R})$. By Poppenberg et al. (2002), $N : X \rightarrow \mathbb{R}$ given by

$$N(u) = \int \dot{u}^2 u^2$$

is of class C^1 ,

$$\langle N'(u), v \rangle = 2 \int (\dot{u}^2 uv + u^2 \dot{u} \dot{v}).$$

2. Proof of Thm 1

We denote $X = H^1(\mathbb{R})$. By Poppenberg et al. (2002), $N : X \rightarrow \mathbb{R}$ given by

$$N(u) = \int \dot{u}^2 u^2$$

is of class C^1 ,

$$\langle N'(u), v \rangle = 2 \int (\dot{u}^2 u v + u^2 \dot{u} \dot{v}).$$

Therefore, $\Phi : X \rightarrow \mathbb{R}$

$$\Phi(u) = \frac{1}{2} \int (\dot{u}^2 + V(x)u^2) + \int \dot{u}^2 u^2 - \int F(x, u)$$

2. Proof of Thm 1

We denote $X = H^1(\mathbb{R})$. By Poppenberg et al. (2002), $N : X \rightarrow \mathbb{R}$ given by

$$N(u) = \int \dot{u}^2 u^2$$

is of class C^1 ,

$$\langle N'(u), v \rangle = 2 \int (\dot{u}^2 u v + u^2 \dot{u} \dot{v}).$$

Therefore, $\Phi : X \rightarrow \mathbb{R}$

$$\begin{aligned} \Phi(u) &= \frac{1}{2} \int (\dot{u}^2 + V(x)u^2) + \int \dot{u}^2 u^2 - \int F(x, u) \\ &= \frac{1}{2} \int (\dot{u}^2 + V(x)u^2) + N(u) - \int F(x, u), \end{aligned}$$

is of class C^1 as well,

2. Proof of Thm 1

We denote $X = H^1(\mathbb{R})$. By Poppenberg et al. (2002), $N : X \rightarrow \mathbb{R}$ given by

$$N(u) = \int \dot{u}^2 u^2$$

is of class C^1 ,

$$\langle N'(u), v \rangle = 2 \int (\dot{u}^2 u v + u^2 \dot{u} \dot{v}).$$

Therefore, $\Phi : X \rightarrow \mathbb{R}$

$$\begin{aligned} \Phi(u) &= \frac{1}{2} \int (\dot{u}^2 + V(x)u^2) + \int \dot{u}^2 u^2 - \int F(x, u) \\ &= \frac{1}{2} \int (\dot{u}^2 + V(x)u^2) + N(u) - \int F(x, u), \end{aligned}$$

is of class C^1 as well, with derivative

$$\langle \Phi'(u), v \rangle = \int (\dot{u} \dot{v} + V(x)u v) + \langle N'(u), v \rangle - \int f(x, u) v.$$

2. Proof of Thm 1

We denote $X = H^1(\mathbb{R})$. By Poppenberg et al. (2002), $N : X \rightarrow \mathbb{R}$ given by

$$N(u) = \int \dot{u}^2 u^2$$

is of class C^1 ,

$$\langle N'(u), v \rangle = 2 \int (\dot{u}^2 u v + u^2 \dot{u} \dot{v}).$$

Therefore, $\Phi : X \rightarrow \mathbb{R}$

$$\begin{aligned} \Phi(u) &= \frac{1}{2} \int (\dot{u}^2 + V(x)u^2) + \int \dot{u}^2 u^2 - \int F(x, u) \\ &= \frac{1}{2} \int (\dot{u}^2 + V(x)u^2) + N(u) - \int F(x, u), \end{aligned}$$

is of class C^1 as well, with derivative

$$\langle \Phi'(u), v \rangle = \int (\dot{u} \dot{v} + V(x)u v) + \langle N'(u), v \rangle - \int f(x, u) v.$$

Critical points of Φ are weak solutions of the problem (3).

Lem 1. If $u_n \rightarrow u$ in X ,

Lem 1. If $u_n \rightarrow u$ in X , then

$$\overline{\lim}_{n \rightarrow \infty} (\langle N'(u_n), u \rangle - 4N(u_n)) \leq 0.$$

Lem 1. If $u_n \rightarrow u$ in X , then

$$\overline{\lim}_{n \rightarrow \infty} (\langle N'(u_n), u \rangle - 4N(u_n)) \leq 0. \quad (6)$$

Pf. The inequality (6) is a consequence of

$$\frac{1}{2} \langle N'(u_n), u \rangle = \int (\dot{u}_n^2 u_n u + u_n^2 \dot{u}_n \dot{u})$$

Lem 1. If $u_n \rightarrow u$ in X , then

$$\overline{\lim}_{n \rightarrow \infty} (\langle N'(u_n), u \rangle - 4N(u_n)) \leq 0. \quad (6)$$

Pf. The inequality (6) is a consequence of

$$\begin{aligned} \frac{1}{2} \langle N'(u_n), u \rangle &= \int (\dot{u}_n^2 u_n u + u_n^2 \dot{u}_n \dot{u}) \\ &\leq 2 \int \dot{u}_n^2 u_n^2 + o(1) = 2N(u_n) + o(1), \end{aligned}$$

which has been proven in ([Chen, 2014](#), Eqn (3.16)).

Lem 1. If $u_n \rightarrow u$ in X , then

$$\overline{\lim}_{n \rightarrow \infty} (\langle N'(u_n), u \rangle - 4N(u_n)) \leq 0. \quad (6)$$

Pf. The inequality (6) is a consequence of

$$\begin{aligned} \frac{1}{2} \langle N'(u_n), u \rangle &= \int (\dot{u}_n^2 u_n u + u_n^2 \dot{u}_n \dot{u}) \\ &\leq 2 \int \dot{u}_n^2 u_n^2 + o(1) = 2N(u_n) + o(1), \end{aligned}$$

which has been proven in (Chen, 2014, Eqn (3.16)).

Lem 2. Suppose $g \in C(\mathbb{R}^N \times \mathbb{R})$, $|g(x, t)| \leq \Lambda|t|$ for $(x, t) \in \mathbb{R}^N \times \mathbb{R}$.

Lem 1. If $u_n \rightarrow u$ in X , then

$$\overline{\lim}_{n \rightarrow \infty} (\langle N'(u_n), u \rangle - 4N(u_n)) \leq 0. \quad (6)$$

Pf. The inequality (6) is a consequence of

$$\begin{aligned} \frac{1}{2} \langle N'(u_n), u \rangle &= \int (\dot{u}_n^2 u_n u + u_n^2 \dot{u}_n \dot{u}) \\ &\leq 2 \int \dot{u}_n^2 u_n^2 + o(1) = 2N(u_n) + o(1), \end{aligned}$$

which has been proven in (Chen, 2014, Eqn (3.16)).

Lem 2. Suppose $g \in C(\mathbb{R}^N \times \mathbb{R})$, $|g(x, t)| \leq \Lambda|t|$ for $(x, t) \in \mathbb{R}^N \times \mathbb{R}$. If $u_n \rightarrow u$ in $H^1(\mathbb{R}^N)$, then for all $\phi \in H^1(\mathbb{R}^N)$ we have

$$\int g(x, u_n) \phi \rightarrow \int g(x, u) \phi.$$

Lem 1. If $u_n \rightarrow u$ in X , then

$$\overline{\lim}_{n \rightarrow \infty} (\langle N'(u_n), u \rangle - 4N(u_n)) \leq 0. \quad (6)$$

Pf. The inequality (6) is a consequence of

$$\begin{aligned} \frac{1}{2} \langle N'(u_n), u \rangle &= \int (\dot{u}_n^2 u_n u + u_n^2 \dot{u}_n \dot{u}) \\ &\leq 2 \int \dot{u}_n^2 u_n^2 + o(1) = 2N(u_n) + o(1), \end{aligned}$$

which has been proven in (Chen, 2014, Eqn (3.16)).

Lem 2. Suppose $g \in C(\mathbb{R}^N \times \mathbb{R})$, $|g(x, t)| \leq \Lambda|t|$ for $(x, t) \in \mathbb{R}^N \times \mathbb{R}$. If $u_n \rightarrow u$ in $H^1(\mathbb{R}^N)$, then for all $\phi \in H^1(\mathbb{R}^N)$ we have

$$\int g(x, u_n) \phi \rightarrow \int g(x, u) \phi. \quad (7)$$

Rem 4. By Brézis & Lieb (1983) we get

$$g(u_n) \rightarrow g(u) \quad \text{in } L^2(\mathbb{R}^N),$$

which implies (7).

Pf (Without using B-L). Since g is of linear growth and $\{u_n\}$ is bounded,
$$\beta := \sup_n |g(x, u_n) - g(x, u)|_2 < \infty.$$

Pf (Without using B-L). Since g is of linear growth and $\{u_n\}$ is bounded,
$$\beta := \sup_n |g(x, u_n) - g(x, u)|_2 < \infty.$$

Given $\varepsilon > 0$, choose $R > 0$ such that

$$\int_{|x| \geq R} \phi^2 \leq \varepsilon^2.$$

Pf (Without using B-L). Since g is of linear growth and $\{u_n\}$ is bounded,
$$\beta := \sup_n |g(x, u_n) - g(x, u)|_2 < \infty.$$

Given $\varepsilon > 0$, choose $R > 0$ such that

$$\int_{|x| \geq R} \phi^2 \leq \varepsilon^2.$$

Using Hölder inequality we have

$$\left| \int g(x, u_n) \phi - \int g(x, u) \phi \right|$$

Pf (Without using B-L). Since g is of linear growth and $\{u_n\}$ is bounded,
$$\beta := \sup_n |g(x, u_n) - g(x, u)|_2 < \infty.$$

Given $\varepsilon > 0$, choose $R > 0$ such that

$$\int_{|x| \geq R} \phi^2 \leq \varepsilon^2.$$

Using Hölder inequality we have

$$\begin{aligned} & \left| \int g(x, u_n) \phi - \int g(x, u) \phi \right| \\ & \leq \int_{|x| \geq R} |g(x, u_n) - g(x, u)| |\phi| + \int_{|x| < R} |g(x, u_n) - g(x, u)| |\phi| \end{aligned}$$

Pf (Without using B-L). Since g is of linear growth and $\{u_n\}$ is bounded,
$$\beta := \sup_n |g(x, u_n) - g(x, u)|_2 < \infty.$$

Given $\varepsilon > 0$, choose $R > 0$ such that

$$\int_{|x| \geq R} \phi^2 \leq \varepsilon^2.$$

Using Hölder inequality we have

$$\begin{aligned} & \left| \int g(x, u_n) \phi - \int g(x, u) \phi \right| \\ & \leq \int_{|x| \geq R} |g(x, u_n) - g(x, u)| |\phi| + \int_{|x| < R} |g(x, u_n) - g(x, u)| |\phi| \\ & \leq |g(x, u_n) - g(x, u)|_2 \left(\int_{|x| \geq R} \phi^2 \right)^{1/2} \end{aligned}$$

Pf (Without using B-L). Since g is of linear growth and $\{u_n\}$ is bounded,
$$\beta := \sup_n |g(x, u_n) - g(x, u)|_2 < \infty.$$

Given $\varepsilon > 0$, choose $R > 0$ such that

$$\int_{|x| \geq R} \phi^2 \leq \varepsilon^2.$$

Using Hölder inequality we have

$$\begin{aligned} & \left| \int g(x, u_n) \phi - \int g(x, u) \phi \right| \\ & \leq \int_{|x| \geq R} |g(x, u_n) - g(x, u)| |\phi| + \int_{|x| < R} |g(x, u_n) - g(x, u)| |\phi| \\ & \leq |g(x, u_n) - g(x, u)|_2 \left(\int_{|x| \geq R} \phi^2 \right)^{1/2} + \int_{|x| < R} |g(x, u_n) - g(x, u)| |\phi| \end{aligned}$$

Pf (Without using B-L). Since g is of linear growth and $\{u_n\}$ is bounded,
$$\beta := \sup_n |g(x, u_n) - g(x, u)|_2 < \infty.$$

Given $\varepsilon > 0$, choose $R > 0$ such that

$$\int_{|x| \geq R} \phi^2 \leq \varepsilon^2.$$

Using Hölder inequality we have

$$\begin{aligned} & \left| \int g(x, u_n) \phi - \int g(x, u) \phi \right| \\ & \leq \int_{|x| \geq R} |g(x, u_n) - g(x, u)| |\phi| + \int_{|x| < R} |g(x, u_n) - g(x, u)| |\phi| \\ & \leq |g(x, u_n) - g(x, u)|_2 \left(\int_{|x| \geq R} \phi^2 \right)^{1/2} + \int_{|x| < R} |g(x, u_n) - g(x, u)| |\phi| \\ & \leq \beta \varepsilon + \int_{|x| < R} |g(x, u_n) - g(x, u)| |\phi|. \end{aligned}$$

Pf (Without using B-L). Since g is of linear growth and $\{u_n\}$ is bounded,
$$\beta := \sup_n |g(x, u_n) - g(x, u)|_2 < \infty.$$

Given $\varepsilon > 0$, choose $R > 0$ such that

$$\int_{|x| \geq R} \phi^2 \leq \varepsilon^2.$$

Using Hölder inequality we have

$$\begin{aligned} & \left| \int g(x, u_n) \phi - \int g(x, u) \phi \right| \\ & \leq \int_{|x| \geq R} |g(x, u_n) - g(x, u)| |\phi| + \int_{|x| < R} |g(x, u_n) - g(x, u)| |\phi| \\ & \leq |g(x, u_n) - g(x, u)|_2 \left(\int_{|x| \geq R} \phi^2 \right)^{1/2} + \int_{|x| < R} |g(x, u_n) - g(x, u)| |\phi| \\ & \leq \beta \varepsilon + \int_{|x| < R} |g(x, u_n) - g(x, u)| |\phi|. \end{aligned}$$

Since $H^1(\mathbb{R}^N) \hookrightarrow L^2_{\text{loc}}(B_R)$ we deduce $g(x, u_n) \rightarrow g(x, u)$ in $L^2(B_R)$.

Pf (Without using B-L). Since g is of linear growth and $\{u_n\}$ is bounded,

$$\beta := \sup_n |g(x, u_n) - g(x, u)|_2 < \infty.$$

Given $\varepsilon > 0$, choose $R > 0$ such that

$$\int_{|x| \geq R} \phi^2 \leq \varepsilon^2.$$

Using Hölder inequality we have

$$\begin{aligned} & \left| \int g(x, u_n) \phi - \int g(x, u) \phi \right| \\ & \leq \int_{|x| \geq R} |g(x, u_n) - g(x, u)| |\phi| + \int_{|x| < R} |g(x, u_n) - g(x, u)| |\phi| \\ & \leq |g(x, u_n) - g(x, u)|_2 \left(\int_{|x| \geq R} \phi^2 \right)^{1/2} + \int_{|x| < R} |g(x, u_n) - g(x, u)| |\phi| \\ & \leq \beta \varepsilon + \int_{|x| < R} |g(x, u_n) - g(x, u)| |\phi|. \end{aligned}$$

Since $H^1(\mathbb{R}^N) \hookrightarrow L^2_{\text{loc}}(B_R)$ we deduce $g(x, u_n) \rightarrow g(x, u)$ in $L^2(B_R)$. Hence

$$\overline{\lim}_{n \rightarrow \infty} \left| \int g(x, u_n) \phi - \int g(x, u) \phi \right| \leq \beta \varepsilon.$$

Let E^- be the negative subspace of $S - h$ and $E^+ = (E^-)^\perp$.

Let E^- be the negative subspace of $S - h$ and $E^+ = (E^-)^\perp$. Then $\dim E^- < \infty$ and there is an equivalent norm $\|\cdot\|$ on $X = E^- \oplus E^+$

Let E^- be the negative subspace of $S - h$ and $E^+ = (E^-)^\perp$. Then $\dim E^- < \infty$ and there is an equivalent norm $\|\cdot\|$ on $X = E^- \oplus E^+$ such that

$$\int (\dot{u}^2 + V(x)u^2 - hu^2) = \|u^+\|^2 - \|u^-\|^2,$$

Let E^- be the negative subspace of $S - h$ and $E^+ = (E^-)^\perp$. Then $\dim E^- < \infty$ and there is an equivalent norm $\|\cdot\|$ on $X = E^- \oplus E^+$ such that

$$\int (\dot{u}^2 + V(x)u^2 - hu^2) = \|u^+\|^2 - \|u^-\|^2,$$

where u^\pm are the orthogonal projections of u on E^\pm .

Let E^- be the negative subspace of $S - h$ and $E^+ = (E^-)^\perp$. Then $\dim E^- < \infty$ and there is an equivalent norm $\|\cdot\|$ on $X = E^- \oplus E^+$ such that

$$\int (\dot{u}^2 + V(x)u^2 - hu^2) = \|u^+\|^2 - \|u^-\|^2,$$

where u^\pm are the orthogonal projections of u on E^\pm .

Lem 3. Under conditions (V_1) , (f_1) and (f_3^*) , Φ satisfies the $(PS)_c$ condition for all $c \in \mathbb{R}$.

Let E^- be the negative subspace of $S - h$ and $E^+ = (E^-)^\perp$. Then $\dim E^- < \infty$ and there is an equivalent norm $\|\cdot\|$ on $X = E^- \oplus E^+$ such that

$$\int (\dot{u}^2 + V(x)u^2 - hu^2) = \|u^+\|^2 - \|u^-\|^2,$$

where u^\pm are the orthogonal projections of u on E^\pm .

Lem 3. Under conditions (V_1) , (f_1) and (f_3^*) , Φ satisfies the $(PS)_c$ condition for all $c \in \mathbb{R}$.

Pf. Under (V_1) , (f_1) and the following condition weaker than (f_3^*) :

(f_3) There exists $h \in (0, \lambda_\infty)$ such that $F(x, t) \leq \frac{1}{2}ht^2$ for all $(x, t) \in \mathbb{R} \times \mathbb{R}$,

Let E^- be the negative subspace of $S - h$ and $E^+ = (E^-)^\perp$. Then $\dim E^- < \infty$ and there is an equivalent norm $\|\cdot\|$ on $X = E^- \oplus E^+$ such that

$$\int (\dot{u}^2 + V(x)u^2 - hu^2) = \|u^+\|^2 - \|u^-\|^2,$$

where u^\pm are the orthogonal projections of u on E^\pm .

Lem 3. Under conditions (V_1) , (f_1) and (f_3^*) , Φ satisfies the $(PS)_c$ condition for all $c \in \mathbb{R}$.

Pf. Under (V_1) , (f_1) and the following condition weaker than (f_3^*) :

(f_3) There exists $h \in (0, \lambda_\infty)$ such that $F(x, t) \leq \frac{1}{2}ht^2$ for all $(x, t) \in \mathbb{R} \times \mathbb{R}$,
it has been shown by Wang & Yang (2015) that Φ is coercive.

Let E^- be the negative subspace of $S - h$ and $E^+ = (E^-)^\perp$. Then $\dim E^- < \infty$ and there is an equivalent norm $\|\cdot\|$ on $X = E^- \oplus E^+$ such that

$$\int (\dot{u}^2 + V(x)u^2 - hu^2) = \|u^+\|^2 - \|u^-\|^2,$$

where u^\pm are the orthogonal projections of u on E^\pm .

Lem 3. Under conditions (V_1) , (f_1) and (f_3^*) , Φ satisfies the $(PS)_c$ condition for all $c \in \mathbb{R}$.

Pf. Under (V_1) , (f_1) and the following condition weaker than (f_3^*) :

(f_3) There exists $h \in (0, \lambda_\infty)$ such that $F(x, t) \leq \frac{1}{2}ht^2$ for all $(x, t) \in \mathbb{R} \times \mathbb{R}$,

it has been shown by Wang & Yang (2015) that Φ is coercive.

Let $\{u_n\}$ be a $(PS)_c$ sequence of Φ , that is

$$\Phi(u_n) \rightarrow c, \quad \Phi'(u_n) \rightarrow 0.$$

By the coerciveness of Φ , $\{u_n\}$ is bounded in X .

Let E^- be the negative subspace of $S - h$ and $E^+ = (E^-)^\perp$. Then $\dim E^- < \infty$ and there is an equivalent norm $\|\cdot\|$ on $X = E^- \oplus E^+$ such that

$$\int (\dot{u}^2 + V(x)u^2 - hu^2) = \|u^+\|^2 - \|u^-\|^2,$$

where u^\pm are the orthogonal projections of u on E^\pm .

Lem 3. Under conditions (V_1) , (f_1) and (f_3^*) , Φ satisfies the $(PS)_c$ condition for all $c \in \mathbb{R}$.

Pf. Under (V_1) , (f_1) and the following condition weaker than (f_3^*) :

(f_3) There exists $h \in (0, \lambda_\infty)$ such that $F(x, t) \leq \frac{1}{2}ht^2$ for all $(x, t) \in \mathbb{R} \times \mathbb{R}$,

it has been shown by Wang & Yang (2015) that Φ is coercive.

Let $\{u_n\}$ be a $(PS)_c$ sequence of Φ , that is

$$\Phi(u_n) \rightarrow c, \quad \Phi'(u_n) \rightarrow 0.$$

By the coerciveness of Φ , $\{u_n\}$ is bounded in X .

We may assume $u_n \rightarrow u$ in X . Since $\dim E^- < \infty$, we have $u_n^- \rightarrow u^-$ and $\|u_n^-\| \rightarrow \|u^-\|$.

Since $\Phi'(u_n) \rightarrow 0$, we have

$$o(1) = o(\|u_n\|) = \langle \Phi'(u_n), u_n \rangle$$

Since $\Phi'(u_n) \rightarrow 0$, we have

$$\begin{aligned} o(1) &= o(\|u_n\|) = \langle \Phi'(u_n), u_n \rangle \\ &= \int (\dot{u}_n^2 + V(x)u_n^2 - hu_n^2) + \langle N'(u_n), u_n \rangle - \int (f(x, u_n)u_n - hu_n^2) \end{aligned}$$

Since $\Phi'(u_n) \rightarrow 0$, we have

$$\begin{aligned} o(1) &= o(\|u_n\|) = \langle \Phi'(u_n), u_n \rangle \\ &= \int (\dot{u}_n^2 + V(x)u_n^2 - hu_n^2) + \langle N'(u_n), u_n \rangle - \int (f(x, u_n)u_n - hu_n^2) \\ &= \|u_n^+\|^2 - \|u_n^-\|^2 + 4N(u_n) + \int (hu_n^2 - f(x, u_n)u_n). \end{aligned}$$

Since $\Phi'(u_n) \rightarrow 0$, we have

$$\begin{aligned} o(1) &= o(\|u_n\|) = \langle \Phi'(u_n), u_n \rangle \\ &= \int (\dot{u}_n^2 + V(x)u_n^2 - hu_n^2) + \langle N'(u_n), u_n \rangle - \int (f(x, u_n)u_n - hu_n^2) \\ &= \|u_n^+\|^2 - \|u_n^-\|^2 + 4N(u_n) + \int (hu_n^2 - f(x, u_n)u_n). \end{aligned} \tag{8}$$

Applying [Lem 2](#) to $g : (x, t) \mapsto ht - f(x, t)$ yields

$$\int (hu_n u - f(x, u_n)u) = \int (hu^2 - f(x, u)u) + o(1).$$

Since $\Phi'(u_n) \rightarrow 0$, we have

$$\begin{aligned} o(1) &= o(\|u_n\|) = \langle \Phi'(u_n), u_n \rangle \\ &= \int (\dot{u}_n^2 + V(x)u_n^2 - hu_n^2) + \langle N'(u_n), u_n \rangle - \int (f(x, u_n)u_n - hu_n^2) \\ &= \|u_n^+\|^2 - \|u_n^-\|^2 + 4N(u_n) + \int (hu_n^2 - f(x, u_n)u_n). \end{aligned} \quad (8)$$

Applying [Lem2](#) to $g : (x, t) \mapsto ht - f(x, t)$ yields

$$\int (hu_n u - f(x, u_n)u) = \int (hu^2 - f(x, u)u) + o(1).$$

Therefore

$$o(1) = \langle \Phi'(u_n), u \rangle$$

Since $\Phi'(u_n) \rightarrow 0$, we have

$$\begin{aligned} o(1) &= o(\|u_n\|) = \langle \Phi'(u_n), u_n \rangle \\ &= \int (\dot{u}_n^2 + V(x)u_n^2 - hu_n^2) + \langle N'(u_n), u_n \rangle - \int (f(x, u_n)u_n - hu_n^2) \\ &= \|u_n^+\|^2 - \|u_n^-\|^2 + 4N(u_n) + \int (hu_n^2 - f(x, u_n)u_n). \end{aligned} \quad (8)$$

Applying [Lem2](#) to $g : (x, t) \mapsto ht - f(x, t)$ yields

$$\int (hu_n u - f(x, u_n)u) = \int (hu^2 - f(x, u)u) + o(1).$$

Therefore

$$\begin{aligned} o(1) &= \langle \Phi'(u_n), u \rangle \\ &= \|u^+\|^2 - \|u^-\|^2 + \langle N'(u_n), u \rangle + \int (hu^2 - f(x, u)u) + o(1). \end{aligned}$$

Since $\Phi'(u_n) \rightarrow 0$, we have

$$\begin{aligned} o(1) &= o(\|u_n\|) = \langle \Phi'(u_n), u_n \rangle \\ &= \int (\dot{u}_n^2 + V(x)u_n^2 - hu_n^2) + \langle N'(u_n), u_n \rangle - \int (f(x, u_n)u_n - hu_n^2) \\ &= \|u_n^+\|^2 - \|u_n^-\|^2 + 4N(u_n) + \int (hu_n^2 - f(x, u_n)u_n). \end{aligned} \quad (8)$$

Applying [Lem2](#) to $g : (x, t) \mapsto ht - f(x, t)$ yields

$$\int (hu_n u - f(x, u_n)u) = \int (hu^2 - f(x, u)u) + o(1).$$

Therefore

$$\begin{aligned} o(1) &= \langle \Phi'(u_n), u \rangle \\ &= \|u^+\|^2 - \|u^-\|^2 + \langle N'(u_n), u \rangle + \int (hu^2 - f(x, u)u) + o(1). \end{aligned} \quad (9)$$

From (8), (9), [Lem1](#), and

$$\int (hu^2 - f(x, u)u) \leq \lim_{n \rightarrow \infty} \int (hu_n^2 - f(x, u_n)u_n), \quad ((f_3^*) \text{ \& Fatou})$$

Since $\Phi'(u_n) \rightarrow 0$, we have

$$\begin{aligned} o(1) &= o(\|u_n\|) = \langle \Phi'(u_n), u_n \rangle \\ &= \int (\dot{u}_n^2 + V(x)u_n^2 - hu_n^2) + \langle N'(u_n), u_n \rangle - \int (f(x, u_n)u_n - hu_n^2) \\ &= \|u_n^+\|^2 - \|u_n^-\|^2 + 4N(u_n) + \int (hu_n^2 - f(x, u_n)u_n). \end{aligned} \quad (8)$$

Applying [Lem2](#) to $g : (x, t) \mapsto ht - f(x, t)$ yields

$$\int (hu_n u - f(x, u_n)u) = \int (hu^2 - f(x, u)u) + o(1).$$

Therefore

$$\begin{aligned} o(1) &= \langle \Phi'(u_n), u \rangle \\ &= \|u^+\|^2 - \|u^-\|^2 + \langle N'(u_n), u \rangle + \int (hu^2 - f(x, u)u) + o(1). \end{aligned} \quad (9)$$

From (8), (9), [Lem1](#), and

$$\int (hu^2 - f(x, u)u) \leq \lim_{n \rightarrow \infty} \int (hu_n^2 - f(x, u_n)u_n), \quad ((f_3^*) \text{ \& Fatou})$$

as well as $\|u_n^-\| \rightarrow \|u^-\|$,

we deduce

$$\begin{aligned}\overline{\lim}_{n \rightarrow \infty} \|u_n^+\|^2 &= \|u^+\|^2 + \int (hu^2 - f(x, u)u) \\ &\quad + \overline{\lim}_{n \rightarrow \infty} \left([\langle N'(u_n), u \rangle - 4N(u_n)] - \int (hu_n u - f(x, u_n)u) \right)\end{aligned}$$

we deduce

$$\begin{aligned}\overline{\lim}_{n \rightarrow \infty} \|u_n^+\|^2 &= \|u^+\|^2 + \int (hu^2 - f(x, u)u) \\ &\quad + \overline{\lim}_{n \rightarrow \infty} \left([\langle N'(u_n), u \rangle - 4N(u_n)] - \int (hu_n u - f(x, u_n)u) \right) \\ &\leq \|u^+\|^2 + \int (hu^2 - f(x, u)u) - \underline{\lim}_{n \rightarrow \infty} \int (hu_n^2 - f(x, u_n)u_n)\end{aligned}$$

we deduce

$$\begin{aligned}\overline{\lim}_{n \rightarrow \infty} \|u_n^+\|^2 &= \|u^+\|^2 + \int (hu^2 - f(x, u)u) \\ &\quad + \overline{\lim}_{n \rightarrow \infty} \left([\langle N'(u_n), u \rangle - 4N(u_n)] - \int (hu_n u - f(x, u_n)u) \right) \\ &\leq \|u^+\|^2 + \int (hu^2 - f(x, u)u) - \underline{\lim}_{n \rightarrow \infty} \int (hu_n^2 - f(x, u_n)u_n) \\ &\leq \|u^+\|^2.\end{aligned}$$

we deduce

$$\begin{aligned}\overline{\lim}_{n \rightarrow \infty} \|u_n^+\|^2 &= \|u^+\|^2 + \int (hu^2 - f(x, u)u) \\ &\quad + \overline{\lim}_{n \rightarrow \infty} \left([\langle N'(u_n), u \rangle - 4N(u_n)] - \int (hu_n u - f(x, u_n)u) \right) \\ &\leq \|u^+\|^2 + \int (hu^2 - f(x, u)u) - \underline{\lim}_{n \rightarrow \infty} \int (hu_n^2 - f(x, u_n)u_n) \\ &\leq \|u^+\|^2.\end{aligned}$$

This and the weak lower semi-continuity of norm functional $u \mapsto \|u\|$, i.e.,

$$\|u^+\|^2 \leq \underline{\lim}_{n \rightarrow \infty} \|u_n^+\|^2, \quad (10)$$

yields $\|u_n^+\| \rightarrow \|u^+\|$.

we deduce

$$\begin{aligned}\overline{\lim}_{n \rightarrow \infty} \|u_n^+\|^2 &= \|u^+\|^2 + \int (hu^2 - f(x, u)u) \\ &\quad + \overline{\lim}_{n \rightarrow \infty} \left([\langle N'(u_n), u \rangle - 4N(u_n)] - \int (hu_n u - f(x, u_n)u) \right) \\ &\leq \|u^+\|^2 + \int (hu^2 - f(x, u)u) - \underline{\lim}_{n \rightarrow \infty} \int (hu_n^2 - f(x, u_n)u_n) \\ &\leq \|u^+\|^2.\end{aligned}$$

This and the weak lower semi-continuity of norm functional $u \mapsto \|u\|$, i.e.,

$$\|u^+\|^2 \leq \underline{\lim}_{n \rightarrow \infty} \|u_n^+\|^2, \quad (10)$$

yields $\|u_n^+\| \rightarrow \|u^+\|$. Therefore $\|u_n\| \rightarrow \|u\|$ and $u_n \rightarrow u$ in X .

we deduce

$$\begin{aligned}
 \overline{\lim}_{n \rightarrow \infty} \|u_n^+\|^2 &= \|u^+\|^2 + \int (hu^2 - f(x, u)u) \\
 &\quad + \overline{\lim}_{n \rightarrow \infty} \left([\langle N'(u_n), u \rangle - 4N(u_n)] - \int (hu_n u - f(x, u_n)u) \right) \\
 &\leq \|u^+\|^2 + \int (hu^2 - f(x, u)u) - \underline{\lim}_{n \rightarrow \infty} \int (hu_n^2 - f(x, u_n)u_n) \\
 &\leq \|u^+\|^2.
 \end{aligned}$$

This and the weak lower semi-continuity of norm functional $u \mapsto \|u\|$, i.e.,

$$\|u^+\|^2 \leq \underline{\lim}_{n \rightarrow \infty} \|u_n^+\|^2, \quad (10)$$

yields $\|u_n^+\| \rightarrow \|u^+\|$. Therefore $\|u_n\| \rightarrow \|u\|$ and $u_n \rightarrow u$ in X .

Pf of Thm 1. Let X^\pm be \pm -spaces of S .

we deduce

$$\begin{aligned}
 \overline{\lim}_{n \rightarrow \infty} \|u_n^+\|^2 &= \|u^+\|^2 + \int (hu^2 - f(x, u)u) \\
 &\quad + \overline{\lim}_{n \rightarrow \infty} \left([\langle N'(u_n), u \rangle - 4N(u_n)] - \int (hu_n u - f(x, u_n)u) \right) \\
 &\leq \|u^+\|^2 + \int (hu^2 - f(x, u)u) - \underline{\lim}_{n \rightarrow \infty} \int (hu_n^2 - f(x, u_n)u_n) \\
 &\leq \|u^+\|^2.
 \end{aligned}$$

This and the weak lower semi-continuity of norm functional $u \mapsto \|u\|$, i.e.,

$$\|u^+\|^2 \leq \underline{\lim}_{n \rightarrow \infty} \|u_n^+\|^2, \quad (10)$$

yields $\|u_n^+\| \rightarrow \|u^+\|$. Therefore $\|u_n\| \rightarrow \|u\|$ and $u_n \rightarrow u$ in X .

Pf of Thm 1. Let X^\pm be \pm -spaces of S . As $u \rightarrow 0$,

Clark (1972/73)

$$\Phi(u) = \frac{1}{2} \int (\dot{u}^2 + V(x)u^2) + \int \dot{u}^2 u^2 - \int F(x, u)$$

we deduce

$$\begin{aligned}
 \overline{\lim}_{n \rightarrow \infty} \|u_n^+\|^2 &= \|u^+\|^2 + \int (hu^2 - f(x, u)u) \\
 &\quad + \overline{\lim}_{n \rightarrow \infty} \left([\langle N'(u_n), u \rangle - 4N(u_n)] - \int (hu_n u - f(x, u_n)u) \right) \\
 &\leq \|u^+\|^2 + \int (hu^2 - f(x, u)u) - \underline{\lim}_{n \rightarrow \infty} \int (hu_n^2 - f(x, u_n)u_n) \\
 &\leq \|u^+\|^2.
 \end{aligned}$$

This and the weak lower semi-continuity of norm functional $u \mapsto \|u\|$, i.e.,

$$\|u^+\|^2 \leq \underline{\lim}_{n \rightarrow \infty} \|u_n^+\|^2, \quad (10)$$

yields $\|u_n^+\| \rightarrow \|u^+\|$. Therefore $\|u_n\| \rightarrow \|u\|$ and $u_n \rightarrow u$ in X .

Pf of Thm 1. Let X^\pm be \pm -spaces of S . As $u \rightarrow 0$, Clark (1972/73)

$$\Phi(u) = \frac{1}{2} \int (\dot{u}^2 + V(x)u^2) + \int \dot{u}^2 u^2 - \int F(x, u) = \frac{1}{2} \int (\dot{u}^2 + V(x)u^2) + o(\|u\|^2),$$

we deduce

$$\begin{aligned}
 \overline{\lim}_{n \rightarrow \infty} \|u_n^+\|^2 &= \|u^+\|^2 + \int (hu^2 - f(x, u)u) \\
 &\quad + \overline{\lim}_{n \rightarrow \infty} \left([\langle N'(u_n), u \rangle - 4N(u_n)] - \int (hu_n u - f(x, u_n)u) \right) \\
 &\leq \|u^+\|^2 + \int (hu^2 - f(x, u)u) - \underline{\lim}_{n \rightarrow \infty} \int (hu_n^2 - f(x, u_n)u_n) \\
 &\leq \|u^+\|^2.
 \end{aligned}$$

This and the weak lower semi-continuity of norm functional $u \mapsto \|u\|$, i.e.,

$$\|u^+\|^2 \leq \underline{\lim}_{n \rightarrow \infty} \|u_n^+\|^2, \quad (10)$$

yields $\|u_n^+\| \rightarrow \|u^+\|$. Therefore $\|u_n\| \rightarrow \|u\|$ and $u_n \rightarrow u$ in X .

Pf of Thm 1. Let X^\pm be \pm -spaces of S . As $u \rightarrow 0$, Clark (1972/73)

$$\Phi(u) = \frac{1}{2} \int (\dot{u}^2 + V(x)u^2) + \int \dot{u}^2 u^2 - \int F(x, u) = \frac{1}{2} \int (\dot{u}^2 + V(x)u^2) + o(\|u\|^2),$$

So Φ has a [loc link](#) at $u = 0$, hence has **3** critical points (Liu (1989)).

3. Proof of Thm 2

Let X^- be the negative subspace of S and $X^+ = (X^-)^\perp$.

3. Proof of Thm 2

Let X^- be the negative subspace of S and $X^+ = (X^-)^\perp$. Then $\dim X^- < \infty$ and there is an equivalent norm $\|\cdot\|$ on X such that

$$\int (\dot{u}^2 + V(x)u^2) = \|u^+\|^2 - \|u^-\|^2,$$

where u^\pm are the orthogonal projections of u on X^\pm .

3. Proof of Thm 2

Let X^- be the negative subspace of S and $X^+ = (X^-)^\perp$. Then $\dim X^- < \infty$ and there is an equivalent norm $\|\cdot\|$ on X such that

$$\int (\dot{u}^2 + V(x)u^2) = \|u^+\|^2 - \|u^-\|^2,$$

where u^\pm are the orthogonal projections of u on X^\pm . Hence

$$\Phi(u) = \frac{1}{2} (\|u^+\|^2 - \|u^-\|^2) + N(u) - \int F(x, u).$$

3. Proof of Thm 2

Let X^- be the negative subspace of S and $X^+ = (X^-)^\perp$. Then $\dim X^- < \infty$ and there is an equivalent norm $\|\cdot\|$ on X such that

$$\int (\dot{u}^2 + V(x)u^2) = \|u^+\|^2 - \|u^-\|^2,$$

where u^\pm are the orthogonal projections of u on X^\pm . Hence

$$\Phi(u) = \frac{1}{2} (\|u^+\|^2 - \|u^-\|^2) + N(u) - \int F(x, u).$$

Lem 4. Under the conditions (V_1) , (f_1) , (f_4) and (f_5) , Φ satisfies the $(PS)_c$ condition for all $c \in \mathbb{R}$.

3. Proof of Thm 2

Let X^- be the negative subspace of S and $X^+ = (X^-)^\perp$. Then $\dim X^- < \infty$ and there is an equivalent norm $\|\cdot\|$ on X such that

$$\int (\dot{u}^2 + V(x)u^2) = \|u^+\|^2 - \|u^-\|^2,$$

where u^\pm are the orthogonal projections of u on X^\pm . Hence

$$\Phi(u) = \frac{1}{2} (\|u^+\|^2 - \|u^-\|^2) + N(u) - \int F(x, u).$$

Lem 4. Under the conditions (V_1) , (f_1) , (f_4) and (f_5) , Φ satisfies the $(PS)_c$ condition for all $c \in \mathbb{R}$.

Pf. Let $\{u_n\}$ be a $(PS)_c$ sequence of Φ .

3. Proof of Thm 2

Let X^- be the negative subspace of S and $X^+ = (X^-)^\perp$. Then $\dim X^- < \infty$ and there is an equivalent norm $\|\cdot\|$ on X such that

$$\int (\dot{u}^2 + V(x)u^2) = \|u^+\|^2 - \|u^-\|^2,$$

where u^\pm are the orthogonal projections of u on X^\pm . Hence

$$\Phi(u) = \frac{1}{2} (\|u^+\|^2 - \|u^-\|^2) + N(u) - \int F(x, u).$$

Lem 4. Under the conditions (V_1) , (f_1) , (f_4) and (f_5) , Φ satisfies the $(PS)_c$ condition for all $c \in \mathbb{R}$.

Pf. Let $\{u_n\}$ be a $(PS)_c$ sequence of Φ .

Step 1. If $\|u_n\| \rightarrow \infty$. Let $v_n = \|u_n\|^{-1}u_n$,

3. Proof of Thm2

Let X^- be the negative subspace of S and $X^+ = (X^-)^\perp$. Then $\dim X^- < \infty$ and there is an equivalent norm $\|\cdot\|$ on X such that

$$\int (\dot{u}^2 + V(x)u^2) = \|u^+\|^2 - \|u^-\|^2,$$

where u^\pm are the orthogonal projections of u on X^\pm . Hence

$$\Phi(u) = \frac{1}{2} (\|u^+\|^2 - \|u^-\|^2) + N(u) - \int F(x, u).$$

Lem 4. Under the conditions (V_1) , (f_1) , (f_4) and (f_5) , Φ satisfies the $(PS)_c$ condition for all $c \in \mathbb{R}$.

Pf. Let $\{u_n\}$ be a $(PS)_c$ sequence of Φ .

Step 1. If $\|u_n\| \rightarrow \infty$. Let $v_n = \|u_n\|^{-1}u_n$, then $v_n \rightarrow v$ in X ,

$$v_n^- \rightarrow v^- \quad \text{in } X$$

because $\dim X^- < \infty$.

If $v = 0$, then $v^- = 0$, $\|v_n^+\|^2 - \|v_n^-\|^2 \geq \frac{1}{2}$ for large n .

If $v = 0$, then $v^- = 0$, $\|v_n^+\|^2 - \|v_n^-\|^2 \geq \frac{1}{2}$ for large n . By (f_4)

$$1 + c + \|u_n\| \geq \Phi(u_n) - \frac{1}{4} \langle \Phi'(u_n), u_n \rangle$$

If $v = 0$, then $v^- = 0$, $\|v_n^+\|^2 - \|v_n^-\|^2 \geq \frac{1}{2}$ for large n . By (f_4)

$$\begin{aligned} 1 + c + \|u_n\| &\geq \Phi(u_n) - \frac{1}{4} \langle \Phi'(u_n), u_n \rangle \\ &= \frac{1}{4} \|u_n\|^2 (\|v_n^+\|^2 - \|v_n^-\|^2) + \int \left(\frac{1}{4} f(x, u_n) u_n - F(x, u_n) \right) \end{aligned}$$

If $v = 0$, then $v^- = 0$, $\|v_n^+\|^2 - \|v_n^-\|^2 \geq \frac{1}{2}$ for large n . By (f_4)

$$\begin{aligned} 1 + c + \|u_n\| &\geq \Phi(u_n) - \frac{1}{4} \langle \Phi'(u_n), u_n \rangle \\ &= \frac{1}{4} \|u_n\|^2 (\|v_n^+\|^2 - \|v_n^-\|^2) + \int \left(\frac{1}{4} f(x, u_n) u_n - F(x, u_n) \right) \\ &\geq \frac{1}{8} \|u_n\|^2, \end{aligned}$$

If $v = 0$, then $v^- = 0$, $\|v_n^+\|^2 - \|v_n^-\|^2 \geq \frac{1}{2}$ for large n . By (f_4)

$$\begin{aligned} 1 + c + \|u_n\| &\geq \Phi(u_n) - \frac{1}{4} \langle \Phi'(u_n), u_n \rangle \\ &= \frac{1}{4} \|u_n\|^2 (\|v_n^+\|^2 - \|v_n^-\|^2) + \int \left(\frac{1}{4} f(x, u_n) u_n - F(x, u_n) \right) \\ &\geq \frac{1}{8} \|u_n\|^2, \end{aligned}$$

contradicting $\|u_n\| \rightarrow \infty$.

If $v = 0$, then $v^- = 0$, $\|v_n^+\|^2 - \|v_n^-\|^2 \geq \frac{1}{2}$ for large n . By (f_4)

$$\begin{aligned} 1 + c + \|u_n\| &\geq \Phi(u_n) - \frac{1}{4} \langle \Phi'(u_n), u_n \rangle \\ &= \frac{1}{4} \|u_n\|^2 (\|v_n^+\|^2 - \|v_n^-\|^2) + \int \left(\frac{1}{4} f(x, u_n) u_n - F(x, u_n) \right) \\ &\geq \frac{1}{8} \|u_n\|^2, \end{aligned}$$

contradicting $\|u_n\| \rightarrow \infty$.

If $v \neq 0$, by Fatou's lemma (5) implies

$$\int \frac{F(x, u_n)}{\|u_n\|^4} \geq \int_{v \neq 0} \frac{F(x, u_n)}{\|u_n\|^4} v_n^4 \rightarrow +\infty.$$

If $v = 0$, then $v^- = 0$, $\|v_n^+\|^2 - \|v_n^-\|^2 \geq \frac{1}{2}$ for large n . By (f_4)

$$\begin{aligned} 1 + c + \|u_n\| &\geq \Phi(u_n) - \frac{1}{4} \langle \Phi'(u_n), u_n \rangle \\ &= \frac{1}{4} \|u_n\|^2 (\|v_n^+\|^2 - \|v_n^-\|^2) + \int \left(\frac{1}{4} f(x, u_n) u_n - F(x, u_n) \right) \\ &\geq \frac{1}{8} \|u_n\|^2, \end{aligned}$$

contradicting $\|u_n\| \rightarrow \infty$.

If $v \neq 0$, by Fatou's lemma (5) implies

$$\int \frac{F(x, u_n)}{\|u_n\|^4} \geq \int_{v \neq 0} \frac{F(x, u_n)}{\|u_n\|^4} v_n^4 \rightarrow +\infty. \quad (11)$$

Using $|N(u)| \leq B\|u\|^4$ we get a contradiction:

$$\frac{c-1}{\|u_n\|^4} \leq \frac{\Phi(u_n)}{\|u_n\|^4}$$

If $v = 0$, then $v^- = 0$, $\|v_n^+\|^2 - \|v_n^-\|^2 \geq \frac{1}{2}$ for large n . By (f_4)

$$\begin{aligned} 1 + c + \|u_n\| &\geq \Phi(u_n) - \frac{1}{4} \langle \Phi'(u_n), u_n \rangle \\ &= \frac{1}{4} \|u_n\|^2 (\|v_n^+\|^2 - \|v_n^-\|^2) + \int \left(\frac{1}{4} f(x, u_n) u_n - F(x, u_n) \right) \\ &\geq \frac{1}{8} \|u_n\|^2, \end{aligned}$$

contradicting $\|u_n\| \rightarrow \infty$.

If $v \neq 0$, by Fatou's lemma (5) implies

$$\int \frac{F(x, u_n)}{\|u_n\|^4} \geq \int_{v \neq 0} \frac{F(x, u_n)}{\|u_n\|^4} v_n^4 \rightarrow +\infty. \quad (11)$$

Using $|N(u)| \leq B\|u\|^4$ we get a contradiction:

$$\frac{c-1}{\|u_n\|^4} \leq \frac{\Phi(u_n)}{\|u_n\|^4} \leq \frac{\|u_n^+\|^2 - \|u_n^-\|^2}{2\|u_n\|^4} + B - \int \frac{F(x, u_n)}{\|u_n\|^4} \rightarrow -\infty.$$

Step 2. Assume $u_n \rightarrow u$. We show that $\|u_n\| \rightarrow \|u\|$.

Step 2. Assume $u_n \rightarrow u$. We show that $\|u_n\| \rightarrow \|u\|$. Since $\dim X^- < \infty$, we have $\|u_n^-\| \rightarrow \|u^-\|$.

Step 2. Assume $u_n \rightarrow u$. We show that $\|u_n\| \rightarrow \|u\|$. Since $\dim X^- < \infty$, we have $\|u_n^-\| \rightarrow \|u^-\|$. Noting

$$\langle N'(u_n), u_n - u \rangle = 4N(u_n) - \langle N'(u_n), u \rangle,$$

Step 2. Assume $u_n \rightarrow u$. We show that $\|u_n\| \rightarrow \|u\|$. Since $\dim X^- < \infty$, we have $\|u_n^-\| \rightarrow \|u^-\|$. Noting

$$\langle N'(u_n), u_n - u \rangle = 4N(u_n) - \langle N'(u_n), u \rangle,$$

by direct computation we deduce

$$0 = \langle \Phi'(u_n), u_n - u \rangle + o(1)$$

Step 2. Assume $u_n \rightarrow u$. We show that $\|u_n\| \rightarrow \|u\|$. Since $\dim X^- < \infty$, we have $\|u_n^-\| \rightarrow \|u^-\|$. Noting

$$\langle N'(u_n), u_n - u \rangle = 4N(u_n) - \langle N'(u_n), u \rangle,$$

by direct computation we deduce

$$\begin{aligned} 0 &= \langle \Phi'(u_n), u_n - u \rangle + o(1) \\ &= (\|u_n^+\|^2 - \|u_n^-\|^2) - (\|u^+\|^2 - \|u^-\|^2) \\ &\quad + 4N(u_n) - \langle N'(u_n), u \rangle - \int f(x, u_n)(u_n - u). \end{aligned}$$

Step 2. Assume $u_n \rightarrow u$. We show that $\|u_n\| \rightarrow \|u\|$. Since $\dim X^- < \infty$, we have $\|u_n^-\| \rightarrow \|u^-\|$. Noting

$$\langle N'(u_n), u_n - u \rangle = 4N(u_n) - \langle N'(u_n), u \rangle,$$

by direct computation we deduce

$$\begin{aligned} 0 &= \langle \Phi'(u_n), u_n - u \rangle + o(1) \\ &= (\|u_n^+\|^2 - \|u_n^-\|^2) - (\|u^+\|^2 - \|u^-\|^2) \\ &\quad + 4N(u_n) - \langle N'(u_n), u \rangle - \int f(x, u_n)(u_n - u). \end{aligned}$$

Now, applying [Lem1](#), using $\|u_n^-\| \rightarrow \|u^-\|$ and condition (f_5) we get

$$\overline{\lim}_{n \rightarrow \infty} \|u_n^+\|^2 = \|u^+\|^2 + \overline{\lim}_{n \rightarrow \infty} \left([\langle N'(u_n), u \rangle - 4N(u_n)] + \int f(x, u_n)(u_n - u) \right)$$

Step 2. Assume $u_n \rightarrow u$. We show that $\|u_n\| \rightarrow \|u\|$. Since $\dim X^- < \infty$, we have $\|u_n^-\| \rightarrow \|u^-\|$. Noting

$$\langle N'(u_n), u_n - u \rangle = 4N(u_n) - \langle N'(u_n), u \rangle,$$

by direct computation we deduce

$$\begin{aligned} 0 &= \langle \Phi'(u_n), u_n - u \rangle + o(1) \\ &= (\|u_n^+\|^2 - \|u_n^-\|^2) - (\|u^+\|^2 - \|u^-\|^2) \\ &\quad + 4N(u_n) - \langle N'(u_n), u \rangle - \int f(x, u_n)(u_n - u). \end{aligned}$$

Now, applying **Lem1**, using $\|u_n^-\| \rightarrow \|u^-\|$ and condition (f_5) we get

$$\begin{aligned} \overline{\lim}_{n \rightarrow \infty} \|u_n^+\|^2 &= \|u^+\|^2 + \overline{\lim}_{n \rightarrow \infty} \left([\langle N'(u_n), u \rangle - 4N(u_n)] + \int f(x, u_n)(u_n - u) \right) \\ &\leq \|u^+\|^2. \end{aligned}$$

Hence $\|u_n^+\| \rightarrow \|u^+\|$.

Step 2. Assume $u_n \rightarrow u$. We show that $\|u_n\| \rightarrow \|u\|$. Since $\dim X^- < \infty$, we have $\|u_n^-\| \rightarrow \|u^-\|$. Noting

$$\langle N'(u_n), u_n - u \rangle = 4N(u_n) - \langle N'(u_n), u \rangle,$$

by direct computation we deduce

$$\begin{aligned} 0 &= \langle \Phi'(u_n), u_n - u \rangle + o(1) \\ &= (\|u_n^+\|^2 - \|u_n^-\|^2) - (\|u^+\|^2 - \|u^-\|^2) \\ &\quad + 4N(u_n) - \langle N'(u_n), u \rangle - \int f(x, u_n)(u_n - u). \end{aligned}$$

Now, applying **Lem1**, using $\|u_n^-\| \rightarrow \|u^-\|$ and condition (f_5) we get

$$\begin{aligned} \overline{\lim}_{n \rightarrow \infty} \|u_n^+\|^2 &= \|u^+\|^2 + \overline{\lim}_{n \rightarrow \infty} \left([\langle N'(u_n), u \rangle - 4N(u_n)] + \int f(x, u_n)(u_n - u) \right) \\ &\leq \|u^+\|^2. \end{aligned}$$

Hence $\|u_n^+\| \rightarrow \|u^+\|$. Noting $\|u_n^-\| \rightarrow \|u^-\|$, we conclude $\|u_n\| \rightarrow \|u\|$.

Lem 5. Assume (V_1) , (f_1) , (f_4) , there exists $A < \inf_{B_2} \Phi$,

Lem 5. Assume $(V_1), (f_1), (f_4)$, there exists $A < \inf_{B_2} \Phi$, s.t. if $\Phi(u) \leq A$, then

$$\left. \frac{d}{dt} \right|_{t=1} \Phi(tu) < 0.$$

Lem 5. Assume (V_1) , (f_1) , (f_4) , there exists $A < \inf_{B_2} \Phi$, s.t. if $\Phi(u) \leq A$, then

$$\left. \frac{d}{dt} \right|_{t=1} \Phi(tu) < 0.$$

Pf. Otherwise, there is a sequence $\{u_n\} \subset X$ such that $\Phi(u_n) \leq -n$ but

$$\langle \Phi'(u_n), u_n \rangle = \left. \frac{d}{dt} \right|_{t=1} \Phi(tu_n) \geq 0.$$

Lem 5. Assume (V_1) , (f_1) , (f_4) , there exists $A < \inf_{B_2} \Phi$, s.t. if $\Phi(u) \leq A$, then

$$\left. \frac{d}{dt} \right|_{t=1} \Phi(tu) < 0.$$

Pf. Otherwise, there is a sequence $\{u_n\} \subset X$ such that $\Phi(u_n) \leq -n$ but

$$\langle \Phi'(u_n), u_n \rangle = \left. \frac{d}{dt} \right|_{t=1} \Phi(tu_n) \geq 0. \quad (12)$$

Using (f_4) , we deduce

$$\|u_n^+\|^2 - \|u_n^-\|^2 \leq (\|u_n^+\|^2 - \|u_n^-\|^2) + \int [f(x, u_n)u_n - 4F(x, u_n)]$$

Lem 5. Assume (V_1) , (f_1) , (f_4) , there exists $A < \inf_{B_2} \Phi$, s.t. if $\Phi(u) \leq A$, then

$$\left. \frac{d}{dt} \right|_{t=1} \Phi(tu) < 0.$$

Pf. Otherwise, there is a sequence $\{u_n\} \subset X$ such that $\Phi(u_n) \leq -n$ but

$$\langle \Phi'(u_n), u_n \rangle = \left. \frac{d}{dt} \right|_{t=1} \Phi(tu_n) \geq 0. \quad (12)$$

Using (f_4) , we deduce

$$\begin{aligned} \|u_n^+\|^2 - \|u_n^-\|^2 &\leq (\|u_n^+\|^2 - \|u_n^-\|^2) + \int [f(x, u_n)u_n - 4F(x, u_n)] \\ &= 4\Phi(u_n) - \langle \Phi'(u_n), u_n \rangle \end{aligned}$$

Lem 5. Assume (V_1) , (f_1) , (f_4) , there exists $A < \inf_{B_2} \Phi$, s.t. if $\Phi(u) \leq A$, then

$$\left. \frac{d}{dt} \right|_{t=1} \Phi(tu) < 0.$$

Pf. Otherwise, there is a sequence $\{u_n\} \subset X$ such that $\Phi(u_n) \leq -n$ but

$$\langle \Phi'(u_n), u_n \rangle = \left. \frac{d}{dt} \right|_{t=1} \Phi(tu_n) \geq 0. \quad (12)$$

Using (f_4) , we deduce

$$\begin{aligned} \|u_n^+\|^2 - \|u_n^-\|^2 &\leq (\|u_n^+\|^2 - \|u_n^-\|^2) + \int [f(x, u_n)u_n - 4F(x, u_n)] \\ &= 4\Phi(u_n) - \langle \Phi'(u_n), u_n \rangle \leq -4n. \end{aligned}$$

Lem 5. Assume (V_1) , (f_1) , (f_4) , there exists $A < \inf_{B_2} \Phi$, s.t. if $\Phi(u) \leq A$, then

$$\left. \frac{d}{dt} \right|_{t=1} \Phi(tu) < 0.$$

Pf. Otherwise, there is a sequence $\{u_n\} \subset X$ such that $\Phi(u_n) \leq -n$ but

$$\langle \Phi'(u_n), u_n \rangle = \left. \frac{d}{dt} \right|_{t=1} \Phi(tu_n) \geq 0. \quad (12)$$

Using (f_4) , we deduce

$$\begin{aligned} \|u_n^+\|^2 - \|u_n^-\|^2 &\leq (\|u_n^+\|^2 - \|u_n^-\|^2) + \int [f(x, u_n)u_n - 4F(x, u_n)] \\ &= 4\Phi(u_n) - \langle \Phi'(u_n), u_n \rangle \leq -4n. \end{aligned} \quad (13)$$

Let $v_n = \|u_n\|^{-1}u_n$, then $v_n^- \rightarrow v^-$ in X ($\dim X^- < \infty$), $v^- \neq 0$.

Lem 5. Assume (V_1) , (f_1) , (f_4) , there exists $A < \inf_{B_2} \Phi$, s.t. if $\Phi(u) \leq A$, then

$$\left. \frac{d}{dt} \right|_{t=1} \Phi(tu) < 0.$$

Pf. Otherwise, there is a sequence $\{u_n\} \subset X$ such that $\Phi(u_n) \leq -n$ but

$$\langle \Phi'(u_n), u_n \rangle = \left. \frac{d}{dt} \right|_{t=1} \Phi(tu_n) \geq 0. \quad (12)$$

Using (f_4) , we deduce

$$\begin{aligned} \|u_n^+\|^2 - \|u_n^-\|^2 &\leq (\|u_n^+\|^2 - \|u_n^-\|^2) + \int [f(x, u_n)u_n - 4F(x, u_n)] \\ &= 4\Phi(u_n) - \langle \Phi'(u_n), u_n \rangle \leq -4n. \end{aligned} \quad (13)$$

Let $v_n = \|u_n\|^{-1}u_n$, then $v_n^- \rightarrow v^-$ in X ($\dim X^- < \infty$), $v^- \neq 0$. Hence

$$\int \frac{f(x, u_n)u_n}{\|u_n\|^4} \geq 4 \int \frac{F(x, u_n)}{\|u_n\|^4} \rightarrow +\infty.$$

Lem 5. Assume (V_1) , (f_1) , (f_4) , there exists $A < \inf_{B_2} \Phi$, s.t. if $\Phi(u) \leq A$, then

$$\left. \frac{d}{dt} \right|_{t=1} \Phi(tu) < 0.$$

Pf. Otherwise, there is a sequence $\{u_n\} \subset X$ such that $\Phi(u_n) \leq -n$ but

$$\langle \Phi'(u_n), u_n \rangle = \left. \frac{d}{dt} \right|_{t=1} \Phi(tu_n) \geq 0. \quad (12)$$

Using (f_4) , we deduce

$$\begin{aligned} \|u_n^+\|^2 - \|u_n^-\|^2 &\leq (\|u_n^+\|^2 - \|u_n^-\|^2) + \int [f(x, u_n)u_n - 4F(x, u_n)] \\ &= 4\Phi(u_n) - \langle \Phi'(u_n), u_n \rangle \leq -4n. \end{aligned} \quad (13)$$

Let $v_n = \|u_n\|^{-1}u_n$, then $v_n^- \rightarrow v^-$ in X ($\dim X^- < \infty$), $v^- \neq 0$. Hence

$$\int \frac{f(x, u_n)u_n}{\|u_n\|^4} \geq 4 \int \frac{F(x, u_n)}{\|u_n\|^4} \rightarrow +\infty. \quad (\text{see (11)})$$

$$0 \leq \frac{\langle \Phi'(u_n), u_n \rangle}{\|u_n\|^4}$$

Lem 5. Assume (V_1) , (f_1) , (f_4) , there exists $A < \inf_{B_2} \Phi$, s.t. if $\Phi(u) \leq A$, then

$$\left. \frac{d}{dt} \right|_{t=1} \Phi(tu) < 0.$$

Pf. Otherwise, there is a sequence $\{u_n\} \subset X$ such that $\Phi(u_n) \leq -n$ but

$$\langle \Phi'(u_n), u_n \rangle = \left. \frac{d}{dt} \right|_{t=1} \Phi(tu_n) \geq 0. \quad (12)$$


Using (f_4) , we deduce

$$\begin{aligned} \|u_n^+\|^2 - \|u_n^-\|^2 &\leq (\|u_n^+\|^2 - \|u_n^-\|^2) + \int [f(x, u_n)u_n - 4F(x, u_n)] \\ &= 4\Phi(u_n) - \langle \Phi'(u_n), u_n \rangle \leq -4n. \end{aligned} \quad (13)$$

Let $v_n = \|u_n\|^{-1}u_n$, then $v_n^- \rightarrow v^-$ in X ($\dim X^- < \infty$), $v^- \neq 0$. Hence

$$\int \frac{f(x, u_n)u_n}{\|u_n\|^4} \geq 4 \int \frac{F(x, u_n)}{\|u_n\|^4} \rightarrow +\infty. \quad (\text{see (11)})$$

$$0 \leq \frac{\langle \Phi'(u_n), u_n \rangle}{\|u_n\|^4} \leq \frac{\|u_n^+\|^2 - \|u_n^-\|^2}{\|u_n\|^4} + \frac{\langle N'(u_n), u_n \rangle}{\|u_n\|^4} - \int \frac{f(x, u_n)u_n}{\|u_n\|^4} \rightarrow -\infty.$$




$$\varphi : X \setminus B_1 \rightarrow \Phi_A,$$

$$\varphi(v) = \begin{cases} T_{v/\|v\|} \frac{v}{\|v\|} & \text{if } \Phi(v) > A \\ v & \text{if } \Phi(v) \leq A \end{cases}$$

Pf of Thm 2. For $u \in \partial B_1$, it is clear that

$$\Phi(tu) \rightarrow -\infty \quad \text{as } t \rightarrow +\infty.$$




$$\varphi : X \setminus B_1 \rightarrow \Phi_A,$$

$$\varphi(v) = \begin{cases} T_{v/\|v\|} \frac{v}{\|v\|} & \text{if } \Phi(v) > A \\ v & \text{if } \Phi(v) \leq A \end{cases}$$

Pf of Thm2. For $u \in \partial B_1$, it is clear that

$$\Phi(tu) \rightarrow -\infty \quad \text{as } t \rightarrow +\infty.$$

Thus $\exists 1 T_u > 1$ s.t. $\Phi(T_u u) = A$.



$$\varphi : X \setminus B_1 \rightarrow \Phi_A,$$


$$\varphi(v) = \begin{cases} T_{v/\|v\|} \frac{v}{\|v\|} & \text{if } \Phi(v) > A \\ v & \text{if } \Phi(v) \leq A \end{cases}$$

Pf of Thm2. For $u \in \partial B_1$, it is clear that

$$\Phi(tu) \rightarrow -\infty \quad \text{as } t \rightarrow +\infty.$$

Thus $\exists 1 T_u > 1$ s.t. $\Phi(T_u u) = A$. By Lem5

$$\left. \frac{d}{ds} \right|_{s=T_u} \Phi(su)$$



$$\varphi : X \setminus B_1 \rightarrow \Phi_A,$$


$$\varphi(v) = \begin{cases} T_{v/\|v\|} \frac{v}{\|v\|} & \text{if } \Phi(v) > A \\ v & \text{if } \Phi(v) \leq A \end{cases}$$

Pf of Thm2. For $u \in \partial B_1$, it is clear that

$$\Phi(tu) \rightarrow -\infty \quad \text{as } t \rightarrow +\infty.$$

Thus $\exists 1 T_u > 1$ s.t. $\Phi(T_u u) = A$. By Lem5

$$\left. \frac{d}{ds} \right|_{s=T_u} \Phi(su) = \frac{1}{T_u} \left. \frac{d}{dt} \right|_{t=1} \Phi(t \cdot T_u u) < 0.$$



$$\varphi : X \setminus B_1 \rightarrow \Phi_A,$$

$$\varphi(v) = \begin{cases} T_{v/\|v\|} \frac{v}{\|v\|} & \text{if } \Phi(v) > A \\ v & \text{if } \Phi(v) \leq A \end{cases}$$


Pf of Thm2. For $u \in \partial B_1$, it is clear that

$$\Phi(tu) \rightarrow -\infty \quad \text{as } t \rightarrow +\infty.$$

Thus $\exists T_u > 1$ s.t. $\Phi(T_u u) = A$. By Lem5

$$\left. \frac{d}{ds} \right|_{s=T_u} \Phi(su) = \frac{1}{T_u} \left. \frac{d}{dt} \right|_{t=1} \Phi(t \cdot T_u u) < 0.$$

Following Wang (1991), by IFT $u \mapsto T_u$ is continuous,



$$\varphi : X \setminus B_1 \rightarrow \Phi_A,$$

$$\varphi(v) = \begin{cases} T_{v/\|v\|} \frac{v}{\|v\|} & \text{if } \Phi(v) > A \\ v & \text{if } \Phi(v) \leq A \end{cases}$$


Pf of Thm2. For $u \in \partial B_1$, it is clear that

$$\Phi(tu) \rightarrow -\infty \quad \text{as } t \rightarrow +\infty.$$

Thus $\exists T_u > 1$ s.t. $\Phi(T_u u) = A$. By Lem5

$$\left. \frac{d}{ds} \right|_{s=T_u} \Phi(su) = \frac{1}{T_u} \left. \frac{d}{dt} \right|_{t=1} \Phi(t \cdot T_u u) < 0.$$

Following Wang (1991), by IFT $u \mapsto T_u$ is continuous, and we construct a deformation $\varphi : X \setminus B_1 \rightarrow \Phi_A$,



$$\varphi : X \setminus B_1 \rightarrow \Phi_A,$$

$$\varphi(v) = \begin{cases} T_{v/\|v\|} \frac{v}{\|v\|} & \text{if } \Phi(v) > A \\ v & \text{if } \Phi(v) \leq A \end{cases}$$

Pf of Thm2. For $u \in \partial B_1$, it is clear that


$$\Phi(tu) \rightarrow -\infty \quad \text{as } t \rightarrow +\infty.$$

Thus $\exists 1 T_u > 1$ s.t. $\Phi(T_u u) = A$. By Lem5

$$\left. \frac{d}{ds} \right|_{s=T_u} \Phi(su) = \frac{1}{T_u} \left. \frac{d}{dt} \right|_{t=1} \Phi(t \cdot T_u u) < 0.$$

Following Wang (1991), by IFT $u \mapsto T_u$ is continuous, and we construct a deformation $\varphi : X \setminus B_1 \rightarrow \Phi_A$, and deduce (critical groups)

$$C_i(\Phi, \infty) = H_i(X, \Phi_A)$$



$$\varphi : X \setminus B_1 \rightarrow \Phi_A,$$

$$\varphi(v) = \begin{cases} T_{v/\|v\|} \frac{v}{\|v\|} & \text{if } \Phi(v) > A \\ v & \text{if } \Phi(v) \leq A \end{cases}$$

Pf of Thm2. For $u \in \partial B_1$, it is clear that


$$\Phi(tu) \rightarrow -\infty \quad \text{as } t \rightarrow +\infty.$$

Thus $\exists 1 T_u > 1$ s.t. $\Phi(T_u u) = A$. By [Lem5](#)

$$\left. \frac{d}{ds} \right|_{s=T_u} \Phi(su) = \frac{1}{T_u} \left. \frac{d}{dt} \right|_{t=1} \Phi(t \cdot T_u u) < 0.$$

Following [Wang \(1991\)](#), by IFT $u \mapsto T_u$ is continuous, and we construct a deformation $\varphi : X \setminus B_1 \rightarrow \Phi_A$, and deduce (critical groups)

$$C_i(\Phi, \infty) = H_i(X, \Phi_A) = H_i(X, X \setminus B_1) = 0 \quad \text{for } i \in \mathbb{N}_0.$$



$$\varphi : X \setminus B_1 \rightarrow \Phi_A,$$

$$\varphi(v) = \begin{cases} T_{v/\|v\|} \frac{v}{\|v\|} & \text{if } \Phi(v) > A \\ v & \text{if } \Phi(v) \leq A \end{cases}$$

Pf of Thm2. For $u \in \partial B_1$, it is clear that

$$\Phi(tu) \rightarrow -\infty \quad \text{as } t \rightarrow +\infty.$$

Thus $\exists T_u > 1$ s.t. $\Phi(T_u u) = A$. By [Lem5](#)


$$\left. \frac{d}{ds} \right|_{s=T_u} \Phi(su) = \frac{1}{T_u} \left. \frac{d}{dt} \right|_{t=1} \Phi(t \cdot T_u u) < 0.$$

Following [Wang \(1991\)](#), by IFT $u \mapsto T_u$ is continuous, and we construct a deformation $\varphi : X \setminus B_1 \rightarrow \Phi_A$, and deduce (critical groups)

$$C_i(\Phi, \infty) = H_i(X, \Phi_A) = H_i(X, X \setminus B_1) = 0 \quad \text{for } i \in \mathbb{N}_0.$$

Since Φ has a loc link at $u = 0$, by [Liu \(1989\)](#) (for $\ell = \dim X^-$)

$$C_\ell(\Phi, 0) \neq 0.$$



$$\varphi : X \setminus B_1 \rightarrow \Phi_A,$$

$$\varphi(v) = \begin{cases} T_{v/\|v\|} \frac{v}{\|v\|} & \text{if } \Phi(v) > A \\ v & \text{if } \Phi(v) \leq A \end{cases}$$

Pf of Thm2. For $u \in \partial B_1$, it is clear that

$$\Phi(tu) \rightarrow -\infty \quad \text{as } t \rightarrow +\infty.$$

Thus $\exists 1 T_u > 1$ s.t. $\Phi(T_u u) = A$. By Lem5

$$\left. \frac{d}{ds} \right|_{s=T_u} \Phi(su) = \frac{1}{T_u} \left. \frac{d}{dt} \right|_{t=1} \Phi(t \cdot T_u u) < 0.$$

Following Wang (1991), by IFT $u \mapsto T_u$ is continuous, and we construct a deformation $\varphi : X \setminus B_1 \rightarrow \Phi_A$, and deduce (critical groups)

$$C_i(\Phi, \infty) = H_i(X, \Phi_A) = H_i(X, X \setminus B_1) = 0 \quad \text{for } i \in \mathbb{N}_0.$$

Since Φ has a loc link at $u = 0$, by Liu (1989) (for $\ell = \dim X^-$)

$$C_\ell(\Phi, 0) \neq 0. \quad \text{Hence } C_\ell(\Phi, 0) \neq C_\ell(\Phi, \infty),$$

Applying Bartsch & Li (1997), Φ has a crt pt $u \neq 0$.

References

- Poppenberg M, Schmitt K, Wang ZQ (2002). [On the existence of soliton solutions to quasilinear Schrödinger equations](#). Calc. Var. Partial Differential Equations, 14(3) 329–344.
- Liu Jq, Wang Yq, Wang ZQ (2003). [Soliton solutions for quasilinear Schrödinger equations. II](#). J. Differential Equations, 187(2) 473–493.
- Colin M, Jeanjean L (2004). [Solutions for a quasilinear Schrödinger equation: a dual approach](#). Nonlinear Anal., 56(2) 213–226.
- Shen Z, Han Z (2015). [Existence of solutions to quasilinear Schrödinger equations with indefinite potential](#). Electron. J. Differential Equations, No. 91, 9.
- Liu S, Zhou J (2018). [Standing waves for quasilinear Schrödinger equations with indefinite potentials](#). J. Differential Equations, 265(9) 3970–3987.
- Silva ED, Silva JS (2019). [Quasilinear Schrödinger equations with nonlin-](#)

earities interacting with high eigenvalues. J. Math. Phys., 60(8) 081504, 24.

Wang DB, Yang K (2015). Existence of solutions for a class of quasilinear Schrödinger equations on \mathbb{R} . Bound. Value Probl., 2015:215, 5.

Chen S, Wang C (2014). Existence of multiple nontrivial solutions for a Schrödinger-Poisson system. J. Math. Anal. Appl., 411(2) 787–793.

Benci V, Fortunato D (1998). An eigenvalue problem for the Schrödinger-Maxwell equations. Topol. Methods Nonlinear Anal., 11(2) 283–293.

Chen H, Liu S (2015). Standing waves with large frequency for 4-superlinear Schrödinger-Poisson systems. Ann. Mat. Pura Appl. (4), 194(1) 43–53.

Liu S, Wu Y (2017). Standing waves for 4-superlinear Schrödinger-Poisson systems with indefinite potentials. Bull. Lond. Math. Soc., 49(2) 226–234.

Bartsch T, Liu Z, Weth T (2004). Sign changing solutions of superlinear Schrödinger equations. Comm. Partial Differential Equations, 29(1-2) 25–42.

- Chen J (2014). [Multiple positive solutions to a class of modified nonlinear Schrödinger equations](#). J. Math. Anal. Appl., 415(2) 525–542.
- Brézis H, Lieb E (1983). [A relation between pointwise convergence of functions and convergence of functionals](#). Proc. Amer. Math. Soc., 88(3) 486–490.
- Clark DC (1972/73). [A variant of the Lusternik-Schnirelman theory](#). Indiana Univ. Math. J., 22 65–74.
- Liu JQ (1989). [The Morse index of a saddle point](#). Systems Sci. Math. Sci., 2(1) 32–39.
- Wang ZQ (1991). [On a superlinear elliptic equation](#). Ann. Inst. H. Poincaré Anal. Non Linéaire, 8(1) 43–57.
- Bartsch T, Li S (1997). [Critical point theory for asymptotically quadratic functionals and applications to problems with resonance](#). Nonlinear Anal., 28(3) 419–441.

Thank you!

<http://lausb.github.io>