

Gagliardo-Nirenberg-Sobolev inequality: an induction proof

The Author

Abstract. An induction proof of the G-N-S inequality is presented, which is much simpler and transparent than other proofs found in the literature.

MSC: 46E35

Let $p \in [1, n)$, then there is a constant $C_{n,p} > 0$ such that

$$\left(\int_{\mathbb{R}^n} |u|^{p^*} \right)^{1/p^*} \leq C_{n,p} \left(\int_{\mathbb{R}^n} |\nabla u|^p \right)^{1/p} \quad \text{for all } u \in C_0^1(\mathbb{R}^n), \quad (1)$$

where $p^* = np/(n-p)$ is the critical Sobolev exponent.

This is the Gagliardo-Nirenberg-Sobolev inequality, which is very fundamental in the theory of Sobolev spaces and partial differential equations. It is well known that (1) follows from applying the result for the case $p = 1$ to $|u|^\gamma$ with some suitable exponent γ . Traditionally (see e.g. [2, Page 277–278]), the inequality (1) for $p = 1$ is proved by integrating

$$|u(x)|^{n/(n-1)} \leq \prod_{i=1}^n \left(\int_{-\infty}^{\infty} |\partial_i u(x_1, \dots, x_{i-1}, y_i, x_{i+1}, \dots, x_n)| dy_i \right)^{1/(n-1)}$$

with respect to x_1 and applying the extended Hölder inequality, then repeating this procedure with respect to x_2, x_3, \dots, x_n successively. In every step the extended Hölder inequality is applied. This tedious procedure is not very transparent, and is not easy to follow (especially for the beginning graduate students).

Observe that the inequality (for $p = 1$) is a proposition about the dimension n . Therefore it should be natural to prove the result by induction on n . Surprisingly, we could not find such a proof in the literature. The purpose of this note is to present the induction proof, which is very simple and transparent.

Theorem 1. Let $n > 1$, $u \in C_0^1(\mathbb{R}^n)$, then

$$\int_{\mathbb{R}^n} |u|^{n/(n-1)} \leq \left(\int_{\mathbb{R}^n} |\nabla u| \right)^{n/(n-1)}. \quad (2)$$

Proof. Our induction proof starts with $n = 2$, which can be found in [1, Page 27]. We present it here for the reader's convenience. Since $u \in C_0^1(\mathbb{R}^n)$, we have

$$|u(x)| = \left| \int_{-\infty}^{x_1} \partial_1 u(s, x_2) ds \right| \leq \int_{-\infty}^{\infty} |\nabla u(s, x_2)| ds.$$

Replacing the right hand side by $\int_{-\infty}^{\infty} |\nabla u(x_1, s)| ds$, the inequality is also true. Thus

$$|u(x)|^2 \leq \left(\int_{-\infty}^{\infty} |\nabla u(s, x_2)| ds \right) \left(\int_{-\infty}^{\infty} |\nabla u(x_1, s)| ds \right).$$

Integrating both sides over \mathbb{R}^2 gives (2) for $n = 2$.

Assume that the result is true for $n = m$, namely

$$\int_{\mathbb{R}^m} |u|^{m/(m-1)} \leq \left(\int_{\mathbb{R}^m} |\nabla u| \right)^{m/(m-1)} \quad \text{for } u \in C_0^1(\mathbb{R}^m). \quad (3)$$

To consider the case that $n = m + 1$, we write points in \mathbb{R}^{m+1} by (x, t) , being $x \in \mathbb{R}^m$ and $t \in \mathbb{R}$. Then for $u \in C_0^1(\mathbb{R}^{m+1})$,

$$|u(x, t)| = \left| \int_{-\infty}^t \partial_{m+1} u(x, s) \, ds \right| \leq \int_{-\infty}^t |\nabla u(x, s)| \, ds,$$

and

$$\int_{\mathbb{R}^m} |u(x, t)| \, dx \leq \int_{\mathbb{R}^m} dx \int_{-\infty}^{\infty} |\nabla u(x, s)| \, ds = \int_{\mathbb{R}^{m+1}} |\nabla u|. \quad (4)$$

Here we write the differentials like ds to indicate with respect to which variable we are integrating; if we are integrating with respect to all variables, we omit the differentials (for example, $dx ds$ is omitted in the final integral in (4)).

Using the Hölder inequality, (3) with u replaced by $u(\cdot, t) \in C_0^1(\mathbb{R}^m)$, and (4), we have

$$\begin{aligned} \int_{\mathbb{R}^m} |u(x, t)|^{(m+1)/m} \, dx &= \int_{\mathbb{R}^m} |u|^{1/m} |u| \, dx \\ &\leq \left(\int_{\mathbb{R}^m} |u| \, dx \right)^{1/m} \left(\int_{\mathbb{R}^m} |u|^{m/(m-1)} \, dx \right)^{(m-1)/m} \\ &\leq \left(\int_{\mathbb{R}^{m+1}} |\nabla u| \right)^{1/m} \left(\int_{\mathbb{R}^m} |\tilde{\nabla} u(x, t)| \, dx \right), \end{aligned} \quad (5)$$

where $\tilde{\nabla} u(x, t)$ is the gradient of $u(\cdot, t)$ at x . Obviously $|\tilde{\nabla} u(x, t)| \leq |\nabla u(x, t)|$, thus integrating both sides with respect to t over \mathbb{R} yields

$$\begin{aligned} \int_{\mathbb{R}^{m+1}} |u|^{(m+1)/m} &= \int_{-\infty}^{\infty} dt \int_{\mathbb{R}^m} |u(x, t)|^{(m+1)/m} \, dx \\ &\leq \left(\int_{\mathbb{R}^{m+1}} |\nabla u| \right)^{1/m} \left[\int_{-\infty}^{\infty} dt \int_{\mathbb{R}^m} |\nabla u(x, t)| \, dx \right] \\ &= \left(\int_{\mathbb{R}^{m+1}} |\nabla u| \right)^{(m+1)/m}, \end{aligned}$$

we have obtained (2) for $n = m + 1$. So (2) is true for all $n \geq 2$. ■

Remark. Using the same idea, a stronger Gagliardo-Nirenberg-Sobolev inequality

$$\int_{\mathbb{R}^n} |u|^{n/(n-1)} \leq \left(\prod_{i=1}^n \int_{\mathbb{R}^n} |\partial_i u| \right)^{1/(n-1)} \quad (S_n)$$

can be proved. To derive (S_{m+1}) from (S_m) , we replace all the ∇u before (5) by $\partial_{m+1} u$, then apply the induction assumption (S_m) to $\int |u|^{m/(m-1)}$ in (5). When integrating with

respect to t we need the extended Hölder inequality, which is not needed in our proof of Theorem 1.

In [3], combining a simple iteration scheme with this stronger version of the Gagliardo-Nirenberg-Sobolev inequality, the author gives a new proof of the L^∞ -embedding

$$\|u\|_{L^\infty(\mathbb{R}^n)} \leq C_p (\|u\|_{L^p(\mathbb{R}^n)} + \|\nabla u\|_{L^p(\mathbb{R}^n)}),$$

where $p > n$, $u \in C_0^1(\mathbb{R}^n)$.

REFERENCES

1. Chen, W., Li, C. (2010). *Methods on nonlinear elliptic equations*. AIMS Series on Differential Equations & Dynamical Systems. Springfield, MO: American Institute of Mathematical Sciences (AIMS).
2. Evans, L. C. (2010). *Partial differential equations*, 2nd ed. Graduate Studies in Mathematics. Providence, RI: American Mathematical Society.
3. Porretta, A. (2020). A note on the Sobolev and Gagliardo-Nirenberg inequality when $p > N$, *Adv. Non-linear Stud.* 20(2): 361–371. doi.org/10.1515/ans-2020-2086