

Lab 9

Lauren Waters (A16326738)

```
candy_fine <- "candy-data.csv"
candy = read.csv(candy_fine, row.names=1)
head(candy)
```

	chocolate	fruity	caramel	peanutyalmondy	nougat	crispedricewafer
100 Grand	1	0	1	0	0	1
3 Musketeers	1	0	0	0	1	0
One dime	0	0	0	0	0	0
One quarter	0	0	0	0	0	0
Air Heads	0	1	0	0	0	0
Almond Joy	1	0	0	1	0	0

	hard	bar	pluribus	sugarpercent	pricepercent	winpercent
100 Grand	0	1	0	0.732	0.860	66.97173
3 Musketeers	0	1	0	0.604	0.511	67.60294
One dime	0	0	0	0.011	0.116	32.26109
One quarter	0	0	0	0.011	0.511	46.11650
Air Heads	0	0	0	0.906	0.511	52.34146
Almond Joy	0	1	0	0.465	0.767	50.34755

Q1. How many different candy types are in this dataset?

```
nrow(candy)
```

```
[1] 85
```

Q2. How many fruity candy types are in the dataset?

```
sum(candy$fruity)
```

```
[1] 38
```

Q.3 What is your favorite candy in the dataset and what is it's winpercent value?

```
candy["100 Grand", ]$winpercent
```

```
[1] 66.97173
```

Q4. What is the winpercent value for “Kit Kat”?

```
candy["Kit Kat", ]$winpercent
```

```
[1] 76.7686
```

Q5. What is the winpercent value for “Tootsie Roll Snack Bars”?

```
candy["Tootsie Roll Snack Bars", ]$winpercent
```

```
[1] 49.6535
```

```
library("skimr")
skim(candy)
```

Table 1: Data summary

Name	candy
Number of rows	85
Number of columns	12
Column type frequency:	
numeric	12
Group variables	None

Variable type: numeric

skim_variable	n_missing	complete_rate	mean	sd	p0	p25	p50	p75	p100	hist
chocolate	0	1	0.44	0.50	0.00	0.00	0.00	1.00	1.00	
fruity	0	1	0.45	0.50	0.00	0.00	0.00	1.00	1.00	

skim_variable	n_missing	complete_rate	mean	sd	p0	p25	p50	p75	p100	hist
caramel	0	1	0.16	0.37	0.00	0.00	0.00	0.00	1.00	
peanutyalmondy	0	1	0.16	0.37	0.00	0.00	0.00	0.00	1.00	
nougat	0	1	0.08	0.28	0.00	0.00	0.00	0.00	1.00	
crispedricewafer	0	1	0.08	0.28	0.00	0.00	0.00	0.00	1.00	
hard	0	1	0.18	0.38	0.00	0.00	0.00	0.00	1.00	
bar	0	1	0.25	0.43	0.00	0.00	0.00	0.00	1.00	
pluribus	0	1	0.52	0.50	0.00	0.00	1.00	1.00	1.00	
sugarpercent	0	1	0.48	0.28	0.01	0.22	0.47	0.73	0.99	
pricepercent	0	1	0.47	0.29	0.01	0.26	0.47	0.65	0.98	
winpercent	0	1	50.32	14.71	22.45	39.14	47.83	59.86	84.18	

Q6. Is there any variable/column that looks to be on a different scale to the majority of the other columns in the dataset?


The winpercent is not within a 0-1 scale.

Q7. What do you think a zero and one represent for the candy\$chocolate column?

A 1 represents that the candy is chocolate, and a 0 represents that there is no chocolate in the candy.

Q8. Plot a histogram of winpercent values

```
hist(candy$winpercent)
```




Q9. Is the distribution of winpercent values symmetrical?

The graph is slightly skewed.

Q10. Is the center of the distribution above or below 50%?

```
hist(candy$winpercent)
```

Histogram of candy\$winpercent



Below

Q.11 On average is chocolate candy higher or lower ranked than fruit candy?

```
choco.inds <- as.logical(candy$chocolate)
choco.win <- candy[choco.inds, "winpercent"]
mean(choco.win)
```

```
[1] 60.92153
```

```
fruity.inds <- as.logical(candy$fruity)
fruity.win <- candy[fruity.inds, "winpercent"]
mean(fruity.win)
```

```
[1] 44.11974
```

On average, chocolate is ranked higher.

Q12. Is this difference statistically significant?

```
t.test(choco.win, fruity.win)
```

Welch Two Sample t-test

```
data: choco.win and fruity.win
t = 6.2582, df = 68.882, p-value = 2.871e-08
alternative hypothesis: true difference in means is not equal to 0
95 percent confidence interval:
 11.44563 22.15795
sample estimates:
mean of x mean of y
 60.92153  44.11974
```

Because the P-value is so small, it's significant.

Q13. What are the five least liked candy types in this set?

```
head(candy[order(candy$winpercent),], n=5)
```

	chocolate	fruity	caramel	peanut	almond	nougat
Nik L Nip	0	1	0		0	0
Boston Baked Beans	0	0	0		1	0
Chiclets	0	1	0		0	0
Super Bubble	0	1	0		0	0
Jawbusters	0	1	0		0	0

	crisped	rice	wafer	hard	bar	pluribus	sugar	percent	price	percent
Nik L Nip				0	0	0	1	0.197		0.976
Boston Baked Beans				0	0	0	1	0.313		0.511
Chiclets				0	0	0	1	0.046		0.325
Super Bubble				0	0	0	0	0.162		0.116
Jawbusters				0	1	0	1	0.093		0.511

	winpercent
Nik L Nip	22.44534
Boston Baked Beans	23.41782
Chiclets	24.52499
Super Bubble	27.30386
Jawbusters	28.12744

Q14. What are the top 5 all time favorite candy types out of this set?

```
head(candy[order(-candy$winpercent),], n=5)
```

	chocolate	fruity	caramel	peanut	almond	nougat
Reese's Peanut Butter cup	1	0	0		1	0
Reese's Miniatures	1	0	0		1	0
Twix	1	0	1		0	0
Kit Kat	1	0	0		0	0
Snickers	1	0	1		1	1


	crisped	rice	wafers	hard	bar	pluribus	sugar
Reese's Peanut Butter cup		0	0	0		0	0.720
Reese's Miniatures		0	0	0		0	0.034
Twix		1	0	1		0	0.546
Kit Kat		1	0	1		0	0.313
Snickers		0	0	1		0	0.546

	price	percent	winpercent
Reese's Peanut Butter cup	0.651	84.18029	
Reese's Miniatures	0.279	81.86626	
Twix	0.906	81.64291	
Kit Kat	0.511	76.76860	
Snickers	0.651	76.67378	

Q15. Make a first barplot of candy ranking based on winpercent values.


```
library(ggplot2)

ggplot(candy) +
  aes(winpercent, rownames(candy)) +
  geom_col()
```




Q16. This is quite ugly, use the `reorder()` function to get the bars sorted by winpercent?

```
ggplot(candy) +  
  aes(winpercent, reorder(rownames(candy), winpercent)) +  
  geom_col()
```

```
mycols <- rep("gray", nrow(candy))
mycols[ as.logical(candy$chocolate) ] <- "chocolate"
mycols[ as.logical(candy$fruity) ] <- "pink"
mycols[ as.logical(candy$bar)] <- "brown"

ggplot(candy) +
  aes(winpercent, reorder(rownames(candy),winpercent)) +
  geom_col(fill=mycols)
```



Q17. What is the worst ranked chocolate candy?

Sixlets


Q18. What is the best ranked fruity candy?

Starburst

```
library(ggrepel)

# How about a plot of price vs win
ggplot(candy) +
  aes(winpercent, pricepercent, label=rownames(candy)) +
  geom_point(col=mycols) +
  geom_text_repel(col=mycols, size=3.3, max.overlaps = 5)
```

Warning: ggrepel: 65 unlabeled data points (too many overlaps). Consider increasing max.overlaps



Q19. Which candy type is the highest ranked in terms of winpercent for the least money - i.e. offers the most bang for your buck?

Reese's miniatures

Q20. What are the top 5 most expensive candy types in the dataset and of these which is the least popular?

```
ord <- order(candy$pricepercent, decreasing = TRUE)
head( candy[ord,c(11,12)], n=5 )
```


	pricepercent	winpercent
Nik L Nip	0.976	22.44534
Nestle Smarties	0.976	37.88719
Ring pop	0.965	35.29076
Hershey's Krackel	0.918	62.28448
Hershey's Milk Chocolate	0.918	56.49050

Nik L Nip

```
library(corrplot)
```

corrplot 0.92 loaded

```
cij <- cor(candy)
corrplot(cij)
```



Q22. Examining this plot what two variables are anti-correlated (i.e. have minus values)?

chocolate and fruity

Q23. Similarly, what two variables are most positively correlated?

chocolate and winpercent/bar (other than chocolate-chocolate, fruity-fruity, ...).


```
pca <- prcomp(candy, scale=TRUE)
summary(pca)
```

Importance of components:

	PC1	PC2	PC3	PC4	PC5	PC6	PC7
Standard deviation	2.0788	1.1378	1.1092	1.07533	0.9518	0.81923	0.81530
Proportion of Variance	0.3601	0.1079	0.1025	0.09636	0.0755	0.05593	0.05539
Cumulative Proportion	0.3601	0.4680	0.5705	0.66688	0.7424	0.79830	0.85369

	PC8	PC9	PC10	PC11	PC12
Standard deviation	0.74530	0.67824	0.62349	0.43974	0.39760
Proportion of Variance	0.04629	0.03833	0.03239	0.01611	0.01317
Cumulative Proportion	0.89998	0.93832	0.97071	0.98683	1.00000


```
plot(pca$x[,1:2], col=mycols, pch=16)
```



```
my_data <- cbind(candy, pca$x[,1:3])

p <- ggplot(my_data) +
  aes(x=PC1, y=PC2,
      size=winpercent/100,
      text=rownames(my_data),
      label=rownames(my_data)) +
  geom_point(col=mycols)
```

```
p
```




```
library(ggrepel)

p + geom_text_repel(size=3.3, col=mycols, max.overlaps = 7) +
  theme(legend.position = "none") +
  labs(title="Halloween Candy PCA Space",
        subtitle="Colored by type: chocolate bar (dark brown), chocolate other (light brown)",
        caption="Data from 538")
```

Warning: ggrepel: 59 unlabeled data points (too many overlaps). Consider increasing max.overlaps


Halloween Candy PCA Space

Colored by type: chocolate bar (dark brown), chocolate other (light brown),



Data from 538

```
par(mar=c(8,4,2,2))
barplot(pca$rotation[,1], las=2, ylab="PC1 Contribution")
```



Q24. What original variables are picked up strongly by PC1 in the positive direction? Do these make sense to you?

Fruity candies are strongly positive (which makes sense when considering fruity candies like skittles, nerds, ...).