
Unique HE breakpoints (T. timococcum + T. zhukovskyi)
Non-breakpoint windows within HE active regions around telomere
a Am
n = 616 unique pairs (both species) vs 35,982 non-breakpoint windows
Mann–Whitney p = 4.0e-75; permutation p < 1e-4
Density, HE breakpoints
Density, non-breakpoint windows
K-mer variations per 100-kb window
b At
n = 616 unique pairs (both species) vs 26,112 non-breakpoint windows
Mann–Whitney p = 6.8e-59; permutation p < 1e-4
Density, HE breakpoints
Density, non-breakpoint windows
K-mer variations per 100-kb window
c Am
K-mer variations per 100-kb window
***
n.s.
***
n = 35,982
mean = 1,077
SE = ±1.2
n = 487
mean = 923
SE = ±7.6
n = 129
mean = 904
SE = ±13.2
Non-breakpoint HE active region
T. timococcum HE breakpoint windows
T. zhukovskyi HE breakpoint windows
d At
K-mer variations per 100-kb window
***
n.s.
***
n = 26,112
mean = 1,087
SE = ±1.3
n = 487
mean = 953
SE = ±8.1
n = 129
mean = 953
SE = ±11.3
Non-breakpoint HE active region
T. timococcum HE breakpoint windows
T. zhukovskyi HE breakpoint windows