

Subject: Our ep3d sequence from Bay 2
Date: Saturday, October 24, 2026 at 14:16:23 Eastern Daylight Time
From: Nasr, Shahin,Ph.D.
To: Pizzuti, Alessandra (PSYCHOLOGY), Huber, Laurentius
Attachments: Martinos_ep3d.pdf, image.png

Hi Alessandra,

A copy of our ep3d protocol, generated by Renzo, is attached to this email.

Renzo is very familiar with our slice-positioning. But as a reminder, we do it almost like this:



** In our first attempt in Bay 2 we noticed pulasatility in the scans. Later @Huber, Laurentius found the source and fixed the issue. I highly suggest eyeballing the scan to make sure that the problem is not back in Bay 3. If it was back, don't continue the scan.

I prefer having a TR=3 s for both rs-FC and my visual stimulation runs.

For rs-FC, #TR=128 is enough. We can use the same run for tSNR measurements.
For motion-selectivity test, we aim for #TR=88.

So, the optimum day will include these scans:

- Localizer/advanced-shimming (if available)

- Mesovein
- Motion-selectivity (0.8mm) -> 6 runs
- Flashing-checker board (0.3mm) -> ??? Runs (skip this step if you think we need more than 2-3 runs)
- Rs-FC (0.8mm) -> 3 runs
- TAKE ME OUT!

Please let me know if you need any further info.

Bests,

Shahin Nasr, Ph. D.
Presidential Early Career Award for Scientists and Engineers 2025
Principal Investigator, MesoVision Laboratory
Assistant Professor of Radiology
Mass General/Martinos Center for Biomedical Imaging
Harvard Medical School
https://mesovision.martinos.org/

Mailing address: 149 13th Street, Suite 2301, Charlestown MA 02129