

First Order Logic of Sparse Random Hyper-Graphs

Lázaro Alberto Larrauri Borroto
Supervisor: Marc Noy Serrano

July 6, 2019

1 First Order Logic of Sparse Graphs

2 General relational structures

Contents

1 First Order Logic of Sparse Graphs

2 General relational structures

The first order language of graphs.

- Variables x_1, \dots, x_n, \dots
- Connectives \wedge, \vee , equality symbol $=$, and negation symbol \neg .
- Quantifiers \forall, \exists .
- A binary relation symbol R
- Vertices.
- “And”, “or”, “equals”, “not”.
- “For all”, “there exists”.
- Edges $x \sim y$.

$$\forall x_1, x_2 R(x_1, x_2) \implies \exists x_3 (\neg(x_3 = x_1) \wedge \neg(x_3 = x_2) \wedge R(x_1, x_3))$$

The binomial model

The binomial model of random graphs $G(n, p)$ is a discrete probability space where we assign to each graph $G = ([n], E)$ the probability

$$\Pr(G) = p^{|E|} \cdot (1 - p)^{\binom{n}{2} - |E|}.$$

Lynch's theorem

Theorem (Lynch, 1992)

Let φ be a sentence in the F.O. language of graphs. Then the map $F_\varphi : [0, \infty) \rightarrow \mathbb{R}$ given by

$$F_\varphi(\beta) = \lim_{n \rightarrow \infty} \Pr(G(n, \beta/n) \models \varphi)$$

is well defined and admits an analytic extension to \mathbb{C} .

Overview of the proof

Some properties of $G(n, \beta/n)$:

- The number of cycles of length $3, 4 \dots, r$ are asymptotically distributed like independent Poisson variables.
- Small cycles are a.a.s far away.
- Fixed vertices are a.a.s far away.
- The ball of a given radius centered in fixed vertex is a.a.s a tree. Any tree occurs with a positive probability.

Overview of the proof

For each fixed quantifier rank k :

- (1) It is given a finite classification of "small" uni-cycles.
- (2) It is shown that the rank k type of random graph G in $G(n, \beta/n)$ a.a.s depends exclusively on the number of "small" uni-cycles belonging to each class.
- (3) The asymptotic distribution of those quantities is obtained.

Contents

1 First Order Logic of Sparse Graphs

2 General relational structures

Edge sets

Definition

Fix

- natural numbers $n, a \in \mathbb{N}$, with $a \geq 2$
- a subgroup of the symmetric group $\Phi \leq S_a$
- and a subset of pairs $A \subset ([a] \times [a]) \setminus \Delta_a$.

The **total edge set** $\mathcal{H}_{(a,\Phi,A)}(n)$ of size a , symmetry group Φ and restrictions A , on n elements is the set

$$\mathcal{H}_{(a,\Phi,A)}(n) = ([n]^a / \Phi) \setminus R,$$

where

$$R = \{ [x_1, \dots, x_a] \in [n]^a / \Phi \mid x_i = x_j \text{ for some } (i, j) \in A \}$$

Graphs

Definition

An (hyper)-graph $([n], H_1, \dots, H_c)$ with edge colors $1, \dots, c$, sizes a_1, \dots, a_c , symmetry groups Φ_1, \dots, Φ_c and restrictions A_1, \dots, A_c consists of

- The vertex set $[n]$ for some natural number n .
- For $i = 1, \dots, c$, a “colored” edge set $H_i \subseteq \mathcal{H}_{(a_i, \Phi_i, A_i)}(n)$ whose elements have color i .

The first order language

Consider the first order purely relational language \mathcal{L} whose signature consists of the relation symbols R_1, \dots, R_c with arities a_1, \dots, a_c .

A graph $G = ([n], H_1, \dots, H_c)$ is a \mathcal{L} -structure in the following way:

- The universe of G is its vertex set, $[n]$.
- For each $1 \leq i \leq c$,

$$(x_1, \dots, x_{a_i}) \in R_i^G \iff [x_1, \dots, x_{a_i}] \in H_i.$$

The first order language

By definition, a graph $G = ([n], H_1, \dots, H_n)$ satisfies, for each $1 \leq i \leq c$:

- Symmetry formulas:

$$S_g := (R_i(x_1 \dots, x_{a_i}) \iff R_i(x_{g(1)} \dots, x_{g(a_i)})),$$

where g is an element from Φ_i .

- Anti-reflexivity formulas:

$$A_{i,(j,l)} := (R_i(x_1 \dots, x_{a_i}) \implies \neg(x_j = x_l)),$$

where $(j, l) \in A_i$.

The random model

The random model $HG(n, p_1, \dots, p_c)$ is a discrete probability space where for each graph $G = ([n], H_1, \dots, H_c)$,

$$\Pr(G) = \prod_{i=1}^c p_i^{|H_i|} \cdot (1 - p_i)^{|\mathcal{H}_{(a_i, \Phi_i, A_i)}(n)| - |H_i|}.$$

We consider the case where for each $1 \leq i \leq c$, $p_i(n) = \beta_i / n^{a_i - 1}$.

Let $\beta = (\beta_1, \dots, \beta_c) \in [0, \infty)^c$. We abbreviate

$$HG(n, p(n, \beta)) := HG(n, \beta_1 / n^{a_1 - 1}, \dots, \beta_c / n^{a_c - 1}).$$

The theorem

For each $1 \leq i \leq c$ let $p_i(n, \beta_i) = \beta_i / n^{a_i-1}$. Let $\beta = (\beta_1, \dots, \beta_c)$ and $p(n, \beta) = (p_1(n, \beta_1), \dots, p_c(n, \beta_c))$.