

¹ Observation of the Higgs boson in the WW^*
² channel and search for Higgs boson pair
³ production in the $b\bar{b}b\bar{b}$ channel with the
⁴ ATLAS detector

⁵ A DISSERTATION PRESENTED
⁶ BY
⁷ TOMO LAZOVICH
⁸ TO
⁹ THE DEPARTMENT OF PHYSICS

¹⁰ IN PARTIAL FULFILLMENT OF THE REQUIREMENTS
¹¹ FOR THE DEGREE OF
¹² DOCTOR OF PHILOSOPHY
¹³ IN THE SUBJECT OF
¹⁴ PHYSICS

¹⁵ HARVARD UNIVERSITY
¹⁶ CAMBRIDGE, MASSACHUSETTS
¹⁷ MAY 2016

¹⁸ ©2016 – TOMO LAZOVICH

¹⁹ ALL RIGHTS RESERVED.

²⁰ **Observation of the Higgs boson in the WW^* channel and search
²¹ for Higgs boson pair production in the $b\bar{b}b\bar{b}$ channel with the
²² ATLAS detector**

²³ ABSTRACT

²⁴ This dissertation presents the observation and measurement of the Higgs boson in the $H \rightarrow WW^* \rightarrow$
²⁵ $\ell\nu\ell\nu$ channel at $\sqrt{s} = 7$ TeV and $\sqrt{s} = 8$ TeV and a search for Higgs pair production in the $HH \rightarrow$
²⁶ $b\bar{b}b\bar{b}$ channel at $\sqrt{s} = 13$ TeV with the ATLAS detector in pp collisions at the Large Hadron Collider.

²⁷ First, the discovery of a particle consistent with the Higgs boson in 4.8 fb^{-1} at $\sqrt{s} = 7$ TeV and
²⁸ 5.8 fb^{-1} at $\sqrt{s} = 8$ TeV is discussed. Then, the measurement of the Higgs boson signal strength
²⁹ and cross section in both the gluon fusion and vector boson fusion (VBF) production modes using
³⁰ 20.3 fb^{-1} of $\sqrt{s} = 8$ TeV data combined with 4.8 fb^{-1} of 7 TeV data is shown. The combined signal
³¹ strength is measured to be $\mu = 1.09^{+0.23}_{-0.21}$. The total observed significance of the $H \rightarrow WW^*$ process
³² is observed to be 6.1σ (with 5.8σ expected). Advanced methods for background reduction and estima-
³³ tion, particularly in same-flavor lepton final states, are shown. The VBF signal strength is measured to
³⁴ be $\mu_{\text{VBF}} = 1.27^{+0.53}_{-0.45}$ with an observed significance of 3.2σ (with 2.7σ expected). In the VBF chan-
³⁵ nel, a selection requirement based method, the precursor to the final multivariate technique used for the
³⁶ result, is detailed.

³⁷ Finally, a search for Higgs pair production in the $b\bar{b}b\bar{b}$ final state with 3.2 fb^{-1} at $\sqrt{s} = 13$ TeV is
³⁸ presented. A particular focus is placed on a tailored signal region for resonant production of Higgs pairs
³⁹ at high masses. No significant excesses are observed, and upper limits on cross sections are placed for
⁴⁰ spin-2 Randall Sundrum gravitons (RSG) and narrow spin-0 resonances. The cross section of $\sigma(pp \rightarrow$
⁴¹ $G_{\text{KK}}^* \rightarrow hh \rightarrow b\bar{b}b\bar{b})$ with $k/\bar{M}_{\text{Pl}} = 1$ is constrained to be less than 70 fb for masses in the range
⁴² $600 < m_{G_{\text{KK}}^*} < 3000$ GeV. The cross section upper limits for $\sigma(pp \rightarrow H \rightarrow hh \rightarrow b\bar{b}b\bar{b})$ ranges
⁴³ from 30 to 300 fb in the mass range of $500 < m_H < 3000$ GeV.

Contents

44

45	o INTRODUCTION	I
46	I Theoretical and Experimental Background	5
47	I THE PHYSICS OF THE HIGGS BOSON	6
48	1.1 The Standard Model of Particle Physics	6
49	1.2 Electroweak Symmetry Breaking and the Higgs	8
50	1.3 Higgs Boson Production and Decay	II
51	1.4 Higgs Pair Production in the Standard Model	16
52	1.5 Higgs Pair Production in Theories Beyond the Standard Model	17
53	1.6 Conclusion	21
54	2 THE ATLAS DETECTOR AND THE LARGE HADRON COLLIDER	23
55	2.1 The Large Hadron Collider	24
56	2.2 The ATLAS Detector	26
57	2.3 The ATLAS Muon New Small Wheel Upgrade	37
58	2.4 Object Reconstruction in ATLAS	41
59	II Observation and measurement of Higgs boson decays to WW^* in LHC	
60	Run 1 at $\sqrt{s} = 7$ and 8 TeV	50
61	3 $H \rightarrow WW^* \rightarrow \ell\nu\ell\nu$ ANALYSIS STRATEGY	51
62	3.1 Introduction	51
63	3.2 The $H \rightarrow WW^* \rightarrow \ell\nu\ell\nu$ signal in ATLAS	52
64	3.3 Background processes	53
65	3.4 Shared signal region selection requirements	56
66	3.5 Background reduction in same-flavor final states	60
67	3.6 Parameters of interest and statistical treatment	64
68	4 THE DISCOVERY OF THE HIGGS BOSON AND THE ROLE OF THE $H \rightarrow WW^* \rightarrow \ell\nu\ell\nu$ CHANNEL	70
69	4.1 Introduction	70

71	4.2	Data and simulation samples	71
72	4.3	$H \rightarrow WW \rightarrow e\nu\mu\nu$ search	71
73	4.4	$H \rightarrow \gamma\gamma$ search	76
74	4.5	$H \rightarrow ZZ \rightarrow 4\ell$ search	77
75	4.6	Combined results	79
76	4.7	Conclusion	81
77	5	EVIDENCE FOR VECTOR BOSON FUSION PRODUCTION OF $H \rightarrow WW^* \rightarrow \ell\nu\ell\nu$	83
78	5.1	Introduction	83
79	5.2	Data and simulation samples	84
80	5.3	Object selection	88
81	5.4	Analysis selection	91
82	5.5	Background estimation	100
83	5.6	Systematic uncertainties	110
84	5.7	Results	113
85	6	COMBINED RUN I $H \rightarrow WW^* \rightarrow \ell\nu\ell\nu$ RESULTS	118
86	6.1	Introduction	118
87	6.2	Results of dedicated gluon fusion $H \rightarrow WW^* \rightarrow \ell\nu\ell\nu$ analysis	119
88	6.3	Signal strength measurements in ggF and VBF production	121
89	6.4	Measurement of Higgs couplings to vector bosons and fermions	126
90	6.5	Higgs production cross section measurement	126
91	6.6	Conclusion	128
92	III	Search for Higgs pair production in the $HH \rightarrow b\bar{b}b\bar{b}$ channel in LHC	
93		Run 2 at $\sqrt{s} = 13$ TeV	130
94	7	SEARCH FOR HIGGS PAIR PRODUCTION IN BOOSTED $b\bar{b}b\bar{b}$ FINAL STATES	131
95	7.1	Introduction	131
96	7.2	Motivation	132
97	7.3	Data and simulation samples	134
98	7.4	Event reconstruction and object selection	136
99	7.5	Event selection	139
100	7.6	Data-driven background estimation	144
101	7.7	Systematic uncertainties	150
102	7.8	Results	153

103	8 COMBINED LIMITS FROM BOOSTED AND RESOLVED SEARCHES	155
104	8.1 Introduction	155
105	8.2 Resolved results	156
106	8.3 Search technique and results	156
107	8.4 Limit setting	157
108	IV Looking ahead	161
109	9 CONCLUSION	162
110	APPENDIX A <i>b</i>-TAGGING PERFORMANCE AT HIGH p_T	166
111	A.1 Changes in MV2 score at high p_T	166
112	A.2 Effect of multiple <i>b</i> -quarks inside one jet	168
113	A.3 Changes in track quality at high p_T	170
114	REFERENCES	173

Listing of figures

116	I.1	The particles of the Standard Model and their properties [6].	7
117	I.2	The four most common Higgs boson production modes at the LHC: (a) gluon-gluon fusion, (b) vector boson fusion, (c) $W/Z + H$ production, (d) $t\bar{t}H$ production	11
118	I.3	Higgs production cross sections as a function of center of mass energy (\sqrt{s}) at a pp collider [18].	12
119	I.4	Higgs boson branching ratios as a function of m_H [18].	15
120	I.5	The two leading diagrams for Standard Model di-Higgs production at the LHC: (a) box diagram, (b) Higgs self coupling.	16
121	I.6	Diagrams with new vertices for non-resonant Higgs pair production arising in composite Higgs models.	18
122	I.7	Generic Feynman diagram for resonant Higgs pair production in BSM theories.	18
123	I.8	Branching ratios for a spin-2 Randall-Sundrum graviton as a function of mass computed in MadGraph with the CP3-Origins implementation [25, 31, 32].	19
124	I.9	$\sigma \times \text{BR}(HH)$ for Randall-Sundrum gravitons as a function of mass computed in MadGraph with the CP3-Origins implementation [25, 31, 32].	20
125	I.10	Randall-Sundrum graviton width as a function of mass computed in MadGraph with the CP3-Origins implementation [25, 31, 32]	20
126	I.11	Branching ratios for heavy Higgs H in Type I (left) and Type II (right) 2HDM models with $\tan \beta = 1.5$ and $\cos(\beta - \alpha) = 0.1(0.01)$ for Type I (Type II) [29].	22
135	2.1	A schematic view of the LHC ring [38]	24
136	2.2	A full diagram of the ATLAS detector [34]	27
137	2.3	The ATLAS coordinate system	28
138	2.4	Layout of the ATLAS Inner Detector system [42]	29
139	2.5	Layout of the ATLAS calorimeter system [34]	31
140	2.6	Layout of the ATLAS muon system [34]	33
141	2.7	Predicted field integral as a function of $ \eta $ for the ATLAS magnet system [34]	35
142	2.8	ATLAS trigger rates for Level-1 triggers as a function of instantaneous luminosity in 2012 and 2015 operation. These are single object triggers for electromagnetic clusters (EM), muons (MU), jets (J), missing energy (XE), and τ leptons (TAU). The threshold of the trigger is given in the name in GeV. [44]	36
144			

146	2.9	Instantaneous luminosity as a function of time for data recorded by ATLAS at different center of mass energies [45, 46]	37
148	2.10	MDT tube hit (solid) and segment (dashed) efficiency as a function of hit rate per tube [47]	38
149	2.11	Trigger rate as a function of p_T threshold with and without the NSW upgrade [47]	39
150	2.12	Illustrations of the geometry (left) and operating principle (right) of the micromegas detector [47]	40
152	2.13	Geometry of the sTGC detector [47]	40
153	2.14	Illustration of particle interactions in ATLAS [50]	42
154	2.15	Electron performance: (a) reconstruction efficiency as a function of electron E_T [52] (b) energy resolution in simulation as a function of $ \eta $ for different energy electrons [53] . . .	43
155	2.16	Muon performance: (a) reconstruction efficiency as a function of muon p_T (b) dimuon mass resolution as a function of average p_T [54]	44
158	2.17	Jet energy response after calibration as a function of true p_T in simulation [58]	46
159	2.18	Summary of the inputs to the MV2 b -tagging algorithm	47
160	2.19	Light jet rejection (1/efficiency) vs. b -jet efficiency for MV1 and its input algorithms (a) [59] and MV2 (b) [60] in simulated $t\bar{t}$ events. The numbers in the algorithm names in (b) refer to the fraction of charm events used in the MV2 training.	48
163	2.20	Resolution of E_T^{miss} components as a function of $\sum E_T$ before pileup suppression with different pileup techniques [62]	49
165	3.1	Branching ratios for a WW system. q refers to quarks. ℓ can be either an electron or muon, and the leptonic branching ratios of the τ are included. For example, the $\ell\nu qq$ final state includes one W decaying to $e\nu$, $\mu\nu$, or $\tau\nu$. τ_h refer to hadronic decays of the τ	53
168	3.2	Feynman diagram for Standard Model WW production	54
169	3.3	Feynman diagrams for top pair production (left) and Wt production (right)	54
170	3.4	An example Feynman diagram of W +jets production	55
171	3.5	An example Feynman diagram of Z +jets production	56
172	3.6	An illustration of the unique analysis signal regions [63]. The most sensitive regions for both gluon fusion and vector boson fusion production are underlined.	57
174	3.7	A graphical illustration of the $E_{T,\text{rel}}^{\text{miss}}$ calculation	58
175	3.8	Predicted backgrounds (compared with data) as a function of n_j (a and b) and n_b (c) after pre-selection requirements	59
177	3.9	An event display of a Z/γ^* + jets event illustrating the effect of pileup interactions	61
178	3.10	The RMS of different missing transverse momentum definitions as a function of the average number of interactions per bunch crossing	61
180	3.11	The difference between the true and reconstructed values of the missing transverse momentum (a) and m_T (b) in a gluon fusion signal sample	63

182	3.12	Comparison of f_{recoil} distributions for $Z/\gamma^* + \text{jets}$, $H \rightarrow WW^*$, and other backgrounds with real neutrinos.	64
183			
184	3.13	Signal significance as a function of required value for f_{recoil} and $p_{\text{T},\text{rel}}^{\text{miss}(\text{trk})}$ in the ggF $H \rightarrow WW^*$ with $n_j = 0$	65
185			
186	4.1	Jet multiplicity distribution in data and MC after applying lepton, jet, and $E_{\text{T},\text{rel}}^{\text{miss}}$ selections. The WW and top backgrounds have been normalized using control samples, and the hashed band indicates the total uncertainty on the prediction. [1]	72
187			
188	4.2	Comparison of m_{T} between data and simulation in the $n_j = 0$ WW (a) and $n_j = 1$ top (b) control samples [1]	74
189			
190	4.3	m_{T} distribution in the $H \rightarrow WW \rightarrow e\nu\mu\nu$ $n_j \leq 1$ channels for 8 TeV data [1].	76
191			
192	4.4	Diphoton mass spectrum in 7 and 8 TeV data. Panel a) shows the unweighted data distribution superimposed on the background fit, while panel c) shows the data where each event category is weighted by its signal to background ratio. Panels b) and d) show the respective distributions with background subtracted [1].	77
193			
194	4.5	Four lepton invariant mass spectrum ($m_{4\ell}$) in 7 and 8 TeV data compared to background estimate. A 125 GeV SM Higgs signal is shown in blue [1].	78
195			
196	4.6	Local p_0 distribution as a function of hypothesized Higgs mass for the $H \rightarrow ZZ^* \rightarrow 4\ell$ (a), $H \rightarrow \gamma\gamma$ (b), and $H \rightarrow WW^* \rightarrow \ell\nu\ell\nu$ (c) channels. Dashed curves show expected results, while solid curves show observed. Red curves are from 7 TeV data, blue curves from 8 TeV, and black curved combined [1].	80
197			
198	4.7	Combined 95% CL limits (a), local p_0 values (b), and signal strength measurement (c) as a function of Higgs mass [1].	81
199			
200	4.8	Comparison of measured signal strength μ for a 126 GeV Higgs in the 7 and 8 TeV datasets [1].	82
201			
202	4.9	Two dimensional likelihood as a function of signal strength μ and Higgs mass m_H [1].	82
203			
204	5.1	A comparison of the subleading lepton p_{T} spectrum between VBF $H \rightarrow WW^*$ production and $t\bar{t}$ background	85
205			
206	5.2	Leading jet η in VBF $H \rightarrow WW^*$ (red) and $t\bar{t}$ (black)	93
207			
208	5.3	Distributions of (a) m_{jj} , (b) Δy_{jj} , (c) $C_{\ell 1}$, and (d) $\Sigma m_{\ell j}$, for the VBF analysis. The top panels compare simulation and data, while the bottom panels show normalized distributions for all background processes and signal [63].	95
209			
210	5.4	A cartoon of the WW final state. Momenta are represented with thin arrows, spins with thick arrows. [63]	96
211			
212	5.5	Event display of a VBF candidate event [63].	98
213			
214	5.6	Distributions of $m_{\ell\ell}$ (top left), $\Delta\phi_{\ell\ell}$ (top right), and m_{T} (bottom), Higgs topology variables used in the selection requirements of the cut-based signal region and as inputs to the BDT result. These are plotted after all of the BDT pre-training selection cuts [63].	99
215			
216			
217			

218	5.7	Distributions of m_{jj} (top left), Δy_{jj} (top right), $\sum C_\ell$ (bottom), VBF topology variables used in the selection requirements of the cut-based signal region and as inputs to the BDT result. These are plotted after all of the BDT pre-training selection cuts [63].	100
219			
220	5.8	Distributions of m_{jj} (a) and O_{BDT} (b) in the VBF $n_b = 1$ top CR [63].	103
221			
222	5.9	Comparison of m_{jj} shape in a same flavor $Z \rightarrow \ell\ell$ control region and the VBF cut-based signal region.	104
223			
224	5.10	General illustration of the ABCD region definitions for $Z/\gamma^* \rightarrow \ell\ell$ background estimation.	105
225			
226	5.11	Distribution of $m_{\text{T}2}$ in the WW validation region of the VBF analysis [63].	107
227			
228	5.12	Extrapolation factors for the $W + \text{jets}$ estimate derived for muons (a) and electrons (b) as a function of lepton p_{T} [63].	109
229			
230	5.13	Background composition in final VBF signal region [63].	110
231			
232	5.14	Variations in the top background extrapolation factor in the cut-based analysis due to PDF uncertainties, binned in m_{T}	112
233			
234	5.15	Variations in the top background extrapolation factor in the cut-based analysis due to QCD scale uncertainties, binned in m_{T}	112
235			
236	5.16	Post-fit distributions in the cut-based VBF analysis. Panel (a) shows the one-dimensional m_{T} distribution, while (b) shows the data candidates split into the bins of m_{T} and m_{jj} used in the final fit [63].	115
237			
238	5.17	Postfit distributions in the BDT VBF analysis [63].	116
239			
240	5.18	Overlap between cut-based and BDT VBF signal region candidates in the m_{jj} - m_{T} plane. .	117
241			
242	6.1	Post-fit m_{T} distribution in the $n_j \leq 1$ regions for the same flavor ($ee/\mu\mu$) final states [63]. .	120
243			
244	6.2	Post-fit m_{T} distribution in the $n_j \leq 1$ regions [63].	122
245			
246	6.3	Best fit signal strength $\hat{\mu}$ as a function of hypothesized m_H [63].	123
247			
248	6.4	Local p_0 as a function of m_H [63].	124
249			
246	6.5	Likelihood as a function of $\mu_{\text{VBF}}/\mu_{\text{ggF}}$ [63].	124
247			
248	6.6	Likelihood scan as a function of μ_{VBF} and μ_{ggF} [63].	125
249			
246	6.7	Likelihood scan as a function of κ_F and κ_V [63].	127
247			
248	6.8	Comparison of signal strength measurements in different Higgs decay channels on ATLAS [92]. .	129
249			
246	7.1	Parton luminosity ratios as a function of resonance mass M_X for 13/8 TeV and 7/8 TeV [93]. .	132
247			
248	7.2	Summary of HH branching ratios [94].	133
249			
246	7.3	Minimum ΔR between B decay vertices for different RSG masses in a $G_{\text{KK}}^* \rightarrow HH \rightarrow 4b$ sample with $c = 1$	134
247			

250	7.4 Trigger efficiency for events passing all signal region selections as a function of mass in	
251	$G_{\text{KK}}^* \rightarrow HH \rightarrow 4b$ samples with $c = 1$ [103]. In the trigger names, “j” refers to a jet	
252	or jets. “ht” refers to H_T , the scalar sum of transverse momenta in the event. “bloose”	
253	refers to a loose b -tagging requirement applied to the jet. “aor” refers to anti- k_T jets with	
254	$R = 1.0$. The numbers at the end are the thresholds on the given quantity in GeV.	136
255	7.5 Comparison of untrimmed and trimmed jet masses for large radius jets in a RSG sample	
256	with $m_{G_{\text{KK}}^*} = 1$ TeV. JES (JMS) refers to the standard jet energy (mass) scale calibration	
257	for ATLAS [58].	137
258	7.6 Efficiency of finding two b -jets from each Higgs in an RSG event using calorimeter jets	
259	with $R = 0.3$ or different track jet radii [107]	138
260	7.7 Illustration of the boosted selection requirements on Higgs candidates. Each large-radius	
261	calorimeter jet (Higgs candidate) must contain two track jets	140
262	7.8 Estimated significance as a function of signal mass for RSG $c = 1$ models in the 3 b (a) and	
263	4 b (b) regions for different b -tagging efficiency working points	141
264	7.9 Acceptance \times efficiency as a function of mass for (a) RSG and (b) narrow heavy scalar	
265	signal models [110].	142
266	7.10 Efficiency of requiring 3 or 4 b -tagged track jets vs. RSG mass. The efficiency quoted is	
267	relative to the previous selection requirements (rather than an absolute efficiency).	143
268	7.11 MV2c2o b -tagging efficiency for each of the four track jets in the boosted 4 b selection as a	
269	function of RSG mass for $c = 1$ models.	144
270	7.12 M_J^{sublead} vs. M_J^{lead} in a 2 b -tag data sample. The signal region is defined by the inner black	
271	contour ($X_{hh} < 1.6$) and the sideband region is defined by the outer contour ($R_{hh} >$	
272	35.8 GeV). The region between the black contours is the control region. The mass region	
273	which is enriched in $t\bar{t}$ background is also shown for illustration. [110]	145
274	7.13 An illustration of the data-driven background estimation technique for the boosted analysis	147
275	7.14 Leading large-R jet mass in the 3 b (a) and 4 b (b) sideband regions. The multijet and $t\bar{t}$	
276	backgrounds are estimated using the data-driven methods described above. Because their	
277	normalizations are derived in the sideband region, the total background normalization is	
278	constrained by default to match the normalization of the data [110].	148
279	7.15 Di-jet invariant mass ($M_{2,J}$) in the 3 b (a) and 4 b (b) control regions. The multijet and $t\bar{t}$	
280	backgrounds are estimated using the data-driven methods described above [110].	149
281	7.16 Di-jet invariant mass ($M_{2,J}$) in the 3 b (a) and 4 b (b) signal regions. The multijet and $t\bar{t}$	
282	backgrounds are estimated using the data-driven methods described above. In the 3 b re-	
283	gion, a graviton signal with $m_{G_{\text{KK}}^*} = 1.8$ TeV and $c = 1$ is overlaid, with the cross section	
284	multiplied by a factor of 50 so that the signal is visible. In the 4 b region, signals with	
285	$m_{G_{\text{KK}}^*} = 1.0$ TeV and $m_{G_{\text{KK}}^*} = 1.5$ TeV are overlaid, both with $c = 1$ and the yields	
286	multiplied by factors of 2 and 5 respectively [110].	154

287	8.1	Di-jet invariant mass (M_{2J}) in the resolved signal region. A graviton signal with $m_{G_{KK}^*} =$		
288		800 GeV and $c = 1$ is overlaid. [110].	157	
289	8.2	Expected and observed upper limit as a function of mass for G_{KK}^* in the RSG model with		
290		(a) $c = 1$ and (b) $c = 2$, as well as (c) H with fixed $\Gamma_H = 1$ GeV, at the 95% confidence		
291		level in the CL_s method. [110]	160	
292	9.1	Combined ATLAS and CMS measurements in Run 1 for (a) Higgs signal strength in gluon		
293		fusion and VBF and (b) Higgs couplings normalized to their SM predictions	164	
294	9.2	Discovery significance for RSG models at the HL-LHC in three different budget scenar-		
295		ios [117]. Systematic uncertainties on the background prediction (σ_B) of 2.5% and 5.0%		
296		are both tested.	165	
297	A.1	p_T of the leading track jet in the leading calorimeter jet for different signal masses in RSG		
298		$c = 1$ models	167	
299	A.2	MV _{2c2o} score for the leading track jet (a) and subleading track jet (b) of the leading calorime-		
300		ter jet for different signal masses in RSG $c = 1$ models	167	
301	A.3	IP _{3D} log-likelihood ratio ($\log(p_b/p_u)$) of the leading track jet in the leading calorimeter		
302		jet for different signal masses in RSG $c = 1$ models	168	
303	A.4	Mass (a) and number of tracks (b) for the secondary vertices computed with the SV1 algo-		
304		rithm. When no secondary vertex is found, the quantities are assigned to default negative		
305		values.	169	
306	A.5	Mass (a) and number of tracks (b) for vertices computed with the JetFitter algorithm.		
307		When no vertices are found, the quantities are assigned to default negative values.	169	
308	A.6	MV _{2c2o} score (a) and SV1 mass (b) for leading track jets with two truth b quarks ($n_{tb,lead} =$		
309		2) compared to those with only one truth b ($n_{tb,lead} = 1$).	170	
310	A.7	Track fit χ^2/n_{DOF} (a) and number of pixel detector hits (b) for the leading track of the		
311		leading track jet in different mass RSG $c = 1$ samples	171	
312	A.8	MV _{2c2o} score (a) and SV1 mass (b) for leading track jets whose leading track jet has at least		
313		four pixel hits ($N_{pix} \geq 4$) compared to those which do not ($N_{pix} < 4$).	172	

Listing of tables

314

315	1.1	Production cross sections for a 125 GeV Higgs boson at $\sqrt{s} = 8$ TeV with scale and PDF uncertainties [18].	13
316	1.2	Theoretical branching ratios for a 125 GeV Higgs boson, quoted as a percentage of the total width of the Higgs. Uncertainties shown are relative to the branching ratio value [18].	15
317	1.3	Possible channels for Higgs searches. Checkmarks denote the most sensitive production modes for each decay channel [5].	16
318	1.4	Production cross sections for pair production of a 125 GeV Higgs boson at $\sqrt{s} = 14$ TeV with total uncertainty [20]. The uncertainties include QCD scale and PDF variations as well as uncertainties on α_S	17
321	2.1	Evolution of LHC machine conditions [40, 41]	26
322	2.2	Performance requirements for the ATLAS detector [34].	37
323	2.3	Signal efficiencies for WH production with $H \rightarrow b\bar{b}$ and $H \rightarrow WW^* \rightarrow \mu\nu qq$ under different trigger configurations [47].	41
324	3.1	A summary of backgrounds to the $H \rightarrow WW^* \rightarrow \ell\nu\ell\nu$ signal	56
325	4.1	Monte carlo generators used to model signal and background for the Higgs search [1].	71
326	4.2	Normalization factors (ratio of data and MC yields in a control sample) for the Standard Model WW and top backgrounds in the $H \rightarrow WW^* \rightarrow \ell\nu\ell\nu$ analysis [1]. Only statistical uncertainties are shown.	74
327	4.3	Data and expected yields for signal and background in the final $H \rightarrow WW^* \rightarrow \ell\nu\ell\nu$ signal region. Uncertainties shown are both statistical and systematic. [1]	75
328	4.4	Summary of the expected and observed significance and measured signal strengths in the combined 7 and 8 TeV datasets for the Higgs discovery analysis [1].	79
337	5.1	Single lepton triggers used for electrons and muons. A logical “or” of the triggers listed for each lepton type is taken. Units are in GeV, and the i denotes an isolation requirement in the trigger.	85
338	5.2	Di-lepton triggers used for different flavor combinations. The two thresholds listed refer to leading and sub-leading leptons, respectively. The di-muon trigger only requires a single lepton at level-1.	85

343	5.3	Trigger efficiency for signal events and relative gain of adding a dilepton trigger on top of the single lepton trigger selection. The first lepton is the leading, while the second is the sub-leading. Efficiencies shown here are for the ggF signal in the $n_j = 0$ category but are comparable for the VBF signal.	86
344			
345	5.4	Monte Carlo samples used to model the signal and background processes [63].	87
346			
347	5.5	p_T dependent isolation requirements for muons. Muons are required to have the amount of calorimeter or track based cone sums be less than this fraction of their p_T	89
348			
349	5.6	p_T dependent requirements for electrons. Electrons are required to have the amount of calorimeter or track based cone sums be less than this fraction of their E_T	90
350			
351	5.7	Summary of event selection for the $n_j \geq 2$ VBF analysis in the 8 TeV cut-based analy- sis [63].	97
352			
353	5.8	Background composition after each requirement in the $n_j \geq 2$ VBF analysis in the 8 TeV cut-based analysis [63].	97
354			
355	5.9	Top normalization factors computed at each stage of the cut-based selection. Uncertainties are statistical only.	102
356			
357	5.10	Top normalization factors computed for each bin of O_{BDT} . Uncertainties are statistical only.	102
358			
359	5.11	$Z/\gamma^* \rightarrow \tau\tau$ correction factors for the VBF cut-based analysis. Uncertainties are statistical only.	105
360			
361	5.12	$Z/\gamma^* \rightarrow \ell\ell$ normalization factors for cut-based and BDT analyses. Uncertainties are statistical only.	106
362			
363	5.13	Systematic uncertainties for various processes in the cut-based VBF analysis, given in units of % change in yield. Values are given for the low m_{jj} signal region.	III
364			
365	5.14	Composition of the post-fit uncertainties (in %) on the total signal (N_{sig}), total back- ground (N_{bkg}), and individual background yields in the VBF analysis [63].	II4
366			
367	5.15	Event selection for the VBF BDT analysis. The event yields in (a) are shown after the pre- selection and the additional requirements applied before the BDT classification (see text). The event yields in (b) are given in bins in O_{BDT} after the classification [63].	II4
368			
369			
370			
371	6.1	Post-fit yields in ggF dedicated signal regions for the $ee/\mu\mu$ final states [63].	II9
372	6.2	All signal regions definitions input into final statistical fit [63].	II21
373	6.3	Post-fit yields in the both ggF and VBF dedicated signal regions with all lepton flavor final states combined [63].	II21
374			
375	7.1	Summary of requirements on objects used in the $X \rightarrow HH \rightarrow b\bar{b}b\bar{b}$ search	II39
376	7.2	Effect of boosted selection on data, RSG signal models, $t\bar{t}$, and $Z + \text{jets}$. The numbers from simulation are normalized with the MC generator cross section and do not take into account the data driven estimates described in section 7.6 [III].	II44
377			
378			

379	7.3	Mass region definitions used for background estimation	146
380	7.4	Parameters derived for exponential fit to background M_{2J} shape in the $3b$ and $4b$ signal	
381		regions [111]	148
382	7.5	The number of events in data and predicted background events in the boosted 3-tag and	
383		4-tag sideband and control regions. The uncertainties shown are statistical only. [110] . . .	149
384	7.6	Summary of systematic uncertainties in the total background and signal event yields (ex-	
385		pressed in %) in the boosted 3-tag and 4-tag signal regions. Systematic uncertainties on	
386		the signal normalization are shown for models with $m_{G_{KK}^*} = 1.5$ TeV and both $c = 1$	
387		and $c = 2$ as well as a narrow width heavy scalar.	151
388	7.7	Alternate fit functions used to model the M_{2J} distribution in the QCD multijet back-	
389		ground. In the equations, $x = M_{2J}/\sqrt{s}$	152
390	7.8	Observed yields in the 3-tag and 4-tag signal regions for the boosted analysis compared to	
391		the predicted number of background events Errors correspond to the total uncertainties	
392		in the predicted event yields. The yields for a graviton with $m_{G_{KK}^*} = 1$ TeV and $c = 1$	
393		are also shown. [110]	153
394	8.1	Observed yields in the resolve selection 4-tag signal region compared to the predicted num-	
395		ber of background events Errors correspond to the total uncertainties in the predicted	
396		event yields. The yields for a graviton with $m_{G_{KK}^*} = 800$ GeV and $c = 1$ are also	
397		shown. [110]	156

Acknowledgments

400 I have been a member of the Harvard ATLAS group for many years now, first as an undergraduate and
401 then as a graduate student. As a result, I have had the privilege of interacting with many amazing people
402 there over the years and have accumulated a large list of folks to thank.

403 First and foremost, I must thank the two people who have effectively been my academic parents since I
404 started in the Harvard group: João Guimarães da Costa and Melissa Franklin. Melissa Franklin and João
405 Guimarães da Costa. They have both been so important to both my academic and personal development
406 that I can't even put one before the other. João has been an excellent PhD advisor, showing me how to
407 look at the big picture and helping me navigate the sometimes complicated politics of ATLAS. He got me
408 started on my first projects with ATLAS as a young college sophomore before there was even beam in the
409 LHC (go cosmic ray muons!). He has also been a constant source of advice and support, even when we
410 have been on different continents. Melissa gave me my start in HEP as a summer student on CDF and
411 has been an unbelievable mentor throughout my time at Harvard. I still remember our weekly chalkboard
412 particle physics lessons after that first summer. She also graciously took me on as a co-advisee after João
413 moved on to his new position at IHEP. I am incredibly lucky to have had both of them as advisors.

414 Another mentor who was essential to my development as a graduate student is Paolo Giromini. His un-
415 canny knowledge and intuition about detectors is unmatched and I am very grateful to have had the chance
416 to work with him on the micromegas for the ATLAS New Small Wheel upgrade project. I owe essentially
417 all my practical knowledge about detectors (and building things in general) to him. I also appreciated his
418 unique sense of humor which made sometimes difficult tasks much easier to get through.

419 I am grateful to John Huth and Masahiro Morii for their helpful advice as the other professors in the
420 Harvard ATLAS group as well. I especially thank John for helping me get started on the micromegas
421 trigger project and being a great professor to TF particle physics for. Additionally I thank Howard Georgi
422 for serving as my third committee member and offering me feedback throughout my graduate career.

423 I also owe enormous thanks to Hugh Skottowe, the postdoc that I worked most closely with in my early
424 years as a graduate student. He was always able to help me through complicated tasks in everything from
425 writing code to understanding difficult physics concepts. I particularly enjoyed walking down to his office
426 in Palfrey at random times and talking through whatever problem I was tackling on that day.

427 Alex Tuna, the second postdoc that I worked closely with at Harvard, deserves great thanks as well. He
428 helped me push through to the end of my graduate career and offered great advice along the way.

429 Being at Harvard, I have seen an incredible array of graduate students graduate before me: Ben Smith,
430 Verena Martinez Outschoorn, Srivas Prasad, Michael Kagan, Giovanni Zevi Della Porta, Laura Jeanty,
431 Kevin Mercurio, William Spearman, and Andy Yen. I want to thank them all for showing me what a good
432 physicist looks like and for patiently answering my questions and offering insightful advice about physics
433 and life.

434 Getting through graduate school would not have been possible without the support and friendship of
435 the other students in our group. Thanks to Emma Tolley for geeking out with me about cool comput-
436 ing stuff, going to taste delicious beers with me, and helping start the Palfrey tradition of Taco Tuesdays.
437 Thanks to Brian Clark for being a great friend and housing companion both in Kirkland House and in our
438 tiny summer apartment in Geneva (and thanks to his partner Allison Goff for the same reasons!). Thanks
439 to Siyuan Sun for giving me my first aikido lesson and always being there for great conversations, big and
440 small. Tony (Baojia) Tong deserves special recognition for working with me on the $4b$ analysis and putting
441 up with my sometimes strange requests (and giving me rides to the Val Thoiry Migros so I wouldn't have
442 to pay exorbitant Geneva grocery prices!). Stephen Chan is probably the only student in the group who
443 both understands my references to the Sopranos and makes some of his own. To the younger graduate
444 students - Karri Di Petrillo, Jennifer Roloff, Julia Gonski, and Ann Wang - I want to say thank you for
445 making the group a fun and lively place to be and giving all of us energy that the older graduate students
446 like myself can sometimes lack.

447 I'd like to thank Annie Wei and Gray Putnam, the two undergraduates I have worked with as a graduate
448 student. Their unbelievable intuition and quickness in picking up difficult particle physics concepts is
449 inspiring.

450 I would also like to thank all of the postdocs that I have interacted with in my time in the Harvard group:
451 Kevin Black, Alberto Belloni (who would always ask me “Do you have it?”...I can now say that I do!),
452 Shulamit Moed, Corrinne Mills, Geraldine Conti, David Lopez Mateos, Chris Rogan, Valerio Ippolito,
453 and Stefano Zambito.

454 There are many people on ATLAS who have helped me get to this point as well. In the *WW* group, I
455 have to thank Jonathan Long, Joana Machado Miguens, Ben Cerio, Philip Chang, Bonnie Chow, Richard
456 Polifka, Heberth Torres, Tae Min Hong, and Jennifer Hsu for being wonderful colleagues and making the
457 entire analysis run smoothly. In the *4b* group, I have to thank Qi Zeng, Tony Tong, Alex Tuna, Michael
458 Kagan, Max Bellomo, John Alison, and Patrick Bryant.

459 Kirkland House was my home for the last three years of graduate school and was an wonderful envi-
460 ronment and support system. I want to thank my fellow tutors, especially Brian Clark and Allison Goff
461 (again), Zach Abel, Kelly Bodwin, Alex Lupsasca, John and Pam Park, Luke and Erin Walczewski, and
462 Philip Gant for their friendship and support. I also want to thank Kate Drizos Cavell, Bob Butler, and the
463 Faculty Deans Tom and Verena Conley.

464 There are still a few friends that haven’t been covered yet and deserve great thanks. Jake Connors and
465 Meredith MacGregor have been absolutely wonderful friends and I thank them in particular for the many
466 home-cooked meals and great conversations we’ve had in their apartment. Nihar Shah has been my friend
467 and confidant since we were both wee freshmen in Harvard Yard. Gareth Kafka, though he sits on the
468 “neutrino” side of Palfrey House, has made days there more fun and has also been an enthusiastic partici-
469 pant in the Palfrey Taco Tuesdays.

470 Being at Harvard necessarily means having to navigate through bureaucracy at some point or another.
471 I thank Lisa Cacciabuado, Carol Davis, and Jacob Barandes for always having open doors and being the
472 most kind, helpful people in the Physics department.

473 I thank Venky Narayananuriti for putting on a great SPU course that I was proud to be a part of and
474 TF for. I’d also like to thank Jim Waldo for offering me much advice about working in Computer Science
475 and giving me a fun data project to be a part of in my free time.

476 I grew up in a very tight knit Serbian community on the south side of Chicago which helped make me

477 the person I am today. I would like to thank all of the people at St. Simeon Mirotochivi Serbian Orthodox
478 Church who have always been sources of enthusiasm and support in my life.

479 I would not be here without the unconditional love and constant support and encouragement of my
480 family. To my pokojni Deda Branko and Miloje, my pokojni Baba Milka, and my Baba Desa, I want to
481 say thank you for instilling in me at an early age the love of curiosity and storytelling that I have carried
482 throughout my life. To my sister Angelina, I want to say thank you for always loving me and being my
483 partner in crime throughout our childhoods. To my parents, Miroljub and Nada, Tata and Mama, I really
484 cannot express how grateful I am to you and how much I owe you. As I look back now I see how I am a
485 combination of both of your best qualities and every day I am in situations where I understand more and
486 more the lessons you made sure to teach me and the sacrifices you made to make sure I got the best possible
487 education. I love you all.

488 Finally, I have to thank my soul mate, the one person in my life who understands me more than anyone
489 else, my fiancée Kelly Brock. You are my sounding board, my support system, my cheerleader (figuratively
490 and literally!), my best friend, my role model, and my everything. I would not have gotten through grad-
491 uate school without you and my life would not be the same without you. I cannot wait to start our new
492 lives together as the married doctors, tackling whatever comes our way with the same zeal with which we
493 tackled graduate school. I love you with all my heart and soul.

0

494

495

Introduction

496 The Higgs boson is often described as one of the cornerstones of particle physics. When the Standard
497 Model was first developed as a theory to describe the fundamental particles and forces of nature, physicists
498 were faced with a dilemma. The electroweak theory beautifully characterized both electromagnetism and
499 the weak force with a single underlying framework. However, the mass of the weak W and Z bosons
500 was puzzling given the fact that their electromagnetic counterpart, the photon, is massless. The Higgs
501 mechanism was developed as the leading theory for the origin of this electroweak symmetry breaking. It
502 predicted the existence of an additional spin-0 boson in the Standard Model, the Higgs boson. Generations
503 of collider experiments searched for this elusive particle. This dissertation presents research work on the
504 Higgs boson from its discovery to its use as a tool in the search for physics beyond the Standard Model
505 with the ATLAS detector at the Large Hadron Collider (LHC).

506 One of the first priorities for the LHC when it began colliding proton beams in 2010 was the search
507 for the Higgs boson. This search was initially tackled in the $H \rightarrow WW^* \rightarrow \ell\nu\ell\nu$ channel, followed by

508 the $H \rightarrow \gamma\gamma$ and $H \rightarrow ZZ^* \rightarrow 4\ell$ channels. Each channel has its own merits, but the WW^* mode is
509 particularly suited to searching over a wide range of masses. The $H \rightarrow WW^*$ branching ratio is large and it
510 is the primary decay channel above the $2m_W$ mass threshold. Despite the fact that the full Higgs invariant
511 mass cannot be reconstructed in the $H \rightarrow WW^* \rightarrow \ell\nu\ell\nu$ channel, its signal to background ratio makes
512 it ideal for measurement of Higgs properties such as the production cross section and couplings.

513 In 2012, the ATLAS and CMS experiments announced the discovery of a new particle consistent with
514 the Higgs boson [1, 2]. In ATLAS, this discovery was made with 4.8 fb^{-1} collected at $\sqrt{s} = 7 \text{ TeV}$
515 and 5.8 fb^{-1} at $\sqrt{s} = 8 \text{ TeV}$. This dissertation first presents the search for gluon fusion production
516 of the Higgs in the $H \rightarrow WW^* \rightarrow \ell\nu\ell\nu$ channel, which played an important role in this discovery.
517 Selection requirements which were optimized to maximize the discovery significance in this channel, as
518 well as background estimation procedures, are discussed.

519 After its discovery, interest in the Higgs shifted to focus on the measurement of its properties. As a result,
520 extensions of the initial discovery analysis in larger datasets had two main goals. Improvement of signal to
521 background ratio was important to allow for precision measurements. Also, searches for rarer production
522 modes of the Higgs were a priority. The first such extension presented in this dissertation is a tailored
523 selection for $\ell\nu\ell\nu$ final states with same flavor leptons. Novel variables for the reduction of the $Z+\text{jets}$
524 background that could remain robust under increasing LHC instantaneous luminosities are shown. The
525 second post-discovery result shown is the first evidence of Vector Boson Fusion (VBF) production of the
526 Higgs boson.

527 VBF production of the Higgs boson is particularly interesting in the $H \rightarrow WW^* \rightarrow \ell\nu\ell\nu$ final state.
528 In this combination of production and decay modes, the Higgs boson couples exclusively to W bosons,
529 allowing for precise measurement of the Higgs- W coupling constant. However, it is challenging to observe
530 VBF Higgs production because its cross section at the LHC is an order of magnitude lower than gluon
531 fusion production. The large $H \rightarrow WW^*$ branching ratio thus presents another advantage over other
532 final states. Additionally, VBF production of the Higgs boson creates two forward jets in addition to the
533 Higgs, and these jets can be used to isolate VBF Higgs events from other production modes. The VBF
534 $H \rightarrow WW^* \rightarrow \ell\nu\ell\nu$ analysis first created a selection requirement based signal region using variables

535 constructed specifically for the VBF Higgs production topology. This “cut-based” analysis is presented
536 in detail in this dissertation. These VBF topology variables, once validated in the cut-based analysis, were
537 then input into a multivariate boosted decision tree discriminant to achieve the first evidence of VBF Higgs
538 production with the full 20.3 fb^{-1} of $\sqrt{s} = 8 \text{ TeV}$ data in ATLAS.

539 After a two year shutdown, the LHC restarted in 2015 with a center of mass energy of $\sqrt{s} = 13 \text{ TeV}$.
540 This increase improved the LHC’s ability to probe for physics beyond the Standard Model, and the Higgs
541 sector remained one of the largest regions of unprobed phase space where such new physics could be dis-
542 covered. Production of high mass resonances benefited most from the center of mass energy increase. In
543 particular, the cross section for a generic gluon-initiated 2 TeV resonance increased tenfold with the in-
544 crease from 8 to 13 TeV. Therefore, a natural next step in studies of the Higgs was a search for a new
545 heavy resonance which decays into a pair of Higgs bosons. The final result shown in this dissertation is
546 a search for resonant di-Higgs production in the $X \rightarrow HH \rightarrow b\bar{b}b\bar{b}$ final state with 3.2 fb^{-1} recorded
547 by ATLAS at $\sqrt{s} = 13 \text{ TeV}$. This search has the unique advantage that it can both probe new physics
548 and gain further understanding of the Higgs potential through constraints on SM pair production of the
549 Higgs. It also extends the previous ATLAS results at $\sqrt{s} = 8 \text{ TeV}$ and probes higher mass resonances
550 that were not previously accessible. Additionally, it is an informative precursor to di-Higgs analyses at the
551 future High Luminosity LHC (HL-LHC), where a projected dataset of 3000 fb^{-1} at $\sqrt{s} = 14 \text{ TeV}$ will
552 begin to become sensitive to the SM Higgs self coupling.

553 As mentioned above, this dissertation begins by discussing the discovery of the Higgs and the role of
554 the $H \rightarrow WW^* \rightarrow \ell\nu\ell\nu$ channel. It then presents the first evidence for the VBF production mode using
555 the $H \rightarrow WW^* \rightarrow \ell\nu\ell\nu$ channel with the full ATLAS Run 1 dataset. It also shows the final combined
556 Run 1 measurements of gluon fusion Higgs production from this channel. Finally, it presents a search for
557 Higgs pair production in the $HH \rightarrow b\bar{b}b\bar{b}$ channel. It is organized into four parts.

558 Part 1 presents the theoretical and experimental background required for the subsequent parts. Chap-
559 ter 1 gives an overview of Higgs physics, particularly single and double Higgs production in the Standard
560 Model and beyond. Chapter 2 presents details regarding the Large Hadron Collider and the ATLAS experi-
561 ment. The evolution of machine conditions, descriptions of the ATLAS sub-detectors, and an overview of

object reconstruction in ATLAS are all shown. A brief interlude on the ATLAS Muon New Small Wheel upgrade is also given, as this upgrade has been a focus of my graduate work and will have an important impact on ATLAS' ability to study the Higgs at the High Luminosity LHC.

Part 2 discusses the observation and measurement of the Higgs in the $H \rightarrow WW^* \rightarrow \ell\nu\ell\nu$ channel in the ATLAS Run 1 dataset at $\sqrt{s} = 7$ and 8 TeV. Because I worked in this channel from before the discovery through to the final analysis of the Run 1 dataset, Part 2 is organized in such a way to allow easy presentation of multiple analyses on different subsets of the full Run 1 dataset. Chapter 3 presents a general overview of the $H \rightarrow WW^*$ analysis strategy and defines many of the variables and common elements used in the rest of Part 2. Chapter 4 presents the discovery and subsequent measurements of the Higgs boson, focusing on the role of the WW^* channel in this discovery. Chapter 5 presents the first evidence for the VBF production mode of the Higgs, a result from the WW^* channel in the full Run 1 ATLAS dataset. In this chapter, the focus is mainly on the cut-based VBF analysis. The cut-based analysis was an important first step to the final VBF result which used a boosted decision tree. Where appropriate, connections between the cut-based and BDT analyses are shown and their compatibility is discussed. Finally, the VBF analysis was an important input into the combined Run 1 $H \rightarrow WW^* \rightarrow \ell\nu\ell\nu$ result, which used both the gluon fusion and VBF channels in a combined fit to infer properties of the Higgs, including its couplings to the gauge bosons and its production cross section. This is the topic of Chapter 6.

Part 3 presents a search for Higgs pair production in the $HH \rightarrow b\bar{b}b\bar{b}$ channel. Chapter 7 presents an overview of this search in the boosted regime, where the Higgs pairs are the result of the decay of a heavy resonance. Chapter 8 shows the combined results between the boosted regime and the resolved regime, which is sensitive to lower mass resonances and non-resonant Higgs pair production. Finally, Part 4 presents a conclusion and brief outlook of future Higgs physics with ATLAS.

Part I

Theoretical and Experimental Background

In modern physics, there is no such thing as “nothing.”

Richard Morris

1

587

588

The Physics of the Higgs Boson

589 This chapter presents an overview of the Standard Model of Particle Physics and in particular the physics
590 of the Higgs boson. First, a brief overview of the Standard Model is presented. Then, a description of the
591 Higgs mechanism of electroweak symmetry breaking is given. Next, the physics of single Higgs boson
592 production and decay is described. The Standard Model also allows for production of two Higgs bosons
593 and this is detailed as well. Finally, di-Higgs production in two beyond the Standard Model (BSM) theories
594 - Randall-Sundrum gravitons (RSG) and Two Higgs Doublet Models (2HDM) - is shown.

595 **I.I THE STANDARD MODEL OF PARTICLE PHYSICS**

596 The Standard Model (SM) of Particle Physics is a quantum field theory describing the fundamental
597 particles of nature and the forces that govern their interactions. Several comprehensive pedagogical treat-
598 ments of the SM already exist in the literature [3–8] and this section will not rehash those. Rather, this
599 section presents a brief overview of the SM particles and forces in order to define them for subsequent

600 discussions.

601 The Standard Model consists of two primary categories of fundamental particles: fermions (spin 1/2
602 particles) and bosons (integer spin particles). The SM also describes three forces: electromagnetism, the
603 weak nuclear force, and the strong nuclear force. Gravity is not included in the theory and is largely irrele-
604 vant at the scales currently probed by collider experiments. Within the fermions, there are both quarks
605 (which interact via all three forces) and leptons. The charged leptons interact via electromagnetic and weak
606 interactions, while neutrinos (neutral leptons) interact only via the weak force. Within the bosons, there
607 are the W^\pm and Z bosons (the mediators of the weak force), the gluon (g , the mediator of the strong
608 force), and the photon (γ , the mediator of the electromagnetic force). Finally, there is the Higgs boson,
609 a fundamental spin zero particle resulting from the Higgs mechanism of electroweak symmetry breaking.

610 Figure 1.1 summarizes the fermions and bosons of the SM.




Figure 1.1: The particles of the Standard Model and their properties [6].

611 The Standard Model coalesced into a unified theoretical framework in the 1960s through the work
612 of Glashow, Weinberg, Salam, and others on the theory of electroweak interactions [9–12]. This theory
613 characterized both the electromagnetic and weak interactions as unified under a single gauge symmetry
614 group, namely $SU(2) \times U(1)$. At low enough energy scales (on the order of the W and Z masses), the
615 electroweak symmetry is broken, as evidenced by the fact that the weak bosons have mass while the photon
616 does not. The discovery of the Higgs boson in 2012 confirmed the Higgs mechanism as the most likely

617 candidate for this electroweak symmetry breaking [1, 2]. The complete SM consists of this electroweak
 618 theory combined with the theory of quantum chromodynamics (which models the strong sector as a non-
 619 Abelian $SU(3)$ gauge group) [13].

620 **I.2 ELECTROWEAK SYMMETRY BREAKING AND THE HIGGS**

621 In the Standard Model Lagrangian, it is difficult to include mass terms for the W and Z bosons without
 622 breaking the fundamental gauge symmetry of the Lagrangian. A traditional mass term does not preserve
 623 the $SU(2) \times U(1)$ symmetry. Additionally, scattering of massive W and Z bosons violate unitarity and
 624 these diagrams diverge at high energy scales. In the 1960s, Higgs, Brout, Englert, Guralnik, Kibble, and
 625 Hagen developed a mechanism for spontaneous symmetry breaking via the addition of a complex scalar
 626 doublet to the SM. Three of the four real degrees of freedom of this complex field would go to the lon-
 627 gitudinal modes of the W^\pm and Z , thus allowing them to have mass [14–17]. The remaining degree of
 628 freedom would manifest as an additional scalar, known now as the Higgs boson.

629 The mechanism works by introducing a Lagrangian for the newly introduced field that still respects the
 630 symmetry of the Standard Model inherently, but with a minimum at a non-zero vacuum expectation value
 631 for the field. In this minimum of the potential, the electroweak symmetry is broken. Specifically, consider
 632 a complex scalar doublet Φ with four degrees of freedom, as shown in equation I.1.

$$\Phi = \begin{pmatrix} \phi^+ \\ \phi^0 \end{pmatrix} = \frac{1}{\sqrt{2}} \begin{pmatrix} \phi_1^+ + i\phi_2^+ \\ \phi_1^0 + i\phi_2^0 \end{pmatrix} \quad (\text{I.1})$$

633 The simplest potential of a self-interacting Higgs that still respects the SM symmetry is given in equa-
 634 tion I.2.

$$V(\Phi) = \mu^2 \Phi^\dagger \Phi + \lambda (\Phi^\dagger \Phi)^2 \quad (\text{I.2})$$

635 If the μ^2 term of this potential is positive, then the potential has a minimum at $\Phi = 0$ and the electroweak

⁶³⁶ symmetry is preserved. However, if instead $\mu^2 < 0$, then the minimum is at a finite value of Φ , namely

$$\Phi_{\min} = \frac{1}{\sqrt{2}} \begin{pmatrix} 0 \\ v \end{pmatrix} \quad (1.3)$$

⁶³⁷ where $v = \sqrt{\mu^2/\lambda}$. Because this is the location of the minimum, it corresponds to the vacuum expecta-
⁶³⁸ tion value for the field ($\langle \Phi \rangle = \Phi_{\min}$). The excitations of the Higgs can then be parameterized as

$$\Phi = \frac{1}{\sqrt{2}} \begin{pmatrix} 0 \\ v + H \end{pmatrix} \quad (1.4)$$

⁶³⁹ The full scalar Lagrangian, including the kinetic term, is then given as

$$\mathcal{L}_s = (D^\mu \Phi)^\dagger (D_\mu \Phi) - V(\Phi) \quad (1.5)$$

⁶⁴⁰ where the covariant derivative is defined as

$$D_\mu = \partial_\mu + \frac{ig}{2} \tau^a W_\mu^a + ig' Y B_\mu \quad (1.6)$$

⁶⁴¹ and W^1, W^2, W^3 and B are the $SU(2)$ and $U(1)$ gauge fields of the electroweak theory, respectively. g
⁶⁴² and g' are the corresponding coupling constants. The Pauli matrices are represented with τ . With the
⁶⁴³ scalar Lagrangian in place, the physical gauge fields can then be written as

$$W_\mu^\pm = \frac{1}{\sqrt{2}} (W_\mu^1 \mp i W_\mu^2) \quad (1.7)$$

⁶⁴⁴

$$Z_\mu = \frac{-g' B_\mu + g W_\mu^3}{\sqrt{g^2 + g'^2}} \quad (1.8)$$

⁶⁴⁵

$$A_\mu = \frac{g B_\mu + g' W_\mu^3}{\sqrt{g^2 + g'^2}} \quad (1.9)$$

646 Equation 1.7 corresponds to the charged W^+ and W^- bosons, equation 1.8 corresponds to the neutral Z
 647 boson, and equation 1.9 corresponds to the neutral photon. The masses of the particles also arise from the
 648 Lagrangian. The photon has zero mass, while the masses of the W and Z bosons are given in equation 1.10.

649

$$\begin{aligned} M_W^2 &= \frac{1}{4}g^2v^2 \\ M_Z^2 &= \frac{1}{4}(g^2 + g'^2)v^2 \end{aligned} \quad (1.10)$$

650 The fermion masses also arise through a coupling with the Higgs via the Yukawa interaction (for a detailed
 651 description, see [8]). In this case the coupling between the Higgs and the fermions goes as

$$g_{hf\bar{f}} = \frac{m_f}{v} \quad (1.11)$$

652 The full Lagrangian of Higgs interactions can be written as

$$\mathcal{L}_{\text{Higgs}} = -g_{hf\bar{f}}\bar{f}fh + \frac{g_{hhh}}{6}h^3 + \frac{g_{hhhh}}{24}h^4 + \delta_V V_\mu V^\mu \left(g_{hVV}H + \frac{g_{hhVV}}{2}h^2 \right) \quad (1.12)$$

653 with

$$\begin{aligned} g_{hVV} &= \frac{2m_V^2}{v} & g_{hhVV} &= \frac{2m_V^2}{v^2} \\ g_{hhh} &= \frac{3m_h^2}{v} & g_{hhHH} &= \frac{3m_h^2}{v^2} \end{aligned} \quad (1.13)$$

654 The last term of the Lagrangian appears twice, once for W bosons and once for Z bosons. V refers to
 655 the W^\pm and Z , and $\delta_W = 1$ while $\delta_Z = 1/2$. Phenomenologically, there are a few features of this
 656 Lagrangian that are useful to note. First, note that the Higgs mass is a free parameter of the theory that
 657 must be determined experimentally. Second, note that the coupling of the Higgs to the vector bosons and
 658 fermions scales as a function of the masses of these particles, a fact that is important when considering
 659 both the production and decays of the Higgs. Finally, note the presence of the cubic and quartic Higgs self
 660 interaction terms, which can lead to final states with multiple Higgs bosons produced.

661 1.3 HIGGS BOSON PRODUCTION AND DECAY

662 This section discusses the properties of Higgs production and decay mechanisms. The details presented
663 here will focus on the properties of a 125 GeV Higgs boson, as this is the mass closest to that of the newly
664 discovered Higgs.

665 1.3.1 HIGGS PRODUCTION

666 The Higgs is produced by four main production modes at the Large Hadron Collider - gluon-gluon
667 fusion (ggF), vector boson fusion (VBF), associated production with a W or Z boson, or associated pro-
668 duction with top quarks ($t\bar{t}H$). Figure 1.2 shows the Feynman diagrams for these four modes.




Figure 1.2: The four most common Higgs boson production modes at the LHC: (a) gluon-gluon fusion, (b) vector boson fusion, (c) $W/Z + H$ production, (d) $t\bar{t}H$ production

669 In gluon-gluon fusion, gluons from the incoming protons fuse via a top-quark loop to produce a Higgs.
670 The top quark is the dominant contribution in the loop due to its heavy mass and the fact that the Higgs-
671 fermion coupling constant scales with fermion mass. In vector boson fusion, the incoming quarks each
672 radiate a W or Z boson which fuse to produce the Higgs. This production mode results in a final state
673 with a Higgs boson and two additional jets which tend to be forward because they carry the longitudinal

674 momentum of the incoming partons. The Higgs can also be produced in association with a W or Z boson.
 675 The W/Z is produced normally and then radiates a Higgsⁱ. Finally, the Higgs can be produced in associa-
 676 tion with two top quarks. Each incoming gluon splits into a $t\bar{t}$ pair, and one of the top pairs combines to
 677 create a Higgs. Figure 1.3 shows the production cross section for a 125 GeV Higgs boson in each of these
 modes at a pp collider as a function of center of mass energy. In figure 1.3, note that gluon fusion has the




Figure 1.3: Higgs production cross sections as a function of center of mass energy (\sqrt{s}) at a pp collider [18].

678
 679 largest cross section, while VBF is the second largest at approximately a factor of 10 smaller. The figure also
 680 includes the less commonly studied $b\bar{b}H$ and tH modes. While the $b\bar{b}H$ mode has a larger cross section
 681 than $t\bar{t}H$, it also has larger backgrounds and is thus less sensitive. The tH mode is not as sensitive as $t\bar{t}H$
 682 due to its lower cross section. At $\sqrt{s} = 8$ TeV, ggF production of a 125 GeV Higgs has a cross section of
 683 $19.47^{+1.54}_{-1.67}$ pb, while VBF has a cross section of $1.601^{+0.036}_{-0.035}$ pb [18]. The cross sections of all of the main
 684 Higgs production modes at this center of mass energy, as well as their uncertainties from varying the QCD
 685 renormalization and factorization scales and PDFs, are summarized in table 1.1 for a 125 GeV Higgs. The
 686 relative uncertainty of the gluon fusion mode is larger than the relative uncertainty in the vector boson

ⁱThis mode is also sometimes known as “Higgs-strahlung”.

687 fusion mode due to the fact that gluon fusion production happens through a loop.

Production mode	σ (pb)	QCD scale uncert. (%)	PDF + α_s uncert. (%)
Gluon fusion	19.47	+7.3 / - 8.0	3.1
Vector boson fusion	1.601	+0.3 / - 0.2	2.2
WH	0.7026	+0.6 / - 0.9	2.0
ZH	0.4208	+2.9 / - 2.4	1.7
bbH	0.2021	+20.7 / - 22.3	
$t\bar{t}H$	0.1330	+4.1 / - 9.2	4.3
tH (t -channel)	0.01869	+7.3 / - 16.5	4.6
tH (s -channel)	1.214×10^{-3}	+2.8 / - 2.4	2.8

Table 1.1: Production cross sections for a 125 GeV Higgs boson at $\sqrt{s} = 8$ TeV with scale and PDF uncertainties [18].

688 1.3.2 HIGGS BRANCHING RATIOS

689 The fact that the Higgs couples more strongly to more massive particles is crucial for understanding its
690 branching ratios. The width for Higgs decays to fermions is given by equation 1.14 [5].

$$\Gamma(H \rightarrow f\bar{f}) = \frac{N_c \sqrt{2} G_F m_f^2 m_H}{8\pi} \quad (1.14)$$

691 In this case, N_c is the number of colors, G_F is the Fermi constant, m_f is the mass of the fermion, and
692 m_H is the mass of the Higgs. Note that the width scales with the square of the fermion mass. (This also
693 assumes that the Higgs mass is large enough to decay with both the fermions on shell.)

694 The decay width to WW , in the case where both W bosons are produced on shell ($m_H \geq 2m_W$), is
695 given in equation 1.15 [5].

$$\Gamma(H \rightarrow W^+ W^-) = \frac{\sqrt{2} G_F M_W^2 m_H}{16\pi} \frac{\sqrt{1-x_W}}{x_W} (3x_W^2 - 4x_W + 4) \quad (1.15)$$

696 where m_W is the mass of the W and $x_W = 4M_W^2/m_H^2$. To get the branching ratio to ZZ (in the regime
697 where $m_H \geq 2m_Z$), the equation is divided by 2 to account for identical particles in the final state, and

⁶⁹⁸ x_W is replaced with $x_Z = 4M_Z^2/m_H^2$. This is shown in equation 1.16 [5].

$$\Gamma(H \rightarrow ZZ) = \frac{\sqrt{2}G_F M_Z^2 m_H}{32\pi} \frac{\sqrt{1-x_Z}}{x_Z} (3x_Z^2 - 4x_Z + 4) \quad (1.16)$$

⁶⁹⁹ The more general formula for Higgs branching into WW or ZZ , taking into account the case where one
⁷⁰⁰ or both vector bosons is off-shell, is shown in equation 1.17 [19].

$$\Gamma(H \rightarrow V^*V^*) = \frac{1}{\pi^2} \int_0^{M_H^2} \frac{dq_1^2 M_V \Gamma_V}{(q_1^2 - M_V^2)^2 + M_V^2 \Gamma_V^2} \int_0^{(M_H - q_1)^2} \frac{dq_2^2 M_V \Gamma_V}{(q_2^2 - M_V^2)^2 + M_V^2 \Gamma_V^2} \Gamma_0 \quad (1.17)$$

⁷⁰¹ Here, q_1^2 and q_2^2 are the invariant masses of the virtual gauge bosons, M_V is the W or Z mass, and Γ_V is
⁷⁰² the W or Z width. Γ_0 is the squared matrix element, which is given in equation 1.18 [19].

$$\Gamma_0 = \frac{G_F M_H^3}{8\sqrt{2}\pi} \delta_V \sqrt{\lambda(q_1^2, q_2^2, M_H^2)} \left[\lambda(q_1^2, q_2^2, M_H^2) + \frac{12q_1^2 q_2^2}{M_H^4} \right] \quad (1.18)$$

⁷⁰³ The function λ is defined as $\lambda(x, y, z) = (1 - x/z - y/z)^2 - 4xy/z^2$. The integral in the general
⁷⁰⁴ off-shell boson case is much more difficult to interpret than the simpler on-shell branching ratios, but it
⁷⁰⁵ can be evaluated numerically. These branching ratio formulas can also be visualized as a function of Higgs
⁷⁰⁶ mass, as shown in figure 1.4. There are a few interesting features to note in this figure. First, note that at
⁷⁰⁷ high Higgs masses, once on-shell production of both W and Z bosons is possible, these two decays are
⁷⁰⁸ dominant due to the large masses of the W/Z . Also note that the branching ratio to W s is twice that of
⁷⁰⁹ Z s at these large masses due to the fact that there are two charged W bosons (W^\pm) and only one Z boson².
⁷¹⁰ At 125 GeV, the Higgs is accessible through many different decay modes. The largest branching ratio is
⁷¹¹ the decay $H \rightarrow b\bar{b}$ at 58.24% [18]. This branching is larger than the WW/ZZ decays because one of
⁷¹² the two bosons must be produced off-shell for $m_h = 125$ GeV. The second largest branching ratio is
⁷¹³ to WW^* at 21.37 % (before taking into account the branching ratios of the W). Table 1.2 summarizes
⁷¹⁴ the theoretical branching ratios for a Higgs with a mass of 125 GeV. Note that there is a Higgs branching
⁷¹⁵ ratio to $\gamma\gamma$ even though photons are massless. This decay happens through a loop, which suppresses the

²In the Higgs Lagrangian, this extra symmetry factor is quantified by the δ_V noted in equation 1.12.




Figure 1.4: Higgs boson branching ratios as a function of m_H [18].

⁷¹⁶ branching ratio³.

Decay	Branching ratio (%)	Relative uncertainty (%)
bb	58.24	+0.25 / - 0.25
WW^*	21.37	+0.99 / - 0.99
gg	8.187	+3.40 / - 3.41
$\tau\tau$	6.272	+1.17 / - 1.16
cc	2.891	+1.20 / - 1.20
ZZ^*	2.619	+0.99 / - 0.99
$\gamma\gamma$	0.2270	+1.73 / - 1.72
$Z\gamma$	0.1533	+5.71 / - 5.71
$\mu\mu$	0.02176	+1.23 / - 1.23

Table 1.2: Theoretical branching ratios for a 125 GeV Higgs boson, quoted as a percentage of the total width of the Higgs. Uncertainties shown are relative to the branching ratio value [18].

⁷¹⁷ Note that the branching ratios alone do not tell the full story of which Higgs channels are the most
⁷¹⁸ sensitive. For example, the $H \rightarrow b\bar{b}$ channel in gluon fusion production is incredibly difficult to observe
⁷¹⁹ due to the large QCD dijet background at the LHC. However, in associated production of the Higgs,

³The largest contributions to the loop are the top quark and W boson.

720 where a W or Z gives additional final state particles that can be used to reduce background, a search for
 721 $H \rightarrow b\bar{b}$ can be sensitive. The combinations of production and decay modes that are most commonly
 722 studied at the LHC are summarized in table 1.3 [5].

Decay	Inclusive (incl. ggF)	VBF	WH/ZH	$t\bar{t}H$
$H \rightarrow \gamma\gamma$	✓	✓	✓	✓
$H \rightarrow bb$			✓	✓
$H \rightarrow \tau^+\tau^-$		✓		
$H \rightarrow WW^* \rightarrow \ell\nu\ell\nu$	✓	✓	✓	
$H \rightarrow ZZ \rightarrow 4\ell$	✓			
$H \rightarrow Z\gamma \rightarrow \ell\ell\gamma$	very low			

Table 1.3: Possible channels for Higgs searches. Checkmarks denote the most sensitive production modes for each decay channel [5].

723 1.4 HIGGS PAIR PRODUCTION IN THE STANDARD MODEL

724 The Standard Model also allows for processes that produce two Higgs bosons in the final state, known
 725 as Higgs pair production or di-Higgs production. The two main production mechanisms are shown in
 figure 1.5. The two diagrams in figure 1.5 interfere destructively with one another, resulting in a low overall




Figure 1.5: The two leading diagrams for Standard Model di-Higgs production at the LHC: (a) box diagram, (b) Higgs self coupling.

726 cross section for di-Higgs production at the LHC. Nevertheless, Higgs pair production is quite interesting
 727 to study because it gives direct access to the λ parameter of the Higgs potential, also known as the Higgs
 728 self coupling. The diagram in figure 1.5(b) is sensitive to this coupling through the triple Higgs vertex.

730 One can substitute the gluon fusion production of diagram 1.5(b) with any of the other production
 731 modes previously discussed. These other modes do not suffer from interference with the box diagram in
 732 figure 1.5(a) due to the presence of additional particles in the final state. They still have a lower cross section
 733 than the gluon fusion mode, however. The cross sections for di-Higgs production in the different modes,
 734 as well as their uncertainties, are shown in table 1.4 [20]. These are shown for $\sqrt{s} = 14$ TeV as this is the
 735 expected center of mass energy for the High Luminosity LHC and this energy is more sensitive to di-Higgs
 production. Note that the scale of cross section quoted is now in fb rather than pb.

Production mode	σ (fb)	Total uncert. (%)
Gluon fusion	33.89	+37.2 / - 27.8
Vector boson fusion	2.01	+7.6 / - 5.1
$W H H$	0.57	+3.7 / - 3.3
$Z H H$	0.42	+7.0 / - 5.5
$t \bar{t} H$	1.02	-

Table 1.4: Production cross sections for pair production of a 125 GeV Higgs boson at $\sqrt{s} = 14$ TeV with total uncertainty [20]. The uncertainties include QCD scale and PDF variations as well as uncertainties on α_S .

736

737 1.5 HIGGS PAIR PRODUCTION IN THEORIES BEYOND THE STANDARD MODEL

738 The Higgs pair production cross section in the Standard Model is rather small, and datasets on the
 739 scale of the full 3000 fb^{-1} expected from the High Luminosity LHC will be required to obtain sensitive
 740 measurements of the Higgs self-coupling [20]. However, the discovery of the Higgs also gives particle
 741 physicists a new tool that can be exploited in the search for new physics beyond the Standard Model. In
 742 particular, Higgs pair production is a promising channel in the search for new physics. The cross section for
 743 di-Higgs production can be altered through both resonant and non-resonant production of Higgs pairs. In
 744 non-resonant production, di-Higgs production vertices can arise from the presence of a new strong sector
 745 and additional colored particles [21–23]. Figure 1.6 shows examples of the types of vertices that can arise. In
 746 the resonant case, new heavy particle can decay to Higgs pairs. Such new particles can include heavy Higgs
 747 bosons arising in two Higgs doublet models (2HDM) or Higgs portal models as well as heavy gravitons in
 748 Randall-Sundrum theories [21, 24–30]. Figure 1.7 shows a generic diagram for a heavy resonance decaying

⁷⁴⁹ to two Higgs bosons. In the 2HDM, X corresponds to the heavy CP-even scalar H . In the Randall-Sundrum model, X corresponds to a heavy spin-2 graviton G_{KK}^* . The next sections provide more detail




Figure 1.6: Diagrams with new vertices for non-resonant Higgs pair production arising in composite Higgs models.

⁷⁵⁰ on the phenomenology of resonant Higgs production in Randall-Sundrum and 2HDM models, as these models will later be tested in a dedicated search for resonant production of boosted Higgs pairs.




Figure 1.7: Generic Feynman diagram for resonant Higgs pair production in BSM theories.

⁷⁵²

⁷⁵³ 1.5.1 RANDALL-SUNDRUM GRAVITONS

⁷⁵⁴ The Randall-Sundrum model is a proposed solution to the hierarchy problem that posits a five-dimensional
⁷⁵⁵ warped spacetime that contains two branes: one where the force of gravity is very strong and a second brane
⁷⁵⁶ at the TeV scale corresponding to the known Standard Model sector [24]. In the theory, the branes are
⁷⁵⁷ weakly coupled and the graviton probability function drops exponentially going from the gravity brane
⁷⁵⁸ to the SM brane, rendering gravity weak on the SM brane. The experimental consequence of this theory

759 is a tower of widely spaced (in mass) Kaluza-Klein graviton resonances. In theories where the fermions
 760 are localized to the SM brane, production of gravitons from fermion pairs is suppressed and the primary
 761 mode of production is gluon fusion [25]. These gravitons have a substantial branching fraction to Higgs
 762 pairs, ranging from 6.43% for gravitons with a mass of 500 GeV to 7.66% at 3 TeV. Figure 1.8 shows the
 763 branching ratios of the spin-2 Randall Sundrum graviton (RSG) as a function of its mass. The predomi-
 764 nant decays are to $t\bar{t}$ above the mass threshold for that channel.




Figure 1.8: Branching ratios for a spin-2 Randall-Sundrum graviton as a function of mass computed in MadGraph with the CP3-Origins implementation [25, 31, 32].

765 Randall-Sundrum models have two free parameters - the mass of the graviton and a curvature parameter
 766 k . Typically, rather than k , the theory is parameterized using $c \equiv k/\bar{M}_{\text{pl}}$, where \bar{M}_{pl} is the reduced
 767 Planck mass. The cross section for production of the RSG decreases as a function of mass and is strongly
 768 dependent on the gluon PDF. The increase in center of mass energy from 8 to 13 TeV in LHC Run 2
 769 greatly increases the cross section at higher mass. Figure 1.9 shows the cross section as a function of graviton
 770 mass at $\sqrt{s} = 13$ TeV for RSG models with $c = 1.0$ and $c = 2.0$.

771 Another interesting feature of the theory is that the width of the graviton increases with both c and
 772 $m_{G_{KK}^*}$. Figure 1.10 shows the graviton width for both $c = 1.0$ and $c = 2.0$ as a function of mass. In
 773 $c = 1.0$, the width starts at 8.365 GeV for a mass of 300 GeV and increases to 187.2 GeV at a mass of
 774 3 TeV. Similarly, with $c = 2.0$, the width starts at 33.46 GeV for $m_G = 300$ GeV and increases to




Figure 1.9: $\sigma \times \text{BR}(HH)$ for Randall-Sundrum gravitons as a function of mass computed in MadGraph with the CP3-Origins implementation [25, 31, 32].

775 748.8 GeV at a mass of 3 TeV.




Figure 1.10: Randall-Sundrum graviton width as a function of mass computed in MadGraph with the CP3-Origins implementation [25, 31, 32]

776 1.5.2 TWO HIGGS DOUBLET MODELS

777 In Two Higgs Doublet Models (2HDM), a second complex scalar doublet is added to the Standard
778 Model [27–29]. In this case, all four degrees of freedom in the second doublet correspond to new particles,
779 meaning that there are five total scalars from the two Higgs doublets - h (light CP-even Higgs), H (heavy
780 CP-even Higgs), A (heavy CP-odd Higgs), and H^\pm (charged Higgs). The model is parameterized by two
781 main parameters. The first, $\tan \beta \equiv \frac{v_2}{v_1}$, is the ratio of the vacuum expectation values of the two Higgs
782 doublets (where v_1 corresponds to the v in the SM Higgs model described above). The second parameter
783 is α , a mixing angle between the heavy and light Higgs fields. Models are also often parameterized with
784 $\cos(\beta - \alpha)$ rather than α directly. The limit where $\cos(\beta - \alpha) = 0$ is called the alignment limit, and in
785 this limit the light Higgs h has the same couplings as a Standard Model Higgs.

786 2HDM models are usually separated into two main types - Type I and Type II. In Type I models, the
787 charged fermions only couple to the second Higgs doublet, leading to a fermiophobic light Higgs. In
788 Type II models, up-type quarks couple to the first doublet while down-type quarks couple to the second
789 doublet. One specific realization of a Type II 2HDM is the Minimal Supersymmetric Standard Model
790 (MSSM).

791 Resonant di-Higgs production in 2HDM models can proceed through decays of the heavy CP-even
792 Higgs $H \rightarrow hh$. The branching ratio for $H \rightarrow hh$ depends on the model type as well as the values of
793 $\tan \beta$ and $\cos \beta - \alpha$. Figure 1.II shows the branching ratios as a function of the mass of the heavy scalar
794 H for both Type I and Type II models. Depending on the type of model hh can be a substantial fraction
795 of the decays of H .

796 1.6 CONCLUSION

797 Studying the Higgs sector is essential for understanding the details of how mass arises in the Standard
798 Model and how the electroweak symmetry is broken. The discovery of the Higgs boson also opens the
799 door for its use as a tool to search for new physics, and Higgs pair production is an ideal candidate for
800 this study. Even if no BSM physics is found in Higgs pair production, searches for Higgs pairs will put
801 constraints on the Higgs self coupling and thus improve knowledge of the Standard Model and the details




Figure 1.II: Branching ratios for heavy Higgs H in Type I (left) and Type II (right) 2HDM models with $\tan \beta = 1.5$ and $\cos(\beta - \alpha) = 0.1$ (0.01) for Type I (Type II) [29].

of the Higgs potential.

*The enthusiasm and motivation to explore particle physics
at the high-energy frontier knows no borders between the
nations and regions of the planet.*

Peter Jenni

2

803

804

805

The ATLAS detector and the Large Hadron Collider

806 This chapter presents an overview of the experimental systems used to conduct the measurements pre-
807 sented in this thesis. First, a brief overview of the accelerator, the Large Hadron Collider, will be given.
808 In this section, the accelerator conditions relevant to data-taking are presented as well. Next, an overview
809 of the ATLAS experiment is given. The basics of each sub-detector's role are summarized, as well as the
810 details of the datasets accumulated. Then, a brief interlude on the ATLAS Muon New Small Wheel up-
811 grade is presented. While this new detector does not have a direct impact on any of the datasets taken so
812 far, it will have an impact on future analyses and the work done on it is briefly summarized here. Finally,
813 an overview of object reconstruction in ATLAS is given. While the details of all of the algorithms will not
814 be presented in detail, aspects of the reconstruction performance such as object resolutions are shown as
815 these are relevant to the two studies presented later in this thesis.

816 2.1 THE LARGE HADRON COLLIDER

817 The Large Hadron Collider (LHC) is a proton-proton collider at the CERN laboratory in Geneva,
818 Switzerland [33]. It is designed for a maximum collision center of mass energy of $\sqrt{s} = 14$ TeV and has a
819 circumference of 26.7 kilometers. Four main experiments are located at the interaction points (IP) of the
820 accelerator: ATLAS (A Toroidal LHC Apparatus), CMS (the Compact Muon Solenoid), ALICE (A Large
821 Ion Collider Experiment), and LHCb [34–37]. The studies performed in this thesis were all completed
822 with the ATLAS detector. Figure 2.1 shows a schematic of the LHC ring and the various experiments.




Figure 2.1: A schematic view of the LHC ring [38]

823 One of the most interesting features of the LHC is in its magnet design. Because the tunnel does not
824 have room for separate superconducting magnets for each of the beam pipes, the LHC employs a twin-bore
825 magnet design. Each magnet must hold an 8.3 Tesla magnetic field in order to bend the proton beams at
826 $\sqrt{s} = 14$ TeV. The superconducting magnets are cooled to a temperature of 1.9 Kelvin with superfluid

827 helium.

828 2.1.1 INSTANTANEOUS LUMINOSITY

829 The rate of physics events expected from the accelerator is dependent on the instantaneous luminosity
830 of the machine and the cross section of the physics process, $R_{\text{events}} = L\sigma$. Here, R_{events} is the number
831 of events per second, L is the instantaneous luminosity of the machine, and σ is the cross section for the
832 physics process being measured. The instantaneous luminosity of the LHC is determined by numerous
833 factors related to machine conditions. Equation 2.1 gives the equation for instantaneous luminosity of
834 Gaussian beam profile [38].

$$L = \frac{N_b^2 n_b f_{\text{rev}} \gamma_r}{4\pi \epsilon_n \beta^*} F \quad (2.1)$$

835 The LHC collides protons in bunches, and in the above equation N_b is the number of protons per bunch
836 while n_b is the number of bunches per beam. Nominally, the LHC can hold up to 2808 proton bunches.
837 f_{rev} is the revolution frequency. ϵ_n is the normalized transverse beam emittance, a measurement of the
838 average spread of the particles position-momentum space which has the dimension of length. β^* is the
839 value of the *beta* function for the beam at the interaction point. It relates the emittance to the Gaussian
840 width of the beam with $\sigma_{\text{beam}} = \sqrt{\epsilon \cdot \beta}$. F is a reduction factor that corrects for the fact that the beams
841 are colliding at an angle at the IP.

842 Another way of writing the instantaneous luminosity is shown in equation 2.2. In this case, the instanta-
843 neous luminosity is written as the ratio of the rate of inelastic collisions with the inelastic cross section [39].

844

$$L = \frac{R_{\text{inel}}}{\sigma_{\text{inel}}} = \frac{\mu n_b f_{\text{rev}}}{\sigma_{\text{inel}}} \quad (2.2)$$

845 In this case, μ is the average number of interactions per bunch crossing in the accelerator. μ is a use-
846 ful parameter for characterizing the amount of activity recorded in an experiment. As the instantane-
847 ous luminosity and thus μ increase, there are more interactions per bunch crossing and more activity in the
848 detector. This is often characterized with $\langle \mu \rangle$, the measured per bunch crossing μ value averaged over all
849 bunch crossings. The interactions inside each bunch crossing that are not the main physics process of in-

850 terest are often referred to as “pileup” interactions, and $\langle \mu \rangle$ is a measurement of the level of pileup in the
851 detector.

852 **2.1.2 EVOLUTION OF MACHINE CONDITIONS**

853 This thesis uses datasets taken at three different center of mass energies: $\sqrt{s} = 7\text{ TeV}$ data taken in the
854 year 2011, $\sqrt{s} = 8\text{ TeV}$ data taken in the year 2012, and $\sqrt{s} = 13\text{ TeV}$ data taken in the year 2015. In
855 addition to increasing center of mass energy, the instantaneous luminosity and parameters that determine
856 it were evolving. Table 2.1 summarizes that machine conditions in each of these datasets.

	2011	2012	2015	Design
$\sqrt{s} [\text{TeV}]$	7	8	13	14
Number of bunches	1380	1380	1825	2808
Max. protons per bunch	1.45×10^{11}	1.7×10^{11}	1.2×10^{11}	1.15×10^{11}
Bunch spacing [ns]	50	50	25	25
Max. instantaneous luminosity [$\text{cm}^{-2}\text{s}^{-1}$]	3.7×10^{33}	7.7×10^{33}	5×10^{33}	10^{34}
$\beta^* [\text{m}]$	1.0	0.6	0.8	0.55
$\langle \mu \rangle$	11.6	20.7	13.7	-

Table 2.1: Evolution of LHC machine conditions [40, 41]

857 **2.2 THE ATLAS DETECTOR**

858 The ATLAS detector is a multi-purpose particle detector experiment at the LHC’s Point 1 [34]. It has
859 nearly 4π coverage in solid angle around the interaction point. It consists of an inner detector for mea-
860 suring charged particles, electromagnetic and hadronic calorimeters, and a muon spectrometer. Figure 2.2
861 gives an overview of the detector.

862 **2.2.1 COORDINATE SYSTEM**

863 Before defining the properties of the individual detectors, it is important to establish the coordinate
864 system used. Figure 2.3 shows a schematic of the coordinate system. The azimuthal plane (perpendicular
865 to the beam line) is defined as the x - y plane. The angle in this plane is referred to as ϕ . The angle relative




Figure 2.2: A full diagram of the ATLAS detector [34]

866 to the beam axis is referred to as θ . Rather than using θ directly as a coordinate, the experiment often uses
 867 the pseudorapidity η . η is defined in equation 2.3.

$$\eta = -\ln \left(\tan \left(\frac{\theta}{2} \right) \right) \quad (2.3)$$

868 Pseudorapidity is the massless approximation of rapidity, the angle used to parameterize boosts in spe-
 869 cial relativity. This is important for two reasons. First, it means that differences in η are Lorentz invariant.
 870 Second, particle production is roughly constant in pseudorapidity. Particles with η close to zero are re-
 871 ferred to as “central”, while those at high $|\eta|$ are called “forward”. In general, two main detector topologies
 872 can be seen in figure 2.2. There are “barrel” elements, which surround the beam line cylindrically and are
 873 in the central region of the detector. In the forward region, there are “endcap” regions which are arranged
 874 as disks perpendicular to the beam line.




Figure 2.3: The ATLAS coordinate system

875 **2.2.2 INNER DETECTOR**

876 The ATLAS Inner Detector (ID) system is built for precision tracking of charged particles. It covers
 877 the range $|\eta| < 2.5$. In this range, approximately 1000 particles are generated every bunch crossing in the
 878 detector. This requires having fine granularity to achieve the resolutions required for good momentum
 879 measurement and vertex reconstruction.

880 The ID consists of three sub-components: the pixel detector, semiconductor tracker (SCT), and trans-
 881 sition radiation tracker (TRT). It is surrounded by a solenoid providing a 2 T axial magnetic field which
 882 bends particles in the transverse plane to allow for momentum measurement. Figure 2.4 shows the layout
 883 of each of these components.

884 **PIXEL DETECTOR**

885 The pixel detector is the first detector particles traverse after being generated in proton collisions and
 886 is the most granular detector. Its operation is crucial for precision tracking and vertex reconstruction as



Figure 2.4: Layout of the ATLAS Inner Detector system [42]

well as higher level object reconstruction like tagging of jets from b -quarks. The basic sensing element in this subdetector is a silicon pixel detector. The operating principle for the silicon pixels is that of a p - n junction. When a charged particle passes through, it creates electron-hole pairs that are then separated by the electric field. The sensors are $250\ \mu\text{m}$ thick and use oxygenated n -type wafers with readout pixels on the n^+ side of the detector [34]. Overall, the pixel detector has 1744 sensors and 80.4 million readout channels.

In the barrel region, the pixel detector has three concentric layers of sensors surrounding the beamline. In the endcap region, it consists of disks perpendicular to the beam axis. The detector is segmented in the R - ϕ plane and in z . Usually, three pixel layers are crossed by a charged particle track. The intrinsic accuracies of the sensors are $10\ \mu\text{m}$ in R - ϕ and $115\ \mu\text{m}$ in z (or R for the endcap).

897 INSERTABLE B-LAYER

898 In Run 2, a new innermost pixel layer, known as the insertable B-layer (IBL), was added to the Inner
899 Detector [43]. This layer was added to cope with the higher luminosities planned in LHC Run 2 and at the

900 high luminosity HL-LHC. Additionally it improves tracking position resolution which in turn improves
901 the vertexing and b -tagging capabilities in ATLAS. The detector sits directly on a new beam pipe, only
902 33.25 mm away from the collision points in the azimuthal plane.

903 **SEMICONDUCTOR TRACKER (SCT)**

904 The semiconductor tracker (SCT) consists of silicon microstrips and comprises the next four layers
905 of the ID. This sub-detector has 6.4cm long sensors that are daisy-chained into strips with a strip pitch
906 of $80\ \mu\text{m}$ [34]. Some of the strips have a small stereo angle to allow for measurement of both angular
907 coordinates. In total there are 6.3 million readout channels. The intrinsic accuracies are $17\ \mu\text{m}$ in $R\text{-}\phi$
908 and $580\ \mu\text{m}$ in z (or R in the endcap).

909 **TRANSITION RADIATION TRACKER (TRT)**

910 The transition radiation tracker (TRT) serves two purposes. First, it consists of 4mm diameter straw
911 tubes filled with a 70/27/3% gas mixture of xenon, carbon dioxide, and oxygen to provide tracking of
912 charged particles. Particles typically have 36 TRT straw tube hits per track. The material in between
913 the straws is designed to induce transition radiation which can be useful for particle identification. As
914 particles pass between media with different dielectric constants, they emit transition radiation that can
915 cause additional showers in the TRT. In particular it is useful for discrimination between electrons and
916 pions or other charged hadrons, as the amount of transition radiation is proportional to the Lorentz factor
917 of the particle.

918 **2.2.3 CALORIMETERS**

919 The calorimeter system consists of two main sub-components: a fine granularity electromagnetic calorime-
920 ter tailored for the measurement of photons and electrons and multiple coarser hadronic calorimeters ded-
921 icated to the measurement of hadronic showers [34]. The calorimeter system has broader coverage than
922 the inner detector, covering the region out to $|\eta| < 4.9$. It is also designed to deliver good containment of
923 showers so as to limit leakage into the muon system. Figure 2.5 shows the layout of the calorimeter system.

924 Both the electromagnetic and hadronic calorimeters are sampling calorimeters. They alternate active
925 material for energy measurement with passive material for energy absorption. The materials used for each
926 purpose vary based on the type of calorimeter and its location in the detector.




Figure 2.5: Layout of the ATLAS calorimeter system [34]

927 ELECTROMAGNETIC CALORIMETER

928 The electromagnetic calorimeter (EM calorimeter) use liquid Argon (LAr) as its active material and
929 lead as its passive material. It is arrange in an accordion geometry to increase the absorption area while still
930 allowing it to have no azimuthal cracks (complete symmetry in ϕ). The EM calorimeter is divided into a
931 barrel portion that extends to $|\eta| < 1.475$ and an endcap portion going from $1.375 < |\eta| < 3.2$. The
932 region where these two units overlap is called the “transition region”.

933 In order to provide good containment the calorimeter depth must be optimized. Typically, for elec-
934 tromagnetic calorimeters the depth is measured in radiation lengths. In general, the intensity of a par-
935 ticle beam attenuates exponentially in distance with a constant equal to the radiation length. That is,

936 $I(x) = I_0 e^{-x/X_0}$, where I is the intensity, x is the distance traveled, and X_0 is the radiation length.
937 The ATLAS EM calorimeter is designed to have > 22 radiation lengths in the barrel and > 24 in the
938 endcap [34].

939 **HADRONIC CALORIMETERS**

940 There are three types of hadronic calorimeters present in ATLAS: the tile calorimeter (TileCal), hadronic
941 endcap (HEC), and forward calorimeter (FCal). Each one is optimized for stopping of hadronic showers
942 and the materials chosen are specific to their placement in the detector.

943 The TileCal is a scintillating tile calorimeter placed directly outside the EM calorimeter. It uses steel as
944 the absorber and plastic scintillator tiles as the active material. It has coverage in the barrel at $|\eta| < 1.0$
945 and in the “extended barrel” region of $0.8 < |\eta| < 1.7$.

946 The HEC had two wheels perpendicular to the beam line per endcap and is located directly behind the
947 EM calorimeter endcap modules. The HEC covers the region from $1.5 < |\eta| < 3.2$, overlapping slightly
948 with both the tile calorimeter and the forward calorimeter. Like the EM calorimeter, it uses liquid Argon
949 as the active material, but it uses copper as the absorber.

950 The FCal covers the most forward regions of the calorimeter system, extending to the region of $3.1 <$
951 $|\eta| < 4.9$. It again uses liquid argon as its active material. For absorber, it consists of an innermost module
952 made of copper followed by a module made of tungsten.

953 The hadronic equivalent of radiation length is called the interaction length and is denoted as λ . In the
954 barrel, the hadronic calorimeter depth is approximately 9.7λ , while in the endcap is 10λ . The outer
955 supports contribute an additional 1.3λ . This is been shown to be sufficient to limit punch-through of
956 showers to the muon system [34].

957 **2.2.4 MUON SPECTROMETER**

958 The muon spectrometer is dedicated to measuring the momentum and position of muons. It consists
959 of tracking and trigger chambers which are unique in the barrel and endcap regions. The magnetic field
960 for bending of muons is provided by a system of three large air-core toroid magnets (from which ATLAS

961 derives its name.) These magnets provide 1.5 to 5.5 Tm of bending power at $0 < |\eta| < 1.4$ and approx-
 962 imately 1 to 7.5 Tm in the endcap region of $1.6 < |\eta| < 2.7$. The entire muon system covers the range
 963 $0 < |\eta| < 2.7$. Monitored drift tubes (MDTs) are used for tracking in the barrel and the two outer layers
 964 of the endcap, while cathode strip chambers (CSCs) are used to provide tracking in the innermost endcap
 965 wheel. In the barrel, resistive plate chambers (RPCs) are used as trigger chambers while thin gap chambers
 966 (TGCs) are used in the endcap. Figure 2.6 shows the layout of the ATLAS muon system. The entire muon
 967 system is designed with the specification of providing a 10% momentum resolution for a 1 TeV muon.




Figure 2.6: Layout of the ATLAS muon system [34]

968 MONITORED DRIFT TUBES (MDTs)

969 The monitored drift tubes (MDTs) are aluminum 3cm diameter tubes filled with a 93/7 % mixture of
 970 Argon and CO₂, with trace amounts of water. As a charged particle traverses the tube, it ionizes the gas
 971 and the ions drift to a wire at the center of the tube. The radial distance of traversal of the particle in the

972 tube is determined by the drift time of the electrons, allowing for fine position resolution. The tubes have
973 an average resolution of $80 \mu\text{m}$ per tube and a maximum drift time of approximately 700ns. The tubes
974 are oriented so that they give precision measurement in η and run along ϕ . They cover $|\eta| < 2.7$, except
975 in the innermost layer of the endcap where they only go to $|\eta| < 2.0$ [34].

976 CATHODE STRIP CHAMBERS (CSCs)

977 The cathode strip chambers cover a narrow window of the innermost endcap region at $2.0 < |\eta| <$
978 2.7. In this region the background rates in the cavern are particularly high and the CSCs are designed to
979 handle these higher rates. The CSCs are multiwire proportional chambers with wires pointing in the radial
980 direction (away from the beam pipe). The wire serves as an anode and there are two types of segmented
981 cathode strip, one perpendicular to the wires which gives the precision measurement and one parallel which
982 provides the transverse coordinate. It has an 80/20 gas mixture of Argon and CO₂ [34].

983 RESISTIVE PLATE CHAMBERS (RPCs)

984 The resistive plate chambers (RPCs) are gaseous electrode-plate detectors covering the region $|\eta| <$
985 1.05. They consist of two resistive plates separated by a distance of 2 mm. The gas mixture used is a
986 94.7/5/0.3% mixture of C₂H₂F₄, Iso-C₄H₁₀, and SF₆. It has readout strips with a pitch of 23-35 mm
987 for both η and ϕ measurement and thus provides measurement of the azimuthal coordinate in the barrel
988 that the MDTs do not. The thin gas gap allows for a quick response time which makes it ideal for use in the
989 trigger. There are three layers of RPCs which are referred to as the three trigger stations. They allow for
990 both a low p_T and high p_T trigger. The coincidence of hits in the innermost chambers allows for triggering
991 of muons between 6 and 9 GeV, while the outermost layer allows the trigger to select high momentum
992 tracks in the range of 9 to 35 GeV [34].

993 THIN GAP CHAMBERS (TGCs)

994 The thin gap chambers (TGCs) are multiwire proportional chambers where the wire to cathode dis-
995 tance (1.4mm) is smaller than the wire-to-wire distance (1.8 mm). They contain a gas mixture of CO₂
996 and *n*-pentane and use a high electric field to gain good time resolution. They serve two functions in the

997 end-cap system. First, they serve as the trigger chambers. Second, they also provide azimuthal coordinate
 998 measurement which the MDTs do not. They sit on the inner and middle layers of the endcap. The outer-
 999 most layer's azimuthal coordinate is determined by extrapolation [34].

1000 2.2.5 MAGNET SYSTEM

1001 As mentioned previously, there are two independent magnet systems in ATLAS. The first is a 2 T
 1002 solenoid field in the inner detector which provides bending in the azimuthal plane. The second is an ap-
 1003 proximately 0.5 T toroidal field in the muon system which provides bending in η . Figure 2.7 shows the
 1004 predicted field integral as a function of $|\eta|$ [34].




Figure 2.7: Predicted field integral as a function of $|\eta|$ for the ATLAS magnet system [34]

1005 2.2.6 TRIGGER SYSTEM

1006 The ATLAS trigger system searches for signatures of muons, electrons, photons, hadronically decaying
 1007 τ leptons, and jets in order to save these events for further analysis. The trigger system in ATLAS is de-
 1008 signed to reduce the maximum LHC event rate of 40 MHz to a more reasonable rate that can be recorded.
 1009 The trigger first consists of a fast, hardware based system called the Level-1 (L1) trigger. The L1 trigger
 1010 consists of independent dedicated detector sub-components that can seed regions of interest (RoIs) for
 1011 further analysis downstream. For muons, the RPCs and TGCs are used, while in the calorimeter coarsely
 1012 grained sections of calorimeter cells called towers are used. Once regions of interest are seeded, a software

1013 based system called the High Level Trigger (HLT) is used to reconstruct objects and integrate information
 1014 from different parts of the detector. In Run 1 of ATLAS, the HLT consisted of two separate stages: the
 1015 level 2 (L2) trigger and the event filter (EF).

1016 The maximum trigger rate that the L1 trigger can handle is 75 kHz. In the HLT, the rate of events
 1017 written to disk is approximately 200 Hz. Figure 2.8 shows the trigger rates for different L1 triggers in 2012
 1018 and 2015 for ATLAS [44].




Figure 2.8: ATLAS trigger rates for Level-1 triggers as a function of instantaneous luminosity in 2012 and 2015 operation. These are single object triggers for electromagnetic clusters (EM), muons (MU), jets (J), missing energy (XE), and τ leptons (TAU). The threshold of the trigger is given in name in GeV. [44]

1019 2.2.7 ATLAS DATASETS

1020 ATLAS has collected data at center of mass energies of 7, 8, and 13 TeV. Figure 2.9 shows the integrated
 1021 luminosity as a function of time for each of the three collected datasets. At $\sqrt{s} = 7$ TeV, ATLAS recorded
 1022 5.08 fb^{-1} . Increased instantaneous luminosity in 2012 led to a larger dataset of 21.3 fb^{-1} recorded at
 1023 $\sqrt{s} = 8$ TeV. After Long Shutdown 1 (LS1) of the LHC and a restart in 2015, ATLAS recorded 3.9 fb^{-1}
 1024 of data at $\sqrt{s} = 13$ TeV. [45, 46]

1025 2.2.8 DETECTOR PERFORMANCE

1026 Table 2.2 summarizes the design requirements for each of the different sub-detectors. This table shows
 1027 the energy and momentum resolution of each tracking, calorimetry, and muon measurements.




Figure 2.9: Instantaneous luminosity as a function of time for data recorded by ATLAS at different center of mass energies [45, 46]

	Required resolution
Tracking	$\sigma_{p_T}/p_T = 0.05\% p_T \oplus 1\%$
EM calorimetry	$\sigma_E/E = 10\%/\sqrt{E} \oplus 0.7\%$
Hadronic calorimetry	
Barrel and end-cap	$\sigma_E/E = 50\%/\sqrt{E} \oplus 3\%$
Forward	$\sigma_E/E = 100\%/\sqrt{E} \oplus 10\%$
Muon spectrometer	σ_{p_T}/p_T at $p_T = 1$ TeV

Table 2.2: Performance requirements for the ATLAS detector [34].

2.3 THE ATLAS MUON NEW SMALL WHEEL UPGRADE

As the LHC continues operation, it is scheduled to be upgraded in several phases to allow it to reach higher instantaneous luminosities and thus collect larger datasets. These conditions will open new doors

1031 for study of rare physics processes but will also present interesting challenges that must be faced. ATLAS
 1032 will require new detector technologies to cope with the increased background rates in the cavern in these
 1033 high luminosity conditions. One such upgrade, scheduled to be installed during Long Shutdown 2 (LS2)
 1034 of the LHC in 2018, is the ATLAS Muon New Small Wheel (NSW) upgrade [47]. The NSW will replace
 1035 the innermost end-cap wheel of the muon system with new technologies, as this is the part of the muon
 1036 detector closest to the beam and thus suffers from the highest rates.

1037 2.3.1 MOTIVATION

1038 The motivation of the NSW is two-fold. First, the objective is to alleviate the decreased tracking ef-
 1039 ficiency that comes in a high rate environment. As figure 2.10, at the LHC design luminosity both the
 1040 efficiency of recording hits and reconstructing track segments in the MDTs decreases at the LHC design
 1041 luminosity.




Figure 2.10: MDT tube hit (solid) and segment (dashed) efficiency as a function of hit rate per tube [47]

1042 Second, the NSW will work to alleviate the rate of fake triggers arising in the endcap. Figure 2.11 shows
 1043 the extrapolated trigger rates as a function of the p_T threshold with and without the NSW upgrade. As
 1044 the figure shows, the NSW upgrade will reduce the trigger rate by an order of magnitude compared to the
 1045 current endcap trigger system.




Figure 2.11: Trigger rate as a function of p_T threshold with and without the NSW upgrade [47]

1046 2.3.2 NSW DETECTOR TECHNOLOGIES

1047 The NSW will use two new detector technologies - micromesh gaseous structure detectors (micromegas)
 1048 and small-strip thin gap chambers (sTGCs) [47, 48]. Unlike the previous detectors, both of these detector
 1049 technologies can be used for tracking or trigger. However, the micromegas is more suited to tracking be-
 1050 cause of its good spatial resolution, while the sTGCs have better time resolution and are more suited for
 1051 the trigger. To maintain a fully redundant system, both technologies are used for both purposes.

1052 MICROMEGAS

1053 Micromegas detectors operate using a thin metallic mesh that sits approximately 100 μm away from the
 1054 readout electrodes to create the amplification region. Above this mesh, there is a drift region on the order
 1055 of a few mm in length capped by a drift electrode. As a charged particle traverses the detector, it ionizes gas
 1056 and the electrons drift down towards readout strips. The timing of the drift can be used to reconstruct the
 1057 angle of traversal of the particle. This is illustrated in figure 2.12. The micromegas used in ATLAS will be
 1058 resistive micromegas, where the readout electrodes are topped with resistive strips [49]. This alleviates the

1059 risk of sparking in the large area detectors that ATLAS will use.

1060 In ATLAS, the micromegas drift gap will be 5 mm and the amplification gap will be $128 \mu\text{m}$. They are
1061 filled with the same gas mixture as the MDTs. They will be stacked in an octuplet in an XXUV-UVXX
1062 geometry, where X refers to straight strips and U and V refer to stereo strips at an angle of $\pm 1.5^\circ$. This
1063 arrangement allows for measurement of the azimuthal coordinate and gives a large lever arm between the
1064 straight strips for triggering purposes. Figure 2.12 shows the geometry of a single micromegas detector as
1065 well as its operating principle [47].




Figure 2.12: Illustrations of the geometry (left) and operating principle (right) of the micromegas detector [47]

1066 sTGCs

1067 The sTGCs are similar to the TGCs already described. They consist of gold-plated tungsten wires with a
1068 1.8 mm pitch between two cathode planes 1.4 mm away from the wire plane. One cathode plane consists
1069 of strips with a 3.2 mm pitch (much smaller pitch than the TGCs), while the other consists of coarser
1070 pads that are used for defining regions of interest in the sTGC trigger algorithm. Figure 2.13 shows the
1071 basic detector geometry.




Figure 2.13: Geometry of the sTGC detector [47]

1072 2.3.3 PHYSICS IMPACT

1073 Maintaining low p_T thresholds for muons while still staying within the trigger rate budget at Level 1
1074 (20 kHz) for the muon system is crucial for physics analyses to be successful in high luminosity condi-
1075 tions. One realm where the lepton trigger threshold is especially important is in Higgs physics. In the
1076 $H \rightarrow WW^*$ analysis, one of the W bosons is off shell and tends to decay to soft leptons. In associated
1077 production of a Higgs with a W , the lepton is also important because the lepton provides the main han-
1078 dle which allows the event to be triggered. Table 2.3 shows the impact of increasing the trigger thresholds
1079 on these analyses. It shows that either raising the threshold or using only the barrel both have significant
1080 impacts on the signal efficiency. With the NSW, the signal efficiency is largely maintained and the triggers
1081 can be unprescaled.

Threshold	$H \rightarrow b\bar{b}$ (%)	$H \rightarrow WW^*$ (%)
$p_T > 20 \text{ GeV}$	93	94
$p_T > 40 \text{ GeV}$	61	75
$p_T > 20 \text{ GeV} (\text{barrel only})$	43	72
$p_T > 20 \text{ GeV} (\text{with NSW})$	90	92

Table 2.3: Signal efficiencies for WH production with $H \rightarrow b\bar{b}$ and $H \rightarrow WW^* \rightarrow \mu\nu qq$ under different trigger configurations [47].

1082 2.4 OBJECT RECONSTRUCTION IN ATLAS

1083 ATLAS analyses first start by requiring the presence of certain reconstructed physics objects in the event.
1084 This section will present a brief overview of the algorithms used to reconstruct electrons, muons, jets (in-
1085 cluding b -jets), and missing energy¹. The performance of object reconstruction and measurement will also
1086 be discussed as these are relevant to the analyses presented later. Figure 2.14 gives an overview of the differ-
1087 ent sub-detectors that each type of particle will interact with in ATLAS.

¹Reconstruction algorithms for other objects, such as photons and τ leptons, are not detailed here as these objects are not used in the presented studies.




Figure 2.14: Illustration of particle interactions in ATLAS [50]

1088 2.4.I ELECTRONS

1089 Electrons in ATLAS will leave tracks in the inner detector and energy deposits in the electromagnetic
 1090 calorimeter. The algorithm for recognizing the signature of electrons proceeds in two steps: reconstruction
 1091 and identification.

1092 In reconstruction, an electron candidate is formed by matching EM calorimeter deposits with ID tracks.
 1093 The algorithm first chooses seed clusters in the EM calorimeter by using a sliding window algorithm that
 1094 searches for towers with transverse energy larger than 2.5 GeV. In addition to seed clusters, track candi-
 1095 dates must be identified in the ID. The algorithm selects seed tracks with $p_T > 1$ GeV that do not fit well
 1096 with a pion hypothesis. Once candidate tracks are selected, they are re-fit with a Gaussian Sum Filter (GSF)
 1097 algorithm to estimate electron parameters [51]. Finally, an electron candidate is formed if at least one track
 1098 matches to a seed cluster in the calorimeter. The full details of the reconstruction algorithm can be found
 1099 in reference [52].

1100 Once an electron candidate is present, identification criteria must be applied in order to reject fake elec-

trons from background. Many different variables are used for this identification, most of them related to
 the shower shape in the EM calorimeter and the amount of leakage into the hadronic calorimeter, as well
 as information from the ID and in particular the TRT. There are both cut-based and likelihood-based
 criteria that range from “loose” to “very tight”. For details, see reference [52].
 1101
 1102
 1103
 1104

Figure 2.15 shows the algorithm’s reconstruction efficiency of true electrons for different identification
 criteria as well as the electron energy resolution in simulation [52, 53]. The reconstruction efficiency is
 measured using both Z and J/ψ tag and probe techniques.
 1105
 1106
 1107




Figure 2.15: Electron performance: (a) reconstruction efficiency as a function of electron E_T [52] (b) energy resolution in simulation as a function of $|\eta|$ for different energy electrons [53]

2.4.2 MUONS

The ATLAS detector is designed to stop most particles before they reach the muon spectrometer. Muons,
 however, are minimum ionizing particles, meaning that they will not lose a significant amount of energy
 through interactions with the detector and will thus pass through. Therefore, the muon reconstruction
 works to match tracks in the muon spectrometer with tracks in the inner detector.
 1108
 1109
 1110
 1111
 1112

The first step of reconstruction is to reconstruct local straight line tracks, called segments, in each muon
 chamber. Segments are then fit to larger tracks that traverse the entire muon spectrometer. Such muon
 tracks are referred to as “standalone” tracks (SA) as they only use information from the muon spectrometer.
 1113
 1114
 1115

1116 The standalone tracks are then matched to tracks in the inner detector to form “combined” (CB) muons,
 1117 where the combined ID and MS fit are used to determine the momentum and direction of the muon. To
 1118 improve acceptance, segment-tagged and calorimeter-tagged muons are also reconstructed. In these cases,
 1119 ID tracks are matched to segments in the MS and calorimeter deposits consistent with a minimum ionizing
 1120 particle, respectively. The details of the reconstruction can be found in reference [54].

1121 As with electrons, once muon candidates are reconstructed they have identification criteria applied to
 1122 reduce background. These criteria include the χ^2 match between the ID and MS tracks, the number of
 1123 hits in the ID, overall ID and MS track fit quality, and additional variables [54]. The criteria range from
 1124 “loose” to “tight” as with electrons.

1125 Figure 2.16 shows the muon reconstruction efficiency (measured with Z and J/ψ tag and probe) and
 1126 invariant mass resolution [54].




Figure 2.16: Muon performance: (a) reconstruction efficiency as a function of muon p_T (b) dimuon mass resolution as a function of average p_T [54]

1127 2.4.3 JETS

1128 When a quark or gluon is produced in collisions, it is not measured directly in ATLAS. Rather, due to
1129 QCD effects, it produces a collimated spray of hadrons in the direction of the original parton, which is
1130 known as a jet. Jets are reconstructed in ATLAS using energy deposits in the hadronic calorimeter. The
1131 first step is build “topological clusters” out of energy deposits in calorimeter cells [55, 56]. This is done
1132 using strategy where seed cells are chosen by picking cells whose energy measurements are four times the
1133 amount of noise expected for that cell. Adjacent cells with at least 2σ energy measurements are added to
1134 the cluster, then a final layer of clusters with energy above 0σ are added. Once calorimeter clusters are
1135 formed, they are clustered further into jet candidates using the anti- k_T jet clustering algorithm [57]. This
1136 algorithm uses a parameter R that appears in the denominator of the clustering distance metric and defines
1137 the radial size of the jet in η - ϕ space.

1138 The energy response of the calorimeter must be properly characterized in order to reconstruct jet energy.
1139 Calorimeter clusters can be calibrated either with the EM calibration, where each cluster is assumed to have
1140 come from the energy deposit of an electron or photon, or the LCW calibration, where local cluster weights
1141 are computed to allow for local calibration of clusters as hadronic or electromagnetic. The details of the jet
1142 energy calibration are not detailed here and are discussed in reference [58]. Figure 2.17 shows the jet energy
1143 response after calibration in Monte Carlo as a function of the true p_T of the jet [58].

1144 Analyses often need to know how consistent a particular jet is with the primary vertex of the event in
1145 order to avoid contamination from pileup interactions. One measure of this consistency is known as the
1146 jet vertex fraction (JVF). The JVF is the ratio of tracks associated with a primary vertex to the total number
1147 of tracks inside a jet. Jets from the primary interaction in the event should have a large fraction of tracks
1148 consistent with the primary vertex and therefore have a large JVF value.

1149 2.4.4 b -TAGGING

1150 One important aspect of jet physics is the task of identifying the flavor of parton that produced the
1151 measured jet. While in general this is very difficult, jets from b -quarks offer an interesting case where such
1152 identification is possible. B mesons have a lifetime on the order of 10^{-12} seconds, which makes a $c\tau$ on




Figure 2.17: Jet energy response after calibration as a function of true p_T in simulation [58]

the order of millimeters [6]. This type of displaced decay vertex can be identified in detectors like ATLAS and allows b -jets to be distinguished from other flavors of jets².

ATLAS uses a multivariate machine learning algorithm to identify jets from b -quarks. The inputs to this algorithm are determined from lower level reconstruction algorithms. There are three distinct algorithms that reconstruct variables which are used as input to the multivariate technique.

The first family is referred to as IPxD (where the x can either be 2 or 3). These algorithms use the transverse and longitudinal impact parameters d_0 and z_0 of the tracks inside a jet to determine their consistency with the primary vertex. They use two or three dimensional (hence the x) templates for light flavor, charm, and bottom jets and then evaluate the likelihood of the jet coming from each of these types. The likelihood ratios are used as inputs to the multivariate algorithm.

The next two algorithms used as input are referred to as the secondary vertex (SV) and JetFitter (JF) algorithms. The SV algorithm uses tracks inside the jet to fit for vertices that are displaced from the primary vertex. The JF algorithm attempts to reconstruct the full flight path of the b by looking for multiple displaced vertices along the same line (as B decays often result in subsequent c decays).

In Run 1, the multivariate b -tagging algorithm used a neural network and was referred to as MV1.

²Jets from charm quarks can also be detected in this way but they do not live quite as long so the displacement of the vertex is harder to distinguish

1168 The details of this algorithm and its inputs are given in reference [59]. In Run 2, the number of inputs
 1169 was simplified and a boosted decision tree with 24 input variables was used, referred to as MV2 [60]. The
 1170 MV2 algorithm is a boosted decision tree incorporating twenty-four input variables constructed from three
 1171 lower level input algorithms described above. Figure 2.18 summarizes the inputs to MV2. Figure 2.19 shows
 1172 the performance of each of these algorithms.

IP2D and IP3D (6 inputs)	SV1 (8 inputs)	JetFitter (8 inputs)
$\log(p_b/p_u)$	Mass	Mass
$\log(p_b/p_c)$	Energy fraction	Energy fraction
$\log(p_c/p_u)$	# tracks at vertex	# vertices
	# 2 track vertices	# tracks at vertex
	Lxy	# 1 track vertices
	L3d	# 2 track vertices
	3D significance	3D significance
	ΔR	ΔR
Kinematics (2 inputs)		
p_T		
η		

Figure 2.18: Summary of the inputs to the MV2 b -tagging algorithm

1173 2.4.5 MISSING TRANSVERSE ENERGY

1174 As noted in figure 2.14, neutrinos produced in ATLAS will pass through the detector without interact-
 1175 ing. The only way of detecting the presence of particles like neutrinos (or BSM particles that are long-lived)
 1176 is to use missing transverse momentum. The basic principle of missing transverse energy is to use the mo-
 1177 mentum balance of the incoming protons to infer the presence of missing particles. The net longitudinal
 1178 momentum of the incoming partons that collide is not known (since each carries an unknown fraction of
 1179 the proton's momentum). However, the protons (and thus incoming partons) have no net momentum
 1180 in the plane transverse to the beam line (the x - y) plane. Therefore, if there are no un-measured particles
 1181 in the final state, the transverse momenta of all of the final state particles should balance. The magnitude




Figure 2.19: Light jet rejection ($1/\text{efficiency}$) vs. b -jet efficiency for MV1 and its input algorithms (a) [59] and MV2 (b) [60] in simulated $t\bar{t}$ events. The numbers in the algorithm names in (b) refer to the fraction of charm events used in the MV2 training.

of this imbalance is known as missing transverse momentum ($E_{\text{T}}^{\text{miss}}$).

The basic calculation of missing transverse momentum from calorimeter cells is given in equation 2.4 [61].

$$\begin{aligned} E_x^{\text{miss}} &= - \sum_{i=1}^{N_{\text{cell}}} E_i \sin \theta_i \cos \phi_i \\ E_y^{\text{miss}} &= - \sum_{i=1}^{N_{\text{cell}}} E_i \sin \theta_i \sin \phi_i \end{aligned} \quad (2.4)$$

The $E_{\text{T}}^{\text{miss}}$ calculation is separated into different terms based on the objects that the calorimeter clusters are associated with. This way, each cell's contribution is calibrated appropriately according to the object. This separation of terms is shown in equation 2.5 [61].

$$\begin{aligned} E_{x(y)}^{\text{miss,calo}} &= E_{x(y)}^{\text{miss},e} + E_{x(y)}^{\text{miss},\gamma} + E_{x(y)}^{\text{miss},\tau} + E_{x(y)}^{\text{miss,jets}} \\ &\quad + E_{x(y)}^{\text{miss,softjets}} + E_{x(y)}^{\text{miss},\mu} + E_{x(y)}^{\text{miss,CellOut}} \end{aligned} \quad (2.5)$$

The CellOut term of the above equation corresponds to calorimeter cells with energy deposits that are not associated with other objects. The soft jets term comes from cells associated to jets with p_{T} between 7 and 20 GeV, while the jets term comes from jets with $p_{\text{T}} > 20$ GeV. Because muons do not deposit

¹¹⁹¹ significant energy in the calorimeter, the muon momentum is used for the muon term [61]. The final
¹¹⁹² E_T^{miss} is calculated using equation 2.6.

$$E_T^{\text{miss}} = \sqrt{(E_x^{\text{miss}})^2 + (E_y^{\text{miss}})^2} \quad (2.6)$$

¹¹⁹³ Figure 2.20 shows the resolution of the components of the E_T^{miss} under different pileup suppression techniques [62].




Figure 2.20: Resolution of E_T^{miss} components as a function of $\sum E_T$ before pileup suppression with different pileup techniques [62]

¹¹⁹⁴

1195

Part II

1196

Observation and measurement of Higgs

1197

boson decays to WW^* in LHC Run I at

1198

$$\sqrt{s} = 7 \text{ and } 8 \text{ TeV}$$

*Basic research is what I am doing when I don't know what
I am doing.*

Wernher von Braun

3

1199

1200 $H \rightarrow WW^* \rightarrow \ell\nu\ell\nu$ Analysis Strategy

1201 3.1 INTRODUCTION

1202 This chapter presents an overview of the strategy for searching for a Higgs boson in the $H \rightarrow WW^* \rightarrow$
1203 $\ell\nu\ell\nu$ decay topology. Its purpose is to define in broad terms how the search and measurement are under-
1204 taken, before going into details on the specific sub-categories within the larger analysis. First, the properties
1205 of the Higgs signal are discussed and the associated backgrounds are presented. Next, the observables used
1206 to enhance the signal to background ratio are defined. Finally, the parameters of interest in the search
1207 and measurement will be shown, along with a brief overview of the statistical treatment of the final Higgs
1208 candidates.

1209 Following this chapter, the results of three different studies within the $H \rightarrow WW^* \rightarrow \ell\nu\ell\nu$ channel
1210 are shown. Chapter 4 presents a search for Higgs boson production in gluon fusion mode and the role of
1211 the $H \rightarrow WW^*$ channel in its discovery. Chapter 5 shows the search and first evidence in ATLAS of the
1212 Vector Boson Fusion (VBF) production mode of the Higgs. Finally, chapter 6 shows the combined Run

1213 1 $H \rightarrow WW^*$ results for the measurement of the Higgs cross section and relative coupling strengths to
1214 other SM particles.

1215 3.2 THE $H \rightarrow WW^* \rightarrow \ell\nu\ell\nu$ SIGNAL IN ATLAS

1216 The signal studied in this and subsequent chapters is the Higgs boson in the WW^* final state, where
1217 each W boson subsequently decays into a charged lepton and a neutrino. In the simplest decay path, the
1218 final state consists of two neutrinos and two charged leptons, each of which can be either an electron or a
1219 muon. If a W decays to a τ lepton, only leptonic decays of the τ are considered. The τ lepton produce
1220 additional neutrinos in the final state but still yield two charged leptons (where each lepton is an electron or
1221 muon). Neutrinos are not detected in ATLAS, so the final state ultimately consists of two reconstructed
1222 leptons and missing transverse momentum (denoted as E_T^{miss}). Final states where both of the charged
1223 leptons are electrons or muons are referred to as the “same flavor” ($ee/\mu\mu$) final states, while those with
1224 one electron and one muon are referred to as “different flavor” ($e\mu$ or μe).

1225 There can be additional jets produced in association with the Higgs boson. As described in detail in
1226 Chapter 1, if the Higgs is produced via vector boson fusion production, there will be two additional forward
1227 jets in the event. In gluon fusion, one or more jets can be produced through initial state radiation from
1228 the incoming gluons. Because of the varying background composition as a function of jet multiplicity,
1229 each bin in this variable has its own dedicated requirements applied in the search and measurement. The
1230 $n_j = 0$ and $n_j = 1$ bins are dedicated to gluon fusion production, while the $n_j \geq 2$ bin has separate
1231 dedicated searches for ggF and VBF production.

1232 Figure 3.1 shows the relative branching fractions for the $H \rightarrow WW^*$ process, calculated from the Par-
1233 ticle Data Group values for the W and τ branching ratios [6]. The largest branching ratio corresponds
1234 to both W bosons decaying to quark pairs at 45.44%. The second largest ratio is for one W decaying lep-
1235 tonically and the other decaying to quarks, a branching ratio of 34.18%. In all cases, ℓ denotes either an
1236 electron or muon, and the leptonic branching ratios of the τ are included. For example, the $\ell\nu qq$ final
1237 state includes one W decaying to $e\nu$, $\mu\nu$, or $\tau\nu$. In the case of the $W \rightarrow \tau\nu$ decay, the τ lepton then
1238 decays to an electron or muon via $\tau \rightarrow \nu_\tau \ell \nu_\ell$. Final states with a τ_h refer to hadronic decays of the τ . The




Figure 3.1: Branching ratios for a WW system. q refers to quarks. ℓ can be either an electron or muon, and the leptonic branching ratios of the τ are included. For example, the $\ell\nu qq$ final state includes one W decaying to $e\nu$, $\mu\nu$, or $\tau\nu$. τ_h refer to hadronic decays of the τ .

branching ratio of the $\ell\nu\ell\nu$ final state is 6.43%.

While the $\ell\nu\ell\nu$ final state is not a large fraction of the branching ratio, there are significant advantages to using this channel in an analysis. First, both the $qqqq$ and $\ell\nu qq$ channels suffer from a large QCD multijet background, which is often difficult to model. Second, events in the the $\ell\nu\ell\nu$ channel in data can be triggered more efficiently due to the presence of two leptons.

3.3 BACKGROUND PROCESSES

Many processes from the Standard Model can also produce a final state with two leptons and missing transverse momentum. This section describes the dominant backgrounds to Higgs production and further explains how they can be reduced. Table 3.1 summarizes the different background processes.

3.3.1 STANDARD MODEL WW PRODUCTION

Non-resonant Standard Model diboson production, as shown in figure 3.2, is an irreducible background to Higgs boson production in the WW final state. It produces the same exact final state objects, namely




Figure 3.2: Feynman diagram for Standard Model WW production

leptonically decaying W bosons. There are no additional objects in the final state that allow for background reduction. Therefore the analysis solely relies on the correlations between the leptons to reduce this background.

3.3.2 TOP QUARK PRODUCTION

Top quark production can mimic the Higgs in the WW^* final state as well. Top quarks can be produced either in pairs ($t\bar{t}$ production) or singly (s -channel, t -channel, or associated production Wt). The dominant top background are $t\bar{t}$ and Wt production.

Because top quarks decay via $t \rightarrow Wb$, top pair production can produce a final state with two W bosons that then decay leptonically. In Wt production, there are two real W bosons produced, as with $t\bar{t}$. In both cases, there is at least one b -jet in the final state. By vetoing on the presence of b -jets, these top quark backgrounds can be reduced. Figure 3.3 shows the Feynman diagrams for $t\bar{t}$ and Wt production.




Figure 3.3: Feynman diagrams for top pair production (left) and Wt production (right)

1262 3.3.3 W +JETS BACKGROUND

1263 Single W boson production in association with jets is a unique background to Higgs production. The
1264 other backgrounds considered thus far have all included two prompt leptons, each decaying from a W
1265 boson, in the final state. In W +jets production, however, only one reconstructed lepton originates from
1266 a W . The second reconstructed lepton is either an algorithmic “fake” or the result of non-prompt decays.
1267 In the first case, the lepton is a jet misidentified as a lepton by either the electron or muon reconstruction
1268 algorithms. In the second case, the lepton may be a real lepton but coming from semi-leptonic decays of
1269 particles inside the shower of the jet. This background can be reduced by requiring that the reconstructed
1270 lepton have little activity in the surrounding region of the calorimeter (also known as an “isolation”). Fig-
1271 ure 3.4 shows the Feynman diagram for W +jets production.




Figure 3.4: An example Feynman diagram of W +jets production

1272 3.3.4 Z/γ^* +JETS BACKGROUND

1273 Production of a Z boson or virtual photon (also known as Drell-Yan and denoted with Z/γ^*) in as-
1274 sociation with jets is also a background to Higgs production. The Z boson decays to two leptons of the
1275 same flavor. When the Z/γ^* decays directly to electrons or muons, the background enters the same flavor
1276 final state. When the Z decays to two τ leptons the background can enter the different flavor final state as
1277 well. Figure 3.5 shows the production of a Z in association with one jet. Because there are no neutrinos in
1278 this final state, variables like E_T^{miss} can be used to reduce the background¹.

¹The E_T^{miss} cut is much more effective for the reduction of Z/γ^* production in the same flavor final state. If the background enters the different flavor final state through τ decays, there will be neutrinos present. Other requirements on the lepton invariant mass are made to reduce the $Z/\gamma^* \rightarrow \tau\tau$ background.



Figure 3.5: An example Feynman diagram of Z +jets production

1279 3.3.5 SUBDOMINANT BACKGROUNDS

1280 There are additional processes which contribute to the background composition. These backgrounds
 1281 are subdominant and contribute less to the total background estimate than those discussed previously.
 1282 The first process is referred to as VV or “Other diboson” processes and includes multiple Standard Model
 1283 diboson processes, including WZ , ZZ , $W\gamma$, $W\gamma^*$, and $Z\gamma$ production. Additionally, there is a back-
 1284 ground contribution from QCD multijet production. While the cross section for this process is large, its
 1285 contribution to the WW^* final state is small because two jets must be misidentified as leptons.

Category	Process	Description
SM WW	$WW \rightarrow \ell\nu\ell\nu$	Real leptons and neutrinos
Top quark production	$t\bar{t} \rightarrow WbWb \rightarrow \ell\nu b\bar{b}\nu\bar{b}$	Real leptons, untagged b s
	$tW \rightarrow WbW \rightarrow \ell\nu\ell\nu b$	Real leptons, untagged b
	$t\bar{b}, t\bar{q}\bar{b}$	Untagged b , jet misidentified as lepton
Drell-Yan	$Z/\gamma^* \rightarrow ee, \mu\mu$	“Fake” E_T^{miss}
	$Z/\gamma^* \rightarrow \tau\tau \rightarrow \ell\nu\ell\nu\nu$	Real leptons and neutrinos
Other dibosons	$ZZ \rightarrow \ell\ell\nu\nu$	Real leptons and neutrinos
	$W\gamma^*, WZ \rightarrow \ell\nu\ell\ell, ZZ \rightarrow \ell\ell\ell\ell$	Unreconstructed leptons
	$W\gamma, Z\gamma$	γ reconstructed as e , unreconstructed lepton
W +jets	$Wj \rightarrow \ell\nu j$	Jet reconstructed as lepton
QCD multijet	jj	Jets reconstructed as leptons

Table 3.1: A summary of backgrounds to the $H \rightarrow WW^* \rightarrow \ell\nu\ell\nu$ signal

1286 3.4 SHARED SIGNAL REGION SELECTION REQUIREMENTS

1287 As presented in section 3.2, there are many different combinations of physics objects that can define a
 1288 $H \rightarrow WW^* \rightarrow \ell\nu\ell\nu$ final state. The multiplicity of jets and the flavor combinations of the leptons

1289 both lead to many potential signal regions. Additionally, signal regions can be optimized separately to be
 1290 sensitive to the distinct production modes of the Higgs. Gluon fusion, vector boson fusion, and associated
 1291 production of a Higgs all lead to unique final state topologies. Figure 3.6 delineates the different signal
 1292 regions used in the gluon fusion and vector boson fusion $H \rightarrow WW^*$ analyses. While there are different
 1293 optimizations possible in each signal region, there are also some commonly shared selections that will be
 1294 described here.




Figure 3.6: An illustration of the unique analysis signal regions [63]. The most sensitive regions for both gluon fusion and vector boson fusion production are underlined.

1295 3.4.I EVENT PRE-SELECTION

1296 Before being sorted into the distinct signal regions, basic requirements are applied to the reconstructed
 1297 objects in the event to select Higgs-like event candidates. First, two oppositely charged leptons are required.
 1298 Once the leptons are selected, the last requirement for event pre-selection is the presence of neutrinos. As
 1299 neutrinos cannot be detected directly in ATLAS, E_T^{miss} can be used as a proxy for the combined neutrino
 1300 momentum in the transverse plane.

1301 In general, it is expected that the signal should have a harder E_T^{miss} spectrum than backgrounds, espe-
 1302 cially if these backgrounds do not contain neutrinos in the final state. When using E_T^{miss} , it is possible
 1303 mis-measurements of objects in the detector can lead to imbalances in the transverse plane. When such a
 1304 mis-measurement occurs, the E_T^{miss} vector in the transverse plane will often point in the same direction as
 1305 the mis-measured object. Therefore, a new variable, $E_{T,\text{rel}}^{\text{miss}}$, is used in the pre-selection. $E_{T,\text{rel}}^{\text{miss}}$ is defined
 1306 in equation 3.1.

$$E_{T,\text{rel}}^{\text{miss}} = \begin{cases} E_T^{\text{miss}} \sin \Delta\phi_{\text{near}} & \text{if } \Delta\phi_{\text{near}} < \pi/2 \\ E_T^{\text{miss}} & \text{otherwise,} \end{cases} \quad (3.1)$$

1307 If the closest object to the E_T^{miss} vector is within $\pi/2$ radians in the transverse plane, the E_T^{miss} is projected
 1308 away from this object. Otherwise, the normal E_T^{miss} vector is used. Figure 3.7 shows a graphical illustration
 of this concept.




Figure 3.7: A graphical illustration of the $E_{T,\text{rel}}^{\text{miss}}$ calculation

1309
 1310 Once the lepton and E_T^{miss} pre-selections are made, the analysis is divided into different regions accord-
 1311 ing to jet multiplicity.

1312 3.4.2 JET MULTIPLICITY

1313 Jet multiplicity, denoted as n_j , is used to sub-divide the analysis into distinct signal regions. By creating
 1314 separate signal regions, each bin in jet multiplicity becomes sensitive to different modes of Higgs produc-
 1315 tion and different backgrounds.

1316 For example, the $n_j \geq 2$ region is more sensitive to VBF production because of the two high momen-
 1317 tum jets produced at matrix element level. For gluon fusion production to enter this bin, two initial state
 1318 radiation jets must be emitted.

1319 Figure 3.8 shows the jet multiplicity in both the different flavor and same flavor regions after the pre-
 1320 selection. It also shows the background composition in the bins of n_b . A few trends from this distribution
 1321 are worth noting. The first is that the Drell-Yan background dominates in the same flavor channels for
 1322 $n_j \leq 1$. Second, the top background becomes a clear contributor to the total background for $n_j \geq$
 1323 1. Lastly, the SM WW production dominates in the $n_j = 0$ bin, as it is an irreducible background to
 1324 $H \rightarrow WW^*$ production. Because of these distinct features, each jet multiplicity bin is treated separately.




Figure 3.8: Predicted backgrounds (compared with data) as a function of n_j (a and b) and n_b (c) after pre-selection requirements

1325 **3.5 BACKGROUND REDUCTION IN SAME-FLAVOR FINAL STATES**

1326 As described in section 3.4.2, the background composition of the same flavor final states is different
1327 from that of the different flavor states. In particular, Drell Yan processes play a much larger role because
1328 the Z/γ^* decays to same flavor leptons. Because real neutrinos are absent in the Z/γ^* decays to ee and $\mu\mu$,
1329 a requirement on E_T^{miss} should largely reduce the background. However, as this section will demonstrate,
1330 with increasing pileup conditions the resolution of the calorimeter-based E_T^{miss} degrades greatly. There-
1331 fore, two new variables for Z/γ^* background reduction are constructed and described in this section.

1332 **3.5.1 PILEUP AND E_T^{miss} RESOLUTION**

1333 Secondary interactions of protons in the colliding bunches of the LHC (known as pileup interactions,
1334 described in detail in Chapter 2) deposit energy into the ATLAS calorimeter in addition to the energy that
1335 comes from the hard scatter process of interest. The calculation of E_T^{miss} is fundamentally Poissonian.
1336 Summing up all of the energy deposits in individual calorimeter cells or clusters is similar to a counting
1337 experiment. The error on a mean of N in a Poisson distribution is \sqrt{N} , so the energy resolution scales
1338 as \sqrt{E} . As more energy is deposited in the calorimeter, the E_T^{miss} resolution degrades, meaning that the
1339 E_T^{miss} resolution is particularly sensitive to LHC instantaneous luminosity conditions.

1340 Figure 3.9 shows an event display of a $Z/\gamma^* + \text{jets}$ event candidate with the twenty-five reconstructed
1341 primary vertices. This display illustrates that while the interaction of interest only has tracks coming from
1342 the hardest primary vertex, all of the secondary interactions deposit energy in the calorimeter as well.

1343 Figure 3.10 shows the RMS of the E_T^{miss} distribution in $Z \rightarrow \mu\mu$ events (where there are no real neu-
1344 trinos) as a function of the number of the average number of interactions. Under 2011 LHC conditions,
1345 this RMS was approximately 9 GeV, while under 2012 running conditions the resolution worsened to 12
1346 GeV. The increase in pileup dilutes the E_T^{miss} variable's ability to reduce the Z/γ^* background.

1347 **3.5.2 TRACK-BASED DEFINITIONS OF MISSING TRANSVERSE MOMENTUM**

1348 Because the increasing number of secondary proton-proton interactions degrades calorimeter-based
1349 E_T^{miss} resolution, a new variable using only contributions from the primary interaction vertex is necessary




Figure 3.9: An event display of a Z/γ^* + jets event illustrating the effect of pileup interactions




Figure 3.10: The RMS of different missing transverse momentum definitions as a function of the average number of interactions per bunch crossing

to further reduce the Z/γ^* background. While it is not possible to associate calorimeter energy deposits with a particular vertex, individual charged particle tracks in the Inner Detector are associated to unique vertices. Thus, two track-based definitions of missing transverse momentum , using only tracks coming

1353 from the primary vertex in the event, are used in the analysis. The simplest variable, $p_T^{\text{miss}(\text{trk})}$, is the vec-
 1354 torial sum of the p_T of all of the tracks from the primary vertex and the selected leptons (excluding the
 1355 tracks associated with the selected leptons to avoid double counting). Equation 3.2 defines $p_T^{\text{miss}(\text{trk})}$.

$$p_T^{\text{miss}(\text{trk})} = - \left(\sum_{\text{selected leptons}} p_T + \sum_{\text{other tracks}} p_T \right), \quad (3.2)$$

1356 To further improve the resolution on the missing transverse momentum, the variable p_T^{miss} is used as de-
 1357 fined in equation 3.3. For selected leptons and jets, the nominal p_T measurements are used. Tracks are used
 1358 to estimate the soft component of the missing transverse momentum instead of calorimeter measurements.

1359

$$p_T^{\text{miss}} = - \left(\sum_{\text{selected leptons}} p_T + \sum_{\text{selected jets}} p_T + \sum_{\text{other tracks}} p_T \right), \quad (3.3)$$

1360 Figure 3.10 illustrates that these two new variables accomplish their intended purpose. The resolution as a
 1361 function of mean number of interactions for both $p_T^{\text{miss}(\text{trk})}$ and p_T^{miss} is much flatter than the dependence
 1362 for E_T^{miss} . Figure 3.11a shows the difference between the true and reconstructed values of missing transverse
 1363 momentum using both the track-based p_T^{miss} and calorimeter based E_T^{miss} . The RMS of the distribution
 1364 improves by 3.5 GeV when using p_T^{miss} .

1365 3.5.3 DISTINGUISHING Z/γ^* +JETS AND $H \rightarrow WW^*$ TOPOLOGIES

1366 In addition to measuring missing transverse momentum, another variable can be constructed to exploit
 1367 kinematic and topological differences between the Z/γ^* background and $H \rightarrow WW^*$ signal. Because
 1368 there are no real neutrinos in the final state (in the case of $Z/\gamma^* \rightarrow ee, \mu\mu$ decays), the dilepton system will
 1369 be balanced with the jets produced in the hard scatter. A new variable, f_{recoil} , is constructed to estimate
 1370 the balance between the dilepton system and recoiling jets and is defined in equation 3.4. The transverse
 1371 plane is divided into four sections, or quadrants, with one quadrant centered on the dilepton vector. The
 1372 numerator of f_{recoil} is the magnitude of the vectorial sum of the p_T of jets in the quadrant opposite the
 1373 dilepton system, weighted by each jet's Jet Vertex Fraction (JVF, described in chapter 2). The denominator




Figure 3.11: The difference between the true and reconstructed values of the missing transverse momentum (a) and m_T (b) in a gluon fusion signal sample

¹³⁷⁴ is the magnitude of the dilepton p_T .

$$f_{\text{recoil}} = \left| \sum_{\text{jets } j \text{ in } \wedge} \text{JVF}_j \cdot p_T^j \right| / p_T^{\ell\ell}. \quad (3.4)$$

¹³⁷⁵ Figure 3.12 shows a shape comparison of the f_{recoil} distribution in a simulated $Z/\gamma^* + \text{jets}$ sample, a
¹³⁷⁶ $H \rightarrow WW^*$ signal sample, and other backgrounds that contain real neutrinos. The $Z/\gamma^* + \text{jets}$ events

1377 tend to be more balanced between the dilepton system and recoiling jets, while the processes containing
 1378 real neutrinos are less balanced in the transverse plane. Thus, a requirement on f_{recoil} will reduce the Z/γ^*
 1379 + jets background while maintaining a good signal efficiency.




Figure 3.12: Comparison of f_{recoil} distributions for $Z/\gamma^* + \text{jets}$, $H \rightarrow WW^*$, and other backgrounds with real neutrinos.

1380 3.5.4 OPTIMIZING BACKGROUND REDUCTION SELECTION REQUIREMENTS

1381 The requirements on $p_T^{\text{miss}(\text{trk})}$ and f_{recoil} used to reduce the $Z + \text{jets}$ background must be optimized
 1382 to maximize expected signal significance in the same flavor channels. Figure 3.13 shows an optimization of
 1383 the combination of the two requirements in the gluon fusion zero jet bin. Each bin shows the expected
 1384 signal significance if the $p_{T,\text{rel}}^{\text{miss}(\text{trk})}$ is required to be greater than the left edge of the bin and the f_{recoil} is
 1385 required to be less than the top edge of the bin. The figure shows that the best signal significance comes
 1386 from requiring low values of f_{recoil} (< 0.05) and $p_{T,\text{rel}}^{\text{miss}(\text{trk})}$ values greater than 45 GeV.

1387 3.6 PARAMETERS OF INTEREST AND STATISTICAL TREATMENT

1388 As with any search or measurement, there are particular parameters of the Higgs that the $H \rightarrow WW^*$
 1389 analysis is interested in measuring. In this case, the parameters of interest are the mass of the Higgs boson
 1390 and its production cross section. Because the $H \rightarrow WW^* \rightarrow \ell\nu\ell\nu$ process does not have a closed final




Figure 3.13: Signal significance as a function of required value for f_{recoil} and $p_{T,\text{rel}}^{\text{miss}(\text{trk})}$ in the ggF $H \rightarrow WW^*$ with $n_j = 0$

state, it is not possible to measure the full invariant mass of the particle that may have produced the final state. However, a proxy for the invariant mass is defined using transverse plane information and detailed in section 3.6.1. The second parameter of interest is the ratio of the measured cross section to that expected from the Standard Model Higgs, which is denoted a μ . This is defined in equation 3.5.

$$\mu = \frac{\sigma}{\sigma_{\text{SM}}} \quad (3.5)$$

All of the likelihoods used in the statistical analysis of the final signal region events are parameterized as a function of μ . μ is a natural variable for hypothesis testing, as $\mu = 0$ corresponds to a background only hypothesis and $\mu = 1$ corresponds exactly to a Standard Model Higgs.

3.6.1 TRANSVERSE MASS

The $H \rightarrow WW^* \rightarrow \ell\nu\ell\nu$ analysis cannot reconstruct the full invariant mass of the Higgs because of the neutrinos in the final state. The transverse mass serves as a proxy for the full invariant mass by

1401 exploiting information from the transverse plane. The transverse mass is defined in equation 3.6.

$$m_T = \sqrt{(E_T^{\ell\ell} + p_T^{\text{miss}})^2 - |\vec{p}_T^{\ell\ell} + \vec{p}_T^{\text{miss}}|^2}, \quad (3.6)$$

1402 Here the $E_T^{\ell\ell}$ and $\vec{p}_T^{\ell\ell}$ are the transverse energy and momentum of the dilepton system, while p_T^{miss} is a
1403 proxy for the transverse momentum of the di-neutrino system. The track-based p_T^{miss} is used in the m_T
1404 rather than the calorimeter based E_T^{miss} because it has a better resolution on the true transverse mass.
1405 Figure 3.11b shows the improvement in the RMS of the difference between the true and reconstructed
1406 transverse mass in a ggF signal sample. The RMS improves by 4.7 GeV using p_T^{miss} in the m_T calculation.

1407 **3.6.2 STATISTICAL TREATMENT²**

1408 **LIKELIHOOD FUNCTION**

1409 The statistical analysis of final event candidates is framed as a hypothesis test, where the null hypoth-
1410 esis is background-only (no Standard Model Higgs). The first step in the analysis is to form a likelihood
1411 function for the data. In its simplest form, this likelihood is the probability of observing the number of
1412 events seen in the final signal region given knowledge of the signal strength. Because observation of events
1413 is fundamentally a Poisson counting experiment, this simple likelihood can be expressed as a Poisson prob-
1414 ability of observing N events given a total number of predicted signal and background events. This basic
1415 likelihood is shown in equation 3.7.

$$\mathcal{L}(\mu) = P(N|\mu S + B) \quad (3.7)$$

1416 Here, P is the Poisson probability density function, N is the total number of observed events, μ is the
1417 signal strength, S is the predicted number of signal events, and B is the predicted number of background
1418 events.

1419 In particle physics, certain background estimates are commonly normalized in so-called “control” re-
1420 gions and those predictions are scaled by the same normalization factor in the signal region. This leads to a

²Many thanks to Aaron Armbruster, whose thesis [64] inspired parts of this section.

1421 slightly more complicated likelihood, which is a function of both the signal strength and the background
1422 normalization. This is shown in equation 3.8.

$$\mathcal{L}(\mu, \theta) = P(N|\mu S + \theta B) P(N_{\text{CR}}|\theta B_{\text{CR}}) \quad (3.8)$$

1423 Here, θ serves as a “nuisance parameter”, or a parameter that is not of primary interest but still enters the
1424 likelihood. The second Poisson term enforces that the background normalization be consistent with the
1425 number of observed events in data in the control region, N_{CR} .

1426 So far, these two formulations of likelihoods have assumed a single signal region and do not take into
1427 account any shape information of potential discriminating variables. The $H \rightarrow WW^*$ analysis is divided
1428 into many different categories, the counting experiment described above can be performed in each individ-
1429 ual category. As mentioned in section 3.6.1, the transverse mass is used as the primary discriminating vari-
1430 able in many of the $H \rightarrow WW^*$ sub-analyses. The same counting experiment can be performed in each
1431 bin of the m_T distribution to incorporate some shape information. Thus, the total likelihood becomes a
1432 product over signal regions and bins of the m_T distribution. Finally, there are usually many backgrounds
1433 that are normalized in control regions. The new formulation of the likelihood takes this into account by
1434 including a product over control regions in the second Poisson term. All of these modifications are shown
1435 in equation 3.9.

$$\mathcal{L}(\mu, \theta) = \prod_{\substack{\text{SRs } i \\ \text{bins } b}} P\left(N_{ib} \middle| \mu S_{ib} + \sum_{\text{bkg } k} \theta_k B_{kib}\right) \prod_{\text{CRs } l} P\left(N_l \middle| \sum_{\text{bkg } k} \theta_k B_{kl}\right) \quad (3.9)$$

1436 The final step to obtain the full likelihood used in the analysis is to add nuisance parameters for the
1437 systematic uncertainties. In cases where the uncertainty does not affect the shape of m_T bin-by-bin, each
1438 systematic uncertainty ϵ is allowed to affect the expected event yields through a linear response function
1439 of the nuisance parameter, namely $\nu(\theta) = (1 + \epsilon)\theta$. If instead the uncertainty does affect the shape, the
1440 effect is instead parameterized by $\nu_b(\theta) = 1 + \epsilon_b\theta$. The value of the nuisance parameters for the systematic
1441 uncertainty are constrained with a Gaussian term that is added to the likelihood as well. This is of the form
1442 $g(\delta|\theta) = e^{-(\delta-\theta)^2/2}/\sqrt{2\pi}$, where δ is the central value and θ is a nuisance parameter. Finally, a last term is

¹⁴⁴³ added to account for the statistical uncertainty in the Monte Carlo samples used, which adds an additional
¹⁴⁴⁴ poisson term. The full likelihood used in the final statistical analysis is defined in equation 3.10.

$$\begin{aligned} \mathcal{L}(\mu, \boldsymbol{\theta}) = & \prod_{\substack{\text{SRs i} \\ \text{bins b}}} P \left(N_{ib} \middle| \mu S_{ib} \cdot \prod_{\substack{\text{sig.} \\ r}} \nu_{br}(\theta_r) + \sum_{\text{bkg k}} \theta_k B_{kib} \cdot \prod_{\substack{\text{bkg.} \\ s \\ \text{syst.}}} \nu_{bs}(\theta_s) \right) \\ & \cdot \prod_{\text{CRs l}} P \left(N_l \middle| \sum_{\text{bkg k}} \theta_k B_{kl} \right) \\ & \cdot \prod_{\substack{\text{syst} \\ t}} g(\delta_t | \theta_t) \cdot \prod_{\text{bkg k}} P(\xi_k | \zeta_k \theta_k) \end{aligned} \quad (3.10)$$

¹⁴⁴⁵ The fourth term of the equation quantifies the uncertainty due to finite Monte Carlo sample size. Here,
¹⁴⁴⁶ ξ represents the central value of the background prediction, θ is the associated nuisance parameter, $\zeta =$
¹⁴⁴⁷ $(B/\delta B)^2$, where δB is the statistical uncertainty of B .

¹⁴⁴⁸ The best fit value of the signal strength μ is determined by finding the values of μ and $\boldsymbol{\theta}$ that maximize
¹⁴⁴⁹ the likelihood, while setting $\delta = 0$ and $\xi = \zeta$. Once the likelihood is defined, a test statistic must be built
¹⁴⁵⁰ for use in hypothesis testing.

¹⁴⁵¹ TEST STATISTIC

¹⁴⁵² To distinguish whether the data match a background only or background and signal hypothesis, a test
¹⁴⁵³ statistic must be used. The $H \rightarrow WW^*$ analysis uses the profile likelihood technique [65]. The first step
¹⁴⁵⁴ in formulating this test statistic is to define the profile likelihood ratio, shown in equation 3.11.

$$\lambda(\mu) = \frac{\mathcal{L}(\mu, \hat{\theta}_\mu)}{\mathcal{L}(\hat{\mu}, \hat{\theta})} \quad (3.11)$$

¹⁴⁵⁵ Here $\hat{\theta}_\mu$ is the value of θ that maximizes the likelihood for the choice of μ being tested. Additionally, $\hat{\theta}$
¹⁴⁵⁶ and $\hat{\mu}$ represent the values of θ and μ that gives the overall maximum value of the likelihood.

1457 Once this is defined, a test statistic q_μ is constructed. This is shown in equation 3.12.

$$q_\mu = -2 \ln \lambda(\mu) \quad (3.12)$$

1458 A higher value of q_μ indicates that the data are more incompatible with the hypothesized value of μ , and
1459 q_0 then corresponds to the value of the test statistic for the background only hypothesis. A p_0 value is
1460 then defined to quantify the compatibility between the data and the null hypothesis. The p_0 value is the
1461 probability of obtaining a value of q_0 larger than the observed value, and this is shown in equation 3.13.

$$p_0 = \int_{q_0^{\text{obs}}}^{\infty} f(q_\mu | \mu = 0) dq_\mu \quad (3.13)$$

1462 Here $f(q_\mu)$ is the probability distribution function of the test statistic. Finally, the p_0 value can be con-
1463 verted into a signal significance, using the formula in equation 3.14, or the one-sided tail of the Gaussian
1464 distribution.

$$Z_0 = \sqrt{2} \operatorname{erf}^{-1}(1 - 2p_0) \quad (3.14)$$

1465 The threshold for discovery used in particle physics is $Z_0 \geq 5$, more commonly known as a value of 5σ .

The real voyage of discovery consists not in seeking new landscapes, but in having new eyes.

Marcel Proust

4

1466

1467 The discovery of the Higgs boson and the role 1468 of the $H \rightarrow WW^* \rightarrow \ell\nu\ell\nu$ channel

1469 4.1 INTRODUCTION

1470 This chapter presents the results of the search for the Higgs boson in 4.8 fb^{-1} collected at $\sqrt{s} = 7 \text{ TeV}$
1471 and 5.8 fb^{-1} at $\sqrt{s} = 8 \text{ TeV}$. The results of three searches at $\sqrt{s} = 8 \text{ TeV}$ in the $H \rightarrow WW^* \rightarrow \ell\nu\ell\nu$
1472, $H \rightarrow \gamma\gamma$, and $H \rightarrow ZZ \rightarrow 4\ell$ channels are combined with results of searches at $\sqrt{s} = 7 \text{ TeV}$
1473 in the same search channels (as well as the $H \rightarrow \tau\tau$ production and associated production searches for
1474 $H \rightarrow b\bar{b}$). The results of this combination are a 5.9σ detection of a new particle consistent with a Higgs
1475 boson. Rather than going into detail for all of the different Higgs decay searches, this chapter will discuss
1476 the three most sensitive channels and in particular focus on $H \rightarrow WW^* \rightarrow \ell\nu\ell\nu$. While the focus is on
1477 WW^* , some of the ZZ^* and $\gamma\gamma$ results are shown for completeness. The results not discussed here can

1478 be found in the ATLAS Higgs discovery publication [1].

1479 **4.2 DATA AND SIMULATION SAMPLES**

1480 The data sample used for the following results was taken in 2011 and 2012 at center of mass energies
1481 of 7 and 8 TeV, respectively, with 4.8 fb^{-1} collected at 7 TeV and 5.8 fb^{-1} collected at 8 TeV. Higgs
1482 production in the gluon fusion and vector boson fusion modes is modeled with **POWHEG** for the hard
1483 scattering event and **PYTHIA** for the showering and hadronization. Associated production of a Higgs with a
1484 vector boson or top quarks is modeled via **PYTHIA**. Table 4.1 shows the Monte Carlo generators used for
1485 modeling the signal and background processes relevant for the three analyses to be discussed.

Process	Generator
ggF, VBF H	POWHEG + PYTHIA
$WH, ZH, t\bar{t}H$	PYTHIA
$W + \text{jets}, Z/\gamma^* + \text{jets}$	ALPGEN + HERWIG
$t\bar{t}, tW, tb$	MC@NLO + HERWIG
tqb	ACERMC + PYTHIA
$q\bar{q} \rightarrow WW$	MC@NLO + HERWIG
$gg \rightarrow WW$	GG2WW+ HERWIG
$q\bar{q} \rightarrow ZZ$	POWHEG + PYTHIA
$gg \rightarrow ZZ$	GG2ZZ+ HERWIG
WZ	MADGRAPH+ PYTHIA , HERWIG
$W\gamma + \text{jets}$	ALPGEN + HERWIG
$W\gamma^*$	MADGRAPH+ PYTHIA
$q\bar{q}/gg \rightarrow \gamma\gamma$	SHERPA

Table 4.1: Monte carlo generators used to model signal and background for the Higgs search [1].

1486 **4.3 $H \rightarrow WW \rightarrow e\nu\mu\nu$ SEARCH**

1487 The $H \rightarrow WW \rightarrow e\nu\mu\nu$ search is unique compared to the ZZ and $\gamma\gamma$ channels. The Higgs mass
1488 cannot be fully reconstructed due to the presence of neutrinos in the final state, so the transverse mass m_T
1489 is used as the final discriminating variable. Compared to the other channels, there are more backgrounds
1490 here as well, as discussed in chapter 3. The same flavor final states are excluded from the 8 TeV dataset due

¹⁴⁹¹ to high pileup conditions¹. These final states are later included in results with the full Run 1 dataset, as
¹⁴⁹² discussed in chapters 5 and 6.

¹⁴⁹³ **4.3.1 EVENT SELECTION**

¹⁴⁹⁴ The analysis requires two opposite charge isolated leptons, with the leading (sub-leading) lepton required
¹⁴⁹⁵ to have $p_T > 25(15)$ GeV. The events are separated into different signal regions depending on which
¹⁴⁹⁶ flavor of lepton is leading ($e\mu$ for leading electron, μe for leading muon). Strict lepton quality cuts are
¹⁴⁹⁷ applied to the sample to reduce backgrounds from fake leptons.

¹⁴⁹⁸ Jets are reconstructed with the anti- k_T algorithm with a radius parameter $R = 0.4$. The jets are re-
¹⁴⁹⁹ quired to have $p_T > 25$ GeV and $|eta| < 4.5$, with jets in the tracking volume required to have a jet
¹⁵⁰⁰ vertex fraction of 0.5 and jets in the forward region required to have $p_T > 30$ GeV. The analysis is sepa-
¹⁵⁰¹ rated into three different signal regions based on jet multiplicity: $n_j = 0, 1, \geq 2$.

¹⁵⁰² To indicate the presence of neutrinos in the event, a requirement of $E_{T,\text{rel}}^{\text{miss}} > 25$ GeV is made². This
¹⁵⁰³ requirement significantly reduces the QCD multijet and Z/γ^* + jets backgrounds. Figure 4.1 shows the
¹⁵⁰⁴ distribution of n_j in data and simulation after applying these “pre-selection” requirements.




Figure 4.1: Jet multiplicity distribution in data and MC after applying lepton, jet, and $E_{T,\text{rel}}^{\text{miss}}$ selections. The WW and top backgrounds have been normalized using control samples, and the hashed band indicates the total uncertainty on the prediction. [[1](#)]

¹The less sensitive 7 TeV search result includes both different flavor and same flavor final states.

²For the definition of $E_{T,\text{rel}}^{\text{miss}}$, see chapter 3

1505 Additional selections are applied to require the dilepton topology to correspond to that of a SM Higgs.
1506 The requirements are presented here - more detailed discussion on the motivation for each requirement is
1507 saved for Chapter 5. In all of the jet multiplicity channels, the dilepton system is required to have a small
1508 gap in azimuthal angle, $\Delta\phi_{\ell\ell} < 1.8$. Similarly, the $m_{\ell\ell}$ is required to be less than 50 GeV in the lower jet
1509 multiplicity channels and less than 80 GeV in the $n_j \geq 2$ channel. In the $n_j = 0$ channel, the magnitude
1510 of the dilepton p_T , $p_T^{\ell\ell}$, is required to be greater than 30 GeV.

1511 In the higher jet multiplicity channels ($n_j \geq 1$), the top background is a more important component
1512 and must be reduced. The total transverse momentum p_T^{sum} is thus required to be less than 30 GeV. Ad-
1513 ditionally, the di- τ invariant mass $m_{\tau\tau}$ (dilepton mass computed under the assumption that the neutrinos
1514 from the τ decay are emitted collinear to the charged leptons) is used to reject $Z \rightarrow \tau\tau$ events by requiring
1515 $|m_{\tau\tau} - m_Z| > 25$ GeV. These variables are also discussed in more detail in Chapter 5.

1516 In the $n_j \geq 2$ channel, requirements are made to isolate the VBF contribution to Higgs production.
1517 The kinematics of the two leading jets are used to make these requirements. In particular, the event must
1518 have $\Delta y_{jj} > 3.8$ and $m_{jj} > 500$ GeV, along with a veto on having any additional jets with rapidity
1519 between the two leading jets.

1520 **4.3.2 BACKGROUND ESTIMATION**

1521 The details of the background estimation techniques used in the $H \rightarrow WW^* \rightarrow \ell\nu\ell\nu$ analysis are
1522 discussed in section 5.5. As that section refers to a later iteration of the analysis, a general discussion is given
1523 here for completeness. The dominant backgrounds are SM WW production and top (both pair and
1524 single) production, and these backgrounds have their normalizations estimated from dedicated control
1525 regions while their shapes are taken from simulation.

1526 The control sample for the Standard Model WW background is defined by making the same require-
1527 ments as the signal region with the $m_{\ell\ell}$ requirement inverted (now requiring $m_{\ell\ell} > 80$ GeV) and remov-
1528 ing the $\Delta\phi_{\ell\ell}$ requirement. This creates a control sample that is 70% (40%) pure in the 0(1)-jet region. The
1529 correction to the pure MC-based background estimate is quantified by defining a normalization factor β
1530 which is the ratio of the data yield to the MC yield ($N_{\text{data}}/N_{\text{MC}}$) in this control sample. Table 4.2 shows

1531 the WW normalization factors in the $n_j = 0$ and $n_j = 1$ bins (the $n_j \geq 2$ estimate is taken directly
1532 from MC).

n_j	β_{WW}	β_t
$= 0$	1.06 ± 0.06	1.11 ± 0.06
$= 1$	0.99 ± 0.15	1.11 ± 0.05
≥ 2	-	1.01 ± 0.26

Table 4.2: Normalization factors (ratio of data and MC yields in a control sample) for the Standard Model WW and top backgrounds in the $H \rightarrow WW^* \rightarrow \ell\nu\ell\nu$ analysis [1]. Only statistical uncertainties are shown.

1533 The top background estimate is also computed separately in each jet multiplicity bin. In the $n_j = 0$
1534 channel, the background is first normalized using data after pre-selection requirements with no selection
1535 on n_j . Then, a dedicated b -tagged control sample is used to evaluate the ratio of one-jet to two-jet events in
1536 data. The details of this technique are shown in reference [66]. In the $n_j = 1$ and the $n_j \geq 2$ regions, the
1537 top background is normalized in a control sample where the signal region selections are applied, but the
1538 b -jet veto is reversed and the Higgs topology requirements on $m_{\ell\ell}$ and $\Delta\phi_{\ell\ell}$ are removed. The resulting
1539 normalization factors for these techniques are shown in table 4.2.




Figure 4.2: Comparison of m_T between data and simulation in the $n_j = 0$ WW (a) and $n_j = 1$ top (b) control samples [1]

1540 The control samples which are used for background normalization can also be used to validate the mod-
1541 eling of the m_T distribution for each background. Figure 4.2 shows the comparison between data and MC

1542 for the m_T distribution after correcting the normalization of the backgrounds in the WW and top control
1543 regions. Good agreement between data and simulation is seen in both cases.

1544 The $W + \text{jets}$ background estimate is taken entirely from data using a control sample with one well recon-
1545 structed lepton and one anti-identified lepton. All other backgrounds are taken purely from simulation.

1546 4.3.3 SYSTEMATIC UNCERTAINTIES

1547 The systematic uncertainties that have the largest impact on the analysis are the theoretical uncertainties
1548 associated with the signal cross section, and these are shared with the ZZ^* and $\gamma\gamma$ channels. The uncer-
1549 tainties resulting from variations of the QCD scale are $+7\%/-8\%$ on the final singal yield. Those coming
1550 from variations of the parton distribution function (PDF) used in the simulation add a $\pm 8\%$ uncertainty
1551 on the yield. The uncertainties on the branching ratios of the Higgs are $\pm 5\%$.

1552 The main experimental uncertainties come from variations of the jet energy scale (JES), jet energy reso-
1553 lution, pile-up, E_T^{miss} , b -tagging efficiency, $W + \text{jets}$ background estimate, and integrated luminosity. For
1554 more details, see reference [1].

1555 4.3.4 RESULTS

1556 Table 4.3 shows the signal and background yields in the final signal region after normalizing the back-
1557 grounds according to the methods described above.

	$n_j = 0$	$n_j = 1$	$n_j \geq 2$
Signal	20 ± 4	5 ± 2	0.34 ± 0.07
WW	101 ± 13	12 ± 5	0.10 ± 0.14
Other dibosons	12 ± 3	1.9 ± 1.1	0.10 ± 0.10
$t\bar{t}$	8 ± 2	6 ± 2	0.15 ± 0.10
Single top	3.4 ± 1.5	3.7 ± 1.6	-
$Z/\gamma^* + \text{jets}$	1.9 ± 1.3	0.10 ± 0.10	-
$W + \text{jets}$	15 ± 7	2 ± 1	-
Total background	142 ± 16	26 ± 6	0.35 ± 0.18
Observed in data	185	38	0

Table 4.3: Data and expected yields for signal and background in the final $H \rightarrow WW^* \rightarrow \ell\nu\ell\nu$ signal region.
Uncertainties shown are both statistical and systematic. [1]

1558 Figure 4.3 shows the m_T distribution in the $n_j \leq 1$ channels for 8 TeV data. (No events are observed
 1559 in data in the $n_j \geq 2$ channels in this dataset). The excess shown here relatively flat as a function of
 1560 hypothesized Higgs mass. The combined 7 and 8 TeV data gives an excess with local significance of 2.8σ
 1561 with an expected significance of 2.3σ , corresponding to a μ measurement of 1.3 ± 0.5 .




Figure 4.3: m_T distribution in the $H \rightarrow WW \rightarrow e\nu\mu\nu$ $n_j \leq 1$ channels for 8 TeV data [1].

1562 4.4 $H \rightarrow \gamma\gamma$ SEARCH

1563 The $H \rightarrow \gamma\gamma$ search is in essence a search for a peaked excess above the falling SM diphoton mass
 1564 spectrum, with $m_{\gamma\gamma}$ as the ultimate discriminating variable. Events are selected by requiring two isolated
 1565 photons, with the leading (sub-leading) photon required to have $E_T > 40(30)$ GeV. In the 8 TeV data,
 1566 the photons are required to pass cut-based identification criteria consistent with a photon in the electro-
 1567 magnetic calorimeter and little leakage in the hadronic calorimeter.

1568 The main challenges for this analysis are accurate mass reconstruction and background estimation. In
 1569 order to accurately reconstruct the invariant mass of the di-photon system, both the energy and direction
 1570 of the photons must be measured well. Therefore, the identification of the primary vertex of the hard
 1571 interaction is particularly important, and is done using a multivariate likelihood which combines informa-
 1572 tion about the photon direction and vertex position. The background is modeled with a falling spectrum
 1573 in $m_{\gamma\gamma}$ that is parameterized by different functions depending on the category of the event.

1574 4.4.1 RESULTS

1575 The resulting diphoton mass spectrum is shown in figure 4.4. The best fit mass value in the $\gamma\gamma$ channel
 1576 alone in the combined 7 and 8 TeV data is 126.5 GeV. The local significance at this point is 4.5σ , with
 1577 an expected significance of 2.5σ . Therefore, the measured signal strength μ is 1.8 ± 0.5 in this channel.




Figure 4.4: Diphoton mass spectrum in 7 and 8 TeV data. Panel a) shows the unweighted data distribution superimposed on the background fit, while panel c) shows the data where each event category is weighted by its signal to background ratio. Panels b) and d) show the respective distributions with background subtracted [1].

1578 4.5 $H \rightarrow ZZ \rightarrow 4\ell$ SEARCH

1579 The $H \rightarrow ZZ \rightarrow 4\ell$ analysis searches for a Standard Model Higgs boson decaying to two Z bosons,
 1580 each of which decays to a pair of same flavor, opposite charge isolated leptons. The ultimate discriminating
 1581 variable is $m_{4\ell}$, or the invariant mass of the four selected leptons. The ℓ denotes an e or μ as with the
 1582 $H \rightarrow WW^* \rightarrow \ell\nu\ell\nu$ analysis.

1583 Four distinct signal regions are constructed depending on the flavors of the final state, additionally sep-
 1584 arated by the flavor of the leading lepton pair. These are referred to as $4e$, $2e2\mu$, $2\mu2e$, 4μ .

1585 The main backgrounds in the $H \rightarrow ZZ \rightarrow 4\ell$ search are continuum ZZ^* production, $Z + \text{jets}$ pro-
 1586 duction, and $t\bar{t}$. The $m_{4\ell}$ distribution for background is estimated from simulation. The normalization
 1587 of the SM ZZ^* background is also taken from MC simulation, while the $Z + \text{jets}$ and $t\bar{t}$ normalizations are
 1588 taken from data-driven methods.

1589 **4.5.1 RESULTS**

1590 Figure 4.5 shows the $m_{4\ell}$ spectrum measured in the 7 and 8 TeV datasets. The total number of events
 1591 observed in the window between 120 and 130 GeV is 13, with 6 events in the 4μ channel, 2 events in
 1592 the $4e$ channel, and 5 events in the $2e2\mu/2\mu2e$. The best fit μ value in the combined 7 and 8 TeV data
 1593 occurs at 125 GeV and is measured to be 1.2 ± 0.6 . The observed significance at this mass is 3.6σ , with
 1594 an expected significance of 2.7σ .




Figure 4.5: Four lepton invariant mass spectrum ($m_{4\ell}$) in 7 and 8 TeV data compared to background estimate. A 125 GeV SM Higgs signal is shown in blue [1].

1595 4.6 COMBINED RESULTS

1596 The statistical interpretation of the combined results is undertaken as described in section 3.6.2, with a
 1597 hypothesis test based on a likelihood ratio parameterized by the Higgs signal strength μ . The null hypoth-
 1598 esis corresponds to $\mu = 0$, while the SM Higgs corresponds to $\mu = 1$.

1599 Table 4.4 summarizes the properties of the individual channels as well as the significances of the excesses
 1600 seen. The most significant observed local excess comes from the $\gamma\gamma$ channel. Figure 4.6 shows a compari-
 1601 son of the observed local p_0 values as a function of hypothesized mass for the three different search chan-
 1602 nels. Both the ZZ^* and $\gamma\gamma$ channels have very peaked excesses, while the WW^* excess can be seen as very
 1603 broad because the m_T distribution does not provide detailed information about the true Higgs mass.

Channel	Fit var.	Observed Z_l	Expected Z_l	$\hat{\mu}$
$H \rightarrow ZZ^* \rightarrow 4\ell$	$m_{4\ell}$	3.6	2.7	1.2 ± 0.6
$H \rightarrow \gamma\gamma$	$m_{\gamma\gamma}$	4.5	2.5	1.8 ± 0.5
$H \rightarrow WW^* \rightarrow e\nu\mu\nu$	m_T	2.8	2.3	1.3 ± 0.5
Combined	-	6.0	4.9	1.4 ± 0.3

Table 4.4: Summary of the expected and observed significance and measured signal strengths in the combined 7 and 8 TeV datasets for the Higgs discovery analysis [1].

1604 Figure 4.7 shows the combined exclusion limit, p_0 , and signal strength. The highest local excess comes
 1605 at a value of 126.5 GeV and corresponds to a 6.0σ observed excess.

1606 Figure 4.8 shows a comparison of the measured signal strengths between the different Higgs search
 1607 channels. All measured μ are consistent with unity within their uncertainty, and the combined μ mea-
 1608 surement is 1.4 ± 0.3 .

1609 The likelihood can also be computed in a two-dimensional plane of m_H and μ , and this is shown in
 1610 figure 4.9. The figure shows that while the $\gamma\gamma$ and ZZ^* channels have very good mass resolution, the
 1611 excess in WW^* covers a broad mass range. The banana shape of the WW^* result is due to the fact that
 1612 the excess in this channel can either be explained by increasing the signal strength or by changing the mass
 1613 (and thus the cross section). The two parameters are correlated due to the lack of mass sensitivity in this
 1614 channel.

1615 Because multiple Higgs mass points are searched for, the local significance must be corrected for a look-




Figure 4.6: Local p_0 distribution as a function of hypothesized Higgs mass for the $H \rightarrow ZZ^* \rightarrow 4\ell$ (a), $H \rightarrow \gamma\gamma$ (b), and $H \rightarrow WW^* \rightarrow \ell\nu\ell\nu$ (c) channels. Dashed curves show expected results, while solid curves show observed. Red curves are from 7 TeV data, blue curves from 8 TeV, and black curved combined [1].

¹⁶¹⁶ elsewhere effect to compute a true global significance. The global significance for finding a Higgs anywhere
¹⁶¹⁷ in the mass range of 110 GeV to 600 GeV is 5.1σ . This increases slightly to 5.3σ if only mass range from
¹⁶¹⁸ 110 to 150 GeV.




Figure 4.7: Combined 95% CL limits (a), local p_0 values (b), and signal strength measurement (c) as a function of Higgs mass [1].

¹⁶¹⁹ **4.7 CONCLUSION**

¹⁶²⁰ A search for the production of a Standard Model Higgs boson was conducted in 4.8 fb^{-1} collected at
¹⁶²¹ $\sqrt{s} = 7 \text{ TeV}$ and 5.8 fb^{-1} at $\sqrt{s} = 8 \text{ TeV}$. A new particle consistent with the Higgs boson was observed,
¹⁶²² with a mass of 126.5 GeV and a global (local) significance of $5.1(6.0)\sigma$. This is the first discovery level
¹⁶²³ observation of a particle consistent with the Higgs.




Figure 4.8: Comparison of measured signal strength μ for a 126 GeV Higgs in the 7 and 8 TeV datasets [1].




Figure 4.9: Two dimensional likelihood as a function of signal strength μ and Higgs mass m_H [1].

*The imagination of nature is far, far greater than the
imagination of man.*

Richard Feynman

5

1624

1625 Evidence for Vector Boson Fusion production

1626

$$\text{of } H \rightarrow WW^* \rightarrow \ell\nu\ell\nu$$

1627

5.1 INTRODUCTION

1628 After the discovery of a particle consistent with the Higgs boson, the $H \rightarrow WW^*$ analysis had two main
1629 goals. The first goal was to increase the sensitivity of the analysis to fully confirm that the $H \rightarrow WW^*$
1630 process did indeed exist. The second goal was to characterize the particle as much as possible, including
1631 searching for the lower cross-section production modes, in order to confirm that it was indeed a Higgs
1632 boson. This chapter presents a dedicated search for Vector Boson Fusion (VBF) production of a Higgs
1633 boson decaying via the $H \rightarrow WW^* \rightarrow \ell\nu\ell\nu$ mode. First, basics of the topology of VBF production
1634 are presented. Then, the details of the analysis are shown, including signal region definition, background
1635 estimation techniques, and systematic uncertainties. Finally, the results of the analysis are presented. As

1636 will be shown, this analysis is the first and most sensitive evidence of the VBF production mode of the
1637 Higgs on ATLAS.

1638 In the VBF channel, there are both a selection requirement based signal region analysis (known as the
1639 “cut-based”) and a multivariate analysis which uses a boosted decision tree (known as the BDT analysis).
1640 The focus of this chapter will be on the cut-based signal region, as this is an important component of the
1641 VBF analysis and in particular acts as strong validation for the final BDT result. Connections between the
1642 cut-based and BDT analyses will be discussed where appropriate.

1643 5.2 DATA AND SIMULATION SAMPLES

1644 The results presented here are with 20.3 fb^{-1} taken at $\sqrt{s} = 8 \text{ TeV}$ and 4.5 fb^{-1} taken at $\sqrt{s} = 7 \text{ TeV}$.
1645 The details of the LHC and detector conditions during this period are given in Chapter 2. The trigger
1646 selection defining the dataset is discussed in section 5.2.1. The simulation samples used for signal and back-
1647 ground modeling are given in section 5.2.2.

1648 5.2.1 TRIGGERS

1649 The analysis uses a combination of single lepton and dilepton triggers to allow lowering of the p_T
1650 thresholds and increased signal acceptance. The p_T threshold on the leptons is a particularly important
1651 consideration for this signal. Because the second W produced in the decay can be off-shell, it tends to pro-
1652 duce lower momentum leptons. Thus, being able to lower the p_T threshold while still maintaining a low
1653 background rate is critical. Figure 5.1 shows an example of the subleading lepton p_T for a VBF $H \rightarrow WW^*$
1654 signal compared to the corresponding $t\bar{t}$ background. Note that the lepton p_T spectrum is considerably
1655 softer in the signal sample.

1656 As discussed in Chapter 2, there are multiple levels in the ATLAS trigger system, and there are different
1657 p_T thresholds imposed for the leptons at each level. Additionally, some triggers have a loose selection
1658 on the isolation of the lepton (looser than that applied offline in the analysis object selection). Table 5.1
1659 shows the thresholds used for single lepton triggers, while table 5.2 shows the thresholds coming from
1660 di-lepton triggers. The single lepton trigger efficiency for muons that pass the analysis object selection is



Figure 5.1: A comparison of the subleading lepton p_T spectrum between VBF $H \rightarrow WW^*$ production and $t\bar{t}$ background

1661 70% for muons in the barrel region ($|\eta| < 1.05$) and 90% in the endcap region. The electron trigger
 1662 efficiency increases with electron p_T but the average is approximately 90%. These efficiencies are measured
 1663 by combined performance and trigger signature groups [67, 68].

	Level-1 threshold	High-level threshold
Electron	18	$24i$
	30	60
Muon	15	$24i$
		36

Table 5.1: Single lepton triggers used for electrons and muons. A logical “or” of the triggers listed for each lepton type is taken. Units are in GeV, and the i denotes an isolation requirement in the trigger.

	Level-1 threshold	High-level threshold
ee	10 and 10	12 and 12
$\mu\mu$	15	18 and 8
$e\mu$	10 and 6	12 and 8

Table 5.2: Di-lepton triggers used for different flavor combinations. The two thresholds listed refer to leading and sub-leading leptons, respectively. The di-muon trigger only requires a single lepton at level-1.

1664 The combination of all triggers shown gives good efficiency for signal events. This efficiency is sum-
 1665 marized in table 5.3. The relative improvement in efficiency by adding the dilepton triggers is also shown

1666 in the same table. The largest gain comes in the $\mu\mu$ channel. Overall the trigger selection shows a good
1667 efficiency for $H \rightarrow WW^*$ signal events.

Channel	Trigger efficiency	Gain from 2ℓ trigger
ee	97%	9.1%
$\mu\mu$	89%	18.5%
$e\mu$	95%	8.3%
μe	81%	8.2%

Table 5.3: Trigger efficiency for signal events and relative gain of adding a dilepton trigger on top of the single lepton trigger selection. The first lepton is the leading, while the second is the sub-leading. Efficiencies shown here are for the ggF signal in the $n_j = 0$ category but are comparable for the VBF signal.

1668 **5.2.2 MONTE CARLO SAMPLES**

1669 Modeling of signal and background processes in the signal region, in particular for the m_T distribution,
1670 is an important consideration for the final interpretation of the analysis. Therefore, careful consideration
1671 must be paid to which Monte Carlo (MC) generators are used for specific processes. With the exception of
1672 the $W + \text{jet}$ and multijet backgrounds, the m_T shape used as the final discriminant is taken from simulation.
1673 (Many backgrounds are normalized from data, as described in section 5.5).

1674 Table 5.4 shows the MC generators used for the signal and background processes, as well as their cross
1675 sections. In order to include corrections up to next-to-leading order (NLO) in the QCD coupling constant
1676 α_s , the POWHEG [69] generator is often used. In some cases, only leading order generators like ACERMC
1677 [70] and GG2VV [71] are available for the process in question. If the process requires good modeling for
1678 very high parton multiplicities, the SHERPA [72] and ALPGEN [73] generators are used to provide merged
1679 calculations for five or fewer additional partons. These matrix element level calculations must then be
1680 additionally matched to models of the underlying event, hadronization, and parton shower. There are
1681 four possible generators for this: SHERPA, PYTHIA 6 [74], PYTHIA 8 [75], or HERWIG [76] + JIMMY [77].
1682 The simulation additionally requires an input parton distribution function (PDF). The CTEQ [78] PDFs
1683 are used for SHERPA and POWHEG simulated samples, while CTEQ6Li [79] is used for ALPGEN + HERWIG
1684 and ACERMC simulations. The Drell-Yan samples are reweighted to the MRST [80] PDFs, as these are
1685 found to give the best agreement between data and simulation.

Process	MC generator	$\sigma \cdot \mathcal{B}$ (pb)
Signal		
ggF $H \rightarrow WW^*$	POWHEG +PYTHIA 8	0.435
VBF $H \rightarrow WW^*$	POWHEG +PYTHIA 8	0.0356
VH $H \rightarrow WW^*$	PYTHIA 8	0.0253
WW		
$q\bar{q} \rightarrow WW$ and $qg \rightarrow WW$	POWHEG +PYTHIA 6	5.68
$gg \rightarrow WW$	GG2VV +HERWIG	0.196
$(q\bar{q} \rightarrow W) + (q\bar{q} \rightarrow W)$	PYTHIA 8	0.480
$q\bar{q} \rightarrow WW$	SHERPA	5.68
VBS $WW + 2$ jets	SHERPA	0.0397
Top quarks		
$t\bar{t}$	POWHEG +PYTHIA 6	26.6
Wt	POWHEG +PYTHIA 6	2.35
$t\bar{q}\bar{b}$	ACERMC +PYTHIA 6	28.4
$t\bar{b}$	POWHEG +PYTHIA 6	1.82
Other dibosons (VV)		
$W\gamma$ ($p_T^\gamma > 8$ GeV)	ALPGEN +HERWIG	369
$W\gamma^*$ ($m_{\ell\ell} \leq 7$ GeV)	SHERPA	12.2
WZ ($m_{\ell\ell} > 7$ GeV)	POWHEG +PYTHIA 8	12.7
VBS $WZ + 2$ jets	SHERPA	0.0126
($m_{\ell\ell} > 7$ GeV)		
$Z\gamma$ ($p_T^\gamma > 8$ GeV)	SHERPA	163
$Z\gamma^*$ (min. $m_{\ell\ell} \leq 4$ GeV)	SHERPA	7.31
ZZ ($m_{\ell\ell} > 4$ GeV)	POWHEG +PYTHIA 8	0.733
$ZZ \rightarrow \ell\ell\nu\nu$ ($m_{\ell\ell} > 4$ GeV)	POWHEG +PYTHIA 8	0.504
Drell-Yan		
Z ($m_{\ell\ell} > 10$ GeV)	ALPGEN +HERWIG	16500
VBF $Z + 2$ jets	SHERPA	5.36
($m_{\ell\ell} > 7$ GeV)		

Table 5.4: Monte Carlo samples used to model the signal and background processes [63].

Once the basic hard scattering process is simulated, it must be passed through a detector simulation and additional pile-up events must be overlaid. The pile-up events are modeled with PYTHIA 8, and the ATLAS detector is simulated with GEANT4 [81]. Because of the unique phase space of the $H \rightarrow WW^*$

1689 analysis, events are sometimes filtered at generator level to allow for more efficient generation of relevant
1690 events. The efficiency of the trigger in MC simulation does not always match the measured efficiency in
1691 data, so trigger scale factors are applied to correct the MC efficiency to the data. These are derived by the
1692 combined performance groups [67, 68].

1693 5.3 OBJECT SELECTION

1694 In order to define the signal region, the analysis must first select the objects to be considered. The details
1695 of the object reconstruction algorithms are discussed in Chapter 2, while this section gives specific selection
1696 cuts used in the $H \rightarrow WW^*$ analysis.

1697 The first step in this process is to select a primary vertex candidates. The event’s primary vertex is the
1698 vertex with the largest sum of p_T^2 for associated tracks and is required to have at least three tracks with
1699 $p_T > 450$ MeV. Many of the object selection cuts are then made relative to this chosen primary vertex.

1700 5.3.1 MUONS

1701 The analysis uses combined muon candidates, where a track in the Inner Detector has been matched to
1702 a standalone track in the Muon Spectrometer. The track parameters are combined statistically in the muon
1703 reconstruction algorithm [54]. The muons are required to be within $|\eta| < 2.5$ and have a $p_T > 10$ GeV.
1704 To reduce backgrounds coming from mis-reconstructed leptons, there are requirements on the impact
1705 parameter of the muon relative to the primary vertex. The transverse impact parameter d_0 is required to
1706 be small relative to its estimated uncertainty, the exact cut value being $d_0/\sigma_{d_0} < 3$. The longitudinal
1707 impact parameter z_0 must satisfy $|z_0 \sin \theta| < 1$ mm.

1708 As discussed previously, the muons must also be isolated. There are two types of lepton isolations that
1709 are calculated: track-based and calorimeter-based. For muons, the track-based isolation is defined using the
1710 scalar sum $\sum p_T$ for tracks with $p_T > 1$ GeV (excluding the muon’s track) within a cone of $\Delta R = 0.3$
1711 (0.4) for muon with $p_T > 15$ GeV ($10 < p_T < 15$ GeV). The final isolation requirement is made my
1712 requiring that this scalar sum be no more than a certain fraction of the muon’s p_T . This requirement varies
1713 with muon p_T and the exact cuts are defined in table 5.5.

1714 The calorimeter-based muon isolation is defined using as a $\sum E_T$ calculated from calorimeter cells us-
 1715 ing the same cone size as the track-based isolation but excluding cells with $\Delta R < 0.05$ around the muon.
 1716 This requirement is also defined as a cut on the ratio of the sum to the muon p_T and varies with muon p_T .
 1717 The cut values are also given in table 5.5.
 1718 The isolation requirements loosen as a function of p_T to allow for larger signal acceptance. At low p_T ,
 1719 the isolation is tightened to reduce the $W + \text{jets}$ background which arises from a misidentified lepton.

p_T range (GeV)	Calorimeter isolation	Track isolation
10 – 15	0.06	0.06
15 – 20	0.12	0.08
20 – 25	0.18	0.12
> 25	0.30	0.12

Table 5.5: p_T dependent isolation requirements for muons. Muons are required to have the amount of calorimeter or track based cone sums be less than this fraction of their p_T .

1720 5.3.2 ELECTRONS

1721 Electrons are identified by matching reconstructed clusters in the electromagnetic calorimeter with tracks
 1722 in the inner detector. The electrons are identified using a likelihood based method [51, 52] which takes into
 1723 account the shower shapes in the calorimeter, the matching of tracks to clusters, and the amount of transi-
 1724 tion radiation in the TRT. The electrons are required to have $|\eta| < 2.47$, and candidates in the transition
 1725 region between the barrel and endcap ($1.37 < |\eta| < 1.52$) are excluded. As the muons, the electrons
 1726 are required to have transverse impact parameter significance < 3 , while in the longitudinal direction
 1727 they must have $|z_0 \sin \theta| < 0.4$ mm. Some electron requirements also vary with electron E_T , and these
 1728 requirements are summarized in table 5.6.

1729 The isolation for electrons are defined similarly to the muons but with unique cuts on the objects in-
 1730 cluded. The track-based isolation is defined using tracks with $p_T > 400$ MeV with cone sizes as defined
 1731 previously. The calorimeter-based isolation also uses the same cone size as the muon, but here the cells
 1732 within a 0.125×0.175 area in $\eta \times \phi$ around the electron cluster's barycenter are excluded. The other
 1733 difference with respect to muons is that the denominator of the isolation ratio is the electron's E_T rather
 1734 than p_T . The isolation cuts very with electron E_T and are defined in table 5.6.

1735 The electron is also required to not be consistent with a vertex coming from a photon conversion.

p_T range (GeV)	Quality cut	Calorimeter isolation	Track isolation
10 – 15	Very tight LH	0.20	0.06
15 – 20	Very tight LH	0.24	0.08
20 – 25	Very tight LH	0.28	0.10
> 25	Medium	0.28	0.10

Table 5.6: p_T dependent requirements for electrons. Electrons are required to have the amount of calorimeter or track based cone sums be less than this fraction of their E_T .

1736 5.3.3 JETS

1737 Jets are clustered with the anti- k_T reconstruction algorithm using a radius parameter of $R = 0.4$. They
1738 are required to have a jet vertex fraction (JVF) of at least 50%, meaning that half of the tracks associated with
1739 the jet originated from the primary vertex. Jets with no tracks associated (i.e. those outside the acceptance
1740 of the ID) do not have this requirement applied. Jets are required to have $p_T > 25$ GeV if they are within
1741 the tracking acceptance ($|\eta| < 2.4$). Jets with $2.4 < |\eta| < 4.5$ are required to have $p_T > 30$ GeV.
1742 This tighter requirement reduces jets from pileup in the region where JVF requirements cannot be applied.
1743 The two highest p_T jets in the event are referred to as the “VBF” jets and used to compute various analysis
1744 selections later.

1745 Identification of b -jets is done using the MV1 algorithm and is limited to the acceptance of the ID ($|\eta| <$
1746 2.5). The operating point of MV1 that is used is the one that is 85% efficient for identifying true b -jets. This
1747 operating point has a 10.3% of mis-tagging a light quark jet as a b -jet. In order to improve the rejection of b -
1748 jets, a lower threshold than the nominal p_T threshold described above is used. For the purposes of counting
1749 the number of b -jets, jets with p_T down to 20 GeV are used.

1750 5.3.4 OVERLAP REMOVAL

1751 There are some cases where certain reconstructed objects will overlap and one will have to be chosen
1752 (for example, an electron and a jet in the calorimeter). First, the case of lepton overlap is dealt with. If
1753 an electron candidate extends into the muon spectrometer, it is removed. If a muon or electron have a
1754 $\Delta R < 0.1$, the electron is removed and the muon is kept. If two electron candidates overlap within the

1755 same radius, then the higher E_T electron is kept. Next, the overlap between leptons and jets is considered.
1756 If an electron and jet are within $\Delta R < 0.3$ of one another, the electron is kept and the jet is removed.
1757 However, if a muon and jet overlap within $\Delta R < 0.3$, the jet is kept (as it is likely that the muon is the
1758 result of a semileptonic decay inside the jet).

1759 Once the overlap removal is complete, the final set of objects used in the analysis is defined.

1760 5.4 ANALYSIS SELECTION

1761 The VBF analysis uses two distinct selections. The first is a more standard selection, referred to as “cut-
1762 based”, that applies requirements on the VBF variables and uses m_T as the final discriminating variable.
1763 The second is a looser selection that uses a Boosted Decision Tree (BDT) score as the final discriminator in
1764 order to take advantage of the detailed correlations between the VBF variables. While the BDT analysis is
1765 ultimately more sensitive, the cut-based serves as an important component of the analysis. First, the cut-
1766 based allows for confirming the modeling and validity of many variables used as input to the BDT. Second,
1767 because this is the first use of such an MVA technique in the $H \rightarrow WW^*$ analysis, the cut-based selection
1768 allows confirmation of the final BDT result with a more traditional analysis. The cut-based techniques are
1769 the focus of this chapter, but connections to the BDT result will be illustrated when appropriate.

1770 One important note is that because this analysis is dedicated to the measurement of the VBF pro-
1771 duction mode of the Higgs, events coming from gluon fusion production with the Higgs decaying via
1772 $H \rightarrow WW^* \rightarrow \ell\nu\ell\nu$ are treated as background events. This will be seen throughout the various predic-
1773 tions shown.

1774 5.4.1 COMMON PRE-SELECTION

1775 Both the cut-based and BDT analyses have a common pre-selection that is applied before the main signal
1776 region requirements. The requirements on leptons are common to all n_j bins. The analysis requires two
1777 oppositely charged leptons, with the leading lepton required to have $p_T > 22$ GeV while the subleading
1778 lepton must have $p_T > 10$ GeV. Next, to remove low mass Z/γ^* events, a cut on the dilepton mass
1779 $m_{\ell\ell} > 10$ (12) GeV is applied in the different (same) flavor channel. In the same flavor channels, there is

1780 an additional veto placed on the region around the Z peak, requiring that $|m_{\ell\ell} - m_Z| > 15$ GeV.

1781 There are also requirements on the amount of missing transverse momentum in the event. These are
1782 only applied in the same flavor channels, as in the different flavor channels $t\bar{t}$ is the dominant background
1783 in $n_j \geq 2$. The BDT analysis requires $p_T^{\text{miss}} > 40$ GeV and $E_T^{\text{miss}} > 45$ GeV. The cut-based analysis
1784 must select more tightly on these variables to have maximal sensitivity and thus requires $p_T^{\text{miss}} > 50$ GeV
1785 and $E_T^{\text{miss}} > 55$ GeV.

1786 Finally, because this analysis is focused on VBF, a requirement on the jet multiplicity is placed, with
1787 $n_j \geq 2$. Additionally, the analysis requires that there are no jets identified as b-quarks in the event, or
1788 $n_b = 0$.

1789 5.4.2 CUT-BASED SELECTION

1790 The cut-based selection places sequential requirements on variables reconstructed from the VBF jets in
1791 order to increase the signal to background ratio.

1792 GENERAL BACKGROUND REDUCTION

1793 Top pair production is the primary background in the $n_j \geq 2$ bin. Even though $n_b = 0$ is required, an
1794 additional variable is constructed to further suppress the top background. There is often additional QCD
1795 radiation that accompanies the $t\bar{t}$ system when it is produced. Therefore, a variable which tests for the
1796 presence of this additional radiation, p_T^{sum} , is constructed. It is defined in equation 5.1.

$$p_T^{\text{sum}} = p_T^{\ell\ell} + p_T^{\text{miss}} + \sum p_T^j \quad (5.1)$$

1797 The first cut after pre-selection in the cut-based analysis requires $p_T^{\text{sum}} < 15$ GeV to further suppress $t\bar{t}$
1798 production.

1799 In the different flavor channels, a cut is made to reduce the contamination from $Z \rightarrow \tau\tau$ decays.
1800 The di- τ invariant mass, $m_{\tau\tau}$, is constructed by assuming that the neutrinos from the τ decays were
1801 collinear with the leptons [82]. The analysis requires that this mass not be consistent with a Z by requiring
1802 $m_{\tau\tau} < m_Z - 25$ GeV.

1803 VBF TOPOLOGICAL CUTS

1804 The characteristic feature of VBF production of the Higgs is the presence of two additional forward
1805 jets coming from the incoming partons which radiate the vector bosons that make the Higgs. These jets
1806 are forward because the outgoing partons still carry the longitudinal momentum of the incoming partons.
1807 Figure 5.2 shows the distribution of the η for the leading jet in a VBF event compared to a background top
1808 pair production event. As can be seen, the VBF jets tend to be more forward in η , while the $t\bar{t}$ jets are more
1809 central.




Figure 5.2: Leading jet η in VBF $H \rightarrow WW^*$ (red) and $t\bar{t}$ (black)

1810 Because the cross section for VBF production is an order of magnitude smaller than gluon fusion pro-
1811 duction, these forward jets must be used in order to better reduce background and achieve a good signal to
1812 background ratio. The dedicated VBF search selection requirements are constructed to maximally exploit
1813 the features of the unique VBF topology.

1814 Requirements on the VBF jets are collectively referred to as the “VBF topological cuts”. First, a require-
1815 ment on the dijet invariant mass of the VBF jets, m_{jj} , is placed, requiring $m_{jj} > 600$ GeV. Next, the
1816 event is required to have a large gap in rapidity between the two VBF jets, or $\Delta y_{jj} > 3.6$. Both of these
1817 cuts put tight requirements on the presence of two forward, high p_T jets moving in opposite directions in

1818 the longitudinal plane.

1819 Beyond requiring the presence of the two forward VBF jets, the analysis also vetoes on the presence
1820 of any additional jets that fall between the two VBF jets. This cut is referred to as the central jet veto, or
1821 CJV. Any events with a third jet with $p_T > 20$ GeV whose rapidity is between the region defined by the
1822 two VBF jets are vetoed. This can be expressed in terms of a variable called the jet centrality, defined in
1823 equation 5.2.

$$C_{j3} = \left| \eta_{j3} - \frac{\eta_{j1} + \eta_{j2}}{2} \right| / \frac{|\eta_{j1} - \eta_{j2}|}{2}, \quad (5.2)$$

1824 Here, η_{j1} and η_{j2} are the pseudorapidities of the leading and subleading jets, respectively, while η_{j3} is
1825 the pseudorapidity of the extra jet in the event (if one exists). Intuitively, C_{j3} is zero when η_{j3} is directly
1826 centered between the two jets and unity when η_{j3} is aligned with either of the VBF jets. Thus, the CJV
1827 can be expressed as a requirement that $C_{j3} > 1$.

1828 The decay products of the Higgs tend to be central as well. Thus, the analysis also requires that both
1829 leptons in the analysis fall within the rapidity gap defined by the jets. This cut is referred to as the outside
1830 lepton veto, or OLV. A quantitative way to define the cut is to require that the centrality of each lepton
1831 (defined analogously to that of the third jet in equation 5.2) correspond to the lepton being within the jet
1832 rapidity gap, or $C_\ell < 1$ for both leptons.

1833 Figure 5.3a-c shows the m_{jj} , Δy_{jj} , and $C_{\ell 1}$ variables at the stage where all previous cuts in the sequence
1834 have been made. The agreement between data and Monte Carlo is good, and the bottom panels show their
1835 power in discriminating the VBF signal from the background processes.

1836 The final signal region is also split into two bins of m_{jj} , with the first bin corresponding to $600 \text{ GeV} <$
1837 $m_{jj} < 1 \text{ TeV}$ and the second bin corresponding to $m_{jj} > 1 \text{ TeV}$. The first bin has more statistics but
1838 also a larger contribution from background, while the second bin has lower statistics but a 1:1 signal to
1839 background ratio.

1840 HIGGS TOPOLOGICAL CUTS

1841 The final state leptons will exhibit unique correlations due to the fact that they are arising from the
1842 decay of a spin zero resonance. In particular, the spins of the final state leptons and neutrinos must all




Figure 5.3: Distributions of (a) m_{jj} , (b) Δy_{jj} , (c) $C_{\ell 1}$, and (d) $\Sigma m_{\ell j}$, for the VBF analysis. The top panels compare simulation and data, while the bottom panels show normalized distributions for all background processes and signal [63].




Figure 5.4: A cartoon of the WW final state. Momenta are represented with thin arrows, spins with thick arrows.
[63]

cancel, as shown in figure 5.4. Because the neutrino has a left handed chirality and the anti-neutrino has a right handed chirality (in the massless neutrino approximation), the spin and momentum of the particles will be anti-aligned and aligned, respectively. In the transverse plane, the momenta of all four final state objects must cancel as well. With the constraint of having both the momenta and the spin alignments cancel, the final state kinematics strongly prefer having a small angle between the leptons in the transverse plane (low $\Delta\phi_{\ell\ell}$). This angular correlation will also lead to low values of the di-lepton invariant mass $m_{\ell\ell}$. These unique signal final state kinematic correlations will be exploited to define the ultimate signal region.

The analysis places additional requirements on the final state leptons. Two requirements on dilepton kinematics are made that are common with lower multiplicity jet bins as well. The angle between leptons in the transverse plane, $\Delta\phi_{\ell\ell}$, is required to be less than 1.8 radians. Additionally, the dilepton mass $m_{\ell\ell}$ is required to be less than 50 GeV.

The cut-based analysis uses m_T as the final discriminating variable as in the ggF focused analysis. The optimal number of bins in m_T was found to be three bins, with the bin boundaries at 80 and 130 GeV.

Table 5.7 shows a summary of the data and estimated signal and background yields from simulation as each requirement described above is made. The table shows how the overall signal to background ratio grows through the various selection requirements. Table 5.8 shows the background composition after each selection requirement, illustrating which backgrounds are reduced most by certain requirements. Figure 5.5 shows an ATLAS event display of a candidate event in the final signal region.

Selection	Summary					
	$N_{\text{obs}}/N_{\text{bkg}}$	N_{obs}	N_{bkg}	N_{signal}		
				N_{ggF}	N_{VBF}	N_{VH}
$e\mu$ sample	1.00 ± 0.00	61434	61180	85	32	26
$n_b = 0$	1.02 ± 0.01	7818	7700	63	26	16
$p_T^{\text{sum}} < 15$	1.03 ± 0.01	5787	5630	46	23	13
$m_{\tau\tau} < m_Z - 25$	1.05 ± 0.02	3129	2970	40	20	9.9
$m_{jj} > 600$	1.31 ± 0.12	131	100	2.3	8.2	—
$\Delta y_{jj} > 3.6$	1.33 ± 0.13	107	80	2.1	7.9	—
$C_{j3} > 1$	1.36 ± 0.18	58	43	1.3	6.6	—
$C_{\ell 1} < 1, C_{\ell 2} < 1$	1.42 ± 0.20	51	36	1.2	6.4	—
$m_{\ell\ell}, \Delta\phi_{\ell\ell}, m_T$	2.53 ± 0.71	14	5.5	0.8	4.7	—
$ee/\mu\mu$ sample	0.99 ± 0.01	26949	27190	31	14	10.1
$n_b, p_T^{\text{sum}}, m_{\tau\tau}$	1.03 ± 0.03	1344	1310	13	8.0	4.0
$m_{jj}, \Delta y_{jj}, C_{j3}, C_\ell$	1.39 ± 0.28	26	19	0.4	2.9	0.0
$m_{\ell\ell}, \Delta\phi_{\ell\ell}, m_T$	1.63 ± 0.69	6	3.7	0.3	2.2	0.0

Table 5.7: Summary of event selection for the $n_j \geq 2$ VBF analysis in the 8 TeV cut-based analysis [63].

	Composition of N_{bkg}								
	N_{WW}		N_{top}		N_{misid}		N_{VV}	$N_{\text{Drell-Yan}}$	
	N_{WW}^{QCD}	N_{WW}^{EW}	$N_{t\bar{t}}$	N_t	N_{Wj}	N_{jj}	N_{VV}	$N_{ee/\mu\mu}^{\text{QCD}}$	$N_{\tau\tau}^{\text{EW}}$
$e\mu$ sample	1350	68	51810	2970	847	308	380	51	3260
$n_b = 0$	993	43	3000	367	313	193	273	35	2400
$p_T^{\text{sum}} < 15$	781	38	1910	270	216	107	201	27	2010
$m_{\tau\tau} < m_Z - 25$	484	22	1270	177	141	66	132	7.6	627
$m_{jj} > 600$	18	8.9	40	5.3	1.8	2.4	5.1	0.1	15
$\Delta y_{jj} > 3.6$	11.7	6.9	35	5.0	1.6	2.3	3.3	—	11.6
$C_{j3} > 1$	6.9	5.6	14	3.0	1.3	1.3	2.0	—	6.8
$C_{\ell 1} < 1, C_{\ell 2} < 1$	5.9	5.2	10.8	2.5	1.3	1.3	1.6	—	5.7
$m_{\ell\ell}, \Delta\phi_{\ell\ell}, m_T$	1.0	0.5	1.1	0.3	0.3	0.3	0.6	—	0.5
$ee/\mu\mu$ sample	594	37	23440	1320	230	8.6	137	690	679
$n_b, p_T^{\text{sum}}, m_{\tau\tau}$	229	12.0	633	86	26	0.9	45	187	76
$m_{jj}, \Delta y_{jj}, C_{j3}, C_\ell$	3.1	3.1	5.5	1.0	0.2	0.0	0.7	3.8	0.7
$m_{\ell\ell}, \Delta\phi_{\ell\ell}, m_T$	0.4	0.2	0.6	0.2	0.2	0.0	0.1	1.5	0.3

Table 5.8: Background composition after each requirement in the $n_j \geq 2$ VBF analysis in the 8 TeV cut-based analysis [63].




Figure 5.5: Event display of a VBF candidate event [63].

1861 5.4.3 BDT-BASED SELECTION

1862 The boosted decision tree based analysis takes a different philosophy compared to the cut-based. Rather
 1863 than cutting sequentially on many variables, the BDT analysis uses many of these variables as inputs to
 1864 the BDT and the output BDT score (O_{BDT}) as the final discriminant. The BDT is trained with the
 1865 VBF $H \rightarrow WW^*$ simulation as the signal samples and all other processes as background, including ggF
 1866 $H \rightarrow WW^*$ production. While the BDT based analysis is treated as a separate result, it has significant
 1867 overlap with the cut-based selection.

1868 PRE-TRAINING SELECTION AND BDT INPUTS

1869 Before training, the common pre-selection cuts described in section 5.4.1 are applied. Additionally, the
 1870 central jet veto and outside lepton veto described in section 5.4.2 are applied. The BDT has eight input
 1871 variables, six of which are also variables that are used in the cut-based analysis. The six shared variables

1872 are p_T^{sum} , m_{jj} , Δy_{jj} , $m_{\ell\ell}$, $\Delta\phi_{\ell\ell}$, and m_T . The seventh variable input in the BDT is a combination of
 1873 the variables used to do the OLV in the cut-based analysis. The BDT uses as input the sum of lepton
 1874 centralities, or $\sum C_\ell = C_{\ell 1} + C_{\ell 2}$. The final BDT input variable, $\Sigma m_{\ell j}$, is constructed to account for
 1875 the correlations between the jets and leptons in the event. It is the sum of the invariant masses of all four
 1876 possible lepton-jet combinations.

1877 Figure 5.3d shows the agreement between data and simulation for the $\Sigma m_{\ell j}$ variable, as well as showing
 1878 its discriminating power. Figure 5.6 shows the distributions of the Higgs topological variables that are
 1879 shared between the cut-based and BDT analyses. Figure 5.7 shows the distributions of the VBF topological
 1880 variables shared between the cut-based and BDT analyses. In both cases, the VBF yield has been scaled by
 1881 a factor of 50 to better show the shape difference compared to the backgrounds.




Figure 5.6: Distributions of $m_{\ell\ell}$ (top left), $\Delta\phi_{\ell\ell}$ (top right), and m_T (bottom), Higgs topology variables used in the selection requirements of the cut-based signal region and as inputs to the BDT result. These are plotted after all of the BDT pre-training selection cuts [63].




Figure 5.7: Distributions of m_{jj} (top left), Δy_{jj} (top right), $\sum C_\ell$ (bottom), VBF topology variables used in the selection requirements of the cut-based signal region and as inputs to the BDT result. These are plotted after all of the BDT pre-training selection cuts [63].

1882 5.5 BACKGROUND ESTIMATION

1883 This section describes the procedures used to estimate backgrounds for the VBF analysis in both the
 1884 cut-based and BDT analyses.

1885 5.5.1 GENERAL STRATEGY

1886 Most of the backgrounds in the VBF analysis have shapes estimated from Monte Carlo simulation but
 1887 normalizations derived from control regions in data. In essence, a normalization factor (denoted with β
 1888 or abbreviated as NF) is derived by scaling the MC yield in the control region to the corresponding yield
 1889 in data. Once this factor is derived, it can be used to scale the MC estimate of the background in the signal

1890 region. This is illustrated in equation 5.3.

$$B_{\text{SR}}^{\text{est}} = B_{\text{SR}} \times \frac{N_{\text{CR}}}{B_{\text{CR}}} \equiv B_{\text{SR}} \times \beta \quad (5.3)$$

1891 Here, B denotes the MC yield prediction in the denoted region, while N denotes the observed number of
1892 events in data in the denoted region.

1893 Another way of writing the same equation, in terms of an extrapolation factor α rather than a normal-
1894 ization factor β . The overall calculation is exactly the same. However, when phrased in this way, it shows
1895 how the uncertainty on the background estimation can be reduced. This is shown in equation 5.4.

$$B_{\text{SR}}^{\text{est}} = N_{\text{CR}} \times \frac{B_{\text{SR}}}{B_{\text{CR}}} \equiv N_{\text{CR}} \times \alpha \quad (5.4)$$

1896 Phrased this way, the equation shows that with enough statistics in the control region, a large theoretical
1897 uncertainty on the overall background yield in the signal region can be replaced by a small statistical un-
1898 certainty coming from the number of data events in the CR and a smaller theoretical uncertainty on the
1899 extrapolation from the control region to the signal region.

1900 5.5.2 TOP BACKGROUND

1901 The normalization factor β_t for the top background in the VBF analysis is derived in a region required
1902 to have one b-tagged jet, or $n_b = 1$. In the cut-based analysis, normalization factors are computed at every
1903 stage of the cutflow by applying the appropriate cuts in the CR. These NF are then applied to the $t\bar{t}$ and
1904 single top event yields in the SR. In the BDT analysis, a single normalization factor is computed for each
1905 bin of O_{BDT} after applying the BDT pre-training cuts described previously. The computed normaliza-
1906 tion factors are derived with all flavor combinations combined in order to decrease statistical uncertainty.
1907 Additionally, in the BDT analysis, BDT bins 2 and 3 are merged for the same reason.

1908 Table 5.9 shows the evolution of the β_t through the cut-based selection. Table 5.10 shows the value of
1909 the β_t in each bin of O_{BDT} . In all cases, the computed factors are relatively consistent with unity, with
1910 the largest discrepancy coming in bin 1 of O_{BDT} . The normalization factors in the bins of O_{BDT} are also

¹⁹¹¹ consistent with those derived in teh cut-based sisgnal region, increasing confidence in the BDT estimation.

Cut	β_t
$p_T^{\text{sum}} < 15 \text{ GeV}$	1.03 ± 0.01
$m_{\tau\tau} < m_Z - 25$	1.05 ± 0.01
$m_{jj} > 600 \text{ GeV}$	0.96 ± 0.06
$\Delta y_{jj} > 3.6$	1.02 ± 0.08
CJV	1.13 ± 0.16
OLV	1.01 ± 0.19
$m_{jj} < 1 \text{ TeV}$	0.94 ± 0.19
$m_{jj} > 1 \text{ TeV}$	1.48 ± 0.66

Table 5.9: Top normalization factors computed at each stage of the cut-based selection. Uncertainties are statistical only.

O_{BDT}	β_t
Bin0	1.09 ± 0.02
Bin1	1.58 ± 0.15
Bin2	0.95 ± 0.31
Bin3	0.95 ± 0.31

Table 5.10: Top normalization factors computed for each bin of O_{BDT} . Uncertainties are statistical only.

¹⁹¹² Figure 5.8 shows the m_{jj} and O_{BDT} distributions in the top control region. Overall the modeling looks
¹⁹¹³ consistent with the data.

¹⁹¹⁴ While these normalization factors can be computed and applied to the expected background yields listed
¹⁹¹⁵ in tables like table 5.8, in the end the normalization of the top background is profiled (meaning there is a
¹⁹¹⁶ dedicated Poisson constraint) and allowed to float in the final statistical fit.

¹⁹¹⁷ 5.5.3 $Z/\gamma^* \rightarrow \tau\tau$ BACKGROUND

¹⁹¹⁸ In the different flavor channels, the $Z/\gamma^* \rightarrow \tau\tau$ background is an important one. Di-tau production
¹⁹¹⁹ can produce an $e\mu$ final state if each τ lepton decays to a different flavor lepton.

¹⁹²⁰ In the BDT analysis, a single normalization factor for the background is derived. A control region
¹⁹²¹ is defined using the pre-training selection cuts, except requiring that $|m_{\tau\tau} - m_Z| < 25 \text{ GeV}$ so that
¹⁹²² the region is enriched in $Z/\gamma^* \rightarrow \tau\tau$ background. Additional requirements of $m_{\ell\ell} < 80(75) \text{ GeV}$




Figure 5.8: Distributions of m_{jj} (a) and O_{BDT} (b) in the VBF $n_b = 1$ top CR [63].

in the different (same) flavor channel, as well as $O_{\text{BDT}} > -0.48$ are applied to increase the purity of the region. The final $\beta_{Z/\gamma^* \rightarrow \tau\tau}$ is calculated to be 0.9 ± 0.3 (statistical uncertainty only). Because of the small contribution of this background in the BDT analysis and the large statistical uncertainty, no additional systematics are calculated. The final SR estimate is scaled by this β and not allowed to float in the fit.

The cut-based corrections are a bit more involved because they need to be applied selection by selection, as well as in the final signal region for the fit. The region is defined including all SR cuts up to the $Z/\gamma^* \rightarrow \tau\tau$ veto, which is instead made into a Z mass peak requirement as for the BDT region. The $m_{\ell\ell}$ cut from

1930 the BDT region is included as well. The cut-based approach aims to correct the normalization of the
 1931 $Z/\gamma^* \rightarrow \tau\tau$ background in two ways. First, an overall normalization factor is computed from the control
 1932 region. However, the VBF topological cuts are not included in this region, and applying them as is done in
 1933 the top CR is not feasible due to limited statistics. So, instead, correction factors (CF) to the cut efficiencies
 1934 of the VBF cuts are derived in a same flavor $Z \rightarrow \ell\ell$ control region, which has significantly more statistics.
 1935 The CF is simply the ratio of the cut efficiencies in data and MC derived in this region. In the end, the
 1936 overall background estimate is given by equation 5.5.

$$N_{Z/\gamma^* \rightarrow \tau\tau}^{\text{est}} = B_{Z/\gamma^* \rightarrow \tau\tau}^{\text{SR}} \times \beta_{\tau\tau} \times \frac{\epsilon_{\text{VBF cuts}}^{\text{data}}}{\epsilon_{\text{VBF cuts}}^{\text{MC}}} \quad (5.5)$$

1937 The hypothesis is that while the normalization correction must be derived in a dedicated region, the ef-
 1938 ficiency of the VBF cuts should not be sensitive to the type of Z/γ^* process and thus the larger control
 1939 region can be exploited to derive the CF. Figure 5.9 shows a shape comparison for the m_{jj} variable in
 1940 $Z \rightarrow \tau\tau$ events in the signal region and $Z \rightarrow \ell\ell$ events in the control region. The figure shows that the
 1941 shapes are indeed comparable and thus any CF derived in the same flavor control region can reliably be
 1942 applied in the signal region.




Figure 5.9: Comparison of m_{jj} shape in a same flavor $Z \rightarrow \ell\ell$ control region and the VBF cut-based signal region.

1943 Table 5.11 shows the overall normalization factor $\beta_{\tau\tau}$ and the efficiency correction factors for the various
 1944 VBF topological cuts. In general, the statistical uncertainties on the cut efficiency corrections are quite

¹⁹⁴⁵ good, and the MC tends to underestimate the efficiency of the VBF cuts for the $Z/\gamma^* \rightarrow \tau\tau$ background.
¹⁹⁴⁶ The overall normalization factor is also consistent with that calculated for the BDT analysis.

$\beta_{\tau\tau}$	0.97 ± 0.04
Cut	Correction factors
$m_{jj} > 600 \text{ GeV}$	1.09 ± 0.01
$\Delta y_{jj} > 3.6$	1.14 ± 0.02
CJV	1.20 ± 0.02
OLV	1.17 ± 0.03
$m_{jj} < 1 \text{ TeV}$	1.17 ± 0.06
$m_{jj} > 1 \text{ TeV}$	1.18 ± 0.13

Table 5.II: $Z/\gamma^* \rightarrow \tau\tau$ correction factors for the VBF cut-based analysis. Uncertainties are statistical only.

¹⁹⁴⁷ **5.5.4 $Z/\gamma^* \rightarrow \ell\ell$ BACKGROUND**

¹⁹⁴⁸ In the same flavor channels, the $Z/\gamma^* \rightarrow \ell\ell$ background is dominant and thus must be estimated cor-
¹⁹⁴⁹ rectly. In both the BDT and cut-based analyses, the background is estimated using the so-called “ABCD”
¹⁹⁵⁰ method. The ABCD method creates four different regions by defining cuts on two variables. One of the
¹⁹⁵¹ regions (A) is the signal region, while the other regions are defined by inverting one of both of the cuts.
¹⁹⁵² in this case, the two variables used are $m_{\ell\ell}$ and E_T^{miss} , because inverting either of the SR cuts on these
¹⁹⁵³ variables will give regions rich in the $Z/\gamma^* \rightarrow \ell\ell$ background. Figure 5.10 illustrates the general strategy
¹⁹⁵⁴ for each region.




Figure 5.10: General illustration of the ABCD region definitions for $Z/\gamma^* \rightarrow \ell\ell$ background estimation.

¹⁹⁵⁵ In both of the cut-based and BDT analyses, the Z peak region is defined with $|m_{\ell\ell} - m_Z| < 15 \text{ GeV}$.

1956 In the cut-based analysis, low $m_{\ell\ell}$ corresponds to $m_{\ell\ell} < 50$ GeV (this defines the cut-based SR) while
 1957 in the BDT it is $m_{\ell\ell} < 75$ GeV. In the cut-based, high and low E_T^{miss} are defined as opposite ends of
 1958 the 55 GeV cut applied for the signal region definition. The BDT low E_T^{miss} region is between 25 and
 1959 45 GeV, while the high E_T^{miss} region is $E_T^{\text{miss}} > 45$ GeV.

1960 Once the regions are defined, the final signal region background estimate is done by taking the estimate
 1961 in region B and extrapolating it to the signal region (A) by multiplying it by the ratio of regions C and
 1962 D. Effectively, the Z peak region is used to estimate the efficiency of the E_T^{miss} cut in data, and then this
 1963 efficiency is applied in the low $m_{\ell\ell}$ region. An additional correction is also applied for the non-closure of
 1964 the method in MC. This is summarized in equations 5.6 and 5.7.

$$N_{Z/\gamma^* \rightarrow \ell\ell}^{\text{SR}} = N_{Z/\gamma^* \rightarrow \ell\ell}^B \times \frac{N_{Z/\gamma^* \rightarrow \ell\ell}^C}{N_{Z/\gamma^* \rightarrow \ell\ell}^D} \times f_{\text{corr}} \quad (5.6)$$

$$f_{\text{corr}} = \frac{B_{\text{MC}}^A / B_{\text{MC}}^B}{B_{\text{MC}}^C / B_{\text{MC}}^D} \quad (5.7)$$

1965 Here, the N refer to data yields in each region with the non Z/γ^* backgrounds subtracted, while B
 1966 refer to the Z/γ^* yields in MC in each region.

1967 A normalization factor $\beta_{\ell\ell}$ is computed for each analysis as the ratio of the predicted data yield to the
 1968 MC yield in the SR. The shape of the BDT distribution is taken from data region B, while the shape of
 1969 the m_T distribution in the cut-based analysis is taken from Z/γ^* MC in the SR. The values of the $\beta_{\ell\ell}$ in
 1970 the cut-based and BDT analyses from this method are summarized in table 5.12. They are quite consistent
 1971 with one another within the statistical uncertainties. In the cut-based analysis, the same cut efficiency
 1972 correction factors shown in table 5.11 are also applied (in product with the $\beta_{\ell\ell}$) in the same flavor channels
 1973 to this background, as they were derived in the Z peak region.

	$\beta_{\ell\ell}$
BDT Bin 1	1.01 ± 0.15
BDT Bin 2	0.89 ± 0.28
Cut-based	0.81 ± 0.21

Table 5.12: $Z/\gamma^* \rightarrow \ell\ell$ normalization factors for cut-based and BDT analyses. Uncertainties are statistical only.

1974 5.5.5 WW AND OTHER DIBOSON BACKGROUNDS

1975 The WW and other diboson backgrounds have both their shape and normalization taken from MC
 1976 simulation. They are validated in dedicated control regions and found to agree with data well.

1977 As WW is the largest of these backgrounds and is irreducible, validating the estimate is of particular
 1978 importance. The validation region is constructed by requiring the pre-selection cuts on leptons and $m_{\ell\ell}$,
 1979 $n_b = 0$, and $m_T > 100$ GeV. The m_{T2} variable [83] is an additional discriminant that will isolate
 1980 the WW background, and a requirement of $m_{T2} > 160$ GeV is placed to define the WW validation
 1981 region. This cut gives a 60% purity for the validation region. The derived normalization factor in the
 1982 region is 1.15 ± 0.19 and is thus consistent with unity. Figure 5.11 shows the m_{T2} distribution and how it
 1983 distinguishes the WW background.




Figure 5.11: Distribution of m_{T2} in the WW validation region of the VBF analysis [63].

1984 5.5.6 HIGGS PRODUCTION VIA GLUON-GLUON FUSION

1985 Because this analysis is dedicated to measuring the VBF contribution to Higgs production, the compo-
 1986 nent of Higgs production from gluon-gluon fusion is treated as a background. The shape is taken directly
 1987 from simulation, using the generators described in table 5.4. In the final combined fit of all different signal
 1988 regions, the normalization is controlled by either a combined signal strength parameter μ , which controls
 1989 the normalization of both ggF and VBF production, or a separate parameter μ_{ggF} depending on the in-

1990 terpretation being presented in the final results.

1991 5.5.7 BACKGROUNDS WITH MISIDENTIFIED LEPTONS

1992 As discussed previously, the $W + \text{jets}$ and QCD multijet backgrounds are derived with fully data-driven
1993 methods. These backgrounds do not make a large contribution to the final VBF signal region but their
1994 estimation methods are discussed briefly here.

1995 $W + \text{jets}$ BACKGROUND

1996 The $W + \text{jets}$ background enters the signal region by having one of the jets mis-reconstructed as a lep-
1997 ton. The background is estimated by constructing a control sample with two leptons, where one lepton
1998 passes the usual lepton quality cuts but the second lepton fails one of those cuts (also known as the “anti-
1999 identified” lepton). This control region is rich in the $W + \text{jets}$ contribution because if a second lepton is
2000 reconstructed in a $W + \text{jets}$ event it is likely to be poor quality. The purity of this $W + \text{jets}$ control sample is
2001 85% to 90% depending on the exact configuration of leptons in the final state.

2002 The signal region estimate of $W + \text{jets}$ is estimated by extrapolation from the control sample to the sig-
2003 nal region using extrapolation factors derived in a $Z + \text{jets}$ control sample in data. The extrapolation factor
2004 is the ratio of the number of lepton candidates satisfying all quality criteria to the number of lepton can-
2005 didates anti-identified. This ratio is measured in bins of p_T and η . Thus, the final signal region estimate
2006 (binned as the extrapolation factor is binned) is simply the number of events in the anti-identified lepton
2007 control sample multiplied by the extrapolation factor derived from the $Z + \text{jets}$ control sample. Figure 5.12
2008 shows the extrapolation factors derived for electrons and muons.

2009 QCD MULTIJET BACKGROUND

2010 The method for estimating the multijet background is very similar to the $W + \text{jets}$ estimation method.
2011 The control sample in this case has two anti-identified leptons but otherwise satisfies all signal region re-
2012 quirements. The extrapolation factor is estimated from a multijet sample and applied twice to the control
2013 sample.




Figure 5.12: Extrapolation factors for the $W+jets$ estimate derived for muons (a) and electrons (b) as a function of lepton p_T [63].

2014 5.5.8 BACKGROUND COMPOSITION IN FINAL SIGNAL REGION

2015 After all of these estimation procedures, the final signal region composition can be calculated. The
 2016 estimated yields are all shown in table 5.8. Figure 5.13 shows the relative percentages of the different back-
 2017 ground for the different flavor and same flavor final states. In $e\mu$, the leading backgrounds are top back-
 2018 grounds, ggF Higgs, and SM WW production. In $ee/\mu\mu$, the leading background is Drell-Yan, followed
 2019 by top and ggF Higgs.




Figure 5.13: Background composition in final VBF signal region [63].

2020 5.6 SYSTEMATIC UNCERTAINTIES

2021 There are two main types of systematic uncertainties that are assessed for the analysis. First, theoretical
 2022 uncertainties associated with the various signal and background yield estimates are discussed. Then, exper-
 2023 imental uncertainties due to detector effects are shown. Normalization uncertainties refer to uncertainties
 2024 that affect the cross section of the process in question in the signal region being probed. Shape uncertain-
 2025 ties refer to systematic uncertainties that affect the shape of the final discriminating variable (either m_T or
 2026 O_{BDT}).

2027 5.6.1 THEORETICAL UNCERTAINTIES

2028 There are four main components to theoretical uncertainties assigned to signal and background pro-
 2029 cesses taken from Monte Carlo. Each one is a different source of variation in the overall acceptance for
 2030 that process. The first involves variation of the QCD renormalization and factorization scales used in the
 2031 calculation. In this case, the two scales are varied independently and simultaneously by factors of two high
 2032 or low and quantifying the resulting variation in normalization and shape for the process. This approx-
 2033 imates the correction to the cross section that would come from including the next order of the QCD
 2034 calculation (referred to as scale uncertainty). Next, there is an uncertainty associated with the PDF set
 2035 used in generating the events. The uncertainty eigenvectors for the given PDF set are studied, and the en-

2036 envelope of maximal variation is taken as an uncertainty. Finally, there are two uncertainties associated with
 2037 the choice of MC software (referred to as PDF uncertainty). An uncertainty associated with the generator
 2038 chosen for the hard scattering process is evaluated by keeping the parton showering software constant but
 2039 varying the matrix element generator and taking the maximal variation as an uncertainty (referred to as
 2040 the generator uncertainty). The converse variation can also be done, where the matrix element generator
 2041 remains constant and the generator used for the underlying event/parton shower modeling is varied (re-
 2042 ferred to as the UE/PS uncertainty). In cases where the background is normalized in a control region, the
 2043 systematic uncertainty arises from variations of the extrapolation factor α between the CR and the SR,
 2044 which can affect the normalization of the background in the SR.

2045 There are two additional uncertainties that are applied to the Higgs processes as well. First, there are
 2046 uncertainties assigned to the Higgs total production cross section. Then, there are uncertainties assigned
 2047 based on the fact that the analysis is done in exclusive jet bins and it is possible for signal events to migrate
 2048 from one bin to the next depending on the presence or absence of jets. These are assigned using the Jet
 2049 Veto Efficiency (JVE) procedure [18, 84] for ggF events and the Stewart-Tackmann (ST) method [85] for
 2050 VBF production.

2051 Table 5.13 shows the total theory uncertainties on the backgrounds in the cut-based analysis. These are
 2052 the sum in quadrature of the uncertainties from each of the variations described above.

Process	Theory syst. (%)
ggF H	48
Top	26
QCD WW	37
$Z/\gamma^* \rightarrow \tau\tau$	6.1

Table 5.13: Systematic uncertainties for various processes in the cut-based VBF analysis, given in units of % change
in yield. Values are given for the low m_{jj} signal region.

2053 Figures 5.14 and 5.15 show the variations in the extrapolation factor from the PDF and QCD uncertain-
 2054 ties on the top background estimate, binned in m_T , for the cut-based analysis. In both cases, there was
 2055 no significant shape uncertainty but normalization uncertainties were assigned according to the maximal
 2056 variation. These uncertainties enter into the 26% total uncertainty on top quoted in table 5.13

2057 While the estimate for the same-flavor $Z/\gamma^* \rightarrow \ell\ell$ background is data-driven, there is still a systematic
 2058 uncertainty taken for the non-closure of the method in Monte Carlo. This is taken as the maximum of the
 2059 deviation of the non-closure factor f_{corr} from unity and its uncertainty, or $\max(|1 - f_{\text{corr}}|, \delta f_{\text{corr}})$. For
 2060 the cut-based analysis this non-closure uncertainty 23%, while for the BDT analysis it is 17%.




Figure 5.14: Variations in the top background extrapolation factor in the cut-based analysis due to PDF uncertainties, binned in m_T .




Figure 5.15: Variations in the top background extrapolation factor in the cut-based analysis due to QCD scale uncertainties, binned in m_T .

2061 5.6.2 EXPERIMENTAL UNCERTAINTIES

2062 In this analysis, the theoretical uncertainties end up being the most dominant, but there are some ex-
2063 perimental uncertainties that make a contribution as well. The first is the uncertainty on the measured
2064 integrated luminosity, which affects backgrounds whose normalization is taken from MC and is measured
2065 to be 2.8% in the 8 TeV dataset [86]. The dominant sources of uncertainty overall are uncertainties on the
2066 jet energy scale and resolution and the b -tagging efficiency. Additional sources include lepton uncertain-
2067 ties on identification, resolution, and trigger efficiency, as well as uncertainties on the missing transverse
2068 momentum .

2069 The jet energy scale uncertainty is split into several independent components, including jet-flavor de-
2070 pendent calorimeter response uncertainties, uncertainties on modeling of pile-up interactions, uncertain-
2071 ties on extrapolation from the central to forward detector regions, and MC non-closure [87]. The uncer-
2072 tainty on energy scale for jets used in this analysis ranges from 1% to 7% depending on the jet p_T and η .
2073 The jet energy resolution varies from 5% to 20%, with uncertainties ranging from 2% to 40% (the largest
2074 uncertainties occurring at the selection threshold).

2075 The b -tagging efficiency is independently measured in data samples enriched in dileptonic decays of $t\bar{t}$
2076 events or in events where a muon is reconstructed in the vicinity of a jet [88, 89]. The efficiencies and
2077 their uncertainties are binned in p_T and decomposed into uncorrelated components using an eigenvector
2078 method [90]. Uncertainties on the efficiency range from 1% to 7.8%. The uncertainty on the rate of
2079 misidentification of c -jets as b -jets ranges from 6-14%, while the uncertainty on the rate of light jet mis-
2080 tagging ranges from 9-19% depending on p_T and η .

2081 The total experimental uncertainties on different signal and background components are summarized
2082 in table 5.14. They are compared to the level of other statistical and systematic uncertainties as well. Overall,
2083 the experimental uncertainties are sub-dominant compared to the statistical and theoretical uncertainties.

2084 5.7 RESULTS

2085 While the combined results of all the $H \rightarrow WW^*$ sub-analyses will be discussed in the next chapter,
2086 this section presents the results of the VBF specific analysis and interpretations. As table 5.7 shows, the

Sample	Total error	Stat. error	Expt. syst. err.	Theo. syst. err.
$n_j \geq 2$ VBF-enriched				
N_{sig}	13	—	6.8	12
N_{bkg}	9.2	4.7	6.4	4.5
N_{WW}	32	—	14	28
N_{top}	15	9.6	7.6	8.5
N_{misid}	22	—	12	19
N_{VV}	20	—	12	15
$N_{\tau\tau}$ (DY)	40	25	31	2.9
$N_{ee/\mu\mu}$ (DY)	19	11	15	—

Table 5.14: Composition of the post-fit uncertainties (in %) on the total signal (N_{sig}), total background (N_{bkg}), and individual background yields in the VBF analysis [63].

final cut-based signal region contains 20 events in data with $m_T < 150$ GeV, 14 coming from the $e\mu$ channel and 6 coming from the $ee + \mu\mu$ channel. The BDT analysis has many more candidates due to its looser selection, and the yields in each bin of O_{BDT} are shown in table 5.15.

(a) Before the BDT classification

Selection	Summary						Composition of N_{bkg}										
	$N_{\text{obs}}/N_{\text{bkg}}$	N_{obs}	N_{bkg}	N_{signal}			N_{WW}^{QCD}	N_{WW}^{EW}	N_{tt}	N_t	N_{Wj}	N_{jj}	N_{VV}	$N_{\text{Drell-Yan}}$	$N_{ee/\mu\mu}^{\text{QCD}}$	$N_{\tau\tau}^{\text{QCD}}$	$N_{\tau\tau}^{\text{EW}}$
				N_{ggF}	N_{VBF}	N_{VH}											
$e\mu$ sample	1.04 ± 0.04	718	689	13	15	2.0	90	11	327	42	29	23	31	2.2	130	2	
$ee/\mu\mu$ sample	1.18 ± 0.08	469	397	6.0	7.7	0.9	37	3	132	17	5.2	1.2	10.1	168	23	1	

(b) Bins in O_{BDT}

$e\mu$ sample																
Bin 0 (not used)	1.02 ± 0.04	661	650	8.8	3.0	1.9	83	9	313	40	26	21	28	2.2	126	1
Bin 1	0.99 ± 0.16	37	37	3.0	4.2	0.1	5.0	1.0	17	3.1	3.3	1.8	2.6	—	4.0	0.2
Bin 2	2.26 ± 0.63	14	6.2	1.2	4.2	—	1.5	0.5	1.8	0.3	0.4	0.3	0.8	—	0.3	0.3
Bin 3	5.41 ± 2.32	6	1.1	0.4	3.1	—	0.3	0.2	0.3	0.1	—	—	0.1	—	0.1	0.1
$ee/\mu\mu$ sample																
Bin 0 (not used)	1.91 ± 0.08	396	345	3.8	1.3	0.8	33	2	123	16	4.1	1.1	8.8	137	20.5	0.5
Bin 1	0.82 ± 0.14	53	45	1.5	2.2	0.1	3.0	0.5	10.4	1.8	0.8	0.2	0.9	26	1.7	0.1
Bin 2	1.77 ± 0.49	14	7.9	0.6	2.5	—	0.8	0.3	1.1	0.2	0.2	—	0.3	4.4	0.3	0.1
Bin 3	6.52 ± 2.87	6	0.9	0.2	1.7	—	0.1	0.2	0.2	—	—	—	—	0.7	—	—

Table 5.15: Event selection for the VBF BDT analysis. The event yields in (a) are shown after the pre-selection and the additional requirements applied before the BDT classification (see text). The event yields in (b) are given in bins in O_{BDT} after the classification [63].

Figure 5.16(a) shows the final distribution of data candidates compared to the expected m_T distribution for signal and background. The data are very consistent with a VBF Higgs hypothesis. Figure 5.16(b) shows

2092 where the data candidates fall in the two-dimensional binning of m_T and m_{jj} used in the fit for the cut-
 2093 based analysis.




Figure 5.16: Post-fit distributions in the cut-based VBF analysis. Panel (a) shows the one-dimensional m_T distribution, while (b) shows the data candidates split into the bins of m_T and m_{jj} used in the final fit [63].

2094 Figure 5.17 shows the distributions of O_{BDT} and m_T in the VBF BDT analysis. Again the data are quite
 2095 consistent with a VBF Higgs hypothesis.

2096 Because the cut-based result is used as a validation for the BDT analysis and the two signal regions are
 2097 not fully orthogonal, it is interesting to explore which events overlap between the two analyses. Of the




Figure 5.17: Postfit distributions in the BDT VBF analysis [63].

2098 twenty events in the cut-based signal region, only seven were not selected by the BDT analysis, while the
 2099 other thirteen also enter the BDT signal region. Figure 5.18 shows where the different analysis candidates
 2100 lie in the m_{jj} - m_T plane. This shows clearly that the advantage of the BDT analysis is that it can extract
 2101 signal candidates lower m_{jj} region due to its ability to recognize correlations with other variables.

2102 While the context of these results in the broader $H \rightarrow WW^*$ statistical analysis will be presented in the
 2103 next chapter, the statistical significance of the VBF Higgs result can be shown here. In the BDT analysis, the
 2104 expected signal significance was 2.7σ , while the observed significance was 3.1σ . In the cut-based analysis,
 2105 the expected significance was 2.1σ and the observed significance was 3.0σ . The compatibility between
 2106 these two results can be evaluated by computing the probability of observing a larger difference in Z_0 values
 2107 than the one measured. Using toy Monte Carlo with the ggF signal strength fixed to unity and considering
 2108 only statistical uncertainties, this probability is computed to be 79%, indicating good agreement between
 2109 the analyses. This result represents the first evidence of the vector boson fusion production of a Higgs
 2110 boson.




Figure 5.18: Overlap between cut-based and BDT VBF signal region candidates in the m_{jj} - m_T plane.

*The feeling is less like an ending than just another starting
point.*

Chuck Palahniuk

6

2111

2112

Combined Run I $H \rightarrow WW^* \rightarrow \ell\nu\ell\nu$

2113

results

2114

6.1 INTRODUCTION

2115

In the final statistical analysis of $H \rightarrow WW^* \rightarrow \ell\nu\ell\nu$, the dedicated gluon-gluon fusion and vector boson fusion sensitive signal regions are all combined into a single fit to determine the main parameters of interest, the Higgs signal strength μ and mass m_H . Therefore, while the specific requirements applied for the VBF sensitive analysis are discussed in chapter 5, the final measurement of these parameters can only be discussed in combination with the results of the ggF dedicated analysis. For example, because ggF Higgs production is considered a background in the VBF analysis, the ggF dedicated signal regions can actually constrain the normalization of this background in the VBF dedicated region.

2122

This chapter presents the combined interpretation of results in the $H \rightarrow WW^* \rightarrow \ell\nu\ell\nu$ analysis

2123 for gluon fusion and vector boson fusion Higgs production. First, the results of the dedicated gluon fu-
 2124 sion search are presented. Then, a comparison of the individual production mode signal strengths (μ_{ggF}
 2125 and μ_{VBF} and a measurement of the combined signal strength (μ) are shown. Subsequently, the mea-
 2126 sured values of the Higgs couplings to fermions and vector bosons is presented. Finally, the cross section
 2127 measurement for ggF and VBF production are shown.

2128 6.2 RESULTS OF DEDICATION GLUON FUSION $H \rightarrow WW^* \rightarrow \ell\nu\ell\nu$ ANALYSIS

2129 The details of the dedicated gluon fusion $H \rightarrow WW^* \rightarrow \ell\nu\ell\nu$ analysis are not discussed in this thesis
 2130 and instead left to more comprehensive sources [63]. However, a brief summary of the results is essential
 2131 for describing the measurements of Higgs properties and interpreting the results of the dedicated VBF
 2132 search in a broader context. Additionally, the final Run 1 results on gluon fusion production make use of
 2133 the dedicated variables for same flavor final states developed in section 3.5. The results in the same flavor
 2134 final states will be shown here as well.

2135 6.2.1 RESULTS IN SAME FLAVOR ($ee/\mu\mu$) FINAL STATES

2136 Final states of the $H \rightarrow WW^* \rightarrow \ell\nu\ell\nu$ channel where both leptons have the same flavor ($ee/\mu\mu$)
 2137 were not included in the discovery result due to increased pileup conditions in the $\sqrt{s} = 8$ TeV data.
 2138 Dedicated techniques for background reduction in the same flavor final states were developed, as described
 2139 in section 3.5. The results shown in this section are the first published results using the same flavor channels
 2140 in the $H \rightarrow WW^*$ analysis.

2141 Table 6.1 shows the background estimate, expected signal yield, and event count in data for the same
 2142 flavor channels in the $n_j \leq 1$ signal regions. The dedicated same flavor background techniques allow this
 2143 channel to preserve a signal to background ratio similar to that of the different flavor channels.

	N_{obs}	N_{bkg}	N_{ggF}	N_{VBF}
$n_j = 0$	1108	1040 ± 40	77 ± 15	2.4 ± 1.7
$n_j = 1$	467	427 ± 21	22 ± 6	3.6 ± 1.8

Table 6.1: Post-fit yields in ggF dedicated signal regions for the $ee/\mu\mu$ final states [63].

2144 Figure 6.1 shows the final m_T distribution in data for the $n_j \leq 1$ channels. The data is very consistent
 2145 with the Higgs hypothesis and it can be seen that the same flavor channels are indeed sensitive to gluon
 2146 fusion production of the Higgs.




Figure 6.1: Post-fit m_T distribution in the $n_j \leq 1$ regions for the same flavor ($ee/\mu\mu$) final states [63].

2147 6.2.2 COMBINED GLUON FUSION RESULTS

2148 Table 6.2 shows the individual signal regions that were input into the final statistical fit. The ggF dedi-
 2149 cated bins use m_T as their discriminating variable and are separated into bins of p_T of the subleading
 2150 lepton as well. The VBF dedicated bin uses the O_{BDT} distribution as its final discriminant.

2151 Table 6.3 shows the yields in the various signal regions in both data and expected signal and back-
 2152 grounds. The yields for signal and background are all scaled according to the final normalizations cal-

SR category i			Fit var.
n_j , flavor	$\otimes m_{\ell\ell}$	$\otimes p_T^{\ell 2}$	
$n_j = 0$	$e\mu$	$\otimes [10, 30, 55]$	$\otimes [10, 15, 20, \infty]$
	$ee/\mu\mu$	$\otimes [12, 55]$	$\otimes [10, \infty]$
			m_T
$n_j = 1$	$e\mu$	$\otimes [10, 30, 55]$	$\otimes [10, 15, 20, \infty]$
	$ee/\mu\mu$	$\otimes [12, 55]$	$\otimes [10, \infty]$
			m_T
$n_j \geq 2$ ggF	$e\mu$	$\otimes [10, 55]$	$\otimes [10, \infty]$
$n_j \geq 2$ VBF	$e\mu$	$\otimes [10, 50]$	$\otimes [10, \infty]$
	$ee/\mu\mu$	$\otimes [12, 50]$	$\otimes [10, \infty]$
			O_{BDT}
			O_{BDT}

Table 6.2: All signal regions definitions input into final statistical fit [63].

culated in the fit.

	N_{obs}	N_{bkg}	N_{ggF}	N_{VBF}
$n_j = 0$	3750	3430 ± 90	300 ± 50	8 ± 4
$n_j = 1$	1596	1470 ± 40	102 ± 26	17 ± 5
$n_j \geq 2$, ggF $e\mu$	1017	960 ± 40	37 ± 11	13 ± 1.4
$n_j \geq 2$, VBF	130	99 ± 9	7.7 ± 2.6	21 ± 3

Table 6.3: Post-fit yields in the both ggF and VBF dedicated signal regions with all lepton flavor final states combined [63].

Figure 6.2 shows the final post-fit m_T distribution in the $n_j \leq 1$ regions. The data are very consistent with the hypothesis of ggF Higgs production. These yields are used as input, along with the VBF results in chapter 5, for the physical interpretation of results presented in subsequent sections.

6.3 SIGNAL STRENGTH MEASUREMENTS IN GGF AND VBF PRODUCTION

When all of the signal regions are combined in the fit, there can be a combined measurement of the signal strength as well as the individual ggF and VBF signal strengths. The combined signal strength is the ratio of the sum of the gluon fusion and VBF cross sections to the theory prediction, or a signal strength




Figure 6.2: Post-fit m_T distribution in the $n_j \leq 1$ regions [63].

for the total Higgs production cross section that this analysis is sensitive to. The final measured combined signal strength μ is measured shown in equation 6.1.

$$\begin{aligned}
\mu &= 1.09 \quad {}^{+0.16}_{-0.15} \text{ (stat.)} \quad {}^{+0.08}_{-0.07} \left(\frac{\text{expt}}{\text{syst}} \right) \quad {}^{+0.15}_{-0.12} \left(\frac{\text{theo}}{\text{syst}} \right) \quad \pm 0.03 \left(\frac{\text{lumi}}{\text{syst}} \right) \\
&= 1.09 \quad {}^{+0.16}_{-0.15} \text{ (stat)} \quad {}^{+0.17}_{-0.14} \text{ (syst)} \\
&= 1.09 \quad {}^{+0.23}_{-0.21}.
\end{aligned} \tag{6.1}$$

Figure 6.3 gives the best fit signal strength $\hat{\mu}$ as a function of the hypothesized Higgs mass. The value at 125.36 GeV corresponds to the μ quoted in equation 6.1. This value of the Higgs mass is used because it is the most precise mass measurement from ATLAS, a result of the combined $\gamma\gamma$ and ZZ mass measure-

2166 ments [91].




Figure 6.3: Best fit signal strength $\hat{\mu}$ as a function of hypothesized m_H [63].

2167 As explained in chapter 3, a probability p_0 can be computed using the test statistic q_0 to quantify the
2168 probability that the background could fluctuate to produce an excess at least as large as the one observed
2169 in the data. The local p_0 value is shown in figure 6.4 as a function of m_H . The minimum p_0 value is at
2170 $m_H = 130$ GeV and corresponds to a significance of 6.1σ . The curve is relatively flat and the significance
2171 is the same at 125.36 GeV within the quoted precision. The expected significance for a signal with strength
2172 $\mu = 1.0$ is 5.8σ . This represents the first discovery level significance measurement in the $H \rightarrow WW^* \rightarrow$
2173 $\ell\nu\ell\nu$ analysis.

2174 All the results presented so far in this section have been for the combined gluon fusion and VBF pro-
2175 duction modes. However, each signal strength can be calculated separately in the likelihood as well. There
2176 are two ways to do this. First, the likelihood can be parameterized in terms of a single parameter, the ratio
2177 of the VBF and gluon fusion signal strengths. With this method, the statistical significance of the VBF
2178 Higgs result can be evaluated. Figure 6.5 shows the likelihood as a function of the ratio $\mu_{\text{VBF}}/\mu_{\text{ggF}}$.

2179 The best fit value of the ratio of signal strengths is shown in equation 6.2. Within the quoted uncer-




Figure 6.4: Local p_0 as a function of m_H [63].




Figure 6.5: Likelihood as a function of $\mu_{\text{VBF}} / \mu_{\text{ggF}}$ [63].

tainties, it is consistent with a ratio of unity.

$$\frac{\mu_{\text{VBF}}}{\mu_{\text{ggF}}} = 1.26^{+0.61} (\text{stat.})^{+0.50} (\text{syst.}) = 1.26^{+0.79}_{-0.53} \quad (6.2)$$

2181 The null hypothesis for VBF production corresponds to a ratio of $\mu_{\text{VBF}}/\mu_{\text{ggF}} = 0$. The likelihood in
 2182 figure 6.5 gives a significance of 3.2σ at $\mu_{\text{VBF}}/\mu_{\text{ggF}} = 0$, as quoted in chapter 5.

2183 In addition to the ratio of signal strengths, each signal strength can be varied independently in the like-
 2184 lihood as well. Figure 6.6 shows the two dimensional likelihood scan in the $\mu_{\text{ggF}}-\mu_{\text{VBF}}$ plane. The best fit
 2185 values of the two signal strengths are shown in equation 6.3. Both are consistent with unity within their
 2186 uncertainties.

$$\begin{aligned} \mu_{\text{ggF}} &= 1.02 \pm 0.19^{+0.22}_{-0.18} = 1.02^{+0.29}_{-0.26} \\ \mu_{\text{VBF}} &= 1.27^{+0.44}_{-0.40}^{+0.29}_{-0.21} = 1.27^{+0.53}_{-0.45} \end{aligned} \quad (6.3)$$

(stat.) (syst.)




Figure 6.6: Likelihood scan as a function of μ_{VBF} and μ_{ggF} [63].

2187 **6.4 MEASUREMENT OF HIGGS COUPLINGS TO VECTOR BOSONS AND FERMIONS**

2188 Similar to the parameterization of signal strength, the couplings of the Higgs to fermions and bosons
 2189 can also be parameterized. The parameter of interest in this case is κ , or the ratio of the measured coupling
 2190 to the standard model expectation. Both the fermion and boson couplings have these so-called scale factors,
 2191 κ_F for fermions and κ_V for bosons. Gluon fusion production is sensitive to the fermion couplings through
 2192 the top quark loops in its production, while VBF production is sensitive to the vector boson couplings in
 2193 its production. Both modes are sensitive to the vector boson couplings in their decays. The signal strengths
 2194 will have dependence on the coupling scale factors as described in equation 6.4 [18].

$$\begin{aligned}\mu_{\text{ggF}} &\propto \frac{\kappa_F^2 \cdot \kappa_V^2}{(\mathcal{B}_{H \rightarrow f\bar{f}} + \mathcal{B}_{H \rightarrow gg}) \kappa_F^2 + (\mathcal{B}_{H \rightarrow VV}) \kappa_V^2} \\ \mu_{\text{VBF}} &\propto \frac{\kappa_V^4}{(\mathcal{B}_{H \rightarrow f\bar{f}} + \mathcal{B}_{H \rightarrow gg}) \kappa_F^2 + (\mathcal{B}_{H \rightarrow VV}) \kappa_V^2}.\end{aligned}\quad (6.4)$$

2195 Figure 6.7 shows the two-dimensional likelihood scan of κ_F and κ_V . The best-fit values are given in equa-
 2196 tion 6.5. The best-fit values are consistent with unity within their uncertainties.

$$\begin{aligned}\kappa_F &= 0.93 & {}^{+0.24}_{-0.18} & {}^{+0.21}_{-0.14} & = 0.93 & {}^{+0.32}_{-0.23} \\ \kappa_V &= 1.04 & {}^{+0.07}_{-0.08} & {}^{+0.07}_{-0.08} & = 1.04 & \pm 0.11.\end{aligned}\quad (6.5)$$

(stat.) (syst.)

2197

2198 **6.5 HIGGS PRODUCTION CROSS SECTION MEASUREMENT**

2199 Another measurement that comes naturally from the signal strength numbers quoted earlier is the pro-
 2200 duction cross section at 7 and 8 TeV for both gluon fusion and VBF production. The general equation




Figure 6.7: Likelihood scan as a function of κ_F and κ_V [63].

for calculating the cross section is given in equation 6.6.

$$\begin{aligned}
 (\sigma \cdot \mathcal{B}_{H \rightarrow WW^*})_{\text{obs}} &= \frac{(N_{\text{sig}})_{\text{obs}}}{\mathcal{A} \cdot \mathcal{C} \cdot \mathcal{B}_{WW \rightarrow l\nu l\nu}} \cdot \frac{1}{\int L dt} \\
 &= \hat{\mu} \cdot (\sigma \cdot \mathcal{B}_{H \rightarrow WW^*})_{\text{exp}}
 \end{aligned} \tag{6.6}$$

$(N_{\text{sig}})_{\text{obs}}$ is the number of events observed in data. \mathcal{A} is the geometric and kinematic acceptance of the detector, while \mathcal{C} is the efficiency of the signal region selection for events that are reconstructed in the detector. The branching ratio of a WW system to leptons must also be divided out. The production cross section depends on the center of mass energy and the production mode desired (gluon fusion or VBF), and so three separate cross section measurements are quoted in equation 6.7.

$$\begin{aligned}
 \sigma_{\text{ggf}}^{7\text{TeV}} \cdot \mathcal{B}_{H \rightarrow WW^*} &= 2.0 \pm 1.7 \stackrel{+1.2}{-1.1} = 2.0 \stackrel{+2.1}{-2.0} \text{ pb} \\
 \sigma_{\text{ggf}}^{8\text{TeV}} \cdot \mathcal{B}_{H \rightarrow WW^*} &= 4.6 \pm 0.9 \stackrel{+0.8}{-0.7} = 4.6 \stackrel{+1.2}{-1.1} \text{ pb} \\
 \sigma_{\text{VBF}}^{8\text{TeV}} \cdot \mathcal{B}_{H \rightarrow WW^*} &= 0.51 \stackrel{+0.17}{-0.15} \stackrel{+0.13}{-0.08} = 0.51 \stackrel{+0.22}{-0.17} \text{ pb.}
 \end{aligned} \tag{6.7}$$

(stat.) (syst.)

2207 The predicted cross section values for gluon fusion are 3.3 ± 0.4 pb at 7 TeV and 4.2 ± 0.5 pb at 8 TeV,
2208 consistent with the measured values within their uncertainties. For vector boson fusion, the predicted
2209 cross section is 0.35 ± 0.02 pb, again consistent with the measured value.

2210 **6.6 CONCLUSION**

2211 The combined analysis of the gluon fusion and vector boson fusion processes in $H \rightarrow WW^* \rightarrow \ell\nu\ell\nu$
2212 in the 7 and 8 TeV datasets has yielded the first discovery level significance for Higgs production in this
2213 decay channel. Additionally, precise measurements of the couplings to vector bosons and fermions are
2214 given. Finally, signal strengths and cross sections for each production mode are measured. Figure 6.8 shows
2215 the $H \rightarrow WW^* \rightarrow \ell\nu\ell\nu$ measurements in comparison with other Higgs decay channels in ATLAS. The
2216 measurement of signal strength from this channel remains the most sensitive in both the gluon fusion and
2217 VBF production modes for the Run 1 dataset.

ATLAS

Individual analysis




Figure 6.8: Comparison of signal strength measurements in different Higgs decay channels on ATLAS [92].

Part III

2218

Search for Higgs pair production in the

2219

$HH \rightarrow b\bar{b}b\bar{b}$ channel in LHC Run 2 at $\sqrt{s} =$

2220

13 TeV

2221

Passion is in all great searches and is necessary to all creative endeavors.

W. Eugene Smith

7

2222

2223

Search for Higgs pair production in boosted $b\bar{b}b\bar{b}$ final states

2224

2225

7.1 INTRODUCTION

2226

After the discovery of the Higgs boson in the ATLAS Run 1 dataset and the subsequent measurements of its properties, the Higgs transformed into a potential tool in searches for physics beyond the Standard Model. The pair production cross section of the Higgs can be enhanced through BSM physics. Studying di-Higgs production also probes the Higgs self-coupling, shedding light on the structure of the Higgs potential. This chapter presents a search for resonant production of a Higgs pair in the $X \rightarrow HH \rightarrow b\bar{b}b\bar{b}$ final state in 3.2 fb^{-1} of data collected at $\sqrt{s} = 13 \text{ TeV}$. In particular, this chapter focuses on a search for this final state in the regime where m_X is large ($\gtrsim 1 \text{ TeV}$) and the Higgs bosons in the decay are significantly boosted. A tailored selection for this boosted selection, using novel techniques in jet substructure

and b -tagging, is discussed. Then, the data-driven background estimate is presented. Finally, the results of the search are shown. The signal models used as benchmarks are a spin-2 Randall Sundrum graviton (RSG) and a narrow width spin-0 resonance. These models are described in more detail in Chapter 1. Limits on signal models are reserved for the next chapter where the results of this chapter are combined with the results of a separate selection dedicated to the lower m_X regime.

7.2 MOTIVATION

With the center of mass energy increase from $\sqrt{s} = 8$ TeV to $\sqrt{s} = 13$ TeV, the LHC and ATLAS are able to probe new resonances at higher mass scales than previously accessible in Run 1. This is a powerful motivator for searching for a new resonance in the early 13 TeV data. Figure 7.1 shows the ratios of parton luminosities between 8 and 13 TeV for different resonance masses. For a resonance of $M_X = 2$ TeV, the cross section at $\sqrt{s} = 13$ TeV is roughly a factor of 10 larger than at $\sqrt{s} = 8$ TeV.




Figure 7.1: Parton luminosity ratios as a function of resonance mass M_X for 13/8 TeV and 7/8 TeV [93].

Higgs pair production offers a vast array of unprobed regions of phase space where searches for BSM physics can be made. Chapter 1 discusses some possibilities for both resonant and non-resonant enhance-

ment of the di-Higgs production cross section. Given the increased mass reach of the LHC in Run 2, it is particularly important to focus on resonant searches at high m_X . One consideration when conducting a search in the HH final state is which decay modes of the Higgs to consider. Figure 7.2 shows the branching ratio of the HH final state for different combinations of decays of each individual Higgs. As the largest branching ratio for the 125 GeV Higgs is $H \rightarrow b\bar{b}$, the $HH \rightarrow b\bar{b}b\bar{b}$ branching ratio is also the largest at 33%.




Figure 7.2: Summary of HH branching ratios [94].

At high m_X , the Higgs bosons resulting from the decay of a heavy resonance will have large p_T ¹. The ΔR between the decay products of the Higgs is inversely proportional to the Higgs p_T , as shown in equation 7.1.

$$\Delta R \approx \frac{2m}{p_T} \quad (7.1)$$

Figure 7.3 shows the minimum ΔR between truth level B decay vertices in simulation samples for Randall-Sundrum gravitons of different masses. The figure shows that as the mass of the graviton increases, the ΔR distribution between the b quarks in the Higgs decay tends to shift to lower values. Because of this effect, it is necessary to tailor a selection to target these merged b -jets.

¹In the limit that $m_H \ell \ell m_X$, the Higgs p_T is roughly $m_X/2$.




Figure 7.3: Minimum ΔR between B decay vertices for different RSG masses in a $G_{KK}^* \rightarrow HH \rightarrow 4b$ sample with $c = 1$

2260 7.3 DATA AND SIMULATION SAMPLES

2261 7.3.1 SIGNAL MODELS

2262 While the resonance search is by its nature generic (as it is a simple search for a peak in the $4b$ invariant
 2263 mass spectrum), there are two signal models that the selection requirements have been optimized for.
 2264 The first is Randall-Sundrum (RSG) model, where a tower of massive spin-2 Kaluza-Klein gravitons is
 2265 predicted. The second is a heavy narrow spin-0 resonance, the so-called “heavy Higgs”. This type of res-
 2266 onance arises, for example, in the two Higgs doublet model (2HDM). More details about the physics of
 2267 these models and their motivation is given in chapter 1.

2268 Signal graviton (G_{KK}^*) events are generated at leading order (LO) with **MADGRAPH5 v2.2.2** [95]. The
 2269 PDF set used is the **NNPDF2.3 LO** set [96]. For modeling parton shower and hadronization in jets, **PYTHIA**
 2270 8.186 is used with the A14 tune [75, 97]. The free parameters in the RSG model are the graviton mass
 2271 and the coupling constant $c \equiv k/\bar{M}_{Pl}^2$. Both the production cross section and width of the graviton are
 2272 proportional to c^2 . Samples are generated at both $c = 1$ and $c = 2$ for a variety of mass points between

² k is the curvature constant for the warped extra dimension and \bar{M}_{Pl} is the Planck mass divided by 8π

2273 300 GeV and 3 TeV.

2274 The second signal sample is a heavy spin-0 resonance H with a fixed width of $\Gamma_H = 1$ GeV. This
2275 is generated with **MADGRAPH5** and uses the **CT10** PDF set [78]. The parton shower and hadronization
2276 are handled by **HERWIG ++** with the **CTEQ6L1** PDF set and the **UEEE5** event tune [79, 98, 99]. Because
2277 the width and branching ratios depend on 2HDM parameters, each mass point generated with this fixed
2278 width corresponds to a different point in the 2HDM parameter phase space. Mass points are generated
2279 between 300 GeV and 1 TeV as with the RSG signal samples.

2280 **7.3.2 BACKGROUND SAMPLES**

2281 While the dominant **QCD** multijet background is estimated with a fully data-driven method, the sub-
2282 dominant backgrounds $t\bar{t}$ and Z +jets are modeled with some input from simulation.

2283 $t\bar{t}$ events are simulated at next-to-leading order (NLO) with the **POWHEG-BOX** version 1 generator us-
2284 ing the **CT10** PDF set [100]. The parton shower, hadronization, and underlying event are simulated with
2285 **PYTHIA 6.428** with the **CTEQ6L1** PDF set [74]. The Perugia 2012 tune is used [101]. NNLO **QCD** cor-
2286 rections to the cross sections are computed in **Top++ 2.0** [102]. The top quark mass is set to 172.5 GeV.
2287 The shapes of distributions in $t\bar{t}$ are taken from MC while the normalization is taken from data.

2288 Finally, the Z +jets background is simulated with **PYTHIA 8.186** and the **NNPDF2.3** LO PDF set. This
2289 background is negligible compared to the others and is taken fully from MC.

2290 **7.3.3 DATA SAMPLE AND TRIGGER**

2291 This analysis is done on 3.2 fb^{-1} of data taken in 2015 at $\sqrt{s} = 13$ TeV. The details of the machine
2292 conditions during this time can be found in Chapter 2. Only data which was taken during stable beam con-
2293 ditions with all detectors functioning is used. Events must pass a trigger which requires a single 360 GeV
2294 large radius ($R = 1.0$) jet to be reconstructed in the HLT. Figure 7.4 shows the trigger efficiency for vari-
2295 ous trigger options as a function of graviton mass. Above $m_G > 1$ TeV, the single large radius jet trigger
2296 is 99% efficient for events passing the signal selection.




Figure 7.4: Trigger efficiency for events passing all signal region selections as a function of mass in $G_{\text{KK}}^* \rightarrow HH \rightarrow 4b$ samples with $c = 1$ [103]. In the trigger names, “j” refers to a jet or jets. “ht” refers to H_T , the scalar sum of transverse momenta in the event. “bloose” refers to a loose b -tagging requirement applied to the jet. “a10r” refers to anti- k_T jets with $R = 1.0$. The numbers at the end are the thresholds on the given quantity in GeV.

2297 7.4 EVENT RECONSTRUCTION AND OBJECT SELECTION

2298 The boosted selection first begins by defining a unique set of objects that can be exploited to increase
 2299 signal efficiency in the kinematic regime where the final state b -jets are very merged.

2300 7.4.1 LARGE RADIUS ($R = 1.0$) JETS

2301 The first step towards reconstructing the final state is to define objects that can be used to measure the
 2302 kinematics of the Higgs bosons. In the boosted selection anti- k_T jets with a radius parameter of 1.0 are
 2303 used. These jets are much larger in angular size than the typical $R = 0.4$ jets and are intended to encompass
 2304 both jets resulting from the Higgs decay³. The jets are built from clusters in the calorimeter calibrated with

³This is in contrast to the resolved selection, which uses two $R = 0.4$ anti- k_T jets for each Higgs

local calibration weighting [58].

Because of the large extent of these jets, great care must be taken to remove potential contributions of calorimeter clusters from pile-up. This is done using a technique called jet trimming [104]. With trimming, the constituents of the large radius jet are re-clustered with a smaller radius with the k_T algorithm. Then, these so-called subjets are removed from the larger jet if $p_T^{\text{subjett}}/p_T^{\text{jet}} < f_{\text{cut}}$. In this analysis, the subjet radius is $R = 0.2$ and $f_{\text{cut}} = 0.05$. Trimming has been shown to improve the mass resolution of large radius jets. Figure 7.5 shows the effect of trimming on the large radius jet mass (M_J). Because the large radius jet fully contains the higgs decay products, its invariant mass should correspond to the 125 GeV mass of the Higgs. The trimming algorithm brings the jet mass much closer to the expected Higgs mass and improves the mass resolution.




Figure 7.5: Comparison of untrimmed and trimmed jet masses for large radius jets in a RSG sample with $m_{G_{KK}^*} = 1$ TeV. JES (JMS) refers to the standard jet energy (mass) scale calibration for ATLAS [58].

The large radius jets are required to satisfy $250 < p_T < 1500$ GeV. They must also be within $|\eta| < 2.0$ in order to ensure that the full jet is within the inner detector tracking volume. Finally, they are required to have $M_J > 50$ GeV. The upper p_T cut and lower threshold on mass are applied to correspond to the kinematic range where uncertainties are available in ATLAS calibrations [105, 106].

2319 7.4.2 TRACK JETS AND b -TAGGING

2320 Because the b -jets from boosted Higgs decays are so close together (as illustrated in figure 7.3), narrow ra-
2321 dius jets are required to fully resolve both b -jets. The minimum radius feasible for jets based on calorimeter
2322 deposits is determined by the calorimeter granularity. However, because b -tagging relies on information
2323 from the inner detector, it is possible to define another type of jet that can have a smaller radius and better
2324 b -tagging resolution. These jets are called “track jets” [106, 107].

2325 Track jets are formed by applying the usual anti- k_T clustering algorithm to tracks that are required to be
2326 consistent with the primary vertex. After the jet axis has been determined using these tracks, a second step
2327 of track association is also performed to add tracks that can be useful for b -tagging [107]. In this analysis,
2328 the tracks are clustered with a radius parameter of $R = 0.2$. This radius has been shown to give good
2329 performance in boosted Higgs tagging [106, 107]. Figure 7.6 shows a comparison among different track jet
2330 radii of the efficiency for reconstructing two b -jets from each Higgs in a RSG sample as a function of mass.
Track jets with radius of 0.2 give the best performance, especially at high mass. In this analysis, track jets




Figure 7.6: Efficiency of finding two b -jets from each Higgs in an RSG event using calorimeter jets with $R = 0.3$ or
2331 different track jet radii [107]

2332 are required to have $p_T > 10$ GeV and $|\eta| < 2.5$. They must also have at least two tracks.

2333 **7.4.3 MUONS**

2334 Muons are used in this study to correct the four-momenta of calorimeter jets by accounting for semi-
2335 leptonic b decays. The muons used are combined ID and MS muons which must satisfy tight identification
2336 requirements [54]. The muons must have $p_T > 4 \text{ GeV}$ and $|\eta| < 2.5$. Table 7.1 summarizes the object
2337 requirements described in this section.

	R	p_T	$ \eta $	M
Calorimeter jets	1.0	$250 < p_T < 1500 \text{ GeV}$	< 2.0	$> 50 \text{ GeV}$
Track jets	0.2	$> 10 \text{ GeV}$	< 2.5	-
Muons	-	4 GeV	< 2.5	-

Table 7.1: Summary of requirements on objects used in the $X \rightarrow HH \rightarrow b\bar{b}b\bar{b}$ search

2338 **7.5 EVENT SELECTION**

2339 The first requirement in the boosted selection is for ≥ 2 large radius jets satisfying the selections out-
2340 lined above. The two highest momentum large-R jets in the event are referred to as “Higgs candidates”.
2341 The leading jet is required to have $p_T > 350 \text{ GeV}$.

2342 Track jets satisfying the object selections are matched to Higgs candidate jets via ghost association [108].
2343 Each Higgs candidate must have at least 2 track jets associated with it. These basic requirements are illus-
2344 trated in figure 7.7.

2345 The QCD multijet background produces less central jets than high mass resonances, so there is an ad-
2346 dditional requirement that the two Higgs candidates be close together in η . The large-R jets are required to
2347 satisfy $|\Delta\eta(JJ)| < 1.7$.

2348 **7.5.1 MASS REQUIREMENTS**

2349 The final set of requirements ensures that the Higgs candidates are consistent with expected properties
2350 of the 125.0 GeV Higgs. First, a variable (X_{hh}) is defined to measure the consistency of both of the Higgs




Figure 7.7: Illustration of the boosted selection requirements on Higgs candidates. Each large-radius calorimeter jet (Higgs candidate) must contain two track jets

²³⁵¹ candidate jets with the SM Higgs mass. This is shown in equation 7.2.

$$X_{hh} = \sqrt{\left(\frac{M_J^{\text{lead}} - 124 \text{ GeV}}{0.1 M_J^{\text{lead}}}\right)^2 + \left(\frac{M_J^{\text{sublead}} - 115 \text{ GeV}}{0.1 M_J^{\text{sublead}}}\right)^2} \quad (7.2)$$

²³⁵² The mass values in the X_{hh} formula are optimized to maximize signal efficiency. The sub-leading jet
²³⁵³ typically has a lower mass due to semi-leptonic b decays and final state radiation. X_{hh} effectively acts as
²³⁵⁴ a χ^2 measurement of the consistency of the two Higgs candidate masses with the signal hypothesis. The
²³⁵⁵ denominators of each term ($0.1M$) give the uncertainty on the mass measurement for the large radius jets.
²³⁵⁶ Events are required to satisfy $X_{hh} < 1.6$.

²³⁵⁷ Before making the requirement on X_{hh} , the masses of the Higgs candidates are corrected for semi-

leptonic b decays using muons with the criteria outlined in the previous section. Any muons within a $\Delta R < 0.2$ of a b -tagged track jet (as described in the next section) have their four-momenta added to the four-momentum of the Higgs candidate. This correction does not affect the pre-selection requirements but does affect the X_{hh} requirement and the final invariant mass distribution used.

7.5.2 b -TAGGING REQUIREMENTS

The last requirement applied is on the number of b -tagged track jets. There are two signal regions defined. The first requires exactly four b -tagged track jets, two in each Higgs candidate (known as the $4b$ signal region). At high resonance masses, this requirement is inefficient, so an additional signal region requiring only three b -tagged track jets is also defined (known as the $3b$ signal region). While this has a larger background it is also more efficient for high resonance masses. For both signal regions, the MV2c20 algorithm, where the training sample for the algorithm has 20% charm events is used. More details for this algorithm can be found in Chapter 2.




Figure 7.8: Estimated significance as a function of signal mass for RSG $c = 1$ models in the $3b$ (a) and $4b$ (b) regions for different b -tagging efficiency working points

Once the algorithm is selected, an efficiency working point must also be chosen. This working point defines the efficiency with which true b -jets are tagged and also fixes the overall background rejection of the algorithm. Higher efficiency working points accept more true b -jets but also allow for more background. Five different working points (70%, 77%, 80%, 85%, 90%) are tested. With each working point, the full data driven background estimation method is run to quantify the amount of background that will be

present in the final signal region. The significance is quantified using the median discovery significance for signal and background with Poisson errors, given in equation 7.3 [109].

$$Z = \sqrt{2 \left((s + b) \ln \left(1 + \frac{s}{b} \right) - s \right)} \quad (7.3)$$

Here, s is the expected number of signal events and b is the expected number of background events. This formula is derived using Poisson statistics with errors on both the signal and background. It is used because it is valid in the regime where s and b are of the same order. Note that in the limit where s is much smaller than b , this equation reduces to the more well known s/\sqrt{b} . Figure 7.8 shows the estimated significance as a function of signal mass in RSG $c = 1$ models for the $3b$ and $4b$ signal regions. The 77% working point gives the best performance over a wide range of masses in the $4b$ signal region. As this is the region which contributes the most to the total discovery significance, the 77% efficiency working point is chosen for the analysis.

7.5.3 SELECTION EFFICIENCY




Figure 7.9: Acceptance \times efficiency as a function of mass for (a) RSG and (b) narrow heavy scalar signal models [110].

Figure 7.9 shows the product of acceptance and efficiency as a function of mass for both the RSG and narrow heavy scalar resonance signal models. After $m_X > 1$ TeV, the efficiency of the $4b$ requirement begins to decline. After $m_X > 2$ TeV, the efficiency of requiring two track jets in each Higgs candidate

begins to decline as well. Both of these behaviors illustrate the difficulty of resolving the merged decay products at high mass. Figure 7.10 shows a more detailed comparison of the signal efficiency in the $3b$ vs $4b$ signal regions for the RSG model. The efficiencies shown here are relative to all prior selection requirements. It can be seen there that at high masses the $3b$ signal region is more efficient for signal.




Figure 7.10: Efficiency of requiring 3 or 4 b -tagged track jets vs. RSG mass. The efficiency quoted is relative to the previous selection requirements (rather than an absolute efficiency).

To investigate the degradation of b -tagging efficiency at high p_T , the individual jet tagging efficiencies can be compared as a function of signal mass. This is shown in figure 7.11. The figure shows that the leading jet tagging efficiency in both calorimeter jets degrades heavily, while the sub-lead jet tagging efficiency remains relatively constant. More details on the cause of this degradation are shown in appendix A.

The final discriminating variable used in the boosted analysis is M_{2J} , the invariant mass of the two Higgs candidates. In order to improve the mass resolution, the four-momenta of each Higgs candidate are scaled by m_h/M_J . The effect of this correction is small in the boosted analysis but is done for consistency with the resolved selection. Table 7.2 shows the effect of the selection requirements on signal and background simulations as well as data.




Figure 7.11: MV2c20 b -tagging efficiency for each of the four track jets in the boosted $4b$ selection as a function of RSG mass for $c = 1$ models.

Selection	Data	$m_{G_{KK}^*} = 1\text{TeV}$	$m_{G_{KK}^*} = 2\text{TeV}$	$t\bar{t}$	$Z + \text{jets}$
$N(\text{fiducial large-R jets}) \geq 2$	2202396	23.3	0.48	32345.2	4255.7
leading large-R jet $p_T > 350\text{ GeV}$	1873741	22.9	0.48	26511.7	3649.9
Both large-R jet $m > 50\text{ GeV}$	1854625	21.2	0.47	24369.8	3575.8
Both large-R jet $p_T < 1500\text{ GeV}$	1853601	21.2	0.46	24346.5	3572.9
$ \Delta\eta(JJ) < 1.7$	1435273	20.8	0.44	20751.0	3265.8
≥ 2 track-jets per large-R jet	1224727	19.8	0.40	18234.5	2692.6
3 b -tags, $X_{hh} < 1.6$	316	3.4	0.067	46.7	2.0
4 b -tags, $X_{hh} < 1.6$	20	2.9	0.030	1.4	0.0

Table 7.2: Effect of boosted selection on data, RSG signal models, $t\bar{t}$, and $Z + \text{jets}$. The numbers from simulation are normalized with the MC generator cross section and do not take into account the data driven estimates described in section 7.6 [III].

7.6 DATA-DRIVEN BACKGROUND ESTIMATION

The largest background to this final state is QCD multijet production, constituting 80-90% of the total background. Because of the difficulties in modeling higher order QCD processes, this background is estimated with a fully data-driven method. The only other non-negligible background is $t\bar{t}$, constituting the other 10-20%. Due to the presence of $t\bar{t}$ in the sideband region where the QCD background will be

⁴The $Z + \text{jets}$ background is a sub-percent level contribution

2407 estimated, the normalization of the QCD and $t\bar{t}$ backgrounds are simultaneously estimated.

2408 **7.6.1 MASS REGION DEFINITIONS**

2409 The first step in the data-driven background estimate is to define a sideband mass region where the
2410 background normalization can be derived. Additionally, a control region is defined where the background
2411 estimate can be validated. The control (CR) and sideband (SB) regions are defined using a radial distance
2412 in the two-dimensional large-R jet mass plane, R_{hh} , which is defined in equation 7.4.

$$R_{hh} = \sqrt{(M_J^{\text{lead}} - 124 \text{ GeV})^2 + (M_J^{\text{sublead}} - 115 \text{ GeV})^2} \quad (7.4)$$

2413 Events in the sideband region are required to fail the signal region $X_{hh} < 1.6$ requirement and have
2414 $R_{hh} > 35.8 \text{ GeV}$. The control region consists of those events which are not in the signal or sideband
regions. Figure 7.12 shows the definition of the signal, control, and sideband mass regions. Table 7.3 sum-




Figure 7.12: M_J^{sublead} vs. M_J^{lead} in a 2 b -tag data sample. The signal region is defined by the inner black contour ($X_{hh} < 1.6$) and the sideband region is defined by the outer contour ($R_{hh} > 35.8 \text{ GeV}$). The region between the black contours is the control region. The mass region which is enriched in $t\bar{t}$ background is also shown for illustration. [110]

2415

2416 summarizes the mass region selections for the three different regions used in the analysis.

Region	Requirement	Notes
Signal Region (SR)	$X_{hh} < 1.6$	-
Control Region (CR)	$R_{hh} < 35.8 \text{ GeV}$ and $X_{hh} > 1.6$	Used for validation of background estimates
Sideband Region (SB)	$R_{hh} > 35.8 \text{ GeV}$	Used to derive background normalization

Table 7.3: Mass region definitions used for background estimation

²⁴¹⁷ **7.6.2 BACKGROUND ESTIMATION**

²⁴¹⁸ The method for estimating the background in this analysis is similar to the ABCD method presented in
²⁴¹⁹ Chapter 5. In this case, the two handles used to define different regions for the estimate are the number of
²⁴²⁰ b -tagged track jets and the mass regions. A region requiring exactly two b -tagged track jets in one large-R
²⁴²¹ jet (referred to as the 2-tag or $2b$ region) is defined for use in the background estimate. The number of
²⁴²² expected background events in the $3b$ and $4b$ signal regions is then given by equation 7.5.

$$N_{\text{bkg}}^{3(4)-\text{tag},\text{SR}} = \mu_{\text{Multijet}} N_{\text{Multijet}}^{2-\text{tag},\text{SR}} + \beta_{t\bar{t}} N_{t\bar{t}}^{3(4)-\text{tag},\text{SR}} + N_{Z+\text{jets}}^{3(4)-\text{tag},\text{SR}} \quad (7.5)$$

²⁴²³ In this equation, $N_{\text{bkg}}^{3(4)-\text{tag}}$ is the expected number of background events in the $3b$ or $4b$ signal regions.
²⁴²⁴ $N_{\text{Multijet}}^{2-\text{tag}}$ is the number of multijet events in the 2-tag region. $N_{t\bar{t}}^{3(4)-\text{tag}}$ is the number of $t\bar{t}$ events pre-
²⁴²⁵ dicted in the MC for the $3b$ or $4b$ signal region, and the variable is similarly defined for the $Z+\text{jets}$ back-
²⁴²⁶ ground. The $\beta_{t\bar{t}}$ parameter is a scale factor used to correct the normalization of the $t\bar{t}$ estimate in the signal
²⁴²⁷ region. μ_{Multijet} is an extrapolation factor that is derived in the sideband region and used to estimate the
²⁴²⁸ ratio of 2-tag events to 3(4)-tag events in the signal region. It is defined in equation 7.6.

$$\mu_{\text{Multijet}} = \frac{N_{\text{Multijet}}^{3(4)-\text{tag},\text{SB}}}{N_{\text{Multijet}}^{2-\text{tag},\text{SB}}} = \frac{N_{\text{data}}^{3(4)-\text{tag},\text{SB}} - \beta_{t\bar{t}} N_{t\bar{t}}^{3(4)-\text{tag},\text{SB}} - N_{Z+\text{jets}}^{3(4)-\text{tag},\text{SB}}}{N_{\text{data}}^{2-\text{tag},\text{SB}} - \beta_{t\bar{t}} N_{t\bar{t}}^{2-\text{tag},\text{SB}} - N_{Z+\text{jets}}^{2-\text{tag},\text{SB}}} \quad (7.6)$$

²⁴²⁹ The $t\bar{t}$ scale factor ($\beta_{t\bar{t}}$) and the QCD multijet extrapolation factor (μ_{Multijet}) are estimated together in
²⁴³⁰ a simultaneous fit in the sideband region. Then, the number of events in the 2-tag signal region is used,
²⁴³¹ along with the $t\bar{t}$ estimate in the $3b$ and $4b$ signal regions and μ_{Multijet} , to estimate the total number
²⁴³² of background events in the two final signal regions. The shape of the final discriminant M_{2J} is also

²⁴³³ taken from the 2-tag signal region where there are more statistics. This method is illustrated graphically in
²⁴³⁴ figure 7.13.




Figure 7.13: An illustration of the data-driven background estimation technique for the boosted analysis

²⁴³⁵ In the $3b$ region, the fit yields values of $\mu_{\text{Multijet}} = 0.160 \pm 0.03$ and $\beta_{t\bar{t}} = 1.02 \pm 0.09$. In the $4b$
²⁴³⁶ region, the fit gives $\mu_{\text{Multijet}} = 0.0091 \pm 0.0007$ and $\beta_{t\bar{t}} = 0.82 \pm 0.39$. The uncertainties quoted are
²⁴³⁷ statistical only. The larger uncertainties in the $4b$ values indicate the lower statistics available in that region.

²⁴³⁸ Figure 7.14 shows the distributions of data and background estimates in the $3b$ and $4b$ sideband regions
²⁴³⁹ after the background fit has been done. The normalizations are constrained from the fit to match that of
²⁴⁴⁰ the data, but good modeling of the shape of the mass of the leading large-R jet is seen as well. The shapes
²⁴⁴¹ of the kinematic distributions in the $4b$ region are taken from the $3b$ region due to the better MC statistics
²⁴⁴² in that region.

²⁴⁴³ 7.6.3 BACKGROUND SHAPE FIT

²⁴⁴⁴ As mentioned in the previous section, the background shape in the 3-tag and 4-tag signal regions is
²⁴⁴⁵ taken from the 2-tag signal mass region. Due to the limited statistics available, the background shapes are
²⁴⁴⁶ additionally smoothed after being extrapolated to the 3-tag and 4-tag signal regions. Only the data in the
²⁴⁴⁷ range $900 < M_{2J} < 2000$ GeV is included in the fit due to the limited statistics available above 2 TeV.
²⁴⁴⁸ Both the $t\bar{t}$ and QCD multijet background are independently fit with an exponential shape, $y = e^{ax+b}$.




Figure 7.14: Leading large-R jet mass in the 3b (a) and 4b (b) sideband regions. The multijet and $t\bar{t}$ backgrounds are estimated using the data-driven methods described above. Because their normalizations are derived in the sideband region, the total background normalization is constrained by default to match the normalization of the data [110].

2449 Other shapes are considered and used for the systematic uncertainties. Table 7.4 shows the fit values for
 2450 the parameters. Because both the 3b and 4b QCD shapes come from the 2-tag region, the slopes derived
 2451 are very similar.

	a	b
QCD (4b)	0.00545 ± 0.00021	5.44 ± 0.24
$t\bar{t}$ (4b)	0.00746 ± 0.00021	4.88 ± 0.36
QCD (3b)	0.00545 ± 0.00021	8.30 ± 0.24
$t\bar{t}$ (3b)	0.00746 ± 0.00021	8.58 ± 0.36

Table 7.4: Parameters derived for exponential fit to background M_{2J} shape in the 3b and 4b signal regions [111]

2452 7.6.4 VALIDATION OF BACKGROUND ESTIMATE

2453 The background estimate can be validated by using the method to estimate the number of events in the
 2454 control mass region rather than the signal mass region. Figure 7.15 shows the M_{2J} distribution in the 3b
 2455 and 4b control regions, comparing data and background estimates. In both cases, both the background
 2456 shape and normalization are consistent with the data, indicating good agreement. The ratio of data to the
 2457 background estimates is also fit to a line in the figure to test for any shape difference. The slope of the
 2458 line is within 1σ (from the fit uncertainties) of flat, further indicating that the data is consistent with the
 2459 background estimate in the control region.




Figure 7.15: Di-jet invariant mass (M_{2J}) in the 3b (a) and 4b (b) control regions. The multijet and $t\bar{t}$ backgrounds are estimated using the data-driven methods described above [110].

Table 7.5 shows the yields in data and background estimates in the 3-tag and 4-tag sideband and control regions. Again, here, it can be seen that the total number of predicted background events from the data driven method is consistent with the number of data events in the region.

Sample (3-tag)	Sideband Region	Control Region
Multijet	4328 ± 27	607 ± 10
$t\bar{t}$	683.5 ± 8.1	99.6 ± 3.1
Z+jets	31.8 ± 3.7	7.7 ± 1.8
Total	5043 ± 28	715 ± 11
Data	5043	724
Sample (4-tag)	Sideband Region	Control Region
Multijet	247.4 ± 1.5	34.7 ± 0.6
$t\bar{t}$	28.4 ± 1.5	5.1 ± 0.7
Z+jets	3.4 ± 1.2	0.6 ± 0.5
Total	279.2 ± 2.5	40.3 ± 1.0
Data	279	45

Table 7.5: The number of events in data and predicted background events in the boosted 3-tag and 4-tag sideband and control regions. The uncertainties shown are statistical only. [110]

2463 7.7 SYSTEMATIC UNCERTAINTIES

2464 The systematic uncertainties in this analysis can be divided into two broad categories. The first type
2465 is uncertainties associated with the modeling of the signal processes. The second type of uncertainty is
2466 associated with both the shape and normalization of the background prediction.

2467 7.7.1 SIGNAL MODELING UNCERTAINTIES

2468 The signal modeling uncertainty has three main components: theoretical uncertainty on the acceptance,
2469 experimental uncertainties on the large-R jets, and experimental uncertainties on the track jets related to
2470 b -tagging. In this analysis the experimental uncertainties are the most significant.

2471 The first uncertainty on signal modeling is the theoretical uncertainty on the acceptance. As explained
2472 in section 5.6.1, there are four components to this uncertainty. The first is related to missing higher order
2473 terms from the matrix element calculations which is estimated by varying the QCD renormalization and
2474 factorization scales. The second is uncertainty due to the PDF set used. The third is a generator uncer-
2475 tainty which is estimated by modifying the generator used to model the underlying event and hadroniza-
2476 tion. Finally, there is an uncertainty associated with the modeling of the initial state and final state radia-
2477 tion (ISR/FSR). The total theoretical uncertainty on the signal yield is 3%, and this is dominated by the
2478 ISR/FSR modeling.

2479 There are uncertainties on the large-R jets in both the jet energy scale (JES) and jet energy resolution
2480 (JER) as well as the jet mass scale (JMS) and jet mass resolution (JMR). These are evaluated using $\sqrt{s} =$
2481 8 TeV data from Run 1 of ATLAS and extrapolated to the Run 2 beam and detector conditions using
2482 MC^s. The details of these uncertainties can be found in reference [112].

2483 Uncertainties on the track jets are related to the b -tagging efficiency. The total uncertainty on the signal
2484 yield due to b -tagging is evaluated by propagating variations of the b -tagging efficiency through the boosted
2485 selection requirements. The uncertainties are calculated jet-by-jet and parameterized as a function of b -jet
2486 p_T and η [90]. For high p_T b -jets (with $p_T > 300$ GeV), the uncertainties are extrapolated using MC
2487 simulation from the lower p_T b -jets [113].

^sThe uncertainties are correspondingly larger due to the uncertainty of this extrapolation.

2488 Table 7.6 shows the systematic uncertainties on the signal normalization for models with $m_{G_{\text{KK}}^*} =$
2489 1.5 TeV and both $c = 1$ and $c = 2$ as well as a narrow width heavy scalar. The dominant uncertainty
2490 comes from b -tagging and this uncertainty is larger in the 4-tag region than the 3-tag region.

Source	Background	G_{KK}^*		H
		$c = 1$	$c = 2$	
Luminosity	-	5.0	5.0	5.0
3-tag				
JER	< 1	< 1	< 1	< 1
JES	2	< 1	< 1	< 1
JMR	1	12	12	11
JMS	5	14	13	17
b -tagging	1	23	22	23
Theoretical	-	3	3	3
Multijet Normalization	3	-	-	-
Statistical	2	1	1	1
Total	7	31	30	33
4-tag				
JER	< 1	< 1	< 1	< 1
JES	< 1	< 1	< 1	< 1
JMR	4	12	13	13
JMS	5	13	13	14
b -tagging	2	36	36	36
Theoretical	-	3	3	3
Multijet Normalization	14	-	-	-
Statistical	3	1	1	1
Total	15	42	42	43

Table 7.6: Summary of systematic uncertainties in the total background and signal event yields (expressed in %) in the boosted 3-tag and 4-tag signal regions. Systematic uncertainties on the signal normalization are shown for models with $m_{G_{\text{KK}}^*} = 1.5$ TeV and both $c = 1$ and $c = 2$ as well as a narrow width heavy scalar.

2491 **7.7.2 BACKGROUND UNCERTAINTIES**

2492 Uncertainties on the QCD multijet background normalization and shape are estimated using the con-
2493 trol mass region. As shown previously, the background predictions in the control region match with the

2494 data yields within the statistical uncertainty in both the 3-tag and 4-tag control regions. As an additional
 2495 protection, the statistical uncertainty on the background prediction in the control region is assigned as a
 2496 systematic uncertainty on the normalization of the QCD background.

2497 Additional robustness tests are done by varying the definition of the control mass region and the b -
 2498 tagging requirements used to define the 2-tag sample. In all cases, the effect of the variations is found to be
 2499 within the statistical uncertainties on the background normalization in the control region.

2500 Shape uncertainties on the background are evaluated using two techniques. First, as shown in fig-
 2501 ure 7.15, the ratio between the data and background prediction is fit with a linear function. The uncer-
 2502 tainties on the slope of this fit are assigned as shape uncertainties. An additional uncertainty is assigned by
 2503 using alternate power law fit functions for the smoothing of the background shape. Table 7.7 shows the
 2504 alternate shapes used. The largest difference between the nominal fit function and the alternates, taking
 2505 into account the 1σ uncertainty band on each fit as well, is taken as a shape uncertainty.

Functional Form
$f_1(x) = p_0(1 - x)^{p_1}x^{p_2}$
$f_2(x) = p_0(1 - x)^{p_1}e^{p_2 x^2}$
$f_3(x) = p_0(1 - x)^{p_1}x^{p_2}x$
$f_4(x) = p_0(1 - x)^{p_1}x^{p_2} \ln x$
$f_5(x) = p_0(1 - x)^{p_1}(1 + x)^{p_2}x$
$f_6(x) = p_0(1 - x)^{p_1}(1 + x)^{p_2} \ln x$
$f_7(x) = \frac{p_0}{x}(1 - x)^{p_1-p_2} \ln x$
$f_8(x) = \frac{p_0}{x^2}(1 - x)^{p_1-p_2} \ln x$

Table 7.7: Alternate fit functions used to model the M_{2J} distribution in the QCD multijet background. In the equations, $x = M_{2J}/\sqrt{s}$.

2506 The uncertainties on the $t\bar{t}$ background are obtained by propagating the various experimental variations
 2507 (JES, JER, JMS, JMR, b -tagging) through the analysis selection requirements. Table 7.6 summarizes the
 2508 background uncertainties in the 3-tag and 4-tag regions.

2509 7.8 RESULTS

2510 Table 7.8 shows the observed yields in the 3-tag and 4-tag signal regions for the boosted analysis com-
 2511 pared to the predicted number of background events. In the 3-tag region, 316 events are observed with
 2512 a predicted background of 285 ± 19 . In the 4-tag region, 20 events are observed with a predicted back-
 2513 ground of 14.6 ± 2.4 . Figure 7.16 shows the M_{2J} distribution in the 3-tag and 4-tag regions. There are
 2514 some small excesses in the data, in particular in the 3-tag region around $M_{2J} \approx 900$ GeV and in the region
 2515 of $1.6 < M_{2J} < 2.0$ TeV. The significance of these excesses will be evaluated in the next chapter in the
 2516 statistical combination with the resolved results.

Sample	Signal Region (3-tag)	Signal Region (4-tag)
Multijet	235 ± 14	13.5 ± 2.4
$t\bar{t}$	48 ± 22	1.2 ± 1.0
$Z + \text{jets}$	2.0 ± 2.2	-
Total	285 ± 19	14.6 ± 2.4
Data	316	20
G_{KK}^* (1000 GeV), $c = 1$	3.4 ± 0.9	2.9 ± 1.1

Table 7.8: Observed yields in the 3-tag and 4-tag signal regions for the boosted analysis compared to the predicted number of background events Errors correspond to the total uncertainties in the predicted event yields. The yields for a graviton with $m_{G_{\text{KK}}^*} = 1$ TeV and $c = 1$ are also shown. [110]




Figure 7.16: Di-jet invariant mass (M_{2J}) in the $3b$ (a) and $4b$ (b) signal regions. The multijet and $t\bar{t}$ backgrounds are estimated using the data-driven methods described above. In the $3b$ region, a graviton signal with $m_{G_{KK}^*} = 1.8$ TeV and $c = 1$ is overlaid, with the cross section multiplied by a factor of 50 so that the signal is visible. In the $4b$ region, signals with $m_{G_{KK}^*} = 1.0$ TeV and $m_{G_{KK}^*} = 1.5$ TeV are overlaid, both with $c = 1$ and the yields multiplied by factors of 2 and 5 respectively [110].

*There is no real ending. It's just the place where you stop
the story.*

Frank Herbert

8

2517

2518

Combined limits from boosted and resolved searches

2519

2520

8.1 INTRODUCTION

2521

In order to cover the full mass range of possible resonances decaying to di-Higgs final states, two distinct tailored selections were produced. The resolved selection is more sensitive in the mass range of $400 < m_X < 1100$ GeV while the boosted selection is more sensitive to masses in the range $1100 < m_X < 3000$ GeV. Chapter 7 presents the details of the boosted selection and results. In setting limits on spin-2 Randall-Sundrum graviton (RSG) and narrow width heavy scalar (H) models, the results of the boosted selection are combined with the results of the resolved selection to cover the full mass range.

2527

This chapter presents limits on signal models resulting from the $X \rightarrow HH \rightarrow b\bar{b}b\bar{b}$ search in both the resolved and boosted selections. It first presents a brief overview of the resolved results that go into

2529 the limit setting. Then, an overview of the statistical methods used for the search and limit setting is given.
 2530 Finally, limits on the RSG and heavy scalar models are presented.

2531 **8.2 RESOLVED RESULTS**

2532 The details of the resolved selection will not be presented here and can be found in reference [110]. In
 2533 basic terms, the selection searches for four $R = 0.4$ b-tagged calorimeter jets (where each pair of jets is
 2534 one Higgs candidate). This is distinct from the boosted methodology which searches for merged decay
 2535 products. The backgrounds to the resolved selection are the same as those presented in Chapter 7 for the
 2536 boosted analysis.

2537 Table 8.1 shows the results for data yields and expected background in the resolved signal region. Fig-
 2538 ure 8.1 shows the M_{2J} distribution in the resolved signal region. The total number of events is consistent
 2539 with the prediction and no significant excess is seen. One event in the boosted 4-tag signal is shared with
 2540 the resolved signal region and has a mass of 852 GeV.

Sample	Signal Region Yield
Multijet	43.3 ± 2.3
$t\bar{t}$	4.3 ± 3.0
$Z + \text{jets}$	-
Total	47.6 ± 3.8
Data	46
SM hh	0.25 ± 0.07
$G_{\text{KK}}^*(800 \text{ GeV}), c = 1$	5.7 ± 1.5

Table 8.1: Observed yields in the resolve selection 4-tag signal region compared to the predicted number of background events Errors correspond to the total uncertainties in the predicted event yields. The yields for a graviton with $m_{G_{\text{KK}}^*} = 800 \text{ GeV}$ and $c = 1$ are also shown. [110]

2541 **8.3 SEARCH TECHNIQUE AND RESULTS**

2542 The statistical technique used for the search in this analysis is the same as that used in the $H \rightarrow WW^*$
 2543 analysis presented in section 3.6.2. The test statistic q_0 is used to define the p -values which measure the




Figure 8.1: Di-jet invariant mass (M_{2J}) in the resolved signal region. A graviton signal with $m_{G_{KK}^*} = 800$ GeV and $c = 1$ is overlaid. [110].

compatibility of the data with the background-only hypothesis corresponding to a signal strength $\mu = 0$.

Local p_0 values are computed to quantify the probability that the background could produce a fluctuation greater than or equal to the one observed in the data. In the resolved analysis, no significant excesses are observed. The largest discrepancy with respect to the background only hypothesis occurs near a resonance mass of 900 GeV and is found to be less than 2σ in significance.

In the boosted selection, the largest local excess is a broad excess in the $3b$ signal region that begins near $M_{2J} \approx 1.7$ GeV. Assuming a G_{KK}^* with this mass and $c = 1.0$, the local significance of this excess is 2.0σ .

8.4 LIMIT SETTING

In the absence of any significant excess observed in the data, limits on different signal models can be set. This section describes the limit setting procedure and presents combined results of the resolved and boosted analyses.

8.4.1 LIMIT SETTING PROCEDURE

The procedure used for setting exclusion limits in this analysis is the CL_s method [114]. The first step in setting the limits is to define a test statistic which will be used. For limit setting, the test statistic is shown

2559 in equation 8.1.

$$\tilde{q}_\mu = \begin{cases} -2 \ln \frac{L(\mu, \hat{\theta}(\mu))}{L(0, \hat{\theta}(0))} & \hat{\mu} < 0 \\ -2 \ln \frac{L(\mu, \hat{\theta}(\mu))}{L(\hat{\mu}, \hat{\theta})} & 0 \leq \hat{\mu} < \mu \\ 0 & \hat{\mu} > \mu \end{cases} \quad (8.1)$$

2560 In the above equation, μ is the value of the signal strength under test, $\hat{\mu}$ is the best fit μ , $\hat{\theta}$ is the
2561 best fit value of the nuisance parameters, $\hat{\theta}$ is the best fit value of the nuisance parameters under the fixed
2562 μ value, and L is the Poisson likelihood of the data (as described in section 3.6.2).

2563 The test statistic \tilde{q}_μ is constructed to protect against two interesting corner cases when setting the upper
2564 limit on the cross section. First, it protects against negative signal strengths μ which are unphysical. Second,
2565 it does not count excesses in the data larger than those expected by a signal strength μ as evidence against
2566 the μ hypothesis.

2567 The CL_s statistic is constructed by taking a ratio of two probabilities. CL_{s+b} is the probability that the
2568 signal+background hypothesis would produce a value of the test statistic that is less than or equal to the
2569 observed value¹. CL_b is the probability that the background only hypothesis will produce a value
2570 of the test statistics less than or equal to the observed. The CL_s statistic is then the ratio CL_{s+b}/CL_b . A
2571 95% upper limit on the cross section is set at the value of μ that makes the CL_s statistic less than 5%.

2572 In practice, the limits are computed numerically within an asymptotic approximation for the distribu-
2573 tion of the test statistic \tilde{q}_μ . The details of this approximation can be found in reference [65].

2574 The resolved and boosted analyses are combined using a very simple procedure rather than a full statis-
2575 tical combination. For each mass point tested, the limit which gives the most stringent constraint is used.
2576 This means that for mass points below 1.1 TeV the resolved signal region is used, while at and above this
2577 point the combination of the orthogonal 3b and 4b boosted signal regions is used.

2578 8.4.2 LIMIT SETTING RESULTS

2579 Figure 8.2 shows the combined 95% upper bounds as a function of mass for three different models:
2580 G_{KK}^* with $c = 1$, G_{KK}^* with $c = 2$, and a narrow heavy scalar H .

¹Lower values of \tilde{q}_μ mean better compatibility

2581 The cross section of $\sigma(pp \rightarrow G_{\text{KK}}^* \rightarrow hh \rightarrow b\bar{b}b\bar{b})$ with $c = 1$ is constrained to be less than 70 fb
2582 for masses in the range $600 < m_{G_{\text{KK}}^*} < 3000$ GeV. For the RSG model with $c = 2$, cross sections limits
2583 between 40 fb and 200 fb are set for the mass range of $500 < m_{G_{\text{KK}}^*} < 3000$ GeV. Masses in the range
2584 of $475 < m_{G_{\text{KK}}^*} < 785$ GeV are excluded with $c = 1$ (with an exclusion of the range 465 to 745 GeV
2585 expected). Masses less than 980 GeV are excluded with $c = 2$ (with an exclusion for masses less than
2586 1 TeV expected).

2587 In the heavy Higgs model, the cross section upper limits for $\sigma(pp \rightarrow H \rightarrow hh \rightarrow b\bar{b}b\bar{b})$ ranges from
2588 30 to 300 fb in the mass range of $500 < m_H < 3000$ GeV. The resolved analysis can also set an upper
2589 limit on the Standard Model di-Higgs production cross section discussed in chapter 3. The upper limit on
2590 $\sigma(pp \rightarrow hh \rightarrow b\bar{b}b\bar{b})$ in the Standard Model is constrained to be less than 1.22 pb.




Figure 8.2: Expected and observed upper limit as a function of mass for G^*_{KK} in the RSG model with (a) $c = 1$ and (b) $c = 2$, as well as (c) H with fixed $\Gamma_H = 1$ GeV, at the 95% confidence level in the CL_s method. [110]

2591

Part IV

2592

Looking ahead

9

2593

2594

Conclusion

2595 After being sought for many years at different collider experiments, the Higgs boson was discovered by
2596 the ATLAS and CMS experiments in 2012, confirming the leading theory for the source of electroweak
2597 symmetry breaking and filling in the last missing piece of the Standard Model. After its discovery, mea-
2598 surements of the particle's detailed properties and searches for new particles decaying to Higgs final states
2599 were both extremely important in constraining physics beyond the Standard Model. This dissertation
2600 presented this evolution through two results: the observation and measurement of the Higgs boson in the
2601 $H \rightarrow WW^* \rightarrow \ell\nu\ell\nu$ channel at $\sqrt{s} = 7$ TeV and $\sqrt{s} = 8$ TeV and a search for Higgs pair production
2602 in the $HH \rightarrow b\bar{b}b\bar{b}$ channel at $\sqrt{s} = 13$ TeV with the ATLAS detector in pp collisions at the Large
2603 Hadron Collider.

2604 In the $H \rightarrow WW^* \rightarrow \ell\nu\ell\nu$, results from both the discovery of the Higgs boson and the full ATLAS
2605 Run 1 dataset were presented. The Higgs boson was discovered with a 6.1σ significance in a combination
2606 of the $H \rightarrow \gamma\gamma$, $H \rightarrow ZZ4\ell$, $H \rightarrow WW^* \rightarrow \ell\nu\ell\nu$ with 4.2 fb^{-1} at $\sqrt{s} = 7$ TeV and 5.2 fb^{-1} at

$\sqrt{s} = 8$ TeV. With the full 20.3 fb^{-1} at $\sqrt{s} = 8$ TeV and 4.2 fb^{-1} at $\sqrt{s} = 7$ TeV, ATLAS achieved discovery level significance in the $H \rightarrow WW^*$ channel alone and obtained the first evidence of vector boson fusion production in that channel. The combined signal strength is measured to be $\mu = 1.09^{+0.23}_{-0.21}$. The total observed significance of the $H \rightarrow WW^*$ process is observed to be 6.1σ (with 5.8σ expected). Advanced methods for background reduction and estimation, particularly in same-flavor lepton final states, are shown. The VBF signal strength is measured to be $\mu_{\text{VBF}} = 1.27^{+0.53}_{-0.45}$ with an observed significance of 3.2σ (with 2.7σ expected).

These results required many novel innovations. The increase of pileup interactions in the higher instantaneous luminosity LHC conditions of 2012 led to a degradation of missing transverse momentum resolution. As a result, the prominent $Z/\gamma^* + \text{jets}$ background of the same flavor $H \rightarrow WW^* \rightarrow \ell\nu\ell\nu$ final states increased greatly. New variables, including a track-based missing transverse momentum and a measurement of the balance between the dilepton system and recoiling jets, allowed for significant reduction of this background. In the VBF channel, selections were optimized to exploit the unique VBF final state topology. Incorporating these variables into a boosted decision tree technique allowed the analysis to exceed the 3σ statistical significance threshold.

After the end of Run 1, the results of Higgs measurements from ATLAS were combined with those from CMS to produce the most precise measurements of the Higgs boson so far [115]. Figure 9.1 shows the combination of ATLAS and CMS data for the Higgs signal strength in and coupling measurements. In the signal strength measurements of gluon fusion and vector boson fusion, the $H \rightarrow WW^*$ channel provides the tightest constraints. Additionally, the Higgs coupling to W bosons is the most precisely measured with a relative uncertainty of 10%.

With the discovery of the Higgs firmly established and its properties measured, a natural next step was to search for new physics with Higgs final states. At $\sqrt{s} = 13$ TeV, a search for Higgs pair production in the $b\bar{b}b\bar{b}$ final state with 3.2 fb^{-1} was conducted. A signal region optimized for the boosted final states arising from high mass resonances was constructed. This signal region utilized large-radius calorimeter jets and b -tagging with small radius track jets to maximize the signal acceptance. No significant excesses were observed, and upper limits on cross sections are placed for spin-2 Randall Sundrum gravitons (RSG) and




Figure 9.1: Combined ATLAS and CMS measurements in Run 1 for (a) Higgs signal strength in gluon fusion and VBF and (b) Higgs couplings normalized to their SM predictions

narrow spin-0 resonances. The increase in center of mass energy in Run 2 allowed this analysis to extend upper limits up to 3 TeV, while previous results from ATLAS in Run 1 only quotes limits up to 2 TeV. The cross section of $\sigma(pp \rightarrow G_{KK}^* \rightarrow hh \rightarrow b\bar{b}b\bar{b})$ with $k/\bar{M}_{\text{Pl}} = 1$ is constrained to be less than 70 fb for masses in the range $600 < m_{G_{KK}^*} < 3000$ GeV. For the RSG model with $k/\bar{M}_{\text{Pl}} = 2$, cross sections limits between 40 fb and 200 fb are set for the mass range of $500 < m_{G_{KK}^*} < 3000$ GeV. The cross section upper limits for $\sigma(pp \rightarrow H \rightarrow hh \rightarrow b\bar{b}b\bar{b})$ ranges from 30 to 300 fb in the mass range of $500 < m_H < 3000$ GeV.

While there has been a rigorous program of measurements and searches involving the Higgs, there is still much room for improvement at the High Luminosity LHC (HL-LHC) and beyond. The measured signal strength for VBF production in $H \rightarrow WW^*$ still has a relative error at the level of 40%, largely dominated by statistical uncertainty. Projections for the HL-LHC show that the uncertainty on the VBF signal strength can be reduced to approximately 15% with 3000 fb^{-1} [116, 117]. This uncertainty also

2646 assumes that theoretical uncertainties on the signal, which would be the largest contribution in this dataset,
 2647 remain as they are now. Improvements in the theoretical understanding of the Higgs signal would also
 2648 reduce the signal strength uncertainty dramatically. Such precision measurements allow for measurements
 2649 of the Higgs coupling to vector bosons precise to the few percent level, therefore giving much power to
 2650 constrain or discover new physics.

2651 The prospects for detection of beyond the Standard Model resonant di-Higgs production at the HL-
 2652 LHC are also quite promising. Figure 9.2 shows projections for the discovery significance of RSG signals at
 2653 the HL-LHC in the $X \rightarrow HH \rightarrow b\bar{b}b\bar{b}$ search [117]. In all detector budget scenarios, a 1.5 TeV resonance
 is above or near 5σ significance, while a 2 TeV resonance is between $4-5\sigma$ except for the lowest budget.




Figure 9.2: Discovery significance for RSG models at the HL-LHC in three different budget scenarios [117]. Systematic uncertainties on the background prediction (σ_B) of 2.5% and 5.0% are both tested.

2654
 2655 The Higgs will continue to be an incredibly powerful tool in the understanding of nature at the HL-
 2656 LHC and beyond. Through both precision measurements and searches, the nature of electroweak symme-
 2657 try breaking will be better understood and the potential for the discovery of physics beyond the Standard
 2658 Model has never been greater.

A

2659

2660

b-tagging performance at high p_T

2661 One of the limiting factors of the signal acceptance in the $X \rightarrow HH \rightarrow b\bar{b}b\bar{b}$ search at high resonance
2662 masses is the degradation of the *b*-tagging efficiency for high p_T jets. This appendix presents a study of the
2663 underlying causes of this degradation.

2664 A.I CHANGES IN MV₂ SCORE AT HIGH p_T

2665 The degradation of *b*-tagging at high p_T was studied in particular in the context of RSG models at high
2666 mass. Figure A.I shows the p_T of the leading track jet inside of the leading calorimeter jet in RSG events.
2667 At high $m_{G_{KK}^*}$, the p_T spectrum of track jets is much harder than at lower masses due to the increased
2668 Higgs p_T .

2669 Figure A.2 shows the MV_{2c2o} algorithm score for the leading and subleading track jets inside of the
2670 leading calorimeter jet. In both cases, it can be seen that at higher RSG masses the MV₂ score shifts towards
2671 more background like (negative) values. Additionally, this effect is more pronounced in the leading track




Figure A.1: p_T of the leading track jet in the leading calorimeter jet for different signal masses in RSG $c = 1$ models

2672 jet than the subleading.




Figure A.2: MV2c20 score for the leading track jet (a) and subleading track jet (b) of the leading calorimeter jet for different signal masses in RSG $c = 1$ models

2673 To understand what is causing this change in the MV2c20 score, the same comparisons can be made for
 2674 the input variables of MV2c20. The focus in these comparisons will be on the leading track jet as this is the
 2675 one seen to have the largest difference in MV2 score. Figure A.3 shows the log likelihood ratio $\log(p_b/p_u)$

from the IP₃D (three dimensional impact parameter) algorithm. At higher masses, the IP₃D likelihood ratio distribution does become more background-like.




Figure A.3: IP₃D log-likelihood ratio ($\log(p_b/p_u)$) of the leading track jet in the leading calorimeter jet for different signal masses in RSG $c = 1$ models

Figure A.4 shows the mass and number of tracks at the secondary vertex computed by the SV1 algorithm. When there is no secondary vertex found, the algorithm assigns a default negative value for these quantities. Both of these distributions show that there is a significantly larger fraction of jets where no secondary vertex is found in the high mass samples compared to the $m_{G_{KK}^*} = 1$ TeV sample. The SV1 algorithm's inability to find a secondary vertex could be an important factor in the overall MV₂ score shift, as this eliminates eight of the input variables that would normally contribute information to the algorithm.

Figure A.5 shows the same quantities for the JetFitter algorithm. In this case, there is also a change in the fraction of jets which have their secondary vertices successfully reconstructed, but this change is not as drastic as that seen in SV1. There is also an increase in the number of jets which have high values of mass.

A.2 EFFECT OF MULTIPLE b -QUARKS INSIDE ONE JET

One hypothesis for why the efficiency of b -tagging the leading track jet degrades is that at high masses, the b quarks get close enough together that both of them are inside of the leading track jet. Because MV₂




Figure A.4: Mass (a) and number of tracks (b) for the secondary vertices computed with the SV1 algorithm. When no secondary vertex is found, the quantities are assigned to default negative values.




Figure A.5: Mass (a) and number of tracks (b) for vertices computed with the JetFitter algorithm. When no vertices are found, the quantities are assigned to default negative values.

2690 is not tuned for tagging multiple b quarks inside one jet, the tagging efficiency could degrade. Figure A.6
 2691 shows MV₂ scores and SV₁ mass for cases where there are two b quarks at truth level within the radius of

the leading track jet compared to cases where there is only one true b^l . This figure suggests that the presence




Figure A.6: MV₂c₂₀ score (a) and SV₁ mass (b) for leading track jets with two truth b quarks ($n_{tb,\text{lead}} = 2$) compared to those with only one truth b ($n_{tb,\text{lead}} = 1$).

of two b -quarks inside the leading jet is not the cause of the degradation in efficiency. There is a change in the shape of the MV₂ score distribution, but it is not nearly as pronounced as that seen in A.2 at higher masses. Additionally, the fraction of jets with no secondary vertex found is nearly identical in the track jets with two truth b -quarks.

A.3 CHANGES IN TRACK QUALITY AT HIGH p_T

Another hypothesis for the degradation of the b -tagging efficiency is a decrease in track quality for high p_T b jets. One way to check the overall quality of the tracking inside the jet is to investigate quantities related to the leading track inside of the track jet. Figure A.7 shows the fit χ^2/n_{DOF} and number of hits in the pixel detector for the leading track of the leading track jet. In both cases, the figure shows that in higher mass samples, the quality of the leading track inside of the track jet degrades substantially. The fit quality is lessened and the tracks have less hits in the pixel detector. This is likely due to the fact that at higher p_T ,

¹When two truth b quarks are required in the leading jet, the subleading jet is required to have zero. When one is required for the leading, one is also required for the subleading.




Figure A.7: Track fit χ^2/nDOF (a) and number of pixel detector hits (b) for the leading track of the leading track jet in different mass RSG $c = 1$ samples

the B -hadron will sometimes live long enough to miss the IBL and first pixel layer, thus decreasing the number of hits on the track.

To check whether this is the cause for the shift in the MV_2 score and the higher difficulty in reconstructing secondary vertices, jets whose leading track have at least four pixel hits are compared with those whose tracks have less than four pixel hits. The results for the MV_2 score and SV_1 mass are shown in figure A.8. Track jets where the leading track does not have at least four pixel hits are more likely to not have a secondary vertex reconstructed. Additionally, their MV_{2c2o} score is shifted more significantly to background-like values. This seems to confirm the hypothesis that degrading track quality is responsible for the lowered b -tagging efficiency at high p_T .




Figure A.8: MV₂c₂₀ score (a) and SV₁ mass (b) for leading track jets whose leading track jet has at least four pixel hits ($N_{\text{pix}} \geq 4$) compared to those which do not ($N_{\text{pix}} < 4$).

References

2713

- 2714 [1] Georges Aad et al. Observation of a new particle in the search for the Standard Model Higgs boson
2715 with the ATLAS detector at the LHC. *Phys. Lett.*, B716:1–29, 2012. doi: 10.1016/j.physletb.2012.
2716 08.020.
- 2717 [2] Serguei Chatrchyan et al. Observation of a new boson at a mass of 125 GeV with the CMS experi-
2718 ment at the LHC. *Phys. Lett.*, B716:30–61, 2012. doi: 10.1016/j.physletb.2012.08.021.
- 2719 [3] David Griffiths. *Introduction to elementary particles*. 2008.
- 2720 [4] F. Halzen and Alan D. Martin. *QUARKS AND LEPTONS: AN INTRODUCTORY COURSE
2721 IN MODERN PARTICLE PHYSICS*. 1984. ISBN 0471887412, 9780471887416.
- 2722 [5] Christopher G. Tully. *Elementary particle physics in a nutshell*. 2011.
- 2723 [6] K. A. Olive et al. Review of Particle Physics. *Chin. Phys.*, C38:090001, 2014. doi: 10.1088/1674-1137/
2724 38/9/090001.
- 2725 [7] Matthew D. Schwartz. *Quantum Field Theory and the Standard Model*. Cambridge University
2726 Press, 2014. ISBN 1107034736, 9781107034730. URL <http://www.cambridge.org/us/academic/subjects/physics/theoretical-physics-and-mathematical-physics/quantum-field-theory-and-standard-model>.
- 2729 [8] S. Dawson. Introduction to electroweak symmetry breaking. In *High energy physics and cosmology.
2730 Proceedings, Summer School, Trieste, Italy, June 29-July 17, 1998*, pages 1–83, 1998. URL <http://alice.cern.ch/format/showfull?sysnb=0301862>.
- 2732 [9] S. L. Glashow. Partial Symmetries of Weak Interactions. *Nucl. Phys.*, 22:579–588, 1961. doi: 10.
2733 1016/0029-5582(61)90469-2.
- 2734 [10] Steven Weinberg. A Model of Leptons. *Phys. Rev. Lett.*, 19:1264–1266, 1967. doi: 10.1103/
2735 PhysRevLett.19.1264.
- 2736 [11] A. Salam. *Elementary Particle Theory*. Almqvist and Wiksell, Stockholm, 1968.
- 2737 [12] J. Iliopoulos S.L. Glashow and L. Maiani. *D2:1285*, 1970.
- 2738 [13] R. Keith Ellis, W. James Stirling, and B. R. Webber. *QCD and collider physics*. *Camb. Monogr.
2739 Part. Phys. Nucl. Phys. Cosmol.*, 8:1–435, 1996.

- 2740 [14] P. W. Higgs. Broken symmetries and the masses of gauge bosons. *13*:508, 1964.
- 2741 [15] P. W. Higgs. Spontaneous symmetry breakdown without massless bosons. *145*:1156, 1966.
- 2742 [16] F. Englert and R. Brout. Broken symmetry and the mass of gauge vector mesons. *13*:321, 1964.
- 2743 [17] G. S. Guralnik, C. R. Hagen, and T. W. .B. Kibble. Global conservation laws and massless particles. *Phys. Rev. Lett.*, *13*:585, 1964. doi: [10.1103/PhysRevLett.13.585](https://doi.org/10.1103/PhysRevLett.13.585).
- 2745 [18] LHC Higgs Cross Section Working Group, S. Heinemeyer, C. Mariotti, G. Passarino, and
2746 R. Tanaka (Eds.). Handbook of LHC Higgs Cross Sections: 3, Higgs Properties. 2013.
- 2747 [19] Abdelhak Djouadi. The Anatomy of electro-weak symmetry breaking. I: The Higgs boson in the
2748 standard model. *Phys. Rept.*, *457*:1–216, 2008. doi: [10.1016/j.physrep.2007.10.004](https://doi.org/10.1016/j.physrep.2007.10.004).
- 2749 [20] J. Baglio, A. Djouadi, R. Gröber, M. M. Mühlleitner, J. Quevillon, and M. Spira. The mea-
2750 surement of the Higgs self-coupling at the LHC: theoretical status. *JHEP*, *04*:151, 2013. doi:
2751 [10.1007/JHEP04\(2013\)151](https://doi.org/10.1007/JHEP04(2013)151).
- 2752 [21] Matthew J. Dolan, Christoph Englert, and Michael Spannowsky. New Physics in LHC Higgs boson
2753 pair production. *Phys. Rev.*, *D87*(5):055002, 2013. doi: [10.1103/PhysRevD.87.055002](https://doi.org/10.1103/PhysRevD.87.055002).
- 2754 [22] Roberto Contino, Margherita Ghezzi, Mauro Moretti, Giuliano Panico, Fulvio Piccinini, and An-
2755 drea Wulzer. Anomalous Couplings in Double Higgs Production. *JHEP*, *08*:154, 2012. doi:
2756 [10.1007/JHEP08\(2012\)154](https://doi.org/10.1007/JHEP08(2012)154).
- 2757 [23] R. Grober and M. Mühlleitner. Composite Higgs Boson Pair Production at the LHC. *JHEP*, *06*:
2758 020, 2011. doi: [10.1007/JHEP06\(2011\)020](https://doi.org/10.1007/JHEP06(2011)020).
- 2759 [24] Lisa Randall and Raman Sundrum. A Large mass hierarchy from a small extra dimension. *Phys.*
2760 *Rev. Lett.*, *83*:3370–3373, 1999. doi: [10.1103/PhysRevLett.83.3370](https://doi.org/10.1103/PhysRevLett.83.3370).
- 2761 [25] Kaustubh Agashe, Hooman Davoudiasl, Gilad Perez, and Amarjit Soni. Warped Gravitons at the
2762 LHC and Beyond. *Phys. Rev.*, *D76*:036006, 2007. doi: [10.1103/PhysRevD.76.036006](https://doi.org/10.1103/PhysRevD.76.036006).
- 2763 [26] A. Liam Fitzpatrick, Jared Kaplan, Lisa Randall, and Lian-Tao Wang. Searching for the Kaluza-
2764 Klein Graviton in Bulk RS Models. *JHEP*, *09*:013, 2007. doi: [10.1088/1126-6708/2007/09/013](https://doi.org/10.1088/1126-6708/2007/09/013).
- 2765 [27] Julien Baglio, Otto Eberhardt, Ulrich Nierste, and Martin Wiebusch. Benchmarks for Higgs Pair
2766 Production and Heavy Higgs boson Searches in the Two-Higgs-Doublet Model of Type II. *Phys.*
2767 *Rev.*, *D90*(1):015008, 2014. doi: [10.1103/PhysRevD.90.015008](https://doi.org/10.1103/PhysRevD.90.015008).
- 2768 [28] G. C. Branco, P. M. Ferreira, L. Lavoura, M. N. Rebelo, Marc Sher, and Joao P. Silva. Theory and
2769 phenomenology of two-Higgs-doublet models. *Phys. Rept.*, *516*:1–102, 2012. doi: [10.1016/j.physrep.2012.02.002](https://doi.org/10.1016/j.physrep.2012.02.002).

- 2771 [29] Howard E. Haber and Oscar Stål. New LHC benchmarks for the \mathcal{CP} -conserving two-Higgs-
2772 doublet model. *Eur. Phys. J.*, C75(10):491, 2015. doi: 10.1140/epjc/s10052-015-3697-x.
- 2773 [30] Jose M. No and Michael Ramsey-Musolf. Probing the Higgs Portal at the LHC Through Resonant
2774 di-Higgs Production. *Phys. Rev.*, D89(9):095031, 2014. doi: 10.1103/PhysRevD.89.095031.
- 2775 [31] Johan Alwall, Michel Herquet, Fabio Maltoni, Olivier Mattelaer, and Tim Stelzer. MadGraph
2776 5:Going Beyond. *JHEP*, 1106:128, 2011. doi: 10.1007/JHEP06(2011)128.
- 2777 [32] Oleg Antipin, Tuomas Hapola. CP3 Origins implementation of Randall-Sundrum model. 2013.
2778 URL <http://cp3-origins.dk/research/units/ed-tools>.
- 2779 [33] Lyndon R Evans and Philip Bryant. LHC Machine. *J. Instrum.*, 3:S08001. 164 p, 2008. URL
2780 <https://cds.cern.ch/record/1129806>. This report is an abridged version of the LHC De-
2781 sign Report (CERN-2004-003).
- 2782 [34] ATLAS Collaboration. The ATLAS experiment at the CERN Large Hadron Collider. *JINST*, 3:
2783 S08003, 2008. doi: 10.1088/1748-0221/3/08/S08003.
- 2784 [35] CMS Collaboration. The cms experiment at the cern lhc. *Journal of Instrumentation*, 3(08):S08004,
2785 2008. URL <http://stacks.iop.org/1748-0221/3/i=08/a=S08004>.
- 2786 [36] LHCb Collaoration. The LHCb Detector at the LHC. *JINST*, 3:S08005, 2008. doi: 10.1088/
2787 1748-0221/3/08/S08005.
- 2788 [37] ALICE Collaboration. The alice experiment at the cern lhc. *Journal of Instrumentation*, 3(08):
2789 S08002, 2008. URL <http://stacks.iop.org/1748-0221/3/i=08/a=S08002>.
- 2790 [38] Lyndon Evans. The Large Hadron Collider. In Holstein, BR and Haxton, WC and Jawah-
2791 ery, A, editor, *ANNUAL REVIEW OF NUCLEAR AND PARTICLE SCIENCE, VOL*
2792 *61*, volume 61 of *Annual Review of Nuclear and Particle Science*, pages 435–466. 2011. doi:
2793 {10.1146/annurev-nucl-102010-130438}.
- 2794 [39] ATLAS Collaboration. Luminosity Determination in pp Collisions at $\sqrt{s} = 7$ TeV Using the
2795 ATLAS Detector at the LHC. *Eur. Phys. J.*, C71:1630, 2011. doi: 10.1140/epjc/s10052-011-1630-5.
- 2796 [40] Mike Lamont for the LHC team. The First Years of LHC Operation for Luminosity Production.
2797 International Particle Accelerator Conference, 2013. URL <https://accelconf.web.cern.ch/>
2798 *accelconf/IPAC2013/talks/moyab101_talk.pdf*.
- 2799 [41] Paul Collier for the LHC team. LHC Machine Status. CERN Resource Review Board, 2015. URL
2800 <https://cds.cern.ch/record/2063924/files/CERN-RRB-2015-119.PDF>.

- 2801 [42] Track Reconstruction Performance of the ATLAS Inner Detector at $\sqrt{s} = 13$ TeV. Technical Report ATL-PHYS-PUB-2015-018, CERN, Geneva, Jul 2015. URL <http://cds.cern.ch/record/2037683>.
- 2802
2803
- 2804 [43] M Capeans, G Darbo, K Einsweiller, M Elsing, T Flick, M Garcia-Sciveres, C Gemme, H Pernegger, O Rohne, and R Vuillermet. ATLAS Insertable B-Layer Technical Design Report. Technical Report CERN-LHCC-2010-013. ATLAS-TDR-19, CERN, Geneva, Sep 2010. URL <https://cds.cern.ch/record/1291633>.
- 2805
2806
2807
- 2808 [44] ATLAS Collaboration. ATLAS Trigger Operations Public Results. 2015. URL <https://twiki.cern.ch/twiki/bin/view/AtlasPublic/TriggerOperationPublicResults>.
- 2809
- 2810 [45] ATLAS Collaboration. ATLAS Luminosity Public Results, Run 1. 2012. URL <https://twiki.cern.ch/twiki/bin/view/AtlasPublic/LuminosityPublicResults>.
- 2811
- 2812 [46] ATLAS Collaboration. ATLAS Luminosity Public Results, Run 2. 2015. URL <https://twiki.cern.ch/twiki/bin/view/AtlasPublic/LuminosityPublicResultsRun2>.
- 2813
- 2814 [47] T Kawamoto, S Vlachos, L Pontecorvo, J Dubbert, G Mikenberg, P Iengo, C Dallapiccola, C Amelung, L Levinson, R Richter, and D Lellouch. New Small Wheel Technical Design Report. Technical Report CERN-LHCC-2013-006. ATLAS-TDR-020, CERN, Geneva, Jun 2013. URL <https://cds.cern.ch/record/1552862>. ATLAS New Small Wheel Technical Design Report.
- 2815
2816
2817
2818
- 2819 [48] Y. Giomataris, Ph. Reboursard, J.P. Robert, and G. Charpak. Micromegas: a high-granularity position-sensitive gaseous detector for high particle-flux environments. *Nuclear Instruments and Methods in Physics Research Section A: Accelerators, Spectrometers, Detectors and Associated Equipment*, 376(1):29 – 35, 1996. ISSN 0168-9002. doi: [http://dx.doi.org/10.1016/0168-9002\(96\)00175-1](http://dx.doi.org/10.1016/0168-9002(96)00175-1). URL <http://www.sciencedirect.com/science/article/pii/0168900296001751>.
- 2820
2821
2822
2823
2824
- 2825 [49] T. Alexopoulos, J. Burnens, R. de Oliveira, G. Glonti, O. Pizzirusso, V. Polychronakos, G. Sekhniadze, G. Tsipolitis, and J. Wotschack. A spark-resistant bulk-micromegas chamber for high-rate applications. *Nuclear Instruments and Methods in Physics Research Section A: Accelerators, Spectrometers, Detectors and Associated Equipment*, 640(1):110 – 118, 2011. ISSN 0168-9002. doi: <http://dx.doi.org/10.1016/j.nima.2011.03.025>. URL <http://www.sciencedirect.com/science/article/pii/S0168900211005869>.
- 2826
2827
2828
2829
2830
- 2831 [50] Joao Pequenao and Paul Schaffner. An computer generated image representing how ATLAS detects particles. Jan 2013. URL <https://cds.cern.ch/record/1505342>.
- 2832

- 2833 [51] Improved electron reconstruction in ATLAS using the Gaussian Sum Filter-based model for
 2834 bremsstrahlung. Technical Report ATLAS-CONF-2012-047, CERN, Geneva, May 2012. URL
 2835 <https://cds.cern.ch/record/1449796>.
- 2836 [52] Electron efficiency measurements with the ATLAS detector using the 2012 LHC proton-proton
 2837 collision data. Technical Report ATLAS-CONF-2014-032, CERN, Geneva, Jun 2014. URL
 2838 <https://cds.cern.ch/record/1706245>.
- 2839 [53] Georges Aad et al. Electron and photon energy calibration with the ATLAS detector using LHC
 2840 Run 1 data. *Eur. Phys. J.*, C74(10):3071, 2014. doi: 10.1140/epjc/s10052-014-3071-4.
- 2841 [54] Georges Aad et al. Measurement of the muon reconstruction performance of the ATLAS detector
 2842 using 2011 and 2012 LHC proton–proton collision data. *Eur. Phys. J.*, C74(11):3130, 2014. doi:
 2843 10.1140/epjc/s10052-014-3130-x.
- 2844 [55] W Lampl, S Laplace, D Lelas, P Loch, H Ma, S Menke, S Rajagopalan, D Rousseau, S Snyder,
 2845 and G Unal. Calorimeter Clustering Algorithms: Description and Performance. Technical Re-
 2846 port ATL-LARG-PUB-2008-002, ATL-COM-LARG-2008-003, CERN, Geneva, Apr 2008. URL
 2847 <https://cds.cern.ch/record/1099735>.
- 2848 [56] Georges Aad et al. Topological cell clustering in the ATLAS calorimeters and its performance in
 2849 LHC Run 1. 2016.
- 2850 [57] Matteo Cacciari, Gavin P. Salam, and Gregory Soyez. The Anti-k(t) jet clustering algorithm. *JHEP*,
 2851 04:063, 2008. doi: 10.1088/1126-6708/2008/04/063.
- 2852 [58] Monte Carlo Calibration and Combination of In-situ Measurements of Jet Energy Scale, Jet Energy
 2853 Resolution and Jet Mass in ATLAS. Technical Report ATLAS-CONF-2015-037, CERN, Geneva,
 2854 Aug 2015. URL <http://cds.cern.ch/record/2044941>.
- 2855 [59] Georges Aad et al. Performance of *b*-Jet Identification in the ATLAS Experiment. 2015.
- 2856 [60] Expected performance of the ATLAS *b*-tagging algorithms in Run-2. Technical Report
 2857 ATL-PHYS-PUB-2015-022, CERN, Geneva, Jul 2015. URL <http://cds.cern.ch/record/2037697>.
- 2859 [61] Georges Aad et al. Performance of Missing Transverse Momentum Reconstruction in Proton-
 2860 Proton Collisions at 7 TeV with ATLAS. *Eur. Phys. J.*, C72:1844, 2012. doi: 10.1140/epjc/
 2861 s10052-011-1844-6.
- 2862 [62] Performance of Missing Transverse Momentum Reconstruction in ATLAS studied in Proton-
 2863 Proton Collisions recorded in 2012 at 8 TeV. Technical Report ATLAS-CONF-2013-082, CERN,
 2864 Geneva, Aug 2013. URL <http://cds.cern.ch/record/1570993>.

- 2865 [63] ATLAS Collaboration. Observation and measurement of Higgs boson decays to WW^* with the
 2866 ATLAS detector. *Phys. Rev. D*, 92(012006), 2015.
- 2867 [64] Aaron James Armbruster. Discovery of a Higgs Boson with the ATLAS detector. 2013. CERN-
 2868 THESIS-2013-047.
- 2869 [65] G. Cowan, K. Cranmer, E. Gross, and O. Vitells. Asymptotic formulae for likelihood-based tests
 2870 of new physics. *Eur. Phys. J.*, C 71:1554, 2011. doi: 10.1140/epjc/s10052-011-1554-0.
- 2871 [66] ATLAS Collaboration. Limits on the production of the Standard Model Higgs Boson in pp
 2872 collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector. *Eur. Phys. J.*, C 71:1728, 2011. doi:
 2873 10.1140/epjc/s10052-011-1728-9.
- 2874 [67] ATLAS Collaboration. Performance of the ATLAS muon trigger in pp collisions at $\sqrt{s} = 8$ TeV.
 2875 *Eur. Phys. J. C*, (arXiv:1408.3179. CERN-PH-EP-2014-154):75. 19 p, Aug 2014. URL <https://cds.cern.ch/record/1749694>.
- 2877 [68] ATLAS collaboration. Electron trigger performance in 2012 ATLAS data, 2015. ATLAS-COM-
 2878 DAQ-2015-091.
- 2879 [69] Paolo Nason. A new method for combining NLO QCD with shower Monte Carlo algorithms.
 2880 *JHEP*, 11:040, 2004.
- 2881 [70] B. P. Kersevan and E. Richter-Was. The Monte Carlo event generator AcerMC version 2.0 with
 2882 interfaces to PYTHIA 6.2 and HERWIG 6.5. 2004.
- 2883 [71] Nikolas Kauer and Giampiero Passarino. Inadequacy of zero-width approximation for a light Higgs
 2884 boson signal. 2012.
- 2885 [72] T. Gleisberg, Stefan Hoeche, F. Krauss, M. Schonherr, S. Schumann, et al. Event generation with
 2886 SHERPA 1.1. *JHEP*, 0902:007, 2009. doi: 10.1088/1126-6708/2009/02/007.
- 2887 [73] Michelangelo L. Mangano et al. ALPGEN, a generator for hard multiparton processes in hadronic
 2888 collisions. *JHEP*, 0307:001, 2003. doi: 10.1088/1126-6708/2003/07/001.
- 2889 [74] Torbjorn Sjostrand, Stephen Mrenna, and Peter Z. Skands. PYTHIA 6.4 Physics and Manual.
 2890 *JHEP*, 0605:026, 2006. doi: 10.1088/1126-6708/2006/05/026.
- 2891 [75] Torbjorn Sjostrand, Stephen Mrenna, and Peter Z. Skands. A Brief Introduction to PYTHIA 8.1.
 2892 *Comput.Phys.Commun.*, 178:852–867, 2008. doi: 10.1016/j.cpc.2008.01.036.
- 2893 [76] G. Corcella et al. HERWIG 6: An event generator for hadron emission reactions with interfering
 2894 gluons (including super-symmetric processes). *JHEP*, 01:010, 2001. doi: 10.1088/1126-6708/2001/
 2895 01/010.

- 2896 [77] J. M. Butterworth, Jeffrey R. Forshaw, and M. H. Seymour. Multiparton interactions in photo-
 2897 production at HERA. *Z. Phys.*, C 72:637, 1996. doi: 10.1007/s002880050286.
- 2898 [78] Jun Gao, Marco Guzzi, Joey Huston, Hung-Liang Lai, Zhao Li, et al. The CT10 NNLO Global
 2899 Analysis of QCD. *Phys. Rev.*, D89:033009, 2014. doi: 10.1103/PhysRevD.89.033009.
- 2900 [79] P. M. Nadolsky. Implications of CTEQ global analysis for collider observables. *Phys. Rev.*, D 78:
 2901 013004, 2008. doi: 10.1103/PhysRevD.78.013004.
- 2902 [80] A. Sherstnev and R. S. Thorne. Parton distributions for the LHC. *Eur. Phys. J.*, C 55:553, 2009. doi:
 2903 10.1140/epjc/s10052-008-0610-x.
- 2904 [81] S. Agostinelli et al. GEANT4, a simulation toolkit. *Nucl. Instrum. Meth.*, A 506:250, 2003. doi:
 2905 10.1016/S0168-9002(03)01368-8.
- 2906 [82] R.K. Ellis, I. Hinchliffe, M. Soldate, and J.J. Van Der Bij. Higgs decay to $\tau+\tau$ —a possible signature
 2907 of intermediate mass higgs bosons at high energy hadron colliders. *Nuclear Physics B*, 297(2):221
 2908 – 243, 1988. ISSN 0550-3213. doi: [http://dx.doi.org/10.1016/0550-3213\(88\)90019-3](http://dx.doi.org/10.1016/0550-3213(88)90019-3). URL <http://www.sciencedirect.com/science/article/pii/0550321388900193>.
- 2910 [83] Eilam Gross and Ofer Vitells. Transverse mass observables for charged Higgs boson searches at
 2911 hadron colliders. *Phys. Rev.*, D81:055010, 2010. doi: 10.1103/PhysRevD.81.055010.
- 2912 [84] J. R. Andersen et al. Les Houches 2013: Physics at TeV Colliders: Standard Model Working Group
 2913 Report. 2014.
- 2914 [85] I. Stewart and F. Tackmann. Theory uncertainties for Higgs mass and other searches using jet bins.
 2915 *Phys. Rev.*, D 85:034011, 2012. doi: 10.1103/PhysRevD.85.034011.
- 2916 [86] ATLAS Collaboration. Luminosity Determination in pp Collisions at $\sqrt{s} = 7$ TeV Using the
 2917 ATLAS Detector at the LHC. *Eur. Phys. J.*, C 71:1630, 2011. doi: 10.1140/epjc/s10052-011-1630-5.
- 2918 [87] Jet energy scale and its systematic uncertainty in proton-proton collisions at $\sqrt{s} = 7$ tev with atlas
 2919 data. *ATLAS-CONF-2013-004*, 2013.
- 2920 [88] Calibrating the b -tag efficiency and mistag rate in 35 pb^{-1} of data with the atlas detector. *ATLAS-*
 2921 *CONF-2011-089*, 2011.
- 2922 [89] ATLAS Collaboration. Measurement of the b -tag Efficiency in a Sample of Jets Containing Muons
 2923 with 5 fb^{-1} of Data from the ATLAS Detector. *ATLAS-CONF-2012-043*, 2012. URL <http://cdsweb.cern.ch/record/1435197>.

- 2925 [90] ATLAS Collaboration. Calibration of b -tagging using dileptonic top pair events in a combinatorial
 2926 likelihood approach with the ATLAS experiment. (ATLAS-CONF-2014-004), 2014. URL <http://cds.cern.ch/record/1664335>.
- 2928 [91] Georges Aad et al. Measurement of the Higgs boson mass from the $H \rightarrow \gamma\gamma$ and $H \rightarrow ZZ^* \rightarrow$
 2929 4ℓ channels with the ATLAS detector using 25 fb^{-1} of pp collision data. *Phys. Rev.*, D90(5):052004,
 2930 2014. doi: 10.1103/PhysRevD.90.052004.
- 2931 [92] Georges Aad et al. Measurements of the Higgs boson production and decay rates and coupling
 2932 strengths using pp collision data at $\sqrt{s} = 7$ and 8 TeV in the ATLAS experiment. *Eur. Phys. J.*,
 2933 C76(1):6, 2016. doi: 10.1140/epjc/s10052-015-3769-y.
- 2934 [93] W.J. Stirling. $7/8$ and $13/8$ TeV LHC luminosity ratios. 2013. URL http://www.hep.ph.ic.ac.uk/~wstirlin/plots/lhclumi7813_2013_v0.pdf.
- 2936 [94] J Alison. Experimental Studies of hh. Oct 2014. URL <http://cds.cern.ch/record/1952581>.
- 2937 [95] J. Alwall et al. The automated computation of tree-level and next-to-leading order differential cross
 2938 sections, and their matching to parton shower simulations. *JHEP*, 07:079, 2014.
- 2939 [96] Richard D. Ball et al. Parton distributions with LHC data. *Nucl. Phys. B*, 867:244, 2013.
- 2940 [97] ATLAS Collaboration. ATLAS Run 1 Pythia8 tunes. (ATL-PHYS-PUB-2014-021), Nov 2014.
 2941 URL <https://cds.cern.ch/record/1966419>.
- 2942 [98] M. Bahr et al. Herwig++ Physics and Manual. *Eur. Phys. J. C*, 58:639–707, 2008. doi: 10.1140/
 2943 epjc/s10052-008-0798-9.
- 2944 [99] Stefan Gieseke, Christian Rohr, and Andrzej Siódak. Colour reconnections in Herwig++. *Eur.
 2945 Phys. J. C*, 72:2225, 2012. doi: 10.1140/epjc/s10052-012-2225-5.
- 2946 [100] Simone Alioli, Paolo Nason, Carlo Oleari, and Emanuele Re. A general framework for implement-
 2947 ing NLO calculations in shower Monte Carlo programs: the POWHEG BOX. *JHEP*, 06:043,
 2948 2010.
- 2949 [101] Peter Zeiler Skands. Tuning Monte Carlo Generators: The Perugia Tunes. *Phys. Rev. D*, 82:074018,
 2950 2010. doi: 10.1103/PhysRevD.82.074018.
- 2951 [102] Michal Czakon and Alexander Mitov. Top++: A Program for the Calculation of the Top-Pair
 2952 Cross-Section at Hadron Colliders. 2011.
- 2953 [103] Baojia (Tony) Tong. Private communication.
- 2954 [104] D. Krohn, J. Thaler, and L.-T. Wang. Jet Trimming. *JHEP*, 02:084, 2010. doi: 10.1007/
 2955 JHEP02(2010)084.

- 2956 [105] ATLAS Collaboration. Identification of Boosted, Hadronically Decaying W Bosons and Compar-
 2957 isons with ATLAS Data Taken at $\sqrt{s} = 8$ TeV. 2015.
- 2958 [106] Expected Performance of Boosted Higgs ($\rightarrow b\bar{b}$) Boson Identification with the ATLAS Detector
 2959 at $\sqrt{s} = 13$ TeV. Technical Report ATL-PHYS-PUB-2015-035, CERN, Geneva, Aug 2015. URL
 2960 <https://cds.cern.ch/record/2042155>.
- 2961 [107] Flavor Tagging with Track Jets in Boosted Topologies with the ATLAS Detector. Technical Report
 2962 ATL-PHYS-PUB-2014-013, CERN, Geneva, Aug 2014. URL <https://cds.cern.ch/record/1750681>.
- 2964 [108] Matteo Cacciari and Gavin P. Salam. Pileup subtraction using jet areas. *Phys. Lett. B*, 659:119, 2008.
 2965 doi: 10.1016/j.physletb.2007.09.077.
- 2966 [109] Glen Cowan, Eilam Gross. Discovery significance with statistical uncertainty in the background
 2967 estimate. 2008. URL <http://www.pp.rhul.ac.uk/~cowan/stat/notes/SigCalcNote.pdf>.
- 2969 [110] Search for pair production of Higgs bosons in the $b\bar{b}b\bar{b}$ final state using proton-proton collisions
 2970 at $\sqrt{s} = 13$ TeV with the ATLAS detector. Technical Report ATLAS-CONF-2016-017, CERN,
 2971 Geneva, Mar 2016. URL <https://cds.cern.ch/record/2141006>.
- 2972 [111] Qi Zeng. Private communication.
- 2973 [112] ATLAS Collaboration. Identification of boosted, hadronically-decaying W and Z bosons in
 2974 $\sqrt{s} = 13$ TeV Monte Carlo Simulations for ATLAS. (ATL-PHYS-PUB-2015-033), Aug 2015. URL
 2975 <https://cds.cern.ch/record/2041461>.
- 2976 [113] ATLAS Collaboration. Performance of b -Jet Identification in the ATLAS Experiment. 2015.
- 2977 [114] Alexander L. Read. Presentation of search results: The CL(s) technique. *J. Phys. G*, 28:2693, 2002.
 2978 doi: 10.1088/0954-3899/28/10/313.
- 2979 [115] Measurements of the Higgs boson production and decay rates and constraints on its couplings
 2980 from a combined ATLAS and CMS analysis of the LHC pp collision data at $\sqrt{s} = 7$ and 8 TeV.
 2981 Technical Report ATLAS-CONF-2015-044, CERN, Geneva, Sep 2015. URL <http://cds.cern.ch/record/2052552>.
- 2983 [116] Projections for measurements of Higgs boson signal strengths and coupling parameters with the
 2984 ATLAS detector at a HL-LHC. Technical Report ATL-PHYS-PUB-2014-016, CERN, Geneva,
 2985 Oct 2014. URL <http://cds.cern.ch/record/1956710>.
- 2986 [117] ATLAS Phase-II Upgrade Scoping Document. Technical Report CERN-LHCC-2015-020. LHCC-
 2987 G-166, CERN, Geneva, Sep 2015. URL <http://cds.cern.ch/record/2055248>.



2988

THIS THESIS WAS TYPESET using \LaTeX , originally developed by Leslie Lamport and based on Donald Knuth's \TeX . The body text is set in 11 point Egenolff-Berner Garamond, a revival of Claude Garamont's humanist typeface. The above illustration, *Science Experiment 02*, was created by Ben Schlitter and released under [CC BY-NC-ND 3.0](#). A template that can be used to format a PhD dissertation with this look & feel has been released under the permissive [AGPL](#) license, and can be found online at github.com/asm-products/Dissertate or from its lead author, Jordan Suchow, at suchow@post.harvard.edu.