

# The Higgs boson with the ATLAS experiment at the LHC: Discovery, measurement, and searches for new physics

A DISSERTATION PRESENTED  
BY  
TOMO LAZOVICH  
TO  
THE DEPARTMENT OF PHYSICS

IN PARTIAL FULFILLMENT OF THE REQUIREMENTS  
FOR THE DEGREE OF  
DOCTOR OF PHILOSOPHY  
IN THE SUBJECT OF  
PHYSICS

HARVARD UNIVERSITY  
CAMBRIDGE, MASSACHUSETTS  
MAY 2016

©2014 – TOMO LAZOVICH  
ALL RIGHTS RESERVED.

# The Higgs boson with the ATLAS experiment at the LHC: Discovery, measurement, and searches for new physics

## ABSTRACT

We measured things. And searched for other things. Here is what we found, please let me graduate.

# Contents

|     |  |    |
|-----|--|----|
| 0   | INTRODUCTION   | I  |
| I   | Preliminaries  | 2  |
| 1   | THE STANDARD MODEL AND BEYOND: A THEORETICAL OVERVIEW  | 3  |
| 1.1 | The Standard Model of Particle Physics . . . . .   | 3  |
| 1.2 | Electroweak Symmetry Breaking and the Higgs . . . . .  | 3  |
| 1.3 | Higgs Boson Production and Decay . . . . .   | 3  |
| 1.4 | Physics Beyond the Standard Model . . . . .  | 3  |
| 2   | THE ATLAS DETECTOR AND THE LARGE HADRON COLLIDER   | 4  |
| 2.1 | The Large Hadron Collider . . . . .  | 4  |
| 2.2 | The ATLAS Detector . . . . .   | 4  |
| II  | Observation and measurement of Higgs boson decays to $WW^*$ with the ATLAS detector in LHC Run I at $\sqrt{s} = 7$ and 8 TeV | 5  |
| 3   | $H \rightarrow WW^* \rightarrow \ell\nu\ell\nu$ ANALYSIS STRATEGY  | 6  |
| 3.1 | Introduction . . . . .   | 6  |
| 3.2 | Signal topology . . . . .  | 7  |
| 3.3 | Background processes . . . . .   | 8  |
| 3.4 | Isolating an $H \rightarrow WW^* \rightarrow \ell\nu\ell\nu$ signal . . . . .  | 10 |
| 3.5 | Background reduction in same-flavor final states . . . . .   | 12 |
| 3.6 | Parameters of interest and statistical treatment . . . . .   | 12 |
| 4   | THE DISCOVERY OF THE HIGGS BOSON AND THE ROLE OF THE $H \rightarrow WW^* \rightarrow \ell\nu\ell\nu$ CHANNEL                 | 13 |
| 5   | OBSERVATION OF VECTOR BOSON FUSION PRODUCTION OF $H \rightarrow WW^* \rightarrow \ell\nu\ell\nu$                             | 14 |
| 6   | COMBINED RUN I $H \rightarrow WW^* \rightarrow \ell\nu\ell\nu$ RESULTS   | 15 |

|     |   |    |
|-----|---|----|
| III | Search for Higgs pair production in the $HH \rightarrow b\bar{b}b\bar{b}$ channel in LHC Run 2 at $\sqrt{s} = 13$ TeV | 16 |
| 7   | SEARCH OVERVIEW   | 17 |
| 8   | SEARCH FOR HIGGS PAIR PRODUCTION IN BOOSTED FINAL STATES  | 18 |
| 9   | RESULTS WITH RUN 2 2015 DATASET   | 19 |
| IV  | Looking ahead   | 20 |
| 10  | CONCLUSION  | 21 |
|     | REFERENCES  | 22 |

# Listing of figures

|     |  |    |
|-----|--|----|
| 3.1 | A cartoon of the $WW$ final state. Momenta are represented with thin arrows, spins with thick arrows. <sup>1</sup> . . . . .         | 7  |
| 3.2 | Feynman diagram for Standard Model $WW$ production . . . . .   | 8  |
| 3.3 | Feynman diagrams for top pair production (left) and $Wt$ production (right) . . . . .  | 9  |
| 3.4 | An example Feynman diagram of $W$ +jets production . . . . .   | 10 |
| 3.5 | An example Feynman diagram of $Z$ +jets production . . . . .   | 10 |
| 3.6 | An illustration of the unique analysis signal regions <sup>1</sup> . . . . .   | 12 |
| 3.7 | A comparison of the subleading lepton $p_T$ spectrum between VBF $H \rightarrow WW^*$ production and $t\bar{t}$ background . . . . . | 12 |

# Listing of tables

|     |  |    |
|-----|--|----|
| 3.1 | A summary of backgrounds to the $H \rightarrow WW^* \rightarrow \ell\nu\ell\nu$ signal . . . . . | II |
|-----|--|----|

THIS IS THE DEDICATION.



# Acknowledgments

LOREM IPSUM DOLOR SIT AMET, consectetur adipiscing elit. Morbi commodo, ipsum sed pharetra gravida, orci magna rhoncus neque, id pulvinar odio lorem non turpis. Nullam sit amet enim. Suspendisse id velit vitae ligula volutpat condimentum. Aliquam erat volutpat. Sed quis velit. Nulla facilisi. Nulla libero. Vivamus pharetra posuere sapien. Nam consectetur. Sed aliquam, nunc eget euismod ullamcorper, lectus nunc ullamcorper orci, fermentum bibendum enim nibh eget ipsum. Donec porttitor ligula eu dolor. Maecenas vitae nulla consequat libero cursus venenatis. Nam magna enim, accumsan eu, blandit sed, blandit a, eros.

# 0

## Introduction

# Part I

## Preliminaries

# 1

## The Standard Model and beyond: a theoretical overview

- I.1 THE STANDARD MODEL OF PARTICLE PHYSICS
- I.2 ELECTROWEAK SYMMETRY BREAKING AND THE HIGGS
- I.3 HIGGS BOSON PRODUCTION AND DECAY
- I.4 PHYSICS BEYOND THE STANDARD MODEL

*This is some random quote to start off the chapter.*

Firstname lastname

# 2

## The ATLAS detector and the Large Hadron Collider

### 2.1 THE LARGE HADRON COLLIDER

### 2.2 THE ATLAS DETECTOR

## Part II

Observation and measurement of Higgs  
boson decays to  $WW^*$  with the ATLAS  
detector in LHC Run I at  $\sqrt{s} = 7$  and 8 TeV

*Basic research is what I am doing when I don't know  
what I am doing.*

Wernher von Braun

# 3

## $H \rightarrow WW^* \rightarrow \ell\nu\ell\nu$ Analysis Strategy

### 3.1 INTRODUCTION

This chapter will present an overview of the strategy for searching for a Higgs boson in the  $H \rightarrow WW^* \rightarrow \ell\nu\ell\nu$  decay topology. Its purpose is to present in broad terms how the search and measurement are undertaken, before going into details on the specific sub-categories within the broader analysis.

First, the topology of the signal final state and corresponding backgrounds are presented. Next, an overview of the variables used to reduce the backgrounds and enhance the signal is given. These will be described in general, while specific values of selection cuts and background estimation will be provided in subsequent chapters. Finally, the parameters of interest in the search and measurement will be defined, and a brief overview of the statistical treatment of the final Higgs candidates is shown.



**Figure 3.1:** A cartoon of the  $WW$  final state. Momenta are represented with thin arrows, spins with thick arrows.<sup>1</sup>

### 3.2 SIGNAL TOPOLOGY

The analysis presented here and in subsequent chapters is the study of the Higgs boson in the  $WW$  final state, where each  $W$  boson subsequently decays into a charged lepton and a neutrino. In its simplest form, the final state will then consist of two neutrinos and two charged leptons, each of which can be either an electron or a muon. If one or both of the  $W$ s decay to  $\tau$  leptons, only leptonic decays of the  $\tau$  are considered, leading to additional neutrinos in the final state but still giving two charged leptons as before. Neutrinos are not detected in ATLAS, so the final state ultimately consists of two reconstructed leptons and missing transverse momentum (denoted as  $E_T^{\text{miss}}$ ). Final states where both of the charged leptons are electrons or muons are referred to as the “same flavor” final states, while those with one electron and one muon are referred to as “different flavor”.

The final state leptons will also exhibit unique correlations due to the fact that they are arising from the decay of a spin zero resonance. In particular, the spins of the final state leptons and neutrinos must all cancel, as shown in figure 3.1. Because the neutrino has a left handed helicity and the anti-neutrino has a right handed helicity, the spin and momentum of the particles will be anti-aligned and aligned, respectively. In the transverse plane, the momenta of all four final state objects must cancel as well. With the constraint of having both the momenta and the spin alignments cancel, the final state kinematics strongly prefer having a small angle between the leptons in the transverse plane (low  $\Delta\phi_{\ell\ell}$ ). This angular correlation will also lead to low values of the di-lepton invariant mass  $m_{\ell\ell}$ . These unique signal final state kinematic correlations will be exploited to define the ultimate signal region.




While the basic final state consists of two leptons and  $E_T^{\text{miss}}$ , there can be additional objects as well depending on the production mode of the Higgs. As described in detail in Chapter 1, if the Higgs is produced via vector boson fusion production, there will be two additional forward jets in the event. Even in gluon fusion, one or more jets can be produced through initial state radiation from the incoming gluons. The analysis is separated into different signal regions depending on the number of hard jets reconstructed in the final state as well.

### 3.3 BACKGROUND PROCESSES

Many processes from the Standard Model can also produce a final state with two leptons and missing transverse momentum. This section lists the dominant backgrounds to Higgs production. It gives general descriptions of how the backgrounds mimic Higgs production and how they can be reduced. The details of background estimation and specific cuts are left for later sections. Table 3.1 summarizes the different processes.

#### 3.3.1 STANDARD MODEL WW PRODUCTION



**Figure 3.2:** Feynman diagram for Standard Model WW production

Non-resonant Standard Model diboson production, as shown in figure 3.2, is an irreducible background to Higgs boson production in the WW final state. It produces the same exact final state objects, namely leptonically decaying W bosons. There are no additional objects in the final state that allow for background reduction. Therefore the analysis solely relies on the correlations between the leptons to reduce this background.

### 3.3.2 TOP QUARK PRODUCTION

Production of top quarks, either in pairs ( $t\bar{t}$  production) or singly (e.g.  $Wt$  production), can also mimic Higgs production. Because top quarks decay via  $t \rightarrow Wb$ , top pair production can produce a final state with two  $W$  bosons that then decay leptonically. In this case, however, there are two additional jets from the bottom quarks in the final state. This allows the analysis to veto on the presence of jets identified as originating from a  $b$  in order to reduce the size of the background.

Single top production can occur via  $s$ -channel,  $t$ -channel, or associated production ( $Wt$ ). The mode which most closely resembles the Higgs final state is  $Wt$ . In this case, there are two real  $W$  bosons produced, as with  $t\bar{t}$ . However, the decay of the single top quark will still also produce one  $b$ -jet, meaning a  $b$  veto will reduce this background as well.

Figure 3.3 shows the Feynman diagrams for  $t\bar{t}$  and  $Wt$  production.




Figure 3.3: Feynman diagrams for top pair production (left) and  $Wt$  production (right)

### 3.3.3 $W$ +JETS BACKGROUND

Single  $W$  boson production, in association with jets, is a unique background. The other background considered so far have all included real leptons in the final state. In this case, however, only one real lepton from the decay of a  $W$  exists in the final state. The second reconstructed lepton can arise from two different cases. First, the lepton may truly be an algorithm “fake”, or a jet misidentified as a lepton by either the electron or muon reconstruction algorithms. Second, the lepton may be a real lepton but coming from semi-leptonic decays of particles inside the shower of the jet. This background can be reduced by requiring that the reconstructed lepton have little activity surrounding it in the calorimeter (also known as an “isolated” lepton). Figure 3.4 shows the Feynman diagram for  $W$ +jets production.




Figure 3.4: An example Feynman diagram of  $W^+$  jets production

### 3.3.4 $Z/\gamma^*$ +JETS BACKGROUND

Production of a  $Z/\gamma^*$  in association with jets (also known as Drell-Yan) is also a background to Higgs production. In particular, the same flavor final states have a large  $Z$ +jets background, as the  $Z$  decays into two leptons of the same flavor. (This background also enters the different flavor final state through the leptonic decays of  $Z \rightarrow \tau\tau$ ). Figure 3.5 shows the production of a  $Z$  in association with one jet. Because there are no neutrinos in this final state, variables like  $E_T^{\text{miss}}$  can be used to reduce the background.




Figure 3.5: An example Feynman diagram of  $Z$ +jets production

### 3.3.5 OTHER (SUBDOMINANT) BACKGROUNDS

There are additional processes which contribute to the background composition but are not produced as frequently as those listed already. The first of these are referred to as  $VV$  or “Other diboson” processes and include multiple Standard Model diboson processes, including  $WZ$ ,  $ZZ$ ,  $W\gamma$ ,  $W\gamma^*$ , and  $Z\gamma$  production. Additionally, there is background from QCD multijet production, where two jets are misidentified as leptons.

## 3.4 ISOLATING AN $H \rightarrow WW^* \rightarrow \ell\nu\ell\nu$ SIGNAL

As presented in section 3.2, there are many different combinations of objects that can define a  $H \rightarrow WW^* \rightarrow \ell\nu\ell\nu$  final state. The multiplicity of jets and the flavor combinations of the leptons both lead to a combinatorically large number of potential signal regions. Additionally, signal regions can be optimized separately to be sensitive to the distinct production modes of the Higgs. Gluon fusion, vector

| Category             | Process  | Description  |
|----------------------|--|--|
| SM $WW$              | $WW \rightarrow \ell\nu\ell\nu$  | Real leptons and neutrinos                             |
| Top quark production | $t\bar{t} \rightarrow WbW\bar{b} \rightarrow \ell\nu b\ell\nu\bar{b}$        | Real leptons, untagged $b\bar{s}$                      |
|                      | $tW \rightarrow WbW \rightarrow \ell\nu\ell\nu b$                            | Real leptons, untagged $b$                             |
|                      | $t\bar{b}, tq\bar{b}$  | Untagged $b$ , jet misidentified as lepton             |
| Drell-Yan            | $Z/\gamma^* \rightarrow e\bar{e}, \mu\bar{\mu}$                              | “Fake” $E_T^{\text{miss}}$                             |
|                      | $Z/\gamma^* \rightarrow \tau\bar{\tau} \rightarrow \ell\nu\ell\nu$           | Real leptons and neutrinos                             |
| Other dibosons       | $ZZ \rightarrow \ell\ell\nu\nu$  | Real leptons and neutrinos                             |
|                      | $W\gamma^*, WZ \rightarrow \ell\nu\ell\ell, ZZ \rightarrow \ell\ell\ell\ell$ | Unreconstructed leptons                                |
|                      | $W\gamma, Z\gamma$   | $\gamma$ reconstructed as $e$ , unreconstructed lepton |
| $W$ +jets            | $Wj \rightarrow \ell\nu j$   | Jet reconstructed as lepton                            |
| QCD multijet         | $jj$   | Jets reconstructed as leptons                          |


**Table 3.1:** A summary of backgrounds to the  $H \rightarrow WW^* \rightarrow \ell\nu\ell\nu$  signal

boson fusion, and associated production of a Higgs all lead to unique final state topologies. Figure 3.6 delineates the different signal regions used in the gluon fusion and vector boson fusion  $H \rightarrow WW^*$  analyses. While there are different optimizations possible in each signal region, there are also some commonly shared selections that will be described here.


#### 3.4.1 EVENT PRESELECTION

Before being sorted into the distinct signal regions, basic cuts are applied on the reconstructed objects in the event to select Higgs-like event candidates. First, two oppositely charged leptons are required. The  $p_T$  threshold on the leptons is a particularly important consideration for this signal. Because the second  $W$  produced in the decay can be off-shell, it tends to produce lower momentum leptons. Thus, being able to lower the  $p_T$  threshold while still maintaining a low background rate is critical. Figure 3.7 shows an example of the subleading lepton  $p_T$  for a VBF  $H \rightarrow WW^*$  signal compared to the corresponding  $t\bar{t}$  background.

Once the leptons are selected, additional pre-selection requirements can be made.



**Figure 3.6:** An illustration of the unique analysis signal regions<sup>1</sup>



**Figure 3.7:** A comparison of the subleading lepton  $p_T$  spectrum between VBF  $H \rightarrow WW^*$  production and  $t\bar{t}$  background

### 3.5 BACKGROUND REDUCTION IN SAME-FLAVOR FINAL STATES

### 3.6 PARAMETERS OF INTEREST AND STATISTICAL TREATMENT

# 4

The discovery of the Higgs boson and the  
role of the  $H \rightarrow WW^* \rightarrow \ell\nu\ell\nu$  channel

# 5

Observation of Vector Boson Fusion

production of  $H \rightarrow WW^* \rightarrow \ell\nu\ell\nu$

# 6

Combined Run 1  $H \rightarrow WW^* \rightarrow \ell\nu\ell\nu$  results



## Part III

Search for Higgs pair production in the  
 $HH \rightarrow b\bar{b}b\bar{b}$  channel in LHC Run 2 at  $\sqrt{s} =$   
13 TeV

# 7

## Search overview

# 8

Search for Higgs pair production in boosted  
final states

# 9

Results with Run 2 2015 dataset

## Part IV

### Looking ahead

# 10

## Conclusion

We found the Higgs. Then measured it. Then used it to look for new physics. What a time to be alive!

# References

- [1] Collaboration, A. (2015). Observation and measurement of higgs boson decays to  $ww^*$  with the atlas detector. *Phys. Rev. D*, 92(012006).



**T**HIS THESIS WAS TYPESET using  $\text{\LaTeX}$ , originally developed by Leslie Lamport and based on Donald Knuth's  $\text{\TeX}$ .

The body text is set in 11 point Egenolff-Berner Garamond, a revival of Claude Garamont's humanist typeface. The above illustration, *Science Experiment 02*, was created by Ben Schlitter and released under [CC BY-NC-ND 3.0](#). A template that can be used to format a PhD dissertation with this look & feel has been released under the permissive AGPL license, and can be found online at [github.com/asm-products/Dissertate](https://github.com/asm-products/Dissertate) or from its lead author, Jordan Suchow, at [suchow@post.harvard.edu](mailto:suchow@post.harvard.edu).