The proof of this theorem, following a similar proof by A.V. Skorokhod, is given in Section 5, after we make a crucial step in the next section where we prove, in particular, that for solutions of (1.2), any Borel $f \geq 0$, and $T \in (0, \infty)$

$$E \int_0^T f(t, x_t)\, dt \leq N \|f\|_{p,q}, \tag{3.2}$$

where N is independent of f and (t^0, x^0).

It is worth saying that deciding whether the solutions of (1.2) are weakly unique or not under our conditions is a very challenging open problem even if $\sigma^{ij} \equiv \delta^{ij}$.

Remark 3.1. Theorem 3.1 is also true if $d/p+1/q < 1$. This can be seen from its proof which becomes somewhat more technical in that case because of the form of our main estimate (4.9). Also the main interest in Theorem 3.1 is, of course, the lowest *local* integrability of b, when the condition $d/p + 1/q = 1$ is weaker than $d/p + 1/q < 1$ due to Hölder's inequality.

4. Estimates of the distributions of semimartingales

Here we first prove a version of Lemma 5.1 of [7]. The proof given in [7] uses somewhat advanced knowledge of very powerful results from the theory of fully nonlinear parabolic equations. We give a proof based on a simpler fact which in turn was one of the cornerstones of that theory.

Let $(\Omega, \mathcal{F}, P)$ be a complete probability space, let $\mathcal{F}_t, t \geq 0$, be an increasing family of complete σ-fields $\mathcal{F}_t \subset \mathcal{F}$, $t \geq 0$, let m_t be an $\mathbb{R}^d$-valued continuous local martingale relative to $\mathcal{F}_t$, let A_t be a continuous $\mathcal{F}_t$-adapted nondecreasing process, let B_t be a continuous $\mathbb{R}^d$-valued $\mathcal{F}_t$-adapted process which has finite variation (a.e.) on each finite time interval. Assume that

$$A_0 = 0, \quad m_0 = B_0 = 0, \quad d\langle m \rangle_t \ll dA_t$$

and that we are also given progressively measurable relative to $\mathcal{F}_t$ nonnegative processes r_t and c_t. Finally, take an $\mathcal{F}_0$ measurable $\mathbb{R}^d$-valued x_0 and introduce

$$x_t = x_0 + m_t + B_t, \quad \tau_t = \int_0^t r_s\, dA_s, \quad \phi_t = \int_0^t c_s\, dA_s, \quad a_t^{ij} = \frac{1}{2} \frac{d\langle m^i, m^j \rangle_t}{dA_t}.$$

Lemma 4.1. *Let γ be an $\mathcal{F}_t$-stopping time and set*

$$A = E \int_0^\gamma e^{-\phi_t} \operatorname{tr} a_s\, dA_t, \quad B = E \int_0^\gamma e^{-\phi_t}\, |dB_t|.$$

Then for any Borel $f(t, x) \geq 0$ we have

$$E \int_0^\gamma e^{-\phi_t} (r_t \det a_t)^{1/(d+1)} f(\tau_t, x_t)\, dA_t$$

$$\leq N(d)(B^2 + A)^{d/(2d+2)} \|f\|_{d+1}. \tag{4.1}$$

Proof. Without losing generality we may assume that $A < \infty$ and $B < \infty$. Furthermore, just stopping the processes A_t, m_t, and B_t at time γ, we reduce the general case to the one in which $\gamma = \infty$. In that case we also observe that, as usual, it suffices to prove (4.1) for $f \in C_0^\infty(\mathbb{R}^{d+1})$.

After these reductions we use Theorem 2.2.4 of [9] according to which, for any $\lambda > 0$ on $\mathbb{R}^{d+1}$, there exists a nonnegative function $v(t,x)$ such that

(i) all Sobolev derivatives $\partial_t v$, $D_i v$, $D_{ij} v$ exist and are bounded on $\mathbb{R}^{d+1}$ and $v \le N e^{-|x|/N}$ for all t, x and a constant N;

(ii) for any nonnegative symmetric $d \times d$ matrix α and $r \ge 0$,

$$r\partial_t v + \alpha^{ij} D_{ij} v - \lambda(r + \operatorname{tr}\alpha)v + \sqrt[d+1]{r \det \alpha}\, f \le 0,$$

$$\partial_t v - \lambda v \le 0, \quad (\lambda v \delta_{ij} - D_{ij} v) \ge 0, \quad |Dv| \le \sqrt{\lambda} v \quad (a.e.), \tag{4.2}$$

(iii) for any $y \in \mathbb{R}^d$, $t \in (-\infty, \infty)$, we have

$$v(t,y) e^{-\lambda t} \le N(d) \frac{1}{\lambda^{d/(2d+2)}} I_t, \tag{4.3}$$

where

$$I_t^{d+1} := \int_0^\infty ds \int_{\mathbb{R}^d} e^{-\lambda(d+1)(t+s)} f^{d+1}(t+s,x)\, dx.$$

Take a nonnegative $\zeta \in C_0^\infty(\mathbb{R}^{d+1})$ with unit integral, for $\varepsilon > 0$ denote $\zeta_\varepsilon(t,x) = \varepsilon^{-(d+1)} \zeta(\varepsilon t, \varepsilon x)$ and use the notation $u^{(\varepsilon)} = u * \zeta_\varepsilon$. Then $v^{(\varepsilon)}$ is infinitely differentiable and in light of (4.2), for any nonnegative symmetric $d \times d$ matrix α and $r \ge 0$,

$$r\partial_t v^{(\varepsilon)} + \alpha^{ij} D_{ij} v^{(\varepsilon)} - \lambda(r + \operatorname{tr}\alpha)v^{(\varepsilon)} + \sqrt[d+1]{r \det \alpha}\, f^{(\varepsilon)} \le 0,$$

$$\partial_t v^{(\varepsilon)} - \lambda v^{(\varepsilon)} \le 0, \quad (\lambda v^{(\varepsilon)} \delta_{ij} - D_{ij} v^{(\varepsilon)}) \ge 0, \quad |Dv^{(\varepsilon)}| \le \sqrt{\lambda} v^{(\varepsilon)}. \tag{4.4}$$

Next, by Itô's formula the process

$$v^{(\varepsilon)}(\tau_t, x_t) e^{-\phi_t - \lambda\tau_t} - \int_0^t e^{-\phi_s - \lambda\tau_s} D_i v^{(\varepsilon)}(\tau_s, x_s)\, dB_s^i$$

$$+ \int_0^t e^{-\phi_s - \lambda\tau_s} \big((\lambda r_s + c_s) v^{(\varepsilon)} - r_s \partial_t v^{(\varepsilon)} - a_s^{ij} D_{ij} v^{(\varepsilon)} \big)(\tau_s, x_s)\, dA_s$$

is a local martingale. Here owing to (4.4)

$$\big((\lambda r_s + c_s) v^{(\varepsilon)} - r_s \partial_t v^{(\varepsilon)} - a_s^{ij} D_{ij} v^{(\varepsilon)} \big)\, dA_s - D_i v^{(\varepsilon)}\, dB_s^i$$

$$\ge (r_s \det a_s)^{1/(d+1)} f^{(\varepsilon)}\, dA_s - \lambda \operatorname{tr} a_s v^{(\varepsilon)}\, dA_s - \sqrt{\lambda} v^{(\varepsilon)}\, |dB_s|.$$

Therefore, for

$$M^\varepsilon = \sup_{t \ge 0, x \in \mathbb{R}^d} v^{(\varepsilon)}(t,x) e^{-\lambda t}$$

the process

$$\kappa_t^\varepsilon := v^{(\varepsilon)}(\tau_t, x_t) e^{-\phi_t - \lambda\tau_t} + \int_0^t e^{-\phi_s - \lambda\tau_s} (r_s \det a_s)^{1/(d+1)} f^{(\varepsilon)}(\tau_s, x_s)\, dA_s$$

$$- \int_0^t e^{-\phi_s} \big(\lambda \operatorname{tr} a_s\, dA_s + \sqrt{\lambda}\, |dB_s| \big)\, M^\varepsilon$$

is a local supermartingale. In addition, it is bounded from below by a summable quantity ($A, B < \infty$). Hence, it is a supermartingale and by Fatou's lemma

$$Ev^{(\varepsilon)}(0,x_0) = \kappa_0^\varepsilon \geq E\int_0^\infty e^{-\phi_t - \lambda\tau_s}(r_t \det a_t)^{1/(d+1)} f^{(\varepsilon)}(\tau_t, x_t)\, dA_t$$
$$-M^\varepsilon(\lambda A + \sqrt{\lambda}B).$$

By sending $\varepsilon \downarrow 0$ and using (4.3) and Fatou's lemma once more we obtain

$$E\int_0^\infty e^{-\phi_t - \lambda\tau_s}(r_t \det a_t)^{1/(d+1)} f(\tau_t, x_t)\, dA_t$$
$$\leq N(d)\frac{1}{\lambda^{d/(2d+2)}}\Big(1 + \lambda A + \sqrt{\lambda}B\Big) I_0.$$

We replace here $e^{-\lambda t} f$ by f and arrive at

$$E\int_0^\infty e^{-\phi_t}(r_t \det a_t)^{1/(d+1)} f(\tau_t, x_t)\, dA_t$$
$$\leq N(d)\frac{1}{\lambda^{d/(2d+2)}}\Big(1 + \lambda A + \sqrt{\lambda}B\Big) \|f\|_{d+1}.$$

Now we use the arbitrariness of λ. If $A < B^2$, then for $\lambda = B^{-2}$ we have

$$\frac{1}{\lambda^{d/(2d+2)}}\Big(1 + \lambda A + \sqrt{\lambda}B\Big) \leq 3B^{d/(d+1)} \leq 3(B^2 + A)^{d/(2d+2)}.$$

If $A \geq B^2$ and $A > 0$, then for $\lambda = A^{-1}$ the above inequality between the extreme terms still holds. Finally, if $A = B = 0$, then the left-hand side of (4.1) is zero. The lemma is proved.

Lemma 4.2. *In the notation of Lemma 4.1 for any Borel $f(x) \geq 0$ we have*

$$E\int_0^\gamma e^{-\phi_t}(\det a_t)^{1/d} f(x_t)\, dA_t \leq N(d)(B^2 + A)^{1/2}\|f\|_{L_d(\mathbb{R}^d)}. \tag{4.5}$$

Proof. We follow a probabilistic version of an argument in [12]. We again may concentrate on the case of $A + B < \infty$, $\gamma = \infty$, and $f \in C_0^\infty(\mathbb{R}^d)$. In that case observe that by Theorem 2.2.3 of [9] there exists a nonnegative function $v(x)$ defined on $\mathbb{R}^d$ such that

(a) $v \leq Ne^{-|x|/N}$ for all x and a constant N; the generalized derivatives $D_i v$ and $D_{ij}v$, $i, j = 1, ..., d$, are bounded on $\mathbb{R}^d$;

(b) for any nonnegative symmetric $d \times d$ matrix α (a.e.)

$$-\lambda v \operatorname{tr}\alpha + \alpha^{ij} D_{ij} v + \sqrt[d]{\det \alpha} f \leq 0, \quad |Dv| \leq \sqrt{\lambda} v,$$
$$(\lambda v \delta_{ij} - D_{ij} v) \geq 0; \tag{4.6}$$

(c) for any $x \in \mathbb{R}^d$

$$v(x) \leq N(d)\lambda^{-1/2}\|f\|_{L_d(\mathbb{R}^d)}. \tag{4.7}$$

Then we closely follow the proof of Lemma 4.1. Take a nonnegative $\zeta \in C_0^\infty(\mathbb{R}^d)$ with unit integral, for $\varepsilon > 0$ denote $\zeta_\varepsilon(x) = \varepsilon^{-d}\zeta(\varepsilon x)$ and use