

# **MATH 60604A**

## **Statistical modelling**

### **§ 6c -Random effects**

Léo Belzile

HEC Montréal  
Department of Decision Sciences

# Introduction to random effects models

Random effects give another way of accounting for within-group correlation and allows prediction of group-level effects in addition to population-level effects.

- The main characteristic of the **linear mixed model** is to allow certain variables to have **random effects**, i.e., **to have parameters that vary from one group to another** (from one person to another in repeated measures data).
- While each group is allowed an individual effect, the overall average of these effects is zero.

# Random effects models

- When an explanatory variable is modeled with a random effect, we assume that the total effect of this variable is a combination of
  1. a common effect for the entire population and
  2. a within-group effect.
- For example, when considering repeated measures from the same individuals, the effect of a variable can be split into a common effect for all individuals in the population, and a unique effect for each individual.
- In the example of worker motivation, the effect of years of service could be split up into a common effect for all employees (in all units) and a unique effect in each unit for employees.

# Linear mixed model as hierarchical model

The linear mixed effect model is

$$\begin{aligned} \mathbf{Y}_i \mid \mathcal{B}_i = \mathbf{b}_i &\sim \text{No}_{n_i}(\mathbf{X}_i\boldsymbol{\beta} + \mathbf{Z}_i\mathbf{b}_i, \mathbf{R}_i) \\ \mathcal{B}_i &\sim \text{No}_q(\mathbf{0}_q, \boldsymbol{\Omega}) \end{aligned}$$

- The response for group  $i$ ,  $\mathbf{Y}_i$  follows a multivariate normal distribution given **random effects**  $\mathbf{b}_i$ .
- We term the coefficients  $\boldsymbol{\beta}$  associated to the model matrix  $\mathbf{X}_i$  **fixed effects**.

# Linear mixed models: fixed effects

We can write the linear mixed model as

$$[\mathbf{Y}_i \mid \mathcal{B}_i = \mathbf{b}_i] = \mathbf{X}_i\boldsymbol{\beta} + \mathbf{Z}_i\mathbf{b}_i + \varepsilon_i, \quad i = 1, \dots, m.$$

where

- $\mathbf{Y}_i = (Y_{i1}, \dots, Y_{in_i})^\top$  is the  $n_i$  vector of responses of group  $i$ .
- $\mathbf{X}_i$  is the  $n_i \times (p + 1)$  matrix of explanatory variables for group  $i$ , whose  $i$ th row is  $\mathbf{X}_{ij} = (1, X_{ij1}, \dots, X_{ijp})^\top$ .
  - The first column correspond to the intercept and all its entries are one.
  - The other columns of  $\mathbf{X}_i$  each represent an explanatory variable.
- $\boldsymbol{\beta}$  is a  $(p + 1)$  vector of **fixed** effects parameters.

# Linear mixed models: random effects

We can write the model

$$[\mathbf{Y}_i \mid \mathcal{B}_i = \mathbf{b}_i] = \mathbf{X}_i\boldsymbol{\beta} + \mathbf{Z}_i\mathbf{b}_i + \boldsymbol{\varepsilon}_i, \quad i = 1, \dots, m.$$

where

- $\mathbf{Z}_i$  is a  $n_i \times q$  matrix consisting of a subset of the columns of  $\mathbf{X}_i$ .
  - The columns of  $\mathbf{Z}_i$  are those of the variables with **random effects**.
  - If there are no random effects,  $q = 0$  and we retrieve the linear model.
- $\mathcal{B}_i = \mathbf{b}_i$  is a  $q$  vector of random effects for group  $i$
- $\boldsymbol{\varepsilon}_i$  is the  $n_i$  vector of errors of group  $i$ .

# General form: random effects models

In the linear mixed model, both  $\mathbf{B}_i$  and  $\varepsilon_i$  are random vectors and

- the random effects  $\mathbf{B}_i$  and  $\mathbf{B}_j$  ( $i \neq j$ ) are independent.
- the random effects are independent from the errors
- the  $\varepsilon_i$  are independent from one another and don't depend on the explanatory variables
- both  $\mathbf{B}_i$  and  $\varepsilon_i$  have mean zero, meaning

$$E(\mathbf{B}_i) = \mathbf{0}_{n_i}, \quad E(\varepsilon_i | \mathbf{X}_i) = \mathbf{0}_{n_i}$$

# Conditional and marginal mean and variance

We specify covariance models for the random effects and the errors,

$$\text{Cov}(\mathbf{B}_i) = \mathbf{\Omega}, \quad \text{Cov}(\varepsilon_i) = \mathbf{R}_i, \quad i = 1, \dots, m$$

The **conditional** mean and variance of  $\mathbf{Y}_i$  are

$$\text{E}(\mathbf{Y}_i \mid \mathbf{X}_i, \mathbf{B}_i = \mathbf{b}_i) = \mathbf{X}_i\boldsymbol{\beta} + \mathbf{Z}_i\mathbf{b}_i, \quad \text{Cov}(\mathbf{Y}_i \mid \mathbf{X}_i, \mathbf{B}_i = \mathbf{b}_i) = \mathbf{R}_i$$

whereas the **marginal** mean and variance of  $\mathbf{Y}_i$  are

$$\text{E}(\mathbf{Y}_i \mid \mathbf{X}_i) = \mathbf{X}_i\boldsymbol{\beta}, \quad \text{Cov}(\mathbf{Y}_i \mid \mathbf{X}_i) = \mathbf{\Sigma}_i = \mathbf{Z}_i\mathbf{\Omega}\mathbf{Z}_i^\top + \mathbf{R}_i.$$



# Random effects models

The **parameters** of the models which we will **estimate** are

- the vector of coefficients of the fixed effects,  $\beta$
- the covariance parameters of the errors and of the random effects.
- With a linear mixed model, we can estimate the conditional mean  $E(Y_{ij} \mid \mathbf{X}_i, \mathbf{b}_i)$  which can be thought of as a **prediction** of the value of  $Y_{ij}$  after accounting for the group-specific effects.
- In repeated measures data, this allows us to predict an individual's trajectory while accounting for specific effects of the individual.
- when we add a random effect for the group variable, we can still estimate effects of variables that are fixed within a group.

# Prediction

- We can predict  $\mathcal{B}_i$  by its conditional mean given  $\mathbf{Y}_i$ .
- If the covariance parameters are known,

$$E(\mathcal{B}_i \mid \mathbf{Y}_i) = \mathbf{\Omega} \mathbf{Z}_i^\top \mathbf{\Sigma}_i^{-1} (\mathbf{Y}_i - \mathbf{X}_i \hat{\boldsymbol{\beta}})$$

where

$$\mathbf{\Sigma}_i = \mathbf{Z}_i \mathbf{\Omega} \mathbf{Z}_i^\top + \mathbf{R}_i.$$

- We can plug-in covariance parameters estimates to obtain a prediction of the random effect,

$$\hat{\mathbf{b}}_i = \hat{\mathbf{\Omega}} \mathbf{Z}_i^\top \hat{\mathbf{\Sigma}}_i^{-1} (\mathbf{Y}_i - \mathbf{X}_i \hat{\boldsymbol{\beta}})$$

# Fixed or random effect?

- There is no universal definition of fixed and random effects...
- Loosely speaking, the main difference between fixed and random effects is
  - **fixed effects** for group are used when we have few groups and lots of replicates and we care about the effect of the group (small  $m$ , large  $n_i$ ).
  - **random effects** are used when there are enough levels of the factor group to estimate the variance  $\sigma_b^2$  reliably; we are not interested in the effects per say (large  $m$ , small  $n_i$ ).
- Testing whether a random effect is needed or not is equivalent to testing if the variance of the random effect  $\sigma_b^2 = 0$ ; this is a non-standard testing problem...