# Bayesian modelling

## Computational strategies and probabilistic programming

Léo Belzile

Last compiled Tuesday Oct 13, 2026

# Measuring model adequacy

Consider estimation of the leave-one predictive density under the hypothesis that the likelihood factorizes as

$$L(\boldsymbol{\theta}; \boldsymbol{y}) = \prod_{i=1}^{n} f_i(y_i; \boldsymbol{\theta});$$

where $f_i$ is the density or mass function of the $i$th observation.

Observations are then exchangeable.

# Leave-one-out cross validation

We can measure the predictive performance using the leave-one-out predictive distribution for observation $i$ given all the rest of the observations $\boldsymbol{y}_{-i}$,

$$p(y_i \mid \boldsymbol{y}_{-i}) = \int f(y_i \mid \boldsymbol{\theta}) p(\boldsymbol{\theta} \mid \boldsymbol{y}_{-i}) \mathrm{d}\boldsymbol{\theta}.$$

Naively, this requires running $n$ Markov chains with $n - 1$ observations each.

# Leave-one-out cross-validation (LOO-CV)

With weights $w_i(\boldsymbol{\theta}) = p(\boldsymbol{\theta} \mid \boldsymbol{y}_{-i})/p(\boldsymbol{\theta} \mid \boldsymbol{y}) \overset{\boldsymbol{\theta}}{\propto} 1/f_i(y_i;\boldsymbol{\theta})$ we can write the leave-one-out posterior predictive as

$$\begin{aligned} p(y_i \mid \boldsymbol{y}_{-i}) &= \int_{\boldsymbol{\Theta}} f_i(y_i;\boldsymbol{\theta}) p(\boldsymbol{\theta} \mid \boldsymbol{y}_{-i}) \mathrm{d}\boldsymbol{\theta} \\ &= \int_{\boldsymbol{\Theta}} f(y_i;\boldsymbol{\theta}) w_i(\boldsymbol{\theta}) p(\boldsymbol{\theta} \mid \boldsymbol{y}) \mathrm{d}\boldsymbol{\theta}. \end{aligned}$$

# Importance sampling for LOO-CV

This formula suggests the importance sampling estimator (eq. 25 of Section 7 of Gelfand & Dey (1994))

$$\begin{aligned}\widehat{p}(y_i \mid \boldsymbol{y}_{-i}) &= \frac{\sum_{b=1}^{B} f_i(y_i;\boldsymbol{\theta}_b) w_i(\boldsymbol{\theta}_b)}{\sum_{b=1}^{B} w_i(\boldsymbol{\theta}_b)} \\ &= B\left\{\sum_{b=1}^{B} \frac{1}{f_i(y_i;\boldsymbol{\theta}_b)}\right\}^{-1}.\end{aligned}$$

This estimator takes the form of an harmonic mean, which can have very large variance due to some weights dominating the fit. Vehtari et al. (2017) proposed using Pareto smoothed importance sampling (Vehtari et al., 2024) to avoid overly large weights; this method is implemented in the `loo` **R** package.

# Marginal predictive check

For each posterior draw and each observation, we

- generate a sample from the posterior predictive, resulting in an $n \times B$ matrix.
- Each observation $y_i$ is compared with the simulated samples $y_{i,b}^{\text{rep}}$ for posterior $\boldsymbol{\theta}_b$.
- We compute $\Pr(y_i \geq y_i^{\text{rep}})$ by Monte Carlo, by looking at the sample proportion that are more extreme.

Values close to zero or one may indicate outliers.

# Marginal predictive check with LOO-CV


Suppose we have an $n \times B$ matrix with the posterior predictive draws $y_{i,b}^{\text{rep}}$ associated with each posterior vector $\boldsymbol{\theta}_b$ and the weights $w_{i,b} = 1/f_i(y_i \mid \boldsymbol{\theta}_b)$.

We can then compute the probability integral transform with importance sampling for $i = 1, \ldots, n$:

$$\widehat{p}_i^{\text{loo}} = \Pr(y_i^{\text{rep}} \leq y_i \mid \boldsymbol{y}_{-i}) \approx \frac{\sum_{b=1}^{B} \mathrm{I}(y_i^{\text{rep}} \leq y_i) w_{i,b}}{\sum_{b=1}^{B} w_{i,b}}$$

# Probability-probability plots

The values $\widehat{p}_i^{\text{loo}}$ should be approximately uniform if the model was perfectly calibrated.
We build a P-P plot (ordered values against uniform plotting position) to check this.



# Sources of poor mixing

Slow mixing can be due to the following:

- poor proposals
- strong correlation between posterior parameters
- overparametrization and lack of identifiability

# Computational strategies

These problems can be addressed using one of the following:

- marginalization
- removing redundant parameters or using informative priors
- reparametrization
- clever proposals (adaptive MCMC), see Andrieu & Thoms (2008) and Rosenthal (2011).
- blocking

# Gaussian model with random effects

Consider a hierarchical Gaussian model of the form

$$\boldsymbol{Y} = \mathbf{X}\boldsymbol{\beta} + \mathbf{Z}\boldsymbol{B} + \boldsymbol{\varepsilon}$$

where

- $\mathbf{X}$ is an $n \times p$ design matrix with centered inputs,
- $\boldsymbol{\beta} \sim \mathsf{Gauss}(\mathbf{0}_p, \sigma^2\mathbf{I}_p)$,
- $\boldsymbol{B} \sim \mathsf{Gauss}_q(\mathbf{0}_q, \boldsymbol{\Omega})$ are random effects and
- $\boldsymbol{\varepsilon} \sim \mathsf{Gauss}_n(\mathbf{0}_n, \kappa^2\mathbf{I}_n)$ are independent white noise.

# One-way ANOVA

Consider a one-way ANOVA with $n$ observations from $K$ categories, with observation $i = 1, \ldots, n_k$ from group $k$ having contribution

$$Y_{i,k} \sim \textsf{Gauss}(\mu + \alpha_k, \sigma_y^2)$$
$$\alpha_k \sim \textsf{Gauss}(0, \sigma_\alpha^2).$$

We set an improper prior for the mean $p(\mu) \propto 1$.

If there are $K$ groups, then the group-specific mean is $\mu + \alpha_k$, and there is a redundant parameter. This leads to poor mixing if priors are not informative.

# Removing redundant parameters

We can enforce a sum-to-zero constraint for $\sum_{k=1}^{K} \alpha_k = 0$ and sample $K-1$ parameters for the difference to the global mean.

This yields a degenerate multivariate Gaussian distribution for the vector $\boldsymbol{\alpha} \sim \textsf{Gauss}_K(\mathbf{0}_K, \boldsymbol{\Omega})$ where

$$\boldsymbol{\Omega} = \sigma_\alpha^2 \left(\mathbf{I}_K - \mathbf{1}_K \mathbf{1}_K^\top / K\right).$$

# Marginalization

Given a model $p(\boldsymbol{\theta}, \boldsymbol{Z})$, reduce the dependance by sampling from the marginal

$$p(\boldsymbol{\theta}) = \int p(\boldsymbol{\theta}, \boldsymbol{z}) \mathrm{d}\boldsymbol{z}.$$

This happens for data augmentation, etc., and reduces dependency between parameters, but typically the likelihood becomes more expensive to compute

# Marginalization

We can marginalize over the $K$ vector of $\boldsymbol{\alpha}$ to perform updates of $\mu \mid \sigma_y^2, \sigma_\alpha^2, \boldsymbol{y}$.

We find if $\boldsymbol{\Sigma}_{\sigma_y^2,\sigma_\alpha^2}$ is the marginal covariance matrix of

$$\boldsymbol{Y} \mid \mu, \sigma_y^2, \sigma_\alpha^2 \sim \text{Gauss}_n\left(\mu \mathbf{1}_n, \boldsymbol{\Sigma}\right).$$

Straightforward calculations yield

$$\mu \mid \sigma_y^2, \sigma_\alpha^2, \boldsymbol{y} \sim \text{Gauss}\left(\frac{\boldsymbol{y}^\top \boldsymbol{\Sigma}^{-1} \mathbf{1}_n}{\mathbf{1}_n^\top \boldsymbol{\Sigma}^{-1} \mathbf{1}_n}, \left\{\mathbf{1}_n^\top \boldsymbol{\Sigma}^{-1} \mathbf{1}_n\right\}^{-1}\right).$$

# Marginalization of Gaussian models

We can write

$$\begin{aligned}\boldsymbol{Y} \mid \boldsymbol{\beta}, \boldsymbol{B}, \sigma^2 &\sim \mathsf{Gauss}_n(\mathbf{X}\boldsymbol{\beta} + \mathbf{Z}\boldsymbol{B}, \sigma^2\mathbf{I}_p) \\ \boldsymbol{Y} \mid \boldsymbol{\beta} &\sim \mathsf{Gauss}_n(\mathbf{X}\boldsymbol{\beta}, \mathbf{Q}^{-1}),\end{aligned}$$

where the second line corresponds to marginalizing out the random effects $\boldsymbol{B}$.

# Efficient calculations for Gaussian models

If, as is often the case, $\boldsymbol{\Omega}^{-1}$ and $\mathbf{Z}$ are sparse matrices, the full precision matrix can be efficiently computed using Shermann–Morisson–Woodbury identity as

$$\begin{aligned}\mathbf{Q}^{-1} &= \mathbf{Z}\boldsymbol{\Omega}^{-1}\mathbf{Z}^\top + \kappa^2\mathbf{I}_n, \\ \kappa^2\mathbf{Q} &= \mathbf{I}_n - \mathbf{Z}\boldsymbol{G}^{-1}\mathbf{Z}^\top, \\ \boldsymbol{G} &= \mathbf{Z}^\top\mathbf{Z} + \kappa^2\boldsymbol{\Omega}^{-1}\end{aligned}$$

Section 3.1 of Nychka et al. (2015) details efficient ways of calculating the quadratic form involving $\mathbf{Q}$ and it's determinant.

# Parameter expansion

Add redundant parameter to improve mixing by decorrelating (Liu et al., 1998)

$$Y_{i,k} \sim \textsf{Gauss}(\mu + \xi\eta_k, \sigma_y^2)$$
$$\eta_k \sim \textsf{Gauss}(0, \sigma_\eta^2)$$

so that $\sigma_\alpha = |\xi|\sigma_\eta$.

# Eight school example

Consider the eight school data from Educational Testing Service (ETS) (Gelman et al., 2013). The goal of the experiment was to estimate the effect of coaching on SAT-V scores.

The data consist of estimated treatment mean difference $y_1, \ldots, y_8$ and their standard errors $s_1, \ldots, s_8$ from eight different schools.

Simple sample sizes are large, the average treatment effect (ATE) can be considered approximately Gaussian. We treat the school-specific standard errors $s_j$ as a known quantity.

# Gibbs sampler for eight school problem

The naive Gibbs sampler, which iterates between conditionals of $p(\boldsymbol{\alpha} \mid \boldsymbol{y}, \boldsymbol{s}^2, \mu, \sigma_\alpha^2)$ and $p(\mu \mid \boldsymbol{y}, \boldsymbol{s}^2, \boldsymbol{\alpha})$ leads to strong autocorrelation.

A better strategy is to marginalize over the global mean $\mu$ and simulate from $p(\boldsymbol{\alpha} \mid \boldsymbol{y}, \boldsymbol{s}^2, \sigma_\alpha^2)$ instead. We get then the conditionals

$$\alpha_i \mid y_i, s_i^2, \sigma_\alpha^2 \sim \text{Gauss}\left\{\left(s_i^{-2} + \sigma_\alpha^{-2}\right)^{-1} \frac{y_i}{s_j^2}, \left(s_i^{-2} + \sigma_\alpha^{-2}\right)^{-1}\right\}$$

$$\mu \mid \boldsymbol{y}, \boldsymbol{s}^2, \boldsymbol{\alpha} \sim \text{Gauss}\left(s_\mu^2 \sum_{j=1}^{8} \frac{y_i - \alpha_i}{s_j^2}, s_\mu^2\right)$$

$$\sigma_\alpha^2 \mid \boldsymbol{\alpha} \sim \text{inv. gamma}\left(\frac{n-1}{2}, \frac{\sum_{j=1}^{8} \alpha_j^2}{2}\right)$$

where $s_\mu^{-2} = \sum_{j=1}^{8} s_j^{-2}$.


# Parameter expansion for the eight school problem

The parameter expansion sampler proceeds in exactly the same way with the same conditionals for $\sigma_\xi^2$ and $\boldsymbol{\xi}$ as above; the only additional step is

$$\nu \mid \boldsymbol{y}, \boldsymbol{s}^2, \boldsymbol{\xi} \sim \text{Gauss}\left\{\left(\sum_{j=1}^8 \xi_j^2/s_j^2\right)^{-1} \sum_{j=1}^8 \xi_j(y_j-\mu)/s_j^2, \left(\sum_{j=1}^8 \xi_j^2/s_j^2\right)^{-1}\right\}.$$

After each update, we then save $\alpha_j = \nu\xi_j$ and $\sigma_\alpha = |\nu|\sigma_\xi$, as the other parameters are not identifiable.

# Funnel geometry



# Blocking

Identify groups of strongly correlated parameters and propose a joint update for these.

- The more parameters we propose at the same time, the lower the chance of acceptance
- Often ways to sample these efficiently

# Tokyo rainfall

We consider data from Kitagawa (1987) that provide a binomial time series giving the number of days in years 1983 and 1984 (a leap year) in which there was more than 1mm of rain in Tokyo; see section 4.3.4 of Rue & Held (2005).

We have

- $T = 366$ days and
- $n_t \in \{1, 2\}$ $(t = 1, \ldots, T)$observations in day $t$ and
- $y_t = \{0, \ldots, n_t\}$ the number of days with rain.

# Smoothing probabilities

The objective is to obtain a smoothed probability of rain. The underlying probit model considered takes

$$Y_t \mid n_t, p_t \sim \textsf{binom}(n_t, p_t), \qquad p_t = \Phi(\beta_t).$$

We specify the random effects $\boldsymbol{\beta} \sim \textsf{Gauss}_T(\mathbf{0}, \tau^{-1}\mathbf{Q}^{-1})$, where $\mathbf{Q}$ is a $T \times T$ precision matrix that encodes the local dependence.

# Random walk prior

A circular random walk structure of order 2 is used to model the smooth curves by smoothing over neighbors, and enforces small second derivative. This is a suitable prior because it enforces no constraint on the mean structure.

This amounts to specifying the process with for $t \in \mathbb{N} \mod 366 + 1$

$$\begin{aligned}\Delta^2 \beta_t &= (\beta_{t+1} - \beta_t) - (\beta_t - \beta_{t-1}) \\ &= -\beta_{t-1} + 2\beta_t - \beta_{t+1} \sim \textsf{Gauss}(0, \tau^{-1}).\end{aligned}$$

# Circulant precision matrix

This yields an intrinsic Gaussian Markov random field with a circulant precision matrix $\tau\mathbf{Q}$ of rank $T-1$, where

$$\mathbf{Q} = \begin{pmatrix} 6 & -4 & 1 & 0 & \cdots & 1 & -4 \\ -4 & 6 & -4 & 1 & \ddots & 0 & 1 \\ 1 & -4 & 6 & -4 & \ddots & 0 & 0 \\ \vdots & \ddots & \ddots & \ddots & \ddots & \ddots & \vdots \\ -4 & 1 & 0 & 0 & \cdots & -4 & 6 \end{pmatrix}.$$

Because of the linear dependency, the determinant of $\mathbf{Q}$ is zero.

# Prior draws





Figure 1: Five realizations from the cyclical random walk Gaussian prior of order 2.

# Gibbs sampling for Tokyo data

We can perform data augmentation by imputing Gaussian variables, say $\{z_{t,i}\}$ from truncated Gaussian, where $z_{t,i} = \beta_t + \varepsilon_{t,i}$ and $\varepsilon_{t,i} \sim \textsf{Gauss}(0,1)$ are independent standard Gaussian and

$$z_{t,i} \mid y_{t,i}, \beta_t \sim \begin{cases} \textsf{trunc. Gauss}(\beta_t, 1, -\infty, 0) & y_{t,i} = 0 \\ \textsf{trunc. Gauss}(\beta_t, 1, 0, \infty) & y_{t,i} = 1 \end{cases}$$

# Posterior for Tokyo data

The posterior is proportional to

$$p(\boldsymbol{\beta} \mid \tau) p(\tau) \prod_{t=1}^{T} \prod_{i=1}^{n_t} p(y_{t,i} \mid z_{t,i}) p(z_{t,i} \mid \beta_t)$$

# Data augmentation for Tokyo data

Once we have imputed the Gaussian latent vectors, we can work directly with the values of $z_t = \sum_{i=1}^{n_t} z_{i,t}$

$$
\begin{aligned}
p(\boldsymbol{\beta}, \tau) \propto \tau^{(T-1)/2} \exp\left(-\frac{\tau}{2}\boldsymbol{\beta}^\top \mathbf{Q}\boldsymbol{\beta}\right) \\
\times \exp\left\{-\frac{1}{2}(\boldsymbol{z}/\boldsymbol{n} - \boldsymbol{\beta})^\top \mathrm{diag}(\boldsymbol{n})(\boldsymbol{z}/\boldsymbol{n} - \boldsymbol{\beta})\right\} \\
\times \tau^{a-1}\exp(-\tau b)
\end{aligned}
$$

where $\boldsymbol{z} = (z_1, \ldots, z_T)$, as $n_t^{-1} z_t \sim \textsf{Gauss}(\beta_t, n_t^{-1})$.

# Gibbs for Tokyo data - conditionals


Completing the quadratic form shows that

$$\boldsymbol{\beta} \mid \boldsymbol{z}, \tau \sim \mathsf{Gauss}_T\left(\mathbf{Q}^{\star-1}\boldsymbol{z}, \mathbf{Q}^{\star-1}\right)$$

$$\tau \mid \boldsymbol{\beta} \sim \mathsf{gamma}\left(\frac{T-1}{2} + a, \frac{\boldsymbol{\beta}^\top \mathbf{Q} \boldsymbol{\beta}}{2} + b\right),$$

$$\mathbf{Q}^\star = \{\tau \mathbf{Q} + \mathrm{diag}(\boldsymbol{n})\}$$

# Posterior prediction for probability of rainfall





Posterior prediction for probability of rainfall

# Blocking vs joint update

Compare the following two stategies


- joint update: given $\boldsymbol{z}$ and $\tau$, simulate $\boldsymbol{\beta}$ jointly
- one-parameter at a time: starting from $i \sim \textsf{unif}(\{1, \ldots, 366\})$, get index $t = i \mod 366 + 1$ and simulate $\beta_i \mid \boldsymbol{\beta}_{-i}, \boldsymbol{z}, \tau$ one at a time.

# Blocking strategy




Trace plots for Tokyo with blocking of random effects

# Individual update strategy




Trace plots for Tokyo with random scan Gibbs for random effects

# Lessons from the Tokyo example

What happened?

- there is lower autocorrelation with the joint update (also faster here!) for the $\boldsymbol{\beta}$
- in both cases, $\tau \mid \cdot$ mixes poorly because the values of $\boldsymbol{\beta}$ were sampled conditional on the previous value.

A better avenue would be to use a Metropolis random walk for $\tau^\star$, simulate $\boldsymbol{\beta} \mid \tau^\star$ and propose the joint vector $(\tau^\star, \boldsymbol{\beta}^\star)$ simultaneously.

# One step further

We could also remove the data augmentation step and propose from a Gaussian approximation of the log likelihood, using a Taylor series expansion of the log likelihood about $\boldsymbol{\beta}_{t-1}$

$$\log p(\boldsymbol{\beta} \mid \boldsymbol{y}) \overset{\boldsymbol{\beta}}{\propto} -\frac{\tau}{2}\boldsymbol{\beta}^\top \mathbf{Q}\boldsymbol{\beta} + \sum_{t=1}^{T} \log f(y_t \mid \beta_t)$$

and the $y_t$ are conditionally independent in the likelihood. Refer to Section 4.4.1 of Rue & Held (2005) for more details.

# Technical aside: in sparsity we trust!

It is crucial to exploit the sparsity structure of $\mathbf{Q}$ for efficient calculations of the likelihood

- typically requires re-ordering elements to get a banded precision matrix
- precompute the sparse Cholesky
- compute inverse by solving systems of linear equations; there are dedicated algorithms

# Probabilistic programming


There are several languages and interfaces that implement probabilistic programming where the user has only to specify the likelihood and prior.

Historically, Bugs paved the way to practitioners.

It relies on Gibbs sampling (updating one parameter at the time), but is not actively developed. Still the source of many exercises and inspiration for the syntax of other implementations (e.g., Nimble, JAGS).

# Stan

The programming language Stan is written in C++ and offers cross-platform interfaces.



# Other MCMC software

- Nimble, C++ with **R** interface
- Turing.jl, Julia
- PyMC, Python
- Pigeons, Julia

# Stochastic volatility model

Financial returns $Y_t$ typically exhibit time-varying variability. The **stochastic volatility** model is a parameter-driven model that specifies

$$Y_t = \exp(h_t/2)Z_t$$
$$h_t = \gamma + \phi(h_{t-1} - \gamma) + \sigma U_t$$

where $U_t \overset{\text{iid}}{\sim} \textsf{Gauss}(0, 1)$ and $Z_t \overset{\text{iid}}{\sim} \textsf{Gauss}(0, 1)$.

It is possible to introduce leverage by adding $\textsf{Cor}(Z_t, U_t) = \rho$.

# Summary

In high dimensions, the typical set over which the posterior concentrate can have weird geometry.

Different strategies can improve sampling

- data augmentation can simplify the likelihood, but lead to slow mixing
- parameter expansion adds redundancy (but the parameters are not identifiable)
- marginalization reduces correlation, but the computational cost is higher.

# References


Andrieu, C., & Thoms, J. (2008). A tutorial on adaptive MCMC. *Statistics and Computing*, *18*(4), 343–373. https://doi.org/10.1007/s11222-008-9110-y

Gelfand, A. E., & Dey, D. K. (1994). Bayesian model choice: Asymptotics and exact calculations. *Journal of the Royal Statistical Society: Series B (Methodological)*, *56*(3), 501–514. https://doi.org/10.1111/j.2517-6161.1994.tb01996.x

Gelman, A., Carlin, J. B., Stern, H. S., Dunson, D. B., Vehtari, A., & Rubin, D. B. (2013). *Bayesian data analysis* (3rd ed.). Chapman; Hall/CRC. https://doi.org/10.1201/b16018

Kitagawa, G. (1987). Non-Gaussian state–space modeling of nonstationary time series. *Journal of the American Statistical Association*, *82*(400), 1032–1041. https://doi.org/10.1080/01621459.1987.10478534

Liu, C., Rubin, D. B., & Wu, Y. N. (1998). Parameter expansion to accelerate EM: The PX-EM algorithm. *Biometrika*, *85*(4), 755–770. https://doi.org/10.1093/biomet/85.4.755

Nychka, D., Bandyopadhyay, S., Hammerling, D., Lindgren, F., & Sain, S. (2015). A multiresolution Gaussian process model for the analysis of large spatial datasets. *Journal of Computational and Graphical Statistics*, *24*(2), 579–599.

Rosenthal, J. (2011). Optimal proposal distributions and adaptive MCMC. In S. Brooks, A. Gelman, G. Jones, & X. L. Meng (Eds.), *Handbook of Markov chain Monte Carlo* (pp. 93–112). CRC Press. https://doi.org/10.1201/b10905-5

Rue, H., & Held, L. (2005). *Gaussian Markov random fields: Theory and applications* (p. 280). CRC Press.

Vehtari, A., Gelman, A., & Gabry, J. (2017). Practical Bayesian model evaluation using leave-one-out cross-validation and WAIC. *Statistics and Computing*, *27*(5), 1413–1432. https://doi.org/10.1007/s11222-016-9696-4

Vehtari, A., Simpson, D., Gelman, A., Yao, Y., & Gabry, J. (2024). Pareto smoothed importance sampling. *Journal of Machine Learning Research*, *25*(1), 3491–3548.