

**Vector boson fusion Higgs boson
fiducial and differential cross-sections
measured in the $WW^* \rightarrow \ell\ell\nu\nu$ decay
channel with the ATLAS detector**

A Dissertation

Presented to

The Faculty of the Graduate School of Arts and Sciences

Brandeis University

Department of Physics

Professor Gabriella Sciolla, Advisor

In Partial Fulfillment
of the Requirements for the Degree
Doctor of Philosophy

by

Laura Bergsten

July, 2020

This dissertation, directed and approved by Laura Bergsten's committee, has been accepted and approved by the Graduate Faculty of Brandeis University in partial fulfillment of the requirements for the degree of:

DOCTOR OF PHILOSOPHY

Eric Chasalow, Dean of Arts and Sciences

Dissertation Committee:

Professor Gabriella Sciolla, Chair

Craig Blocker

Alessandro Tricoli

©Copyright by

Laura Bergsten

2020

For Nick

Acknowledgments

Thank all the people

Abstract

**Vector boson fusion Higgs boson fiducial and differential
cross-sections measured in the $WW^* \rightarrow \ell\ell\nu\nu$ decay channel with
the ATLAS detector**

A dissertation presented to the Faculty of
the Graduate School of Arts and Sciences of
Brandeis University, Waltham, Massachusetts

by Laura Bergsten

Contents

Abstract	vii
1 Introduction	1
2 Theory	2
2.1 Standard Model	2
2.2 LHC Physics/Phenomenology	17
2.3 Brief history of SM tests	25
2.4 Measurement motivation	30
3 The LHC and the ATLAS detector	33
3.1 The Large Hadron Collider	33
3.2 A Toroidal LHC Apparatus	36
3.3 The High-Luminosity LHC and Inner Tracker (ITk)	38
4 Tracking and Isolation in ATLAS	50
4.1 Jets	51
4.2 Missing transverse energy	56
4.3 Electrons	60
4.4 Muons	64
5 Event Selection	76
5.1 Data and Monte Carlo samples	76
5.2 Object definitions	79
5.3 Common Observables	84
5.4 Event selection	86
6 Backgrounds and Systematics	98
6.1 Backgrounds	98
6.2 Systematic uncertainties	133

CONTENTS

7	Results	139
7.1	Statistical analysis	139
7.2	Unfolding	141
7.3	Results and future measurements	141
8	Conclusions	142

Chapter 1

Introduction

Introduction to the thesis. Explain what u'll see.

Chapter 2

Theory

The measurement at the heart of this thesis can only be understood within the context of a vast amount of preceding theoretical and experimental work. I have tried to condense and summarize key concepts that will motivate the central measurement's strategy and results. This chapter begins with a brief summary of the Standard Model itself, first describing fundamental particles and their forces before delving into a succinct mathematical formulation. Next, the chapter discusses history of the Standard Model and crucial tests of the theory up until current work at the LHC. The next section outlines some of the recent and current physics analyses at the Large Hadron Collider (LHC) with a focus on Higgs boson measurements. Finally, I'll introduce my thesis' main focus, fiducial and differential cross-section measurements of Vector-Boson-Fusion (VBF) Higgs decaying into two W-bosons.

2.1 Standard Model

The Standard Model is one of the most successful scientific theories to date. Its predictions encompass all of the visible universe and continue to undergo careful testing. The

CHAPTER 2. THEORY

SM combines three forces- electromagnetic, weak, and strong - but it is not complete. One known force, gravity, is not included in the Standard Model and further questions, like an explanation for dark matter and dark energy, remain.

2.1.1 Particles and forces

The particles we define in high energy physics are the most minute portions of observable matter. They are generally considered point-like, have no internal structure, and cannot be further split. Each particle we can define has a unique set of quantum numbers and its own anti-particle (with the same mass and spin, but opposite electrical charge and quantum numbers).

Particles can be sorted into a few distinct groups - first bosons, with integer spin, and fermions, with half-integer spin. Bosons are ‘force carriers’ meaning they are exchanged any time particles interact. Fermions are at the heart of all conventional matter. They can be split further into two categories- leptons and quarks. Quarks have fractional integer charge and interact strongly while leptons have integer charge and interact solely through the weak or electromagnetic forces. Both quarks and leptons are made of three generations of particles, each heavier and more unstable than the next. Charts showing quark/lepton families and their key quantum numbers are shown below in Figure 2.1. Each generation of quarks and leptons contains a particle doublet. Each lepton doublet contains a charged lepton and a neutrino while each quark doublet contains one $+2/3$ charged particle and one with a $-1/3$ charge. Each lepton and quark also has an anti-particle. All conventional, stable matter is made from the first generation of quarks and leptons.

There are four gauge bosons and one scalar boson predicted through the Standard Model. These correspond to three fundamental forces in nature (the fourth, gravity, is so small on the

CHAPTER 2. THEORY

scale of particle interactions as to not be considered). The strongest force on the subatomic scale is the strong force- this is mediated by the gluon- and works primarily to bind quarks together to form composite particles like protons or neutrons. The electromagnetic force is about $60\times$ weaker than the strong force and is mediated by the photon. This force accounts for all electric interactions like that between an electron and an atomic nucleus. Finally, the weak force ($10^4\times$ weaker than the EM) facilitates β -decay and is mediated by massive W and Z bosons. The final boson predicted by the Standard Model is the Higgs boson. The only scalar boson, it has no charge or intrinsic spin. The Higgs gives mass to all other particles through Spontaneous Symmetry Breaking, which will be described in later sections.

Standard Model of Elementary Particles




Figure 2.1: Three generations of quarks and leptons are shown along with all SM bosons [1]

CHAPTER 2. THEORY

Photons are massless, spin-1 particles and mediate all electromagnetic interactions. They couple directly to any particle with electric charge- so quarks, leptons, and W/Z bosons but not neutrinos. Since the photon is massless, the electromagnetic force can operate on infinitely long scales but its force decreases with $1/r^2$.

Gluons are massless particles with no charge and spin-1. They couple to color charges, which are a property only of quarks and gluons. Each quark has one of three colors (RGB) while anti-quarks have "anti" versions of these. Colors are conserved 'charges' similar to electric charge. Quarks are never found alone as they couple so strongly to one another as to be confined in groups of two or three. These groups are "color-confined" meaning the quarks contain colors which add up to a color neutral sum. For instance, a two quark meson $u\bar{u}$ may have colors R and anti-R while a three quark hadron uud (proton) may have colors R, G, and B. Gluons differ from photons in that they are not neutral in the charge they couple to. Gluons have two colors (8 total combinations) and can thus couple to each other. This makes the strong force distinct from the electromagnetic and has implications for long-distance interactions.

W and Z bosons, unlike gluons and photons, are massive. However, like their other gauge boson counterparts, they have spin-1 and mediate a charge (weak). W^\pm mediates charged-current interactions which can violate flavor conservation between quarks and/or leptons and their neutrinos. Z^0 mediates neutral-current interactions which conserve flavor. W^\pm bosons contain electric charge so can interact through EM as well. In addition, W and Z bosons contain weak charge (as do all fermions) so can self-couple as well as couple with all fermions.

The Higgs boson will be further motivated and described in later sections but suffice to say it's a massive spin-0 particle which couples to all particles with mass (including itself). It doesn't mediate any force but is still an integral part of the Standard Model.

CHAPTER 2. THEORY

2.1.2 Gauge Invariance

According to Noether's theorem, for every continuous transformation of a field that leaves the Lagrangian invariant, there is a conserved current. In other words, symmetries found in physical theories lead to conservation laws (and vice-versa). The Standard Model is a gauge theory built on symmetries; all interactions between particles result from requiring the theory to be invariant under local gauge transformations. Each part of the Standard Model- from quantum electrodynamics (QED) to quantum chromodynamics (QCD) - is a gauge theory on its own. Each part has gauge invariance symmetries. In this section I'll step through the basic mathematic formalism for QED, QCD, and the combined electro-weak theory to illustrate the physical ramifications of gauge invariance and set the stage for the Higgs mechanism. The following sections are written with guidance from text [2].

Quantum Electrodynamics

Quantum electrodynamics (QED) is the first, and simplest, physical gauge theory, describing how light and matter interact even under relativistic conditions. The theory produces extremely good agreement with experiment due to the success of perturbative QED calculations. Entire textbooks are dedicated to QED formalism and predictions. Here I will highlight only the effects of its local gauge invariance symmetry and generating the full QED Lagrangian beginning with the Dirac Lagrangian of a free fermion.

First, the Dirac Lagrangian describes a free fermion of mass m

$$\mathcal{L} = i\psi\gamma^\mu\partial_\mu\psi - m\bar{\psi}\psi, \quad (2.1)$$

where ψ is a Dirac spinor and γ^μ represent the Dirac matrices. To demonstrate local gauge

CHAPTER 2. THEORY

invariance we need to transform

$$\psi(x) \rightarrow e^{i\alpha(x)}\psi(x) \quad (2.2)$$

where $\alpha(x)$ depends on space and time arbitrarily. Directly substituting this into our Lagrangian shows that \mathcal{L} is not invariant, and the ∂_μ term breaks this

$$\partial_\mu\psi \rightarrow e^{i\alpha(x)}\partial_\mu\psi + ie^{i\alpha(x)}\partial_\mu\alpha \quad (2.3)$$

In order to mandate the theory is invariant we need to change this term to the “covariant derivative” D_μ which transforms

$$D_\mu\psi \rightarrow e^{i\alpha(x)}D_\mu\psi. \quad (2.4)$$

The “covariant derivative” must contain a vector field A_μ and this field must transform so as to cancel with the unwanted part of the transformed D_μ in order to transform as required by gauge invariance.

$$D_\mu \equiv \partial_\mu - ieA_\mu \quad (2.5)$$

where

$$A_\mu \rightarrow A_\mu + \frac{1}{e}\partial_\mu\alpha \quad (2.6)$$

Now the original Dirac equation is replaced with the following:

$$\mathcal{L} = \bar{\psi}(i\gamma^\mu\partial_\mu - m)\psi + e\bar{\psi}\gamma^\mu\psi A_\mu. \quad (2.7)$$

By requiring local gauge invariance we've introduced a gauge field A_μ which couples to the Dirac particle just as the photon. In fact, if we take this as the photon gauge field and so add a kinetic energy term (which is also local gauge invariant!) we find the Lagrangian of

CHAPTER 2. THEORY

QED.

$$\mathcal{L} = \bar{\psi}(i\gamma^\mu \partial_\mu - m)\psi + e\bar{\psi}\gamma^\mu\psi A_\mu - \frac{1}{4}F_{\mu\nu}F^{\mu\nu} \quad (2.8)$$

One can also see that adding a mass term to the Lagrangian for the new field ($\frac{1}{2}m^2 A_\mu A^\mu$) would break gauge invariance, indicating the photon must be massless. From the free fermion Lagrangian, imposing local gauge invariance leads to the full interacting field theory of QED. This is not a curiosity but an essential component of the theory, and the use of local gauge symmetry in deriving particle interactions does not end here.

Quantum chromodynamics

Quantum chromodynamics differs from QED in a few crucial ways. First, since quark color fields exist the QED $U(1)$ gauge group is replaced with $SU(3)$ and the free Lagrangian contains indices j to denote the three color fields:

$$\mathcal{L} = \bar{q}_j(i\gamma^\mu \partial_\mu - m)q_j. \quad (2.9)$$

QCD also carries three quark flavors, which will be ignored here for simplicity. The QCD group is non-Abelian since not all generators of the group commute with each other. These generators will be defined as T_a where $a = 1, \dots, 8$ and are linearly independent traceless 3×3 matrices (the Gell-Mann matrices λ_a are conventional). The local color phase transformation required is thus

$$q(x) \rightarrow e^{i\alpha_a(x)T_a}q(x) \quad (2.10)$$

We can consider an infinitesimal phase transformation as

$$\begin{aligned} q(x) &\rightarrow [1 + i\alpha_a(x)T_a]q(x), \\ \partial_\mu q &\rightarrow (1 + i\alpha_a T_a)\partial_\mu q + iT_a q \partial_\mu \alpha_a. \end{aligned} \quad (2.11)$$

CHAPTER 2. THEORY

Just as in the QED example, the last line breaks the invariance of \mathcal{L} and we can proceed similarly by introducing a new gauge field (or in this case eight) called G_μ^a which transform

$$G_\mu^a \rightarrow G_\mu^a - \frac{1}{g} \partial_\mu \alpha_a - f_{abc} \alpha_b G_\mu^c \quad (2.12)$$

The last term added here is to cope with the non-Abelian nature of QCD (that not all the generators T_a commute with each other). Just as in QED this invariance forms a covariant derivative:

$$D_\mu = \partial_\mu u + ig T_a G_\mu^a \quad (2.13)$$

Replacing the derivative in our Lagrangian and adding a gauge invariant energy term for each of the G_μ^a fields ($\frac{1}{4} G_{\mu\nu}^a G_{\mu\nu}^a$) yields the final gauge invariant QCD Lagrangian:

$$\mathcal{L} = \bar{q}(i\gamma^\mu \partial_\mu - m)q - g(\bar{q}\gamma^\mu T_a q)G_\mu^a - \frac{1}{4} G_{\mu\nu}^a G_{\mu\nu}^a. \quad (2.14)$$

Just as in the QED case, imposing local color phase invariance produced a new interacting field (actually eight) with a coupling specified as g . These are the gluon fields and just like photons, local gauge invariance requires them to be massless. Unlike the QED case, this Lagrangian's new kinetic term includes self-interaction between the gauge bosons - another key feature of QCD that is mandated by local color phase invariance. Gluons themselves must carry color charge and so self-couple - the structure of these self coupling terms and their single coupling strength g are uniquely determined by gauge invariance.

Electroweak unification

Thus far, I've summarized the theoretical backgrounds for symmetries (and so conserved quantities) in both quantum electrodynamics and chromodynamics. The weak force is the

CHAPTER 2. THEORY

final Standard Model force and weak interactions are mediated by Z and W bosons. Unlike the gluons and photons of QCD and QED, these gauge bosons are massive. This is explained through spontaneous symmetry breaking of the electroweak force, which is outlined in the following section. Assuming that W/Z bosons are massive, the weak force can be combined with QED and a central electroweak force (with its associated symmetries) can be described.

The weak neutral current J_μ^{NC} and weak charged currents (J_μ , J_μ^\dagger) can form a symmetry group for weak interactions. The charged currents correspond to the charged weak interaction through W^\pm bosons while the neutral current is associated with the Z^0 boson.

$$\begin{aligned} J_\mu &= \bar{\nu}_L \gamma_\mu \nu_L, \\ J_\mu^\dagger &= \bar{e}_L \gamma_\mu \nu_L J_\mu^3 = \frac{1}{2} \bar{\nu}_L \gamma_\mu \nu_L - \frac{1}{2} \bar{e}_L \gamma_\mu e_L \end{aligned} \quad (2.15)$$

L denotes that these are left-handed spinors and particle names denote associated Dirac spinors. The charged currents can be written as a doublet using the Pauli spin matrices τ_i where $\tau_\pm = \frac{1}{2}(\tau_1 \pm i\tau_2)$ and

$$\chi_L = \begin{bmatrix} \nu \\ e^- \end{bmatrix} \quad (2.16)$$

as

$$\begin{aligned} J_\mu^+(x) &= \bar{\chi}_L \gamma_\mu \tau_+ \chi_L, \\ J_\mu^-(x) &= \bar{\chi}_L \gamma_\mu \tau_- \chi_L, \\ J_\mu^3(x) &= \bar{\chi}_L \gamma_\mu \frac{1}{2} \tau_i \chi_L \text{ with } i = 1, 2, 3 \end{aligned} \quad (2.17)$$

The corresponding charge $T^i = \int J_0^i(x) d^3x$ can be introduced so we have an $SU(2)_L$ algebra

$$[T^i, T^j] = i\epsilon_{ijk} T^k \quad (2.18)$$

Unfortunately while these currents create an $SU(2)$ group, they don't correspond with the

CHAPTER 2. THEORY

weak neutral current symmetry in an obvious way. Unlike the charged currents, the neutral current has a right handed component. One clear way to resolve this is to add in the electromagnetic current, as its a neutral current with left and right-handed components.

$$j_\mu^{em}(x) = -e_R \bar{\gamma}_\mu e_R - e_L \bar{\gamma}_\mu e_L \quad (2.19)$$

so the the electromagnetic current j_μ can be written using the coupling e

$$j_\mu = ej_\mu^{em} = e\bar{\psi}\gamma_\mu Q\psi \quad (2.20)$$

with Q the charge operator and generator of the $U(1)$ symmetry group of EM. In order to “save” the symmetry of the weak neutral current, we can define an electromagnetic current j_μ^Y , or the weak hypercharge current, that is unchanged by $SU(2)_L$ transformations. We define a weak hypercharge Y and its current j_μ^Y

$$\begin{aligned} Q &= T^3 + \frac{Y}{2}, \\ j_\mu^Y &= \bar{\psi}\gamma_\mu Y\psi \end{aligned} \quad (2.21)$$

The combined current

$$j_\mu^{em} = J_\mu^3 + \frac{1}{2}j_\mu^Y \quad (2.22)$$

now generates the symmetry group $U(1)_\gamma$ and so the electromagnetic interaction and weak interaction are combined into one $SU(2)_L \times U(1)_\gamma$ group. While unified in this way, the two forces still have independent coupling strengths. This brief introduction into electroweak unification is not the complete picture- EM and weak *interactions* still have to be unified. This is simple in the Standard model framework- electroweak currents just have to be coupled to vector bosons. In the electroweak $SU(2)_L \times U(1)_\gamma$ group there is an isotriplet of vector

CHAPTER 2. THEORY

fields W_μ^i coupled with strength g to the weak isospin current J_μ^i while a single vector field B_μ is coupled to the weak hypercharge current j_μ^Y with strength $g'/2$. The electroweak interaction can thus be defined:

$$-ig(J^i)^\mu W_\mu^i - i\frac{g'}{2}(j^Y)^\mu B_\mu. \quad (2.23)$$

This summary of the unified electroweak force will be the starting point for a derivation of the Higgs boson and an explanation for mass of the weak force's vector bosons (and all the fermions). The electroweak theory is unique in its calculability even at higher order scales. Because of this, theoretical uncertainties are relatively low and many deviations from theory could potentially be observed at current energy scales. The measurement central to my thesis probes for such discrepancies in electroweak theory. The mechanisms for this will be explained in the last section in this chapter.

Spontaneous Symmetry Breaking

Unlike QED and QCD, the weak force is mediated by massive gauge bosons. Because of this, we can't apply the same gauge invariance prescription that we did in the last sections. If a mass term is added to the Lagrangian we break the gauge invariance we aimed to find. If we instead ignore the gauge invariance and add a mass term to the Lagrangian, all predictive power of the theory is lost due to unrenormalizable divergences. With "spontaneous symmetry breaking" we can gain massive gauge bosons while maintaining the integrity of the theory. In this section I first describe the "spontaneous symmetry breaking" mechanism in terms of an Abelian theory composed of complex scalar fields to illustrate the overall strategy. This mechanism is then applied to the non-Abelian electroweak theory to gain massive weak gauge bosons $W^{+/-}$ and Z with the Higgs field appearing as a 'spontaneous' result.

CHAPTER 2. THEORY

The Lagrangian for a $U(1)$ gauge symmetry

$$\phi \rightarrow e^{i\alpha(x)}\phi \quad (2.24)$$

As in the QED case, we introduce a gauge field A_μ and covariant derivative $D_\mu = \partial_\mu - ieA_\mu$ to obtain the gauge invariant Lagrangian

$$\mathcal{L} = (\partial^\mu + ieA^\mu)\phi^*(\partial_\mu u - ieA_\mu)\phi - \mu^2\phi^*\phi - \lambda(\phi^*\phi)^2 - \frac{1}{4}F_{\mu\nu}F^{\mu\nu}. \quad (2.25)$$

In this example if $\mu^2 > 0$ we gain back the QED Lagrangian for a charged scalar particle of mass μ - with an additional self-interaction term. However, if we take $\mu^2 < 0$ the potential $V(\phi^*\phi) = \mu^2\phi^*\phi - \lambda(\phi^*\phi)^2$ now has a non-zero vacuum expectation value (v.e.v.) and there's a set of equivalent minima shown in Figure 2.2. Choosing one of these minima spontaneously breaks the potential's rotational symmetry. Next, we can perturbatively expand the field about a minima through




Figure 2.2: Higgs potential when $\mu^2 < 0$, choosing a minima spontaneously breaks the $U(1)$ rotational symmetry [3]

CHAPTER 2. THEORY

$$\phi(x) = \sqrt{\frac{1}{2}[\nu + \eta(x) + i\xi(x)]} \quad (2.26)$$

Substituting this perturbation gives the new Lagrangian

$$\mathcal{L}' = \frac{1}{2}(\partial_\mu \xi)^2 + \frac{1}{2}(\partial_\mu \eta)^2 - \nu^2 \lambda \eta^2 + \frac{1}{2}e^2 \nu_\mu^A A^\mu - e\nu A_\mu \partial^\mu \xi - \frac{1}{4}F_{\mu\nu} F^{\mu\nu} + \text{interaction terms.} \quad (2.27)$$

Three particles seem to emerge here: massless Goldstone boson ξ , massive vector A_μ with $m_A = e\nu$ and a massive scalar η with $m_\eta = \sqrt{2\lambda\nu^2}$. However, the number of particles does not correspond to the polarization degrees of freedom expected. A longitudinal polarization was added, creating an unphysical field, and injecting extra degrees of freedom. To eliminate the unphysical field we can substitute new set of fields:

$$\phi \rightarrow \sqrt{\frac{1}{2}(\nu + h(x))e^{i\theta(x)/\nu}} \quad (2.28)$$

and

$$A_\mu \rightarrow A_\mu + \frac{1}{e\nu} \partial_\mu \theta. \quad (2.29)$$

Introducing these substitutions, the Goldstone boson field disappears and the new Lagrangian is

$$\mathcal{L} = \frac{1}{2}(\partial_\mu h)^2 - \lambda \nu^2 h^2 + \frac{1}{2}e^2 \nu^2 A_\mu^2 - \lambda \nu h^3 - \frac{1}{4}\lambda h^4 + \frac{1}{2}e^2 A_\mu^2 h^2 + \nu e^2 A_\mu^2 h - \frac{1}{4}F_{\mu\nu} F^{\mu\nu}. \quad (2.30)$$

Here the degrees of freedom before our substitutions remains the same and a massive boson A_μ is preserved along with a massive scalar h . The "Higgs mechanism" applied to a scalar field succeeded in creating a massive boson and determined the existence of a massive scalar boson. This same mechanism can be applied in the more complicated Standard Model electroweak field. Through electroweak symmetry breaking we not only gain massive gauge

CHAPTER 2. THEORY

bosons and a massive scalar boson (the Higgs!) but also a way to calculate testable Standard Model predictions for many other quantities. We start with the $SU(2) \times U(1)$ gauge symmetry of electroweak interactions derived in the previous section. In order to gain masses for three gauge bosons and keep the photon massless we need at least 3 degrees of freedom added and a simple choice is the $SU(2)$ doublet of scalar fields ϕ , with four fields in an isospin doublet of weak hypercharge $Y = 1$:

$$\mathcal{L} = (D^\mu \phi)^\dagger (D_\mu \phi) - \mu^2 \phi^\dagger \phi - \lambda (\phi^\dagger \phi)^2 \quad (2.31)$$

where ϕ is a $SU(2)$ doublet of complex scalar fields

$$\phi = \sqrt{\frac{1}{2}} \begin{bmatrix} \phi_1 + i\phi_2 \\ \phi_3 + i\phi_4 \end{bmatrix}. \quad (2.32)$$

Local gauge invariance can be achieved just as in the $U(1)$ case with the covariant derivative, though now a bit more complex:

$$D_\mu = \partial_\mu + ig \frac{\tau_a}{2} W_\mu^a \quad (2.33)$$

with three gauge fields $W_\mu^a(x)$ and $a = 1, 2, 3$. An infinitesimal transformation is defined

$$\phi(x) \rightarrow \phi'(x) = (1 + i\alpha(x) \cdot \tau/2)\phi(x) \quad (2.34)$$

so that we have a Lagrangian potential

$$V(\phi) = \mu^2 \phi^\dagger \phi + \lambda (\phi^\dagger \phi)^2. \quad (2.35)$$

Once again if we choose the conditions $\mu^2 < 0$ and $\lambda > 0$ there is rotational symmetry in

CHAPTER 2. THEORY

our choice of vacuum expectation value. In this case choice of v.e.v. is limited. For the photon to remain massless, the vacuum must be invariant under $U(1)$ (or electromagnetic) transformations, and not be charged in either direction (charge conservation). Thus the chosen minima to spontaneous break electroweak symmetry is

$$\phi_0 = \sqrt{\frac{1}{2}} \begin{bmatrix} 0 \\ \nu \end{bmatrix} \quad (2.36)$$

Next, substituting the vacuum expectation value ϕ_0 for $\phi(x)$ and expanding perturbatively yields

$$\phi(x) \rightarrow \begin{bmatrix} 0 \\ \sqrt{\frac{1}{2}(\nu + H(x))} \end{bmatrix} \quad (2.37)$$

Fully expanding this term in the Lagrangian gives a complex and illuminating result, the Goldstone bosons have been consumed and there is only a Higgs field ($H(x)$) remaining. Next, masses for the vector bosons are found from expanding one key parameter in the Lagrangian

$$|(-ig\frac{\tau}{2} \cdot W_\mu - i\frac{g'}{2}B_\mu)\phi|^2 = \frac{1}{8} \left| \begin{bmatrix} gW_\mu^3 + g'B_\mu & g(W_\mu^1 - iW_\mu^2) \\ g(W_\mu^1 + iW_\mu^2) & igW_\mu^3 + g'B_\mu \end{bmatrix} \begin{bmatrix} 0 \\ \nu \end{bmatrix} \right|^2. \quad (2.38)$$

Expanding further and substituting $W^\pm = (W^1 \pm iW^2)/\sqrt{2}$ gives the result

$$|(-ig\frac{\tau}{2} \cdot W_\mu - i\frac{g'}{2}B_\mu)\phi|^2 = (\frac{1}{2}\nu g)^2 W_\mu^+ W^{-\mu} + \frac{1}{8}(W_\mu^3, B_\mu) \begin{bmatrix} g^2 & -gg' \\ -gg' & g'^2 \end{bmatrix} \begin{bmatrix} W^{\mu 3} \\ B^\mu \end{bmatrix}. \quad (2.39)$$

It is immediately clear that there is a mass-term for the W^\pm , $M_W = \frac{1}{2}\nu g$. Masses for the photon and Z -boson are also apparent after expanding the last final term

$$\frac{1}{8}\nu^2(g^2(W_3^\mu)^2 - 2gg'W_\mu^3B^\mu + g'^2B_\mu^2) = \frac{1}{8}\nu^2(gW_\mu^3 - g'B_\mu)^2 \quad (2.40)$$

CHAPTER 2. THEORY

and using the substitutions

$$\begin{aligned} A_\mu &= \frac{g'W_\mu^3 + gB_\mu}{\sqrt{(g^2 + g'^2)}} \text{ with } M_A = 0, \\ Z_\mu &= \frac{gW_\mu^3 - g'B_\mu}{\sqrt{(g^2 + g'^2)}} \text{ with } M_Z = \frac{1}{2}\nu\sqrt{(g^2 + g'^2)}. \end{aligned} \quad (2.41)$$

Now the Higgs field exists just as in the previous example and the theory contains a massive scalar boson and three massive vector gauge fields - for each of the W^\pm and Z bosons. The Goldstone bosons have been consumed, their degrees of freedom used to give mass to the vector bosons. Choosing a ground state and so breaking the gauge symmetry does not eliminate this symmetry altogether, since the theory is still renormalizable. Fermion masses can also be derived from their interactions with the Higgs boson using this Lagrangian. These derivations can be used to predict masses of bosons and fermions and couplings to the Higgs boson. It's important to note that though the Higgs mechanism gives mass to all fermions and massive gauge bosons, it doesn't determine what the Higgs mass ought to be. This is left as an empirical input to the theory that can then be used to calculate other observables.

The Standard Model has been proven over decades to be an incredibly robust theory and the Large Hadron Collider (LHC) is its key testing ground.

2.2 LHC Physics/Phenomenology

The Large Hadron Collider (LHC) is the foremost Standard Model testing ground and the proton-proton collisions recorded through ATLAS, CMS, ALICE, and LHCb have demonstrated the breadth and accuracy of the theory. Fermion and gauge boson masses and couplings, including the mass of the Higgs boson, have been measured with increasingly

CHAPTER 2. THEORY

high precision. In the next chapter the mechanics of the LHC and ATLAS detector will be discussed, but first here I will introduce the motivations and observations of LHC physics. This section will begin with the mechanics of proton-proton collisions and their decay products, then discuss the concept of decay cross-sections and finally focus more closely on the Higgs boson and its properties.

The LHC was designed with one central goal- discover the missing Standard Model Higgs boson. The protons in the LHC collide at a center-of-mass energy of 13 TeV, but began at half that in 2010. The electroweak symmetry breaking scale was theoretically known to be between 100-1000 GeV and so probing at 7 TeV provided near certainty of finding either the Higgs or an inconsistency in the Standard Model. The motivation for a proton collider was multifaceted. Foremost, using the tunnels built for the electron-positron detector LEP with protons allows the collider to reach higher energies, as protons don't lose energy to synchrotron radiation at the same scale as electrons. However, proton collisions have added complexity from their component quarks. Each parton carries some fraction of the momentum of the proton described by parton distribution functions.

Figure 2.3 shows a proton-proton collision schematic. In this example the hard process comes from the up quark in each proton [4]. “Hardness” refers to the fraction of proton momentum involved in the collision. In contrast, “soft” collisions are those from remaining partons in each proton and usually involve low momentum transfer. These soft collisions are considered the underlying event shown in the figure 2.3. Parton scatter is the most common hard process at the LHC by far due to the high density of gluons in the proton and the scale of QCD couplings above electroweak coupling strength.




Figure 2.3: Example proton-proton collision with quark-gluon scattering and final state jet and Z-boson [4]

Quarks and gluons emitted from the high energy hard scatter do not appear in the detector directly. QCD, in one of its key differences to QED, becomes stronger with larger distances. As a parton reaches high enough energy, it will begin to radiate low energy gluons until resultant partons are able to bind into color-neutral hadrons. These hadrons are seen collimated in groups in the detector as “jets”. The energy and momentum of jets are considered reflections of the intial scattered partons. Various “jet algorithms” can be used to determine initial parton properties as reproducibly and accurately as possible. The jet algorithm used in this analysis will be described in detail in Chapter 3. However, it is important to note that the algorithm used by all LHC experiments- anti- k_t - is collinear and infrared safe, or unaffected by small angle and soft scatterings that occur in a parton shower. Without these qualities, perturbation theory applied to the parton shower would find infinities at high orders.

Cross-sections (denoted σ) measure the probability that a certain process will occur in the collision of two particles, in our case protons. In high energy physics cross-sections are

CHAPTER 2. THEORY

measured in inverse femtobarns. A barn is the cross-sectional area of a uranium nucleus and was named to describe the large target area needed in order to have direct strikes on a nucleus. The name was inspired by the expression "couldn't hit the broad side of a barn". Inverse femtobarns are used to measure the number of particle collision events per femtobarn area of a target and quantifies time-integrated luminosity.

Hard scattering cross-sections in hadron-hadron collisions can be calculated using the QCD factorization theorem, and to leading-order these calculations are relatively simple. In the factorization theorem, developed by Drell and Yan, deep inelastic scattering parton model processes could apply to hadron-hadron collisions. The Drell-Yan process is the production of a massive lepton pair by quark-antiquark annihilation. According to the factorization theorem, a hadronic cross-section $\sigma(AB \rightarrow \mu^+\mu^- + X)$ could be calculated by weighting the Drell-Yan sub-process cross-section $\hat{\sigma}$ for $\bar{q}q \rightarrow \mu^+\mu^-$ with parton distribution functions $f_{q/A}(x)$ which come from deep inelastic scattering [5]:

$$\sigma_{AB} = \int dx_a dx_b f_{a/A}(x_a) f_{b/B}(x_b) \hat{\sigma}_{ab \rightarrow X} \quad (2.42)$$

where X represents the two resulting leptons and ab the two annihilated quarks. This parton model provides good agreement with measured cross sections and so allows understanding of particular hard scattering processes. Predictions for some key Standard Model processes are shown in Figure 2.4. Noting the logarithmic scales it's clear that the Higgs boson of mass 125 GeV is orders of magnitude more numerous at the LHC than the Tevatron and that certain high mass particles like the b quark and W/Z bosons are produced at the LHC at high levels [5]. In addition, the plot shows cross-sections of particular Higgs decay modes. These will be discussed next.

CHAPTER 2. THEORY




Figure 2.4: Predicted Standard Model cross-sections for the Tevatron and LHC [6].

Higgs production at the LHC occurs via four main processes: gluon-gluon fusion, vector-boson fusion, associated production with W/Z bosons, and associated production with top or bottom quarks. The Feynman diagrams for these processes are shown in Figure 2.5.

CHAPTER 2. THEORY




Figure 2.5: Feynman diagrams for the leading Higgs boson production modes at the LHC [7].

The LHC Cross-section Working Group produces predictions on cross-sections, branching ratios, and pseudo-observables for the Higgs boson. Four CERN reports bring together Higgs recommendations for current and planned LHC efforts [8]. Figure 2.6 shows the Higgs cross-section for the main production modes as a function of the Higgs mass and the collision center-of-mass energy. The Standard Model doesn't predict the Higgs mass, but with our experimental knowledge of that m_H is about 125GeV, the calculations can be focussed on the region of interest. Examining cross-section as a function of center-of-mass demonstrates the increase in statistics for events of interest that running the detector at higher energy levels can accomplish. It is also clear that gluon-gluon fusion Higgs production is the leading production mechanism by far. The cross-section is currently known at NNLO in QCD with NLO EW corrections. The second highest production cross-section is from vector-boson-fusion (VBF). As seen in the Feynman diagram, two outgoing quarks are produced in the interaction. These quarks produce two hard jets in the forward region, with the Higgs boson appearing between them. To leading order VBF Higgs production is solely electroweak and QCD corrections (calculated at NLO) have a smaller impact than in ggF. NLO EW corrections are also applied. As a result, VBF theoretical uncertainties are smaller

CHAPTER 2. THEORY

than those on ggF. Vector boson associated Higgs production through a W/Z boson are less common than VBF but also dominated by electroweak processes with a small QCD correction (NNLO). Finally, associated production with top and bottom quarks is shown, though these are quite rare and have high NLO and NNLO QCD corrections.




Figure 2.6: Higgs production cross-sections over Higgs mass at center-of-mass energy 14TeV (left) and over center-of-mass energy for a Higgs mass of 125GeV (right)[8]

Theoretical uncertainties shown as colored bars in 2.6 are calculated from choice of PDFs and renormalization and factorization scales. Parton distribution functions (PDFs) are described in more detail by the PDF4LHC working group [9]. This group performs studies of PDFs and their predictions at the LHC and makes recommendations for methods of estimating PDF uncertainties.

CHAPTER 2. THEORY




Figure 2.7: Predicted branching ratios for the Higgs boson at the LHC as a function of Higgs mass [8].

Since the Higgs boson couplings are directly proportional to the masses of each particle, the Higgs decays most readily into the heaviest particles possible. Figure 2.7 shows key decay mode branching ratios near the experimentally known Higgs mass. While the branching ratios demonstrate the relative abundance of each Higgs decay, these don't translate directly into their ease of discovery or measurement. The current status of Higgs boson coupling and cross-section measurements for each decay mode will be detailed in the next section. The Higgs boson discovery was made through a combination of searches in many channels though dominated by $H \rightarrow ZZ^*$, $H \rightarrow \gamma\gamma$, and $H \rightarrow WW$ [10]. This is because though other decay branching ratios are higher, like $H \rightarrow b\bar{b}$, the backgrounds associated with this decay are much higher. As previously mentioned, proton-proton collisions create large amounts of QCD jets that are difficult to discern from target hard QCD processes. Because of this, Higgs decays to quarks and gluons are particularly difficult and those with leptonic products (ZZ decays to 4 leptons, decays directly to photons, and WW decays to two leptons and two neutrinos) are much easier to reconstruct. As energy and integrated luminosity increased

CHAPTER 2. THEORY

during LHC runs 1 and 2, measurements of even rare and background-heavy Higgs decay channels were able to be made. This thesis focuses on the decay of $H \rightarrow W^+W^- \rightarrow \ell\nu\ell\nu$ through VBF production and in the last section of this chapter I will motivate the choice of these conditions for probing new physics beyond the SM.

2.3 Brief history of SM tests

Here I summarize key moments in the history of the development of the Standard Model and its experimental tests. I use both [11] and [12] as a guide.

The history of the Standard Model could start with any number of physicists well before the formalism of the theory itself. As far back as the fifth century B.C. philosophers posited that matter is composed of discrete “particles” in its most fundamental state. This idea was only tested beginning in the nineteenth century, when chemists were able to detect physical evidence of atoms and their structure. The first of the gauge theories, QED, was invented in the 1930s but only calculated to first order. Renormalization theory was invented simultaneously by Feynman, Schwinger, and Tomonaga in the 1940s. This made calculations of higher order QED results possible. Following this advancement, QED was verified experimentally with incredibly high precision.

Physicists next attempted to understand and formalize the other fundamental forces- strong and weak- in the same way. Symmetries for these theories were not as easy to find as that of QED and in 1954 the first new gauge theory for QCD was proposed by Shaw, Yang, and Mills. Though ultimately incorrect it led to proposals for a gauge theory of weak interactions by Schwinger in 1956 (unifying weak and electromagnetic interactions with photons and massive W^\pm bosons). Glashow added a fourth boson, Z , to the theory, but the problem of a broken symmetry necessary to give mass to the W/Z bosons remained.

CHAPTER 2. THEORY

Spontaneous symmetry breaking was a known and tested concept, but its use in the theory led to unwanted massless “Nambu-Goldstone” bosons. Some thought this problem was inevitable in using SSB in the gauge theory. In 1964 Englert and Brout, then Higgs and a few months later Guralnik, Hagen, and Tribble all published papers with the same conclusion—the Goldstone theorem can be applied to gauge theories and massive bosons can “eat” the Nambu-Goldstone bosons to gain mass. This then creates a new scalar field whose particles are now termed Higgs bosons. A few years later Weinber unified these ideas into the electroweak theory we recognize today. The theory was confirmed many times over at experiments over the next decades.

While electroweak symmetry breaking and its implications were being understood in the 1960s, QCD was gradually being assembled. Experimental discoveries of a host of new particles led to Gell-Man and Zweig’s development of a theory that these were all composed of the same three base particles, “quarks”. Han and Nambu understood that an octet of colored gauge bosons, “gluons”, mediated the strong force through color interactions. In 1973 Gross, Wilczek, and Politzer demonstrated the asymptotic freedom of QCD, its weakness at short distances which allows perturbation theory to calculate high-energy interactions. The Standard Model as it is composed of electroweak dynamics and QCD, has been remarkably predictive but there is much it doesn’t explain. While theoretical physicists work to expand the Standard Model (or replace it entirely), experimentalists search for deviations from Standard Model predictions, which may be the next hint of entirely new physics.

Large amounts of experimental evidence for the predictive power of QED had amassed over time and by the 1970s high energy accelerators at CERN, Fermilab, Brookhaven and SLAC began making first measurements of predicted electroweak and QCD observables. In 1969 physicists at SLAC collided electrons and protons and found that electron scattering behaved as though the proton was made up of point-like particles, quarks. Electroweak

CHAPTER 2. THEORY

theory's predicted new neutral weak current interaction between quarks and leptons was discovered at CERN and then Fermilab in 1973, giving evidence to electroweak unification. The W and Z bosons predicted by the theory with masses $\approx 80\text{GeV}$ were discovered at SPS at CERN. This was the first proton-antiproton collider and with a center-of-mass energy of 540GeV , it was the highest energy collider ever built. The collider was built for the main purpose of finding the predicted W and Z bosons and it succeeded in discovering both in 1983. Higher energy colliders like LEP at CERN and SLC at SLAC were able to produce millions of W and Z bosons and so test electroweak predictions with high precision. The theory continued to prove extremely accurate. QCD remains more difficult to test precisely as calculating QCD parameters theoretically is a computationally difficult task. However, evidence of QCD's accuracy has still accumulated. In 1978, SPEAR at DESY was able to indirectly detect the first gluon. The electron-positron collider produces two quarks that form two jets of particles with equal energy in opposite directions. QCD predicts that there will also be 3-jet events, where a gluon is radiated from one of the scattered quarks and forms the third quark. The strength of the strong interaction was first measured in 1978 and thereafter with more and more precision and has always followed QCD predictions. Quarks beyond the first generation (up, down) were discovered progressively in terms of their mass-strange, charm, bottom, and finally in 1995 the top. Similarly the muon was discovered well before the heavier tau-lepton. After the discovery of the top quark, the last Standard Model particle left undiscovered was the Higgs boson. Its mass is an input rather than a prediction to the Standard Model, but with the large amount of data taken at increasingly high energy colliders, experimentalists and theorists were confident that if it existed, it would be between 100GeV and 1TeV . In order to search for the Higgs boson, the Large Hadron Collider was planned and built at CERN. The collider and its largest all-purpose detector, ATLAS, are discussed in the next chapter.

CHAPTER 2. THEORY

In 2012, the Higgs boson was discovered at the LHC in both the ATLAS and CMS detectors. After just one year of data at a center-of-mass energy of 7-8TeV, combined searches predominantly in the $H \rightarrow ZZ^* \rightarrow \ell\ell\ell\ell$, $H \rightarrow \gamma\gamma$ and $H \rightarrow WW \rightarrow \ell\nu\ell\nu$ channels found a particle compatible with the Standard Model Higgs with a significance of 5.9 standard deviations at a mass close to 125 GeV. The LHC continued data-taking after the discovery with a new goal of measuring this new particle and its properties with accuracy and precision. The center-of-mass energy increased to 13TeV and more than $20\times$ the data used in the first discovery has now been recorded. Measurements of the Higgs boson are now numerous and incredibly precise, but the particle still doesn't deviate significantly from its theoretical expectation. While the Standard Model has proven a successful model of known interactions, there are many phenomena that it does not predict - from dark matter to gravity to neutrino masses. There must be physics beyond this theory and in continuing to probe new aspects of the model, the LHC may find deviations from the known forces or physics beyond the Standard Model.

The mass of the Higgs boson is measured through a combination of decay modes and in combination with CMS to be 125.10 ± 0.14 GeV [1]. The latest combined Higgs cross-section measurement from ATLAS uses data from 2015-2017 and finds the production cross-sections (normalized to their Standard Model predictions) as shown in Figure 2.8. Branching ratios of relevant decay modes are also measured and are shown in Figure 2.9 multiplied by their cross-sections in each relevant production mode. Overall results demonstrate Higgs production and decay agree with SM predictions over a wide array of decay channels and production modes. The differences in theoretical and systematic uncertainty for certain decays (QCD heavy VBF $H \rightarrow b\bar{b}$ in comparison to the leptonically decaying VBF $H \rightarrow WW^*$) show how discernable backgrounds play a major role in the viability of a measurement, even when the branching ratio is high [13].

CHAPTER 2. THEORY




Figure 2.8: Production cross-sectios for ggF, VBF, VH, and $t\bar{t}H + tH$ normalized to their SM predictions. Total, systematic, and statistical uncertainties are shown [13]




Figure 2.9: Branching ratios for measured Higgs decays normalized to their SM predictions. Total, systematic, and statistical uncertainties are shown [13]

CHAPTER 2. THEORY

This thesis details my work on the differential measurement of VBF $H \rightarrow WW^* \rightarrow \ell\nu\ell\nu$ decays. The clear sign of VBF Higgs production (2 jets recoiling in opposite high η directions, combined with the leptonically decaying W bosons) gives the decay a distinctive signature. This channel can be measured with high accuracy which may allow small deviations from theory to be visible. In addition, the differential cross-section measurement can probe higher order perturbative contributions to QCD or EW. Details on the particular motivations for this thesis' analysis will be outlined in the next section. The last ATLAS VBF $H \rightarrow WW$ differential measurement was published in 2016 with 20.3fb^{-1} at center-of-mass energy 8TeV and measured only ggF Higgs boson production differential cross-sections [14]. This new measurement is presented with a $4\times$ higher integrated luminosity and center-of-mass energy 13TeV as well as new methodology for overall analysis strategy in order to take advantage of larger statistics. This is also to be the first ATLAS measurement focussing primarily on VBF Higgs differential cross-sections(?).

2.4 Measurement motivation

There are many reasons why the VBF $H \rightarrow \ell\nu\ell\nu$ channel is particularly interesting to measure precisely. Since the W bosons decay leptonically, they create a clear signal in the detector which allows precise reconstruction of the hard processes. Further, the VBF production mode has a similarly clear signal with the two jets accompanying the decaying W bosons with a high η separation. The pseudorapidity of these jets and the rapidity between them are particularly sensitive to the electroweak symmetry breaking mechanism, so a differential cross-section measurement over these variables (along with other detailed and motivated in further sections) tests the Higgs mechanism. As previously stated, VBF Higgs WW decays have never been measured with the ATLAS detector, so this analysis

CHAPTER 2. THEORY

introduces novel measurements. In Run-1 VBF $H \rightarrow WW^* \rightarrow \ell\nu\ell\nu$ cross-sections were measured with an experimental uncertainty of $\approx 30\%$ [13]. This new measurement uses the entirety of Run-2 data, with a factor of $2.5\times$ higher statistics. This VBF differential $H \rightarrow WW$ measurement uses new analysis techniques, higher statistics, and improvements in theoretical calculations to make an accurate probe of new physics within the electroweak symmetry breaking mechanism.

This thesis focuses on the VBF differential $H \rightarrow WW$ measurement, but this measurement is the first piece to a larger goal beyond the scope of my Ph.D. Our group plans to make a simultaneous measurement of VBF $H \rightarrow WW$ and VBF $W + \text{jets}$ fiducial and differential cross-sections. The VBF $W + jj$ production cross section is sensitive to the WWV vertex. In vector boson scattering, we expect (with minimal assumptions) that new physics ought to be at a scale above the current experiment. Effects from this type of new physics would simultaneously affect HWW and WWV vertices. In particular, dimension-6 operators affect VBF Higgs mechanisms and VBF W production, while the interplay of VBF and VBS mechanisms allows for the probe of dimension-6 and dimension-8 operators. Measuring fiducial and differential cross-section ratios (of VBF Higgs and Wjj) enhances sensitivity to new physics because any new phenomena that simultaneously affect both vertices are suppressed and so sensitivity to the phenomena affecting just the WWV vertex (for example, anomalous quadruple-gauge couplings) is enhanced. VBF Higgs and Wjj productions are also characterized by final-states with two jets with similar kinematics. Because of this, the ratio of cross sections measurement allows for the cancellations of correlated systematic uncertainties. As both process are characterized by large ($10\% - 20\%$) jet-related uncertainties, a reduction via the ratio cancellation would greatly improve the sensitivity to the VBF Higgs process and to the search for new phenomena. This VBF HWW measurement is the first part of the ratio of fiducial differential cross sections for VBF HWW and Wjj , which will

CHAPTER 2. THEORY

set limits on EFT parameters, benefiting from enhanced sensitivity through the cancellation of correlated uncertainties.

Chapter 3

The LHC and the ATLAS detector

3.1 The Large Hadron Collider

The Large Hadron Collider (LHC) is a proton-proton storage ring operating at CERN and for last 13 years it has been the world's highest energy particle collider. The LHC first began operation in 2008 but following a magnetic quench incident it had to be repaired and adjusted. The first data-taking commenced in 2009 [15]. During the 9 years of LHC operation thus far, protons collided with increased center-of-mass energiy, approaching the design energy of 14TeV. Instaneous luminosity has also successively increased, surpassing design instantaneous luminosity of $1 \times 10^{34} \text{cm}^{-2}\text{s}^{-1}$ in 2018 to reach $2 \times 10^{34} \text{cm}^{-2}\text{s}^{-1}$ [16]. The overall data recorded in the ATLAS detector totals more than 10^{16} collisions, and total collisions (not recorded) far surpasses this. Operation of the LHC has led to the most precise measurements of Standard Model constants yet including the coupling of the Higgs boson to bottom quarks [?], W and Z bosons [17], [18], photons[19] and tau leptons [20]. The LHC has also facilitated searches over a wide parameter space, setting confidence level exclusion limits on masses of supersymmetric particles like squarks, gluinos and neutralinos

CHAPTER 3. THE LHC AND THE ATLAS DETECTOR

[21].

The LHC is set to begin Run 3, in which design center-of-mass energy should be reached, in 2021. Following Run 3 detector upgrades will be made during a long shutdown. Then the High-Luminosity LHC (HL-LHC) will begin colliding protons with unprecedented $10 \times$ LHC-luminosity in 2027 [22]. The HL-LHC and its goals will be explained further in later in this chapter. Suffice to say LHC-physics is progressing quickly and promises exciting developments in the near future.

In a brief explanation of the LHC operation, one could begin with the small volume of $\approx 10^{11}$ protons that are accelerated. Linac 2 is the primary source of protons for CERN accelerators and has been since the early 1990s [23]. This injects protons at 50 MeV into the Proton Synchrotron Booster (PSB) where they are further accelerated to 1.4 GeV. Next they enter the Proton Synchrotron (PS) where the protons are separated into bunches with a spacing of 25 ns and are further accelerated to 25 GeV before being extracted to the Super Proton Synchrotron (SPS) where they reach 450 GeV. Finally these bunches of protons enter the LHC and are accelerated to their final energy of 6.5 TeV. Linac 2, PSB, PS, and SPS were all operational accelerators before being involved in the LHC apparatus though each had to be majorly upgraded to handle the energy and beam intensity required for LHC collisions [23].

The LHC layout mimics that of the Large Electron Positron collider (LEP) that was housed in the same tunnels. Figure 3.1 shows the positioning of each experiment on the LHC as well as injection systems and other features. Once proton bunches enter the LHC in two opposing beams they are accelerated with radio frequency (RF) systems. Located at Point 4 in the LHC schematic, the system consists of 16 RF cavities operating at twice the frequency of the SPS injector. RF cavities are metallic chambers containing oscillating electromagnetic fields, in the LHC this oscillation frequency is 400 Mz. The tuning of this

CHAPTER 3. THE LHC AND THE ATLAS DETECTOR

frequency ensures that protons of the ideal energy are not accelerated further and simply maintain their momentum while particles arriving in an RF cavity slightly before or after will be decelerated or accelerated toward the ideal proton energy. This acceleration process can also be used to split beams of protons into discrete bunches, and this is first done with RF cavities in the PS. After proton bunches have circled the LHC approximately 1 million times (15 minutes), peak energy is reached and collisions can commence [24].




Figure 3.1: LHC layout [25]

Superconducting magnets in the LHC main dipoles create a magnetic field of $\approx 8\text{T}$ to bend the proton beams into the circular path of the collider. Figure 3.2 shows the flux in a dipole cross-section. The opposing direction beamlines are shown centered and the flux is shown to be high (and directionally opposed) in the center of each beam. To maintain these fields, the magnets operate at below 1.9K. Pressurized superfluid helium chosen for its low viscosity and high specific heat cools the dipole magnets. Once the two LHC rings are filled from the SPS, center-of-mass energy of the beams increases until it reaches peak energy after about 28 minutes. Finally proton bunches separated by 25ns collide simultaneously in each

detector.




Figure 3.2: Flux within an LHC dipole cross-section [25]

3.2 A Toroidal LHC ApparatuS

The LHC creates proton-proton collisions at the rate and energy necessary for pushing the boundaries of particle physics, but identifying and reconstructing the tracks of such energetic particles is no mean feat. A Toroidal LHC ApparatuS (ATLAS) and the Compact Muon Solenoid (CMS) are multi-purpose detectors built to search for and measure a wide range of particle interactions and properties. Both experiments measured a particle consistent with the Higgs boson in 2012 and their agreement was a key verification of the discovery. The following sections describe each major component of the ATLAS detector so to highlight their necessity in the measurement of $H \rightarrow WW \rightarrow e\nu\mu\nu$.

ATLAS utilizes a coordinate system with its origin at the center of the detector (the “interaction point”) and has a z-axis along the beam pipe. The x-axis points from the interaction point to the center of the LHC ring, and the y-axis points upward. The experiment

CHAPTER 3. THE LHC AND THE ATLAS DETECTOR

uses cylindrical coordinates (r, ϕ) where ϕ is the azimuthal angle around the beam pipe. The pseudorapidity and the transverse momentum are defined in terms of the polar angle θ as $\eta = -\ln(\tan(\theta/2))$ and $p_T = p \sin \theta$.




Figure 3.3: Computer-simulated ATLAS detector schematic [26]

The Inner Detector (ID) detects charged particles with $|\eta| < 2.5$ operating in a 2T solenoidal field. It consists of 3 layers of pixel sensors, 4 layers of silicon strips, and 72 straw layers of transition radiation tracker modules. The ID describes particles closest to the interaction point and locates track parameters with great resolution due to high granularity [26].

The ATLAS detector contains 3 superconducting magnet systems- the barrel toroid, 2 end-cap toroids, and a central solenoid. The central solenoid provides a magnetic field for the inner detector while the toroids create a strong magnetic field for the muon detector. These magnet systems were built to create the largest possible uniform field (for increased momentum resolution on particle tracks) on a large volume enclosing the detector

CHAPTER 3. THE LHC AND THE ATLAS DETECTOR

components. They also need to use as little material as possible so as to not unduly influence particles in the detector. The toroids in the barrel and endcap each have 8 coils and create a 4T magnetic field while the central solenoid creates a 2T magnetic field in the inner detector. Combined the magnet systems contain $>100\text{km}$ of superconducting wire which are cooled to working temperatures below 5K [26].

The Muon Spectrometer precision chambers provide muon momentum measurements at a high resolution over a wide range of p_T . The MS consists of 3 layers of Monitored Drift Tube chambers covering $|\eta| < 2.7$ and an inner layer of Cathode Strip Chambers with $|\eta| > 2.0$. In addition, it includes trigger chambers that contain 3 layers of Resistive Plate Chambers ($|\eta| < 1.05$) and 3 layers of Thin Gap Chambers ($1.05 < |\eta| < 2.4$). As the outermost subdetector, the MS provides precise muon momentum measurements along the muon trajectory and the muon chambers are located with a precision of under $60\ \mu\text{m}$. The MS also contains a system of three superconducting toroidal magnets each with eight coils providing a magnetic field with a bending integral of up to 6 Tm [26].

Calorimeters provide detailed information about the energy deposited as particles pass through. Electromagnetic calorimeters detect and halt the motion of electrons and photons while the hadronic calorimeter does the same for hadrons. Muons and neutrinos are able to pass through the calorimeters to the MS. The electromagnetic and hadronic calorimeters, made of liquid Argon and scintillating tiles respectively, are able to pass information from the location of energy deposits to the various identification and reconstruction algorithms [26].

3.3 The High-Luminosity LHC and Inner Tracker (ITk)

Though the LHC succeeded in its paramount goal of discovering the Higgs boson in 2012, its continuous operation at higher energy and luminosity has led to more rigorous measure-

CHAPTER 3. THE LHC AND THE ATLAS DETECTOR

ments of Higgs boson properties as well as searches for new physics beyond the Standard Model. The LHC has been the leading high energy collider in terms of person power, energy, and scale for over a decade and will continue to be extremely important for understanding theoretical questions in the future. While more data collection is planned in Run-3 starting in 2021, new colliders and detectors take decades to design, develop and build, so the plans for the collider to take its place is well underway. The High Luminosity LHC will operate at an LHC-level energy (14TeV) and begin data-taking in 2026. The HL-LHC will begin with $5 - 7 \times$ the luminosity of the LHC and will have a design instantaneous luminosity of $10 \times$ the LHC, or $12.6 \times 10^{-34} \text{cm}^{-2}\text{s}^{-1}$. This huge increase in number of collisions requires massive upgrades to the LHC including new 11-12T superconducting magnet systems, compact superconducting cavities for beam rotation and phase control, and new technology beam collimation [?]. This massive undertaking has been underway for many years already and has involved laboratories all over the world.

Just as the LHC had to be re-designed and built to create higher luminosity, so too do all the experiments on the LHC have to be redesigned to be able to interpret so many more collisions per second. The detectors must be built to withstand more radiation, as the increased collision rate also means a high radiation rate especially closest to the beamline. They also have to provide greater granularity to be able to reconstruct tracks with good enough resolution that individual tracks can be discriminated. Finally, they have to be able to deal with increased pile-up. Pile-up is caused by high numbers of collisions occurring at each bunch crossing. When there is a large amount of pile-up it becomes difficult to trace which particle tracks come from the same interaction point. Finally, the increased data in and of itself creates a complex problem for the detectors to solve, as briefly discussed previously, the trigger system must quickly choose which collision events may hold interesting events and store these. When there are more events happening near simultaneously these systems must

CHAPTER 3. THE LHC AND THE ATLAS DETECTOR

make these decisions in real-time. New algorithms to speed up this process are necessary when there are an order of magnitude more events to sort through.

Detectors for high energy colliders are not built often- expensive and time-consuming to design and test, they're made to last at least a decade. I was lucky to have the opportunity to work on ATLAS detector research and development during the year I worked at Brookhaven National Laboratory during my Ph.D. Though my thesis isn't directly related to this work, it was formative and extremely interesting, so I'll touch on this work in the next section. Because I worked on the new ATLAS inner detector for the HL-LHC (termed Inner Tracker or ITk) I will discuss solely this sub-detector and the particular role I played in its assembly.

3.3.1 Inner Tracker (ITk)

The Inner Tracker is planned to be an all-silicon detector that will completely replace the current Inner Detector. While the current ID has been extremely successful during Runs 1 and 2 (and will certainly continue to be in Run 3), it does not have the capacity to withstand the radiation and pile-up conditions of the HL-LHC. The ITk is designed to operate for 10 years under an instantaneous luminosity of $7.5 \times 10^{-34} \text{ cm}^{-2} \text{s}^{-1}$ with 25ns between bunch crossings. This will result in $1,000 \text{ fb}^{-1}$ and average pile-up up to $\langle \mu \rangle = 200$ [27]. The current solenoid magnet will remain in the detector with a 2T magnetic field. The ITk will consist of an innermost section with silicon pixels and an outermost section of silicon strips. The pixel detector will contain four barrel layers and six forward region disks, while the strip detector will contain five barrel layers and seven disks. The rapidity range matches the coverage of the Muon Spectrometer with $|\eta| < 2.7$. This layout is shown in ??.




Figure 3.4: ITk layout as defined in ITk Technical Design Report [27]

Building ITk Strip barrel staves

I spent 1.5 years of my Ph.D. at Brookhaven National Laboratory funded through a SCGSR Fellowship and was able to make key contributions to the ITk Strip barrel stave assembly effort. The goal of stave assembly is to precisely calibrate specified positions for silicon modules on carbon fiber stave cores and then glue them in place within a $25\mu\text{m}$ tolerance. Brookhaven National Laboratory is responsible for assembly of half off all ITk staves (200) and their accurate assembly is necessary for the ITk to reduce uncertainty on track positions as well as to ensure a symmetric detector. At Brookhaven I was tasked with co-creating a stave assembly software system through LABView to automatically calibrate required module positions, apply a layer of adhesive gel, and guide a user in accurately placing a module into its specified location. This project was highly collaborative and evolved further after I left the laboratory, but the overall process remains unchanged.

The basic design of the Inner Tracker for both barrel and endcap components is the same - a carbon fiber core (containing titanium cooling pipes) is covered on each side with co-cured kapton service tapes. The carbon fiber core is designed to reduce overall radiation

CHAPTER 3. THE LHC AND THE ATLAS DETECTOR

length and the similar design in the barrel and endcap adds to simplicity. Silicon modules are glued to stave cores. Similar radial silicon strip detectors have been used previously in both ATLAS and CMS but never covering so much fiducial area. The modules consist of one silicon sensor and one or two low-mass PCB’s (hybrid) which host ASICs. Module design has optimized producibility and low cost while maintaining readout goals. Overall module design is the same in barrel and endcap regions, while strip lengths and geometries vary. Components of a short-strip barrel module are shown in [3.5](#).




Figure 3.5: Short-strip barrel module components [\[27\]](#)

Each barrel stave core needs to be “loaded” with 14 modules, as shown in the assembled electrical prototype in [3.6](#). The placement of these modules needs to be accurate to below $25\mu\text{m}$ so that the ITk models accurately demonstrate particle track positions. After the detector is assembled and commisioned tests will be undergo laser alignment to understand the exact positioning of the modules and staves but any module positioning out of specification will have adverse affects on track resolution during operation.

Brookhaven National Laboratory is one of two sites responsible for assembling barrel staves. Assembly procedures have been tested with the production of numerous prototypes including a thermomechanical double-sided stave and a fully operational electrical stave. The thermomechanical prototype was later used for various thermal tests, including IR imaging

CHAPTER 3. THE LHC AND THE ATLAS DETECTOR




Figure 3.6: Electrical stave prototype at Brookhave National Laboratory

(documented in the next section). The electrical stave is currently undergoing tests for full electrical read-out. Stave assembly is composed of three main parts: system calibration, module placement, and survey of results.

Staves at BNL are assembled on a granite table housing an Aerotech XYZ Stage accurate to the micron level. The stage is equipped with a 10-megapixel camera that gives real-time feedback to a nearby computer and a glue dispenser. The stave assembly software system is implemented by a user who interacts with a LabView GUI and monitors progress. The stave is fixed to optical rails drilled into the table. In order to accurately place modules in their correct positions a series of calibrations need to be completed including camera calibrations to test the optimal working point, focus, and pixel-to-micron conversion. Next, the position of the stave with relation to the XYZ stage needs to be calibrated. Transforming coordinates of the XYZ stage to that of the stave and so specified coordinates for modules atop their carbon fiber cores requires locating a fixed point on the stave core as well as the angle of the stave relative to the XYZ stage. Pattern matching algorithms find the exact locations of particular features on the stave core and allow calculation of required positions for all modules based on specifications. Once specified module positions are calculated, calibrations are complete and it is time to apply glue and affix modules.

Next an epoxy (SE4445) is loaded into the glue dispenser on the XYZ-stage which is connected to a vacuum controlled by the LABView software system. The epoxy is automatically dispensed in lines to cover $\approx 60\%$ of area under the module. Then the module

CHAPTER 3. THE LHC AND THE ATLAS DETECTOR

is lifted with a custom-made “pick-up” tool which uses a vacuum applied to module corners to hold the module in place and move it to the needed position along the optical rails. Then using real-time feedback from the software system and its pattern matching algorithm, the user is directed on how to finetune module position using knobs on the “pick-up” tool. Markings etched in the silicon sensor at each corner are used to position the module accurately. The output of the module alignment GUI is shown in 3.7. When the module is within specifications it is lowered into position above the epoxy and held in place for 24 hours until the glue has completely dried.



Figure 3.7: GUI interface showing etched marking on module corner located in real-time to guide user on how to adjust module position

Finally after the glue has set a final survey of module positions is recorded using pattern matching to find positions of etched markings on each module corner. These results are saved into an ITk database and checked for any biases. After module placement on the stave is complete, the loaded stave is moved to another station in the lab for wirebonding so that all data from the modules can be read-out to stave-wide electronics. The module assembly system has been successful at placing modules accurately for all prototypes, achieving specification requirements for almost all modules. Results of the first prototype stave’s

CHAPTER 3. THE LHC AND THE ATLAS DETECTOR

module placement are shown in 3.8. While a few module corners are slightly out of the ideal range, the majority are well within specifications. Throughout prototype assembly issues and inefficiencies were found and corrected. New hardware like an improved glue system and temperature monitoring were also added. The methods described continue to be in use now and will be utilized for the production of 200 ITk staves at BNL.




Figure 3.8: Histograms show difference between ideal and final position of each module corner. Left shows difference from specification in X and right in Y.

IR Testing ITk Strip barrel staves

The first full US stave prototype was the thermo-mechanical stave built in the summer of 2017. Building this stave was the first test of stave assembly procedures and the results showed success. This stave was also assembled to test the thermal and mechanical properties of a fully loaded barrel stave. Multiple studies were conducted including thermal measurements using thermistors and IR imaging, thermal cycling and thermal shock tests, mechanical studies, and bending tests. I will give a short summary on IR imaging tests, as these were another focus of my time at BNL. First, the thermo-mechanical stave consists of 13 modules mounted on each side. The modules used are thermo-mechanical, which means that instead of the usual readout chips their hybrids have copper traces to mimic the power dissipation and location of the chips. The powerboard can vary the TM hybrid power dissi-

CHAPTER 3. THE LHC AND THE ATLAS DETECTOR

pation of each module individually. In addition these modules each have three thermistors—one on the DC-DC converter, and one on each of two hybrids. The stave is powered by a custom End-of-Substructure (EoS) card which is attached to a RaspberryPi and Arduino and can power on or off each module.

Thermal testing of barrel staves had a few main goals: first, to compare with Finite Element Analysis (FEA) simulations and so test that all temperature trends are as we expect, to make sure that individual modules and their slightly modified assembly methods don't exhibit abnormal thermal behavior, and finally to check that loaded staves can cope with potential issues they might face during operation. I will highlight a few key results which demonstrate that each of these goals have been accomplished.

Thermal measurements were taken both through the mounted thermistors on each module and through IR imaging. IR imaging provides information about the entirety of the loaded stave, rather than at just a few module positions so provides a more complete picture. The loaded stave was spray-painted black with a high emissivity, low conductivity black paint since silicon is transparent to the IR camera spectrum ($8\text{-}14\mu\text{m}$). In order to image the entire stave core, the IR camera was attached to rails above and pulled at a constant speed with an external motor as it recorded video. The frames were then stitched together into one image. A section of the painted stave is shown in [3.9](#).

FEA simulations for the thermal performance of a stave were completed by Prof. Graham Beck at Queen Mary University of London. These calculations quickly became intractable if convection was included so conditions of the stave and coolant were adjusted such that convective contributions were minimal or the total electrical power and power absorbed into the coolant were identical. At BNL we adjusted the coolant temperature until we saw this convective power minimization then recorded module temperatures under these conditions. These results were compared to the FEA simulations by averaging hybrid temperatures

CHAPTER 3. THE LHC AND THE ATLAS DETECTOR




Figure 3.9: Portion of the thermo-mechanical stave after spray-painted for increased emissivity

recorded through IR imaging and recording NTC thermistor readings. These comparisons are shown in 3.10. The measurements show very good agreement with FEA calculations within 5% of the expected values.




Figure 3.10: IR measurements, NTC thermistor measurements, and FEA simulations for the TM stave are compared. Agreement with FEA simulation within 5%

During module assembly some slight variations were tested including varying glue thickness below modules, glue curing time, and FEAST versions. Modules with these variations were noted and compared to those without at varying coolant temperatures and output power settings. Overall, no significant differences in module temperature change was observed for any of these assembly modifications. Fully loaded stave thermal properties are thus robust

CHAPTER 3. THE LHC AND THE ATLAS DETECTOR

to assembly modifications like these. Figure 3.11 shows a full IR image of the fully loaded stave. It is clear that there are no obvious module-to-module variations in silicon, hybrid, or FEAST temperature. The module sensors increase in temperature as they get closer to the EoS, which is expected since it dissipates power to the stave.




Figure 3.11: IR image of fully loaded thermo-mechanical stave

The thermo-mechanical stave was pushed to limits beyond what we would expect loaded staves to encounter during operation and never exhibited unexpected behavior. Thermal cycles, thermal shocks and bend tests showed the loaded stave robust to intense temperature variation and that the carbon fiber core is as stiff as it was prior to loading. Another test was how neighboring modules would perform if one module malfunctioned and went offline. Figure 3.12 shows temperature of each module when one of them (fourth to the left) turns off. The rest shows modules continue to operate normally and temperature changes from the unpowered module don't propagate very far. The second image in the figure shows the reverse of the stave when a module is powered off (fourth from the right). The temperature effects are greater directly below than adjacent to which is expected due to material differences.

My experiences on ATLAS detector upgrades for the HL-LHC provided context for the bulk of my thesis research. This project provided me with in depth and hands-on knowledge of the ATLAS detector and its component parts, as well as the scale of effort required to build new detector components. I have an abiding appreciation for the people and technology

CHAPTER 3. THE LHC AND THE ATLAS DETECTOR




Figure 3.12: IR image of fully loaded thermo-mechanical stave

necessary for data-taking at the LHC both of which make measurements like the Higgs cross-section detailed in this thesis possible.

Chapter 4

Tracking and Isolation in ATLAS

The measurement of $H \rightarrow WW^* \rightarrow \ell\nu\ell\nu$ cross-sections relies on the accurate determination of a number of physics objects within the detector primarily jets, missing transverse energy (E_T^{miss}), electrons, and muons. The final state particles of this interaction consist of one electron, one muon, and two neutrinos, which appear as missing transverse energy in the detector since they cannot be directly detected. Because our search focuses on VBF production Higgs, we also require two jets in the final state. Each physics object has a dedicated performance group which is tasked with providing recommendations for reconstruction, identification, isolation and measurements of efficiency, scale, and resolution. Biases or omissions in any of these would severely impact the precision of our analysis and so understanding the uncertainties associated with each reconstructed object is critical. In this chapter I briefly outline the algorithms used for tracking and isolation and determining their relevant uncertainties for jets, missing transverse energy, electrons, and muons. Particular attention will be paid to muons as I spent a significant amount of time working in the Muon Performance Group applying and validating corrections to muon momentum scale and resolution.

4.1 Jets

Quarks and gluons emitted from high energy hard scattering do not appear in the detector directly. As quarks and gluons reach high enough energy, low energy gluons are radiated until partons are able to bind into color-neutral hadrons. These hadrons are seen collimated in groups as “jets”. Their energy and momentum are used in physics analyses as proxies for the initial scattered partons. Jet calibration seeks to determine jet energy scale and jet energy and angular resolution as accurately as possible. Pile-up presents the main difficulty in jet calibration - multiple interactions occurring in the detector at once creates significant, often hadronic, background. The hard interaction of interest must be separated from pile-up background (which is most often soft). During Run-1 the ATLAS experiment reconstructed jets using either only the calorimeter or the tracker, though most often the calorimeter. Topological clusters of calorimeter cells (topo-clusters) were used to trace jet tracks. At the end of Run-1, the jet energy scale (JES) correction factor used to calibrate jets to the particle level was re-calculated using additional track information from the Inner Detector and Muon Spectrometer, which greatly improved jet resolution [28].

Beginning in Run-2, a new algorithm for jet reconstruction took advantage of the improvements shown in Run-1 by including information from the tracker. ‘Particle flow’ uses the tracker’s higher momentum resolution for low-energy charged particles and its greater angular resolution of single charged particles. This is complemented by the calorimeter’s ability to reconstruct both charged and neutral particles and the calorimeter’s higher energy resolution for high energy physics objects. The calorimeter also has an extended acceptance so in the forward region only calorimeter topo-clusters are used. One potential difficulty with the ‘particle flow’ algorithm is the possibility for double counting particles if each tracker reconstructed jet isn’t correctly matched to its calorimeter signal. This is avoided in the

CHAPTER 4. TRACKING AND ISOLATION IN ATLAS

algorithm through the condition that if a particle's track measurement is used, its corresponding energy must be subtracted from the calorimeter measurement. The success of the algorithm in removing only energy deposits from the tracked jet represents a key criteria for its overall performance [29].




Figure 4.1: Flow chart of the particle flow algorithm beginning with track selection and ending with charged particles and changed/unchanged topo-clusters [29].

Topo-clusters are described by two main properties - ϵ , which represents the fraction of true energy deposited in the cluster out of the total true energy deposited in all topo-clusters for that object and ρ , the purity or fraction of a particle's true energy which lies within the topo-cluster. High ρ , high ϵ topo-clusters separate out contributions from different particles and so are easier to apply hadronic shower subtraction [29].

Jet tracks need to meet strict criteria to be considered. These include at least nine hits in silicon detectors, no missing Pixel hits, $\eta < 2.5$, and $40 > p_T > 0.5\text{GeV}$. These are referred to as a “tight” selection. Tracks which obey these criteria and do not deposit enough energy to create topo-clusters are still included. High p_T tracks are excluded because their poor isolation. In addition, tracks which are identified as electrons or muons are excluded [29].

With topo-clusters and tracks assembled, the algorithm next must match each to one another. Topo-clusters are ranked through the distance metric

$$\Delta R' = \sqrt{\left(\frac{\Delta\phi}{\sigma_\phi}\right)^2 + \left(\frac{\Delta\eta}{\sigma_\eta}\right)^2} \quad (4.1)$$

CHAPTER 4. TRACKING AND ISOLATION IN ATLAS

where σ_ϕ and σ_η denote angular topo-cluster widths. A requirement that $E^{clus}/p^{trk} > 0.1$ is applied so that the energy of the topo-cluster must contain more a significant portion of the energy of the track. This requirement rejects about 30–40% of the incorrect topo-clusters at $p_T > 5\text{GeV}$ (this is less pronounced at lower p_T and very rarely rejects the correct cluster). Distance selection is made such that the closest topo-cluster to each track in $\Delta R'$ is taken to be the correct match. This is very successful at $p_T > 5\text{GeV}$. If no topo-cluster is within a cone of $\Delta R' = 1.64$ it is assumed the particle did not form a topo-cluster in the calorimeter [29].

Topo-clusters are thus matched to particle tracks and next the energy the particle deposits must be subtracted from the calorimeter. The average energy deposited by a particle with momentum p^{trk} is given $\langle E_{dep} \rangle = p^{trk} \langle E_{ref}^{clus}/p_{ref}^{trk} \rangle$ where $\langle E_{ref}^{clus}/p_{ref}^{trk} \rangle$ is calculated using single-particle samples without pile-up by summing topo-cluster energies within $\Delta R = 0.4$ about the track position. These are calculated at varying p_T and η values to capture effects from detector geometry and shower development. Particles often split their energy between multiple topo-clusters and this can be determined through the significance of the difference between expected energy and that of the matched topo-cluster. If significance is below -1, a split shower recovery procedure runs: topo-clusters within a cone of $\Delta R = 0.2$ about the track position are considered to be matched to the track. This new full set of matched clusters is considered for energy subtraction [29].

Energy subtraction is performed cell-by-cell (unless $\langle E_{dep} \rangle$ is greater than the energy of the total matched topo-clusters, in which case they are all removed). Rings are formed in η, ϕ about the extrapolated track and are one calorimeter cell wide. The average energy density in each ring is computed and the ring with the highest energy density is subtracted first. This continues to lower density rings until $\langle E_{dep} \rangle$ is reached. Finally, after cell-by-cell subtraction, remnant energy clusters are removed if they're considered purely from shower

CHAPTER 4. TRACKING AND ISOLATION IN ATLAS

fluctuations and otherwise remain. Ideally now the set of selected tracks and remaining topo-clusters together represent the reconstructed event without double counting between the subdetectors. An example display of particle flow events is shown in Figure 4.2




Figure 4.2: Sum of transverse momenta of neutral and charged particle flow objects in an area $\Delta\eta \times \Delta\phi = 0.2 \times 0.2$ from a 2017 event with mean number of interactions per beam crossing $\mu = 38$ [30]

Jet finding algorithms aim to approximate the hadron generators of calorimeter and track-based jets in the detector. They bridge the divide between observable jet objects and theoretical predictions in terms of QCD. There are several jet algorithms in use currently and key properties of all are collinear and infrared safety. This means that neither splitting a jet collinearly nor soft emissions should change jet structure. Without these qualities perturbation theory diverges at high orders. In addition modelling hadronization, the underlying event and pile-up is imprecise, so the ideal theory would be minimally sensitive to these effects. Anti- k_t is the jet algorithm used by ATLAS and is infrared collinear-safe by construction and soft-resilient. Distance between pseudojets i and j , d_{ij} and the distance between pseudojet i and the beam d_{iB} are defined. The algorithm loops through these distances beginning with the smallest distance. If d_{ij} is smallest, pseudojets i and j are combined, if

CHAPTER 4. TRACKING AND ISOLATION IN ATLAS

d_{iB} is smallest then i is a jet and removed from the list. This continues iteratively until all jets are defined. The distances are defined:

$$d_{ij} = \min(k_{ti}^{2p}, k_{tj}^{2p}) \frac{\Delta_{ij}^2}{R^2}, \quad (4.2)$$

$$d_{iB} = k_{ti}^{2p}$$

where $\Delta_{ij}^2 = (y_i - y_j)^2 + (\phi_i - \phi_j)^2$ and k_{ti} is the transverse momentum of particle i. p is a new parameter which signifies the relative power of energy versus geometrical scales (Δ_{ij}). If $p = 1$ the algorithm follows one known as k_t , if $p = 0$ the algorithm is Cambridge/Aachen, and if $p = -1$ the anti- k_t algorithm results. One notable characteristic of this algorithm is its resilience to soft radiation while maintaining collinear safety, which is why it's the prevailing jet algorithm used on the ATLAS experiment [31].

Jet reconstruction and isolation doesn't end with particle flow and the anti- k_t algorithm. Further corrections have to be applied to model data with Monte Carlo simulations. MC simulation are calibrated to better model pile-up and to improve jet angular resolution. Global sequential calibrations are calculated with MC using calorimeter, track, and muon chamber information to improve jet resolution. Following this, data is used to constrain uncertainties with through known dijet samples. These calibrations lead to final measurements of jet energy scale (JES) and jet energy resolution (JER) and provide reconstructed jet events in MC and data as well as a number recommended uncertainties for physics analyses to use. Figures 4.3 show jet energy scale and resolution as a function of p_T for PFlow jets modeling with the anti- k_t algorithm after JES corrections are applied. Fully combined systematic uncertainties are also shown for each [32].

Jet reconstruction and calibration are crucial to this analysis and later chapters will detail how impactful jet energy scale and resolution uncertainties are to the overall precision of the




Figure 4.3: a.) data-to-simulation ratio for average jet p_T as a function of jet p_T . Three techniques shown as well as their their combination (black) and the combination total uncertainty. PFlow Anti- k_t jets with $R=0.4$ and JES correction. b.) relative jet energy resolution as a function of p_T for anti- k_t PFlow jets with $R = 0.4$ in 2017. JES calibrations applied and compared to MC with full systematic uncertainties [30].

$H \rightarrow WW$ cross-section measurement.

4.2 Missing transverse energy

Protons in collider experiments like ATLAS have momentum solely in the plane of the beam. Conservation of momentum implies that in the plane transverse to the beam ($x - y$) the sum of momentum from all interaction by-products is zero. This is most often not the case and the non-zero transverse momentum from any interaction is termed E_T^{miss} . Missing transverse energy is a sign of final state neutrinos, which are massless and solely weakly interact so go undetected within ATLAS. E_T^{miss} could also point to new particles that cannot be directly detected like neutralinos or dark matter particles. E_T^{miss} could also signify interacting particles which evade detection in ATLAS due to detector acceptance or poor reconstruction [33].

CHAPTER 4. TRACKING AND ISOLATION IN ATLAS

Missing transverse energy is determined using a combination of all reconstructed particles in an event. This is challenging because it involves all detector components and final particle types. The Jet/ E_T^{miss} performance group delivers recommendations for multiple E_T^{miss} variables that we use in the $H \rightarrow WW$. This section will discuss E_T^{miss} reconstruction and performance followed by definitions of a few additional E_T^{miss} observables. Reconstructed E_T^{miss} calculations take into account both *hard* and *soft* event signals. Hard-events are composed of fully reconstructed and calibrated particles like electrons, muons, photons, τ -leptons and jets. Hard events are reconstructed charged-particle tracks from the hard-scatter vertex, all objects except those considered *soft*. Reconstruction for all particle types happen independently which means that the same signal may be used to identify two distinct particles. This double-counting is taken into account in E_T^{miss} resolution. At its most basic, E_T^{miss} is defined:

$$E_{x(y)}^{miss} = - \sum_{i \in \text{hard objects}} p_{x(y),i} - \sum_{j \in \text{soft objects}} p_{x(y),j} \quad (4.3)$$

where overall E_T^{miss} is a vector composed of x, y components. In order to avoid double-counting the same detector signal in multiple particle reconstruction algorithms, hard objects are consider in order: electrons, photons, hadronically decaying τ -leptons, and then jets. Muons have little overlap since they are reconstructed using MS and ID tracks (and muons alone leave tracks in the MS). As particles are reconstructed in this sequence, commonly used signals are rejected to avoid overlap. Another key variable is $\sum E_T$ which is defined

$$\sum E_T = \sum_{\text{electrons}} p_T^e + \sum_{\text{photons}} p_T^\gamma + \sum_{\tau\text{-leptons}} p_T^\tau + \sum_{\text{muons}} p_T^\mu + \sum_{\text{jets}} p_T^{jet} - \sum_{\text{unused tracks}} p_T^{track} \quad (4.4)$$

The first five terms show the hard term while the last represents the soft-term. Selections are applied to reconstructed jets and particles to achieve optimal E_T^{miss} for a particular analysis [34]. In the context of $H \rightarrow WW$ we use a “tight” configuration for E_T^{miss} classified by strict

CHAPTER 4. TRACKING AND ISOLATION IN ATLAS

conditions on accepted jets. This working point has the greatest pile-up rejection which is integral to our analysis.

E_T^{miss} reconstruction contains the complexity of each of its component parts and their p_T resolutions all affect total E_T^{miss} resolution. Pile-up and total event activity also play a large role in E_T^{miss} performance. Validations for E_T^{miss} are performed on a variety of observables and MC modelling is compared to reconstructed data. Systematic uncertainties are derived from comparing the reproducibility of these observables and their successful modelling of data. Resolution for reconstructed jets and leptons are also propagated to overall E_T^{miss} uncertainty. E_T^{miss} performance is evaluated using $Z \rightarrow \mu\mu$ or $Z \rightarrow e^-e^+$ and $W \rightarrow e\nu$ events, the first with no genuine E_T^{miss} and the second with significant E_T^{miss} from neutrinos. After specific event selection and E_T^{miss} reconstruction, each of these samples E_T^{miss} distributions is studied and data and MC demonstrated to match within uncertainties. Selected performance observables are shown for the full Run-2 sample set in 4.4. Here $Z \rightarrow e^+e^-$ events are selected and reconstructed track-based soft term E_T^{miss} and E_T^{miss} significance distributions are shown. Data and MC show good agreement.

In this analysis, a number of important E_T^{miss} variables are used. This selection will be further explained in later chapters, but these variables are used to cut on background (particularly from $Z \rightarrow \tau\tau$ events and to select for the signal $H \rightarrow WW \rightarrow \ell\nu\ell\nu$ channel where its two neutrinos require significant E_T^{miss}). Some of the variables used for these selections include track-based E_T^{miss} , track-based soft term E_T^{miss} , E_T^{miss} significance, $m_{\tau\tau}$, and p_T^{tot} . The E_T^{miss} soft term is composed of a track-based and calo-based component. The calorimeter-based component is highly dependent on pile-up and so using a track-based soft term reduces overall pile-up dependence. Track-based E_T^{miss} uses only reconstructed ID tracks from the primary vertex and in our analysis this has been just as powerful discriminant for isolating $Z \rightarrow \tau\tau$ background events from VBF signal as overall E_T^{miss} without the same

CHAPTER 4. TRACKING AND ISOLATION IN ATLAS




Figure 4.4: a.) Track-based Soft Term E_T^{miss} shown for the complete Run-2 dataset with an integrated luminosity of 140 fb^{-1} . Monte Carlo simulations are compared to data for the tight E_T^{miss} working point. b.) Track-based Soft Term E_T^{miss} significance modelling shown in the tight E_T^{miss} working point [30]

pile-up dependence and correlation with other kinematic variables. E_T^{miss} significance is newly defined and recommended variable from the Jet/ E_T^{miss} group. Described further in [33], this discriminates real missing energy from momentum resolution effects using a calculated likelihood. Shown in 4.4, E_T^{miss} significance peaks at low values if E_T^{miss} likely comes from resolution effects and not from a real invisible particle in the event. Our analysis also defines two variables based on $E_T^{miss}:p_T^{tot}$ describes the total transverse momentum from all hard objects in the event and $m_{\tau\tau}$ is defined

$$m_{\tau\tau} = \frac{m_{\ell\ell}}{\sqrt{x_1 * x_2}} \quad (4.5)$$

where

$$\begin{aligned} x_1 &= \frac{p_x^{\ell 0} * p_y^{\ell 1} - p_y^{\ell 0} * p_x^{\ell 1}}{p_y^{\ell 1} * E_{Tx}^{miss} - p_x^{\ell 1} * E_{Ty}^{miss} + p_x^{\ell 0} * p_y^{\ell 1} - p_y^{\ell 0} * p_x^{\ell 1}}, \\ x_2 &= \frac{p_x^{\ell 0} * p_y^{\ell 1} - p_y^{\ell 0} * p_x^{\ell 1}}{p_x^{\ell 1} * E_{Ty}^{miss} - p_y^{\ell 0} * E_{Tx}^{miss} + p_x^{\ell 0} * p_y^{\ell 1} - p_y^{\ell 0} * p_x^{\ell 1}} \end{aligned} \quad (4.6)$$

Each of these variables contributes to our signal region selection and elimination of background. A number of systematic uncertainties from reconstructed E_T^{miss} are defined and used in our analysis, though these are small compared to the uncertainties from jets and other final state particles.

4.3 Electrons

Accurate reconstruction and calibration of electrons within the ATLAS detector is integral to precision measurements, including the $H \rightarrow WW$ measurement in which an electron is required in the signal’s final state. Reconstruction, identification, and energy measurements of electrons and photons are the goals of the Electron and Photon Performance group. This section will summarize each of these calibration processes and their performances with a focus on electrons as they are particularly important in reconstructing final state particles for $H \rightarrow WW$.

Electrons are defined through energy deposits in the calorimeter, called superclusters, each with a matching track from the Inner Detector. Photons are defined strictly through a calorimeter cluster though “converted” photons can be matched to conversion vertexes in the tracker if they are matched to a leptonic initial state. Converted photons constitute a large fraction of reconstructed photons. Figure 4.6 shows the procedure for electron and photon reconstruction. First, topo-clusters in the EM calorimeter and tracks in the ID are

CHAPTER 4. TRACKING AND ISOLATION IN ATLAS

selected and matched together. Topo-clusters are defined based on signal to noise significance in calorimeter cells and calorimeter cell proximity. Standard reconstruction takes place in the inner detector and potential tracks are assigned to topo-clusters if their positions are within a region-of-interest compatible with that topo-clusters EM shower. Next, super-clusters are built from track-matched topo-clusters. Topo-clusters are tested for use as seed cluster candidates which begin super-clusters. Remaining topo-clusters are tested for compatibility as satellite clusters to each seed candidate. The resultant combination of seed and satellite clusters form super-clusters which are defined independently for photons and electrons. Finally, tracks are added to super-clusters, energy calibration and position corrections are applied, and analysis-level electrons and photons are created. Reconstruction efficiency for electrons is quite high, particularly at high p_T . Photon reconstruction efficiency is significantly lower due to their dependence solely on calorimeter clusters [35].




Figure 4.5: Electron and photon reconstruction algorithm [35]

Electron and photon energy resolution is determined from EM calorimeter showers and this is optimized with multivariate regression algorithms. Energy scale is also corrected using

CHAPTER 4. TRACKING AND ISOLATION IN ATLAS

calibration from well-known $Z \rightarrow ee$ decays and verified with other Z -boson decays. Similarly, these calibrations are calculated for photons using $Z \rightarrow \ell\ell\gamma$. Systematic uncertainties that effect these calibrations include passive material between the interaction point and the EM calorimeter and pile-up [35].

While electron and photon objects are already identified and reconstructed further ‘identification’ selections are used to improve electron and photon purity. Prompt electrons are identified with a likelihood discriminant which takes into account track parameters and electromagnetic shower properties. The primary electron track must stretch into the two inner tracking layers closest to the beam line and hit multiple points in the silicon-strip detector. The likelihood discriminant is calculated from probability density functions formed by smoothing histograms of 13 discriminating variables with an adaptive kernel density estimator. These are calculated separately for both the likelihood that an event reconstructs a prompt electron (signal) or that it does not (background). These likelihoods are derived from well-known $Z \rightarrow ee$ and $J/\Psi \rightarrow ee$ events recorded in Run-2. A tag-and-probe method was used to evaluate this likelihood- one electron in each decay must satisfy strict Run-1 likelihood discriminant requirements while the other serves as a probe of the new likelihood dscriiminator. Three electron working points are derived- Loose, Medium, and Tight, each with respectively lower efficiencies and higher purity. Physics analyses use the working points optimized for their analysis. For the $H \rightarrow WW$ measurement we use ‘Medium’ reconstructed electrons which have a lower efficiency than the loose condition but a factor $2\times$ greater rejection of background in electron identification. Figure ?? shows electron identification efficiency for each of the working points in a sample of $Z \rightarrow ee$ events where efficiency is calculated through comparisons to MC simulated $Z \rightarrow ee$ and $J/\Psi \rightarrow ee$ events. The average efficiencies for electroweak processes are 93%, 88% and 80% respectively for Loose, Medium, and Tight operating points and gradually increase at high E_T [35].

CHAPTER 4. TRACKING AND ISOLATION IN ATLAS




Figure 4.6: Electron reconstruction efficiency as a function of E_T and η in $Z \rightarrow ee$ events for Losse, Medium, and Tight electrons [35]

Track hits and calorimeter deposits near reconstructed electrons and muons can bias energy, momentum, and position measurements. Isolation performance are defined for calorimeter clusters by E_T^{cone} , the sum of transverse energy within a cone ΔR near a photon or electron cluster after correcting for leakage and pile-up effects. Track isolation p_T^{cone} is defined as a sum of the transverse momentum of tracks within a cone about the electron track or interpolated photon track. For electrons the distance between nearby decay products is directly related to electron p_T so a variable cone size can used such that

$$\Delta R = \min\left(\frac{10}{p_T}, \Delta R_{max}\right) \quad (4.7)$$

where ΔR_{max} is typically 0.2. Isolation working points strike a balance between efficiency and rejection of misidentified prompt electrons. The gradient working point is used by the $H \rightarrow WW$ analysis and gives an efficiency of 90% at $p_T = 25 \text{ GeV}$ and 99% at $p_T = 60 \text{ GeV}$. These values are reached through cuts on E_T^{cone20} and $p_T^{varcone20}$ derived from $J\Psi \rightarrow ee$ and $Z \rightarrow ee$ MC simulations. Efficiency of different isolation working points for electrons are shown in Figure 4.7. These are shown in Medium identified electrons for $Z \rightarrow ee$ events and

CHAPTER 4. TRACKING AND ISOLATION IN ATLAS

demonstrate the efficiencies possible with different cuts on the calorimeter and track-based isolation variables. The Gradient working point delivers reasonably high efficiency that is stable across η and is coupled with high background rejection of misidentified electrons [35].




Figure 4.7: Electron isolation efficiency shown for four working points in $Z \rightarrow ee$ events as a function of E_T and η [35]

Photon isolation is solely calorimeter-based. $Z \rightarrow \ell\ell\gamma$ events are used for photon isolation efficiency measurements and three working points are available which balance efficiency and background rejection just as in the electron case.

Electron reconstruction, calibration, identification, and isolation are all integral for the $H \rightarrow WW$ analysis in order to accurately identify electrons in the final state of candidate Higgs events. Systematic uncertainties from electron identification and isolation efficiency will be discussed further in later chapters and play a significant role in $H \rightarrow WW$ cross-section measurements.

4.4 Muons

Muons are abundant in the ATLAS detector and help lead to some of the most interesting physics results and analyses produced by the ATLAS experiment including $H \rightarrow WW \rightarrow$

CHAPTER 4. TRACKING AND ISOLATION IN ATLAS

$\ell\nu\ell\nu$ measurements which require a muon in the final state. The Muon Combined Performance group is tasked with producing the most accurate muon data for physics analyses. This includes muon reconstruction, identification, isolation, and analysis of efficiency, as well as muon momentum scale and resolution. The group’s goal is to create a number of “working points” tailored to different types of physics analyses that will isolate, identify, and reconstruct muons in the region of interest to the analyses. The working points are continuously updated and improved before being tested and implemented on different analyses. My work with the MCP group has focused on applying corrections necessary for muon momentum scale at the per mille level and resolution at the percent level in simulation/data. These are derived by a template fit of simulations smeared and corrected by variables in data. These corrections are validated by comparisons to simulations over a variety of variables.

Muon reconstruction is performed independently in the ID and MS and the information from these separate sub-detectors is combined to form full tracks. This section will focus on: 1) reconstruction in the ID, which is the same for any charged particle; 2) reconstruction in the MS, which is particular to muons; and 3) the combined reconstruction, which uses information from both the ID and MS.

In the ID, a pattern recognition algorithm reconstructs particle tracks with an inside-out sequence [36]. A track from a particle traversing the barrel typically has 3 pixel clusters, 8 SCT clusters and more than 30 TRT straw hits. The sequence begins by finding three-dimensional space points from the silicon hits. Each set of three space points which originate in the interaction point are used to trace hits up to the outer edge of the silicon detector. The final track parameters are fit through a collection of hits that extend to the TRT [37].

Muon reconstruction in the MS is not an inside-out procedure like that in the ID. Reconstruction begins with a search for hit patterns in each MS subdetector, which are called segments. The middle of the MS typically exhibits the largest number of trigger hits, there-

CHAPTER 4. TRACKING AND ISOLATION IN ATLAS

fore tracks are built by working out from the center of the MS and connecting segments layer-by-layer. Criteria such as hit multiplicity and fit quality determine track acceptance. At least two segments are needed to build a track. Hits associated with each track candidate are fitted using a global χ^2 fit. A track candidate is accepted if it passes the selection criteria [?].

The combined ID-MS reconstruction uses different algorithms to find different *muon types*. There are four main types outlined below, but preference-in terms of overlap between types-is given to Combined (CB), then Segment-tagged (ST), and finally Calorimeter Tagged (CT) muons. These algorithms have been continuously been improved to yield better precision, speed, and robustness against misidentification [38].

- **Combined muon (CB):** This type combines tracks from the ID and MS detectors using a global refit on all hits (some may be removed or added to improve quality). Most muons are reconstructed using an outside-in method.
- **Segment-tagged muon (ST):** ST muons are assigned an ID track that is associated with at least one local MDT or CSC track after extrapolation. These are used when muons cross only one layer of the MS because of low p_T or regions out of most MS layer boundaries.
- **Calorimeter-tagged muon (CT):** These muons are identified by an ID track that can be matched to a minimum ionizing particle energy deposit in the calorimeter. These muons have the lowest purity but are optimized for $|\eta| < 0.1$ and $1.5 < p_T < 100$ GeV where the MS is only partially instrumented.
- **Extrapolated muon (ME):** These muons are reconstructed in the MS with the addition of silicon points and with a loose requirement that the muon track originated

CHAPTER 4. TRACKING AND ISOLATION IN ATLAS

at the IP. In general, this muon is required to traverse $2 - 3$ layers of MS chambers. These are mainly used to extend acceptance for $2.5 < |\eta| < 2.7$, which is not covered in the ID.

In order to identify muons from other particles (like backgrounds from pion and kaon decays) strict quality requirements must be set to select prompt muons with high efficiency. Ideal signal muons are those that come from W decays (as opposed to light-hadron decays) and originate from the interaction point. We use a few variables to identify muons:

- q/p significance: The absolute value of the difference between the ratio of the charge and momentum of muons in the ID and MS divided by the sum in quadrature of their corresponding uncertainties.
- ρ' : The absolute value of the difference between the p_T measurements in the ID and MS divided by the p_T of the combined track.
- χ^2 : The normalized fit parameter of the combined track.

Specific requirements on the number of hits in the ID and MS assure that inefficiencies are expected and momentum measurements are robust. There are four muon identification selections that each addresses specific needs of physics analyses [38].

- **Loose Muons**: The *Loose* criteria maximizes the reconstruction efficiency, losing very few potential muons, while providing satisfactory tracks. All muon types are used in this criteria.
- **Medium Muons**: *Medium* is the default selection for muons in ATLAS because it minimizes systematic uncertainties associated with reconstruction and calibration. Only CB and ME tracks are used with requirements for over 3 hits in at least two MDT layers in most regions. All *Medium* muons are included in the *Loose* criteria.

CHAPTER 4. TRACKING AND ISOLATION IN ATLAS

- **Tight Muons:** *Tight* selects muons with the highest purity, but sacrifices efficiency. All *Tight* muons are included in the *Medium* selection, but only CB muons with at least two hits in the MS are considered, and the χ^2 value must be less than 8.
- **High- p_T Muons:** *High- p_T* muons have good momentum resolution for tracks with $p_T > 100$ GeV. This is beneficial to searches for high-mass Z' and W' resonances. CB muons in the *Medium* selection with at least 3 hits in 3 MS stations are included.

The $H \rightarrow WW$ analysis uses Tight muons with the added condition that $p_T > 15$ GeV and $\eta < 2.5$ to gain the highest purity possible and eliminate background from misidentified leptons, which constitute a significant background even with this selection.

We measure the muon reconstruction efficiency in two different ways in the regions $|\eta| < 2.5$ and $2.5 < |\eta| < 2.7$. First, in the barrel region, we use the **Tag-and-Probe** method. In this method we select an almost-pure sample of J/ψ and Z decays and require the leading muon to be a *Medium* muon labeled the **tag**. This muon fires the trigger. The subleading muon, the **probe**, must be reconstructed independently. There are three types of probes:

- **ID track:** Allows measurement of MS efficiency and of tracks not accessible to CT muons.
- **CT tracks:** Allows measurement of MS efficiency and has powerful rejection of background (especially at low p_T). This is the most commonly used probe.
- **MS tracks:** Allows measurement of ID and CT efficiency.

To find the overall efficiency of *Medium*, *Tight*, or *High- p_T* muons, we multiply the efficiencies associated with each type of probe. The efficiency $\epsilon(X|CT)$ ($X = Medium / Tight / High- p_T) of reconstructing these muons assuming a reconstructed ID track is measured$

CHAPTER 4. TRACKING AND ISOLATION IN ATLAS

using a CT muon as probe. This result is corrected by the efficiency $\epsilon(\text{ID}|\text{MS})$ of the ID track reconstruction measured using MS probes.

$$\epsilon(\text{X}|\text{ID}) \cdot \epsilon(\text{ID}) = \epsilon(\text{X}|\text{CT}) \cdot \epsilon(\text{ID}|\text{MS}) \quad (\text{X} = \text{Medium}/\text{Tight}/\text{High-}p_T) \quad (4.8)$$

The ID track reconstruction efficiency must be independent from the muon spectrometer track reconstruction ($\epsilon(\text{ID}) = \epsilon(\text{ID}|\text{MS})$). In addition, the use of a CT muon as a probe instead of an ID track must not affect the probability for *Medium*, *Tight*, or *High- p_T* reconstruction ($\epsilon(\text{X}|\text{ID}) = \epsilon(\text{X}|\text{CT})$). These assumptions are largely true with simulations showing some small deviations. These deviations are taken into account when calculating systematic errors.

The reconstruction efficiency of *Loose* muons is measured separately for CT muons within $|\eta| < 0.1$ and all other *Loose* types. The CT muon efficiency is measured using MS probe tracks, and the efficiency of other muons is evaluated similarly to the *Medium*, *Tight*, and *High- p_T* muons using CT probe muons [38]. For $|\eta| > 2.5$, the efficiency is calculated using the ME muons in the **Loose** and **Medium** selections. The number of muons observed in this region is normalized to the number of muons observed in the region $2.2 < |\eta| < 2.5$. A more detailed discussion of the efficiency measurement in this region can be found in Ref [39].

Scale factors are defined as the ratios between the efficiency of data and the efficiency of Monte Carlo simulations. They are used to describe the deviation between simulated and real detector behavior and are used in physics analyses to correct simulations.

Figure 4.8 displays reconstruction efficiency for *Medium* muons over a range of p_T and all other working plotted over η . While *Medium* muons have a higher efficiency than the *Tight* selection used in this analysis, both have an efficiency above 95% for a large range

CHAPTER 4. TRACKING AND ISOLATION IN ATLAS




Figure 4.8: On the left, reconstruction efficiency for *Medium* muons from $Z \rightarrow \mu\mu$ events is displayed as a function of the p_T of the muon in the region $0.1 < |\eta| < 2.5$ with systematic and statistical uncertainties. On the right, muon reconstruction efficiency is shown as a function of η in $Z \rightarrow \mu\mu$ events for muons with $pT > 15$ GeV for *Medium*, *Loose* and *Tight* muons. In both plots the full 139fb^{-1} Run-2 dataset is used[40].

of η and p_T . J/ψ decays probe low p_T muons while Z decays probe muons of a higher p_T allowing a large range to be defined. In addition, MC simulations match data quite well - within 1 – 2%. The only significant loss of efficiency is seen at extremely low η due to criteria excluding ID muons. Overall, the default *Tight* muon selection demonstrates a high reconstruction efficiency.

Isolating muons from heavy particles is one of the keys to understanding the background in many physics analyses. When heavy particles like W , Z , and Higgs bosons decay they often produce muons in isolation. Semileptonic decays, on the other hand, typically produce muons embedded in jets.

The MCP group uses two muon isolation variables: a track-based variable ($p_T^{varcone30}$) and a calorimeter-based variable ($E_T^{topocone20}$). $p_T^{varcone30}$ is defined as the scalar sum of the transverse momenta of tracks with $p_T > 1$ GeV in a cone around the muon of transverse momentum p_T excluding the muon track itself. The cone size is p_T -dependent to improve the performance for muons produced in decays with a large transverse momentum. $E_T^{topocone20}$ is

CHAPTER 4. TRACKING AND ISOLATION IN ATLAS

defined as the sum of the transverse energy of topological clusters in a cone around the muon after subtracting the contribution from the energy deposit of the muon itself and correcting for pile-up effects [41].

Table ?? defines seven isolation selection criteria - called “isolation working points” - that optimize different physics analyses. The *LooseTrackOnly* and *FixedCutTightTrackOnly* working points are defined by cuts on the relative track-based isolation variable. All other working points are defined by cuts applied separately on both relative isolation variables. All cuts are tuned as a function of the η and p_T of the muon to obtain a uniform performance. The target efficiencies of the different working points are described in Table 4.1. The efficiencies for the seven isolation working points are measured in data and simulation using the **Tag-and-Probe** method described previously on $Z \rightarrow \mu\mu$ decays. Figure 4.9 shows the isolation efficiency measured for *Medium* muons in data and simulation as a function of the muon p_T for two different working points. In both the *GradientLoose* and *FixedCutTightTrackOnly* working points, efficiency is above 90% and matches simulation well within errors for muons for higher p_T muons. In the *HWW* analysis, a fixed isolation cuts are optimized independent of Muon Performance working points. Only muons with $p_T > 15$ GeV are considered so the efficiency is optimized for muons which pass this and the *Tight* selection cuts.

The $H \rightarrow WW$ analysis uses muon isolation cuts $p_T^{varcone30}/p_T < 0.06\zeta$ and $p_T^{topocone20}/p_T < 0.09$ which are relatively tight selections and are selected to reduce misidentified lepton backgrounds.

The muon momentum scale and resolution are studied using Z and J/ψ decays. In order to obtain agreement between simulation and data in muon momentum scale to the per mille level and in resolution to the percent level, we need to apply a set of corrections to the simulated muon momentum. After applying the corrections we validate them by comparing

CHAPTER 4. TRACKING AND ISOLATION IN ATLAS

Isolation WP	Discriminating variable(s)	Definition
<i>LooseTrackOnly</i>	$p_T^{\text{varcone}30} / p_T^\mu$	99% efficiency constant in η and p_T
<i>Loose</i>	$p_T^{\text{varcone}30} / p_T^\mu, E_T^{\text{topocone}20} / p_T^\mu$	99% efficiency constant in η and p_T
<i>Tight</i>	$p_T^{\text{varcone}30} / p_T^\mu, E_T^{\text{topocone}20} / p_T^\mu$	96% efficiency constant in η and p_T
<i>Gradient</i>	$p_T^{\text{varcone}30} / p_T^\mu, E_T^{\text{topocone}20} / p_T^\mu$	$\geq 90(99)\%$ efficiency at 25 (60) GeV
<i>GradientLoose</i>	$p_T^{\text{varcone}30} / p_T^\mu, E_T^{\text{topocone}20} / p_T^\mu$	$\geq 95(99)\%$ efficiency at 25 (60) GeV
<i>FixedCutTightTrackOnly</i>	$p_T^{\text{varcone}30} / p_T^\mu$	$p_T^{\text{varcone}30} / p_T^\mu < 0.06$
<i>FixedCutLoose</i>	$p_T^{\text{varcone}30} / p_T^\mu, E_T^{\text{topocone}20} / p_T^\mu$	$p_T^{\text{varcone}30} / p_T^\mu < 0.15, E_T^{\text{topocone}20} / p_T^\mu < 0.30$

Table 4.1: The seven isolation working points are described by their discriminating variables and defining criteria [38].




Figure 4.9: Isolation efficiency for the GradientLoose (left) and FixedCutTrackOnly (right) muon isolation working points. The efficiency is displayed as a function of p_T in $Z \rightarrow \mu\mu$ events. The black markers show efficiency measured in data samples while the red show MC simulations. The bottom panel shows the ratio of the efficiency between the two as well as both statistical and systematic uncertainties [42].

the muon momentum scale and resolution between simulation and data over η , ϕ , and p_T .

We extract the calibration parameters with the transverse momentum of the ID and MS components of a CB track. The corrected transverse momentum is described by the following equation:

CHAPTER 4. TRACKING AND ISOLATION IN ATLAS

$$p_T^{\text{Cor,Det}} = \frac{p_T^{\text{MC,Det}} + \sum_{n=0}^1 s_n^{\text{Det}}(\eta, \phi)(p_T^{\text{MC,Det}})^n}{1 + \sum_{m=0}^2 \Delta r_m^{\text{Det}}(\eta, \phi)(p_T^{\text{MC,Det}})^{m-1} g_m}. \quad (4.9)$$

Here the g_m terms are normally distributed random variables with zero mean and unit width. The Δr and s terms describe momentum resolution smearing and scale corrections applied in specific detector regions, respectively. Both the ID and MS are divided into 18 pseudorapidity regions and the MS is divided into two ϕ bins separating the large and small sectors. Each of these bins leverages different alignment techniques and has different material distributions.

There are two s terms that represent different types of corrections. s_1 corrects for inaccuracy in the description of the magnetic field integral and the detector in the direction perpendicular to the magnetic field. s_0 corrects for the inaccuracy in the simulation of energy loss in the calorimeter and other materials. Since this loss is negligible in the ID, it is only nonzero in the MS [38].

The denominator introduces momentum smearing which broadens the p_T resolution in simulation. The parametrization of the smearing is defined:

$$\frac{\sigma(p_T)}{p_T} = r_0/p_T \oplus r_1 \oplus r_2 \cdot p_T. \quad (4.10)$$

In this equation r_0 is related to the fluctuations in energy loss in the traversed material, r_1 accounts for multiple scattering, local magnetic field inhomogeneities, and local radial displacements of hits, and r_2 describes intrinsic resolution effects caused by the spatial resolution of the hit measurements and by residual misalignment of the MS [38]. Correction parameters are extracted from data using a binned maximum-likelihood fit with templates derived from simulation which compares the invariant mass distributions for J/ψ and Z decay candidates in data and simulation. The muons are carefully selected to be compati-

CHAPTER 4. TRACKING AND ISOLATION IN ATLAS

ble with tracks that start at the interaction point and penetrate both the ID and the MS. Muons are also selected to pass specific momentum and isolation criteria. The dimuon mass distribution of these tracks in data is fitted using a Crystal Ball function convoluted with an exponential background distribution in the ID and MS fits. The background model and its normalization are then used in the template fit. The fits are performed in $\eta - \phi$ regions of fit (ROFs) which compromise regions with uniform features in the ID and MS [38].

From these fits, we can find the smearing terms across all η regions. Once the corrections are applied we can validate that the agreement between data and MC is excellent. This is shown in Figure 4.10. r_0 is set to zero across all η regions since energy loss is negligible in the ID. r_1 and r_2 increase as η increases since spatial resolution decreases and inhomogeneities increase as we move from the barrel to end-cap regions of both the ID and MS. Muon momentum corrections are continuously studied during ATLAS runs to validate muon calibration performance and account for discrepancies.




Figure 4.10: The r -values from each of 10 fits of resolution to p_T for ID muon simulations are shown. Each value corresponds to a particular ROF or η region. These plots show r_1 (left) and r_2 (right) as functions of leading muon η .

In this chapter, I have outlined the procedures for building all the component physics objects we use in the $H \rightarrow WW$ analysis. The successful reconstruction, calibration, and

CHAPTER 4. TRACKING AND ISOLATION IN ATLAS

isolation of jets, missing transverse energy, electrons, and muons is integral to the measurement of VBF Higgs bosons, as they decay to a final state with each of these components. As discussed, the measurement of physics objects includes their systematic and statistical uncertainties all of which are taken into account in our final measurement. The next chapters will use these physics objects and their kinematic variables to understand our background and signal events.

Chapter 5

Event Selection

This chapter steps briefly through some of the concrete object definitions and inputs used in the analysis. First the data and Monte Carlo simulations used for signal and background modelling are summarized. Then the kinematic variables and observables used in event selection and later for the differential measurement are defined. Finally, the signal selection is outlined and its results shown.

5.1 Data and Monte Carlo samples

5.1.1 Data samples

The full Run-2 dataset containing all proton-proton collision data collected from 2015–2018 at $\sqrt{s} = 13\text{TeV}$ with a 25ns bunch spacing configuration are used.

In 2015, 2016, 2017, and 2018 3.86 fb^{-1} , 35.6 fb^{-1} , 46.9 fb^{-1} , and 62.2 fb^{-1} of luminosity were recorded respectively. Peak instantaneous luminosity increased from $5.0 \times 10^{33}\text{ cm}^{-2}\text{s}^{-1}$ in 2015 to $21.4 \times 10^{33}\text{ cm}^{-2}\text{s}^{-1}$ in 2018. Average and peak pile-up also increased from about $\langle\mu\rangle = 13.6$ and 40.5 in 2015 to $\langle\mu\rangle = 37.0$ and 90 in 2018. These pile-up distributions

datasets are shown in 5.1.




Figure 5.1: The luminosity-weighted distribution of the mean number of interactions per crossing is shown for Run-2 pp collision data.

Events are only used if all relevant detector components are operating normally. These events are part of the standard “All Good” Good Run List comprising a total integrated luminosity of 139fb^{-1} and data quality efficiency of 91.5%.

Throughout Run-2 the instantaneous luminosity and so pile-up changed dramatically. This is accounted for in our MC modelling, as described in the next section.

5.1.2 Monte Carlo samples

Monte Carlo samples are generated to compare to data in order to test Standard Model predictions and search for excesses. These simulations are also used to optimize the analysis before applying these techniques to our dataset. Monte Carlo events are fully simulated using the ATLAS detector simulation in the GEANT4 framework [?] and reconstructed with standard ATLAS reconstruction software. Pile-up is simulated as additional pp interactions in a separate simulation step during digitization where minimum bias events are superimposed

CHAPTER 5. EVENT SELECTION

on the simulated signal events. These additional events are added based on that years recorded pile-up to account for dataset differences.

Separate programs are used to generate the hard scattering process and to model the parton showering (PS), hadronization, and the underlying event (UE). The next sections summarize the simulation techniques for our signal vector boson fusion samples, other Higgs production modes, and relevant backgrounds.

The fullset of data is split into three time-based categories. Data-taking conditions in 2015 and 2016 are averaged and described together as mc16a. 2017 and 2018 data-taking conditions are considered separately as mc16d and mc16e respectively.

Vector boson fusion Higgs samples

VBF Higgs events are generated through POWHEG [?] interfaced with PYTHIA 8 with the PDF4LHC15 parton distribution function (PDF) set [?]. Cross sections are calculated with full NLO QCD and EW corrections [? ?] with an approximate NNLO QCD correction applied [?]. These cross-sections as well as associated branching ratios are calculated by the LHC Higgs Cross Section Working Group Yellow Report 4 [?]. Generated events are normalized to calculated cross-sections.

Other production mode Higgs samples

Other Higgs production modes are considered in the analysis including gluon fusion (ggF), associated Higgs boson production (VH , $V = W, Z$) and Higgs boson production in association with a heavy quark pair ($t\bar{t}H$). Though only ggF has an appreciable yield in the signal region, each of these are studied and considered in the analysis. As in VBF Higgs sample, ggF, VH , and $t\bar{t}H$ are produced with PYTHIA8[? ?] for decay, parton shower, hadronisation and multiple parton interactions.

Background samples

Main Standard Model backgrounds include events from production of dibosons, top-quark, $Z+jets$, $W+jets$ and multijets. The WW samples are generated using SHERPA 2.2.2 interfaced with NNPDF3.0 NNLO PDFs and further described in [?]. $Z+jets$ (or Drell-Yan) production is simulated with SHERPA 2.2.1 using the NNPDF3.0 NNLO PDFs with dedicated parton shower tuning developed by Sherpa authors. Top-quark pair production ($t\bar{t}$) is simulated using POWHEG with the POWHEG-Box framework using the NNPDF 3.0 PDFs and interfaced with PYTHIA 8 using NNPDF 2.3 PDFs for parton showering. Single top, mainly Wt , production is generated with POWHEG-Box 2.0 interfaced to PYTHIA 6.428 for parton showering. $Z\gamma$ and $W\gamma$ productions are modeled using SHERPA 2.2.2 at the NLO accuracy for 0- and 1-jet. The $W+jets$ process modeling is based on a data-driven method described in the next chapter. MC samples (with the same MC generators for both $W+jets$ and $Z+jets$ samples) are used to validate the fake estimation and estimate sample composition uncertainties. These processes are generated with POWHEG MiNLO interfaced to PYTHIA 8 with the AZNLO tune.

5.2 Object definitions

This analysis utilizes a number of calibrated physics objects and the accurate use of each determines the precision of our $H \rightarrow WW$ differential cross section measurement. Lepton, jet, and missing transverse energy reconstruction, isolation, and calibration were described in some detail in the previous chapter. This chapter focuses on the particular parameters applied for these physics objects in this analysis. The object definitions are used in accordance with the $H \rightarrow WW$ coupling analysis for consistency where optimization

CHAPTER 5. EVENT SELECTION

focussed on $H \rightarrow WW$ measurements at large. Finally, further observables used in event selection are defined and described here as well.

5.2.1 Lepton

Our choice of lepton identification algorithm impacts both the rejection of fake lepton backgrounds ($W+jets$) and QCD background. Tighter requirements on lepton identification decrease signal efficiency so the optimal criteria balance high signal efficiency and high background rejection. Further studies on lepton identification and isolation criteria can be found in the HWW Coupling Support note [?].

Specific muon and electron requirements will be detailed next. Overall, all events must have at least two leptons tracks and each lepton track must have $p_T > 400\text{MeV}$. Further, leptons all are required to originate at the hard-scatter primary vertex which is defined as the primary vertex with the largest track $\sum p_T$. Leptons must pass two impact parameter selections to ensure they originate at the primary vertex. The longitudinal impact parameter of each lepton track is defined as $|z_0 \sin \theta|$ where z_0 is the impact parameter and θ the track angle. Each lepton longitudinal impact parameter must be less than 0.5mm. The significance of the transverse impact parameter is calculated with respect to the beam line ($|d_0|/\sigma_{d_0}$) and has a requirement of three (five) for muons (electrons) as recommended by the Muon and E/ γ combined performance groups.

Electron

Electron identification utilizes energy deposits in the calorimeter and reconstructed tracks in the inner detector. An electron likelihood combines these quantities and sets recommended working points (*Loose, Medium, Tight*) which correspond to increasing levels of background

CHAPTER 5. EVENT SELECTION

rejection. Further descriptions of the current E/γ combined performance group recommendations and methods can be found here [?].

This analysis uses both the *Medium* and *Tight* identification selections which have 94% and 88% identification efficiency respectively for an electron with $E_T = 100$ GeV. Electrons with $E_T > 25$ GeV must pass the *Medium* identification while those with $15 < E_T < 25$ GeV are required to pass the *Tight* selection. This mimics the selection for the $H \rightarrow WW$ coupling analysis where optimization studies demonstrated its impact on overall significance.

Electrons are identified in the range $|\eta| < 2.47$, where the transition region between barrel and endcaps in the LAr calorimeter ($1.37 < |\eta| < 1.52$) is excluded.

Electron isolation uses two different selections based on electron track p_T . For $p_T > 25$ GeV the *IsoGradient* working point is used. This requires calorimeter and track isolation about cones of $\Delta R = 0.2$ (*topoetcone20*, *ptvarcone20*). Electron isolation is changed to the fixed cut track cone isolation for $p_T < 25$ GeV to further eliminate fake background contributions. These require both track and calorimeter variables to fall below the p_T dependent $0.1143 \times p_T + 92.14$.

The following table summarizes electron selection criteria 5.1.

Table 5.1: Electron selections

p_T range	$ \eta $ range	Electron ID	Isolation	Impact parameter
< 25GeV	0 – 1.37, 1.52 – 2.47	Tight	FixedCutTrackCone40	$ z_0 \sin\theta < 0.5$, $ d_0 /\sigma_{d_0} < 5$
> 25GeV		Medium	IsoGradient	

Muon

As described previously, muons can be reconstructed using inner detector tracks, calorimeter deposits, and muon spectrometer tracks. The muon combined performance group performs an overall fit using each of these physical quantities and recommends six muon iden-

CHAPTER 5. EVENT SELECTION

tification working points- *VeryLoose*, *Loose*, *Medium*, *Tight*, *High- pT* , *Low- pT* . The combination is performed through an overall fit using the hits of the inner detector track, the energy loss in the calorimeter, and the hits of the track in the muon system. Muons in this analysis must fit the *Tight* selection as well as pass cuts $p_T > 15\text{GeV}$ and $|\eta| < 2.5$. Muon identification and isolation criteria match those used in the $H \rightarrow WW$ coupling analysis and optimize background rejection and signal efficiency. Muon isolation uses calorimeter and track isolation with a cone of $\Delta R = 0.2$ (*topoetcone20*, *ptvarcone20*). Muon selection criteria are summarized in the table below [5.2](#).

Table 5.2: Muon selections

p_T range	$ \eta $ range	Muon ID	Calo Isolation	Track Isolation	Impact parameter
$> 15\text{GeV}$	$ \eta < 2.5$	“Tight”	$E_T^{\text{cone}20}/p_T < 0.09$	$p_T^{\text{varcone}30}/p_T < 0.06$	$ z_0 \sin \theta < 0.5$, $ d_0 /\sigma_{d_0} < 3$

5.2.2 Jets

Jets constitute an important part of the analysis both in their number and characteristics. Our signal region selection and most control regions require at least two jets but in order to estimate and reject ggF Higgs background events, regions with less than two jets are also studied. “Tagging” jets refer to those considered in our jet requirement though others may be defined. As detailed in the previous chapter, jets reconstruction uses the anti- k_t algorithm to create jet tracks from calorimeter energy deposits within a cone of $R = 0.4$ and this analysis uses particle flow jet reconstruction.

Jets are required to have $p_T > 30\text{GeV}$ to eliminate high potential for pile-up jets in the full η range, $|\eta| < 4.5$. “Jet vertex tagger” variables suppress pile-up events through combinations of multiple variables into a single jet tagger. For jets with $p_T < 60\text{GeV}$ and $|\eta| < 2.4$, the JVT variable is required to be larger than 0.59. In the forward region a

CHAPTER 5. EVENT SELECTION

designated tagger called “fJVT” ought to be applied. However, the current analysis leaves out this requirement leading to more jet pile-up than expected.

The VBF signal region contains an additional jet observable, central jet veto (CJV), which cuts on events with additional jets (other than the two “tagged” jets) which have $p_T > 20\text{GeV}$ within the rapidity gap between two leading jets. These increase VBF sensitivity through removal of hadronic backgrounds.

***b*-tagged jet**

The Jet/ E_T^{miss} group recommends tools to identify jets formed from bottom quarks. While the previous HWW analysis used the MV2C10 jet tagging algorithm, the current analysis utilizes the DL1 tagger. Both tools use jet kinematics, impact parameters, and secondary vertex variables as input to machine learning classifiers. The DL1 tagger uses a neural network training method (the MV2C10 a boosted decision tree) trained with $t\bar{t}$ signal to discriminate b -quarks from light and c -quarks. The DL1 tagger shows greater b -veto efficiency in isolating top background events and so is used in this analysis. This optimization follows that performed for the HWW coupling measurement [?].

5.2.3 Missing transverse energy

Missing transverse energy is used to both suppress background and build other variables used in signal selection like m_{tt} and p_T^{tot} . E_T^{miss} definitions and calculation is explained in the previous chapter. This analysis uses the “Tight” E_T^{miss} working point which has proven the most robust against increasing pile-up.

5.2.4 Overlap removal

Overlap removal is applied to electrons, muons, and jets following current recommendations and using the official ASG tool [?]. Current removal steps can be summarized as followed based upon [? ?]:

- If a muon and electron share an ID track the electron is removed and if a calo-tagged muon shares an ID track with an electron, the muon is removed.
- If $\Delta R(\text{jet}, e) < 0.2$ the jet is removed (to eliminate overlap with the nearby electron).
The electron is removed if $\Delta R(\text{jet}, e) \geq \min(0.4, 0.04 + 10 \text{ GeV}/p_T^e)$.
- If $\Delta R(\text{jet}, \mu) < 0.2$ and the jet has less than three associated tracks with $p_T > 500\text{MeV}$ the jet is removed. The jet is also removed if the p_T ratio of the muon and jet is larger than 0.5 and the ratio of the muon p_T and the sum of the p_T of all jet tracks with $p_T > 500\text{GeV}$ is larger than 0.7. The muon is removed if $\Delta R(\text{jet}, \mu) \geq \min(0.4, 0.04 + 10 \text{ GeV}/p_T^\mu)$ for any surviving jets.

5.3 Common Observables

Dedicated observables are used in this analysis to reject and isolate backgrounds as well as to enhance VBF signal strength. This section lists and describe key observables used in the analysis.

- $m_{\tau\tau}$: This can be defined as in the previous chapter's E_T^{miss} section, if charged leptons are the products of a pair of τ leptons, the neutrinos are collinear with the charged leptons, and the neutrinos are the only source of the observed E_T^{miss} in the event. We cut on this variable to suppress $Z \rightarrow \tau\tau$ events.

CHAPTER 5. EVENT SELECTION

- N_{b-jet} : Defines number of jets with $p_T > 20$ GeV identified as b-jets from the b-tagged algorithm (DL1) and used to reject $t\bar{t}$ background.
- p_T^{ll} : Topological variable transverse momentum of the dilepton system. This cut targets Drell-Yan/ $Z \rightarrow \tau\tau$ events through a cut at 70 GeV.
- $\Delta\phi_{ll}$: The two leptons from HWW decays tend to be collimated especially compared to non-resonant WW backgrounds. This is due to the spin-zero initial state of the resonant process.
- m_{ll} : The invariant mass of the two leptons from the hard scattering interaction. Similar to p_T^{ll} , targets Drell Yan events and is cut on at 70GeV in the control region.
- m_T : Transverse mass is used as discriminant variable though not directly cut. Defined as

$$m_T = \sqrt{(E_{ll} + E_T^{miss})^2 - |p_{ll} + E_T^{miss}|^2} \quad (5.1)$$

where $E_{ll} = \sqrt{|p_{ll}|^2 + m_{ll}^2}$.

- p_T^{tot} : Total transverse momentum p_T^{tot} , defined in the targeted E_T^{miss} chapter. Aids in rejecting events with significant soft gluon radiation but not high p_T jets.
- ΔY_{jj} : VBF Higgs signal events are characterized by a large separation of the two tagging jets in rapidity. This variable is used as a signal region cut.
- m_{jj} : VBF Higgs events tend to have a high jet invariant mass (m_{jj}), defined by combining mass of tagged jets. This analysis applies a cut on m_{jj} in our signal region.
- η_{lep} centrality: This analysis uses an outside lepton veto defined to reject events with leptons outside the rapidity gap between the two tag jets. This OLV variable is defined

CHAPTER 5. EVENT SELECTION

using η of the tagged jets and leptons as follows:

$$\begin{aligned} \text{OLV}_{l_0} &= 2 \cdot \left| \frac{\eta_{l_0} - \bar{\eta}}{\eta_{j_0} - \eta_{j_1}} \right| \\ \text{OLV}_{l_1} &= 2 \cdot \left| \frac{\eta_{l_1} - \bar{\eta}}{\eta_{j_0} - \eta_{j_1}} \right| \\ \eta_{\text{lep centrality}} &= \text{OLV}_{l_0} + \text{OLV}_{l_1} \end{aligned} \quad (5.2)$$

where $\bar{\eta} = (\eta_{j_0} + \eta_{j_1})/2$ and so for each lepton:

$$\text{OLV}_l \begin{cases} = 0 & : \text{the lepton is right in the middle of the rapidity gap between the two tag jets.} \\ < 1 & : \text{the lepton lies within the rapidity gap between the two tag jets.} \\ > 1 & : \text{the lepton is outside the rapidity gap between the two tag jets.} \end{cases} \quad (5.3)$$

- $\sum_{l,j} M_{lj}$: The sum of the invariant masses of all possible lepton-jet pairs are used to train our signal discriminant as VBF Higgs signal peaks at a higher value than the dominant backgrounds. VBF signal jets tend to be very forward while lepton central so a large separation between lepton and jets is expected as opposed to typical background topologies.

5.4 Event selection

Figure 5.2 shows the Feynman diagram for the vector boson fusion Higgs production mode. Two energetic jets with large separation in rapidity accompany the Higgs and its decay products. In addition, since the Higgs is created through the fusion of two electro-weak bosons, there is no QCD interaction between the jets and the Higgs boson or its decay products. The main backgrounds this analysis seeks to mitigate include top quark

CHAPTER 5. EVENT SELECTION

production, Drell-Yan/ Z , di-bosons (WW , WZ , ZZ , and $W\gamma$), gluon-gluon fusion (ggF) Higgs production, and $W+jets$.




Figure 5.2: Feynman diagram for VBF Higgs production

This analysis controls and estimates these backgrounds with a variety of methods depending on characteristics of each background. Major backgrounds are estimated in CRs and minor backgrounds are estimated using MC simulation. Top quark production is suppressed in the SR by a b -jet veto requirement, and can be studied in a very pure orthogonal set of bins in the signal region defined by a boosted decision tree discriminant. A pure validation region is also defined to cross-check top MC modelling. The Drell-Yan/ Z background is suppressed by kinematic requirements and is constrained in a dedicated and orthogonal CR and ultimately normalized in the signal phase space. The ggF Higgs production mode is particularly difficult to separate from the VBF production mode, therefore several CRs are defined to constrain it in data, and ultimately its normalization is determined in the signal phase space. The $W+jets$ background is estimated with data-driven techniques, as described in the next chapter's section on fake estimation, and then fixed in the simultaneous fit of SR and CRs.

The analysis employs multivariate analysis (MVA), more specifically 2D boosted decision trees trained to discriminate between samples. Specifically there are six trained BDTs used to separate: $WW+top$ and VBF events, ggF events from all other samples in each of their three control regions, $WW+top$ events and VBF , and WW and top events from one another. The use of these BDTs separates the backgrounds from our signal in the signal region and

CHAPTER 5. EVENT SELECTION

maintains discrimination between the two WW and top background samples. The $WW + \text{top}$ vs. VBF BDT is discussed in this section while the others are explained within in the next dedicated background chapter. In addition, two other types of BDT have been tested in this analysis: a $Z + \text{jets}$ vs. VBF BDT and a 3D BDT to separate ggF , WW and VBF events. Results from these tests showed that though these led to minimization of background (in the case of $Z + \text{jets}$ vs. VBF) and good discrimination between key backgrounds in the signal region (in the case of the 3D BDT), when used in the final fit with systematic uncertainties they showed no significant decrease in calculated error in the VBF coupling value. Results from these studies are shown in the Appendix.

Background	Relative size Pre-selection	Relative size Signal Region	Estimation	Generator
WW	$\sim 8\%$	$\sim 20\%$	MC only + BDT	SHERPA 2.2.2 ($gg \rightarrow WW$: SHERPA 2.1)
Top	$\sim 73\%$	$\sim 61\%$	Data+MC (Top CR)	POWHEG+PYTHIA 8
ggF Higgs	$< 1\%$	$\sim 2\%$	MC only + BDT	POWHEG+PYTHIA 8 NNLOPS
$W + \text{jets}$	$\sim 2.5\%$	$\sim 4\%$	Data-driven	-
$Z \rightarrow \tau\tau$	$\sim 16\%$	$\sim 7\%$	Data+MC ($Z \rightarrow \tau\tau$ CR)	SHERPA 2.2.1
$W\gamma, W\gamma^*$	$\sim 1\%$	$\sim 2.5\%$	MC only	SHERPA 2.1/SHERPA 2.2.2

Table 5.3: Each background and its relative size of in SR (after all cuts), estimation method, and generator.

5.4.1 Pre-selection

The VBF differential analysis shares object definitions with the VBF and ggF coupling analysis and pre-selection cuts are based off of those used in the 2016 ggF and VBF $H \rightarrow WW$ 36.1fb^{-1} cross-section paper [?]. The VBF selection utilizes observables defined in the previous section. Cuts applied in the preselection region are listed and described in the table below.

Yields for all samples as well as data for pre-selection cuts are shown in the following table.

CHAPTER 5. EVENT SELECTION

Pre-sel cut	Description
Channel Selection	Fits $e\mu/\mu e$ channel/event weight applied
Trigger Selection	Trigger selected
Trigger Matching	Trigger weight applied
Data driven muon/electron	W+jets split into electron and muon fake flavors
Cut Jet Cleaning	Pass loose jet cleaning
Cut $V\gamma/Z$ +jets overlap	Eliminate any Z +jets events that overlap with $V\gamma$
Only two leptons	Require exactly 2 leptons (in case of fakes make sure 1 ID/1 Anti-ID)
Lead lepton p_T	$p_T^{lead} > 22\text{GeV}$
Sublead lepton p_T	$p_T^{sublead} > 15\text{GeV}$
Opposite sign leptons	Require zero dilepton charge
m_{ll} selection cut	$m_{ll} > 10\text{GeV}$
Fake Factor	W+jets weight applied

Table 5.4: Table describing pre-selection cuts applied in common with VBF and ggF coupling analyses

$\sqrt{s} = 13\text{TeV}, \mathcal{L} = 139\text{fb}^{-1}$ (Full Run 2)	H_{VBF}	H_{soft}	WW	Other VV	Top	Z_{jets}	Mis-Id	Total Bkg	Significance	Data	Data/MC
Channel Selection	904.81 ± 0.92	9240.38 ± 10.13	175857.28 ± 140.42	22386.99 ± 78.25	1709307.70 ± 287.98	655027.79 ± 1233.45	5008168.16 ± 4730.67	7579988.31 ± 4899.95	0.33 ± 0.00	4374979	0.58 ± 0.00
Trigger Selection	904.81 ± 0.92	9240.38 ± 10.13	175857.28 ± 140.42	22386.99 ± 78.25	1709307.70 ± 287.98	655027.79 ± 1233.45	5008168.16 ± 4730.67	7579988.31 ± 4899.95	0.33 ± 0.00	4374979	0.58 ± 0.00
Trigger Matching	882.26 ± 0.90	8940.07 ± 9.82	173094.52 ± 138.41	21501.64 ± 77.38	1678206.02 ± 283.57	624251.33 ± 1168.16	5151805.35 ± 4644.01	7657798.93 ± 4709.69	0.32 ± 0.00	4352644	0.57 ± 0.00
W+jets flavour split muon	882.26 ± 0.90	8940.07 ± 9.82	173094.52 ± 138.41	21501.64 ± 77.38	1678206.02 ± 283.57	624251.33 ± 1168.16	4675498.22 ± 4228.78	7181491.80 ± 4399.19	0.33 ± 0.00	4352644	0.61 ± 0.00
W+jets flavour split electron	882.26 ± 0.90	8940.07 ± 9.82	173094.52 ± 138.41	21501.64 ± 77.38	1678206.02 ± 283.57	624251.33 ± 1168.16	3625576.16 ± 3693.41	6131569.74 ± 3887.35	0.36 ± 0.00	4352644	0.71 ± 0.00
Jet Cleaning	882.26 ± 0.90	8940.07 ± 9.82	173094.52 ± 138.41	21501.64 ± 77.38	1678206.02 ± 283.57	624251.33 ± 1168.16	3625576.16 ± 3693.41	6131569.74 ± 3887.35	0.36 ± 0.00	4352644	0.71 ± 0.00
Overlap: Vgamma/Vjets	882.26 ± 0.90	8940.07 ± 9.82	173094.52 ± 138.41	21501.64 ± 77.38	1678206.02 ± 283.57	624251.33 ± 1168.16	3625576.16 ± 3693.41	6131569.74 ± 3887.35	0.36 ± 0.00	4352644	0.71 ± 0.00
Only two Leptons	880.17 ± 0.89	8934.45 ± 9.82	172870.71 ± 138.36	20467.40 ± 77.14	1662432.27 ± 282.33	622191.46 ± 1160.36	3621793.98 ± 3681.91	6108690.27 ± 3873.99	0.36 ± 0.00	4331979	0.71 ± 0.00
$p_T^{lead} > 22\text{ GeV}$	880.17 ± 0.89	8934.45 ± 9.82	172870.71 ± 138.36	20467.40 ± 77.14	1662432.27 ± 282.33	622191.46 ± 1160.36	3621793.98 ± 3681.91	6108690.27 ± 3873.99	0.36 ± 0.00	4331979	0.71 ± 0.00
$p_T^{sublead} > 15$	879.98 ± 0.89	8933.88 ± 9.82	172854.38 ± 138.35	20409.32 ± 77.10	1661877.68 ± 282.29	622068.28 ± 1160.19	3620023.39 ± 3681.17	6106166.94 ± 3873.23	0.36 ± 0.00	4330240	0.71 ± 0.00
OS Leptons	879.86 ± 0.89	8933.68 ± 9.82	172771.33 ± 134.49	20041.83 ± 38.48	1661388.47 ± 282.25	621966.33 ± 1160.06	3614469.54 ± 3678.75	6099572.24 ± 3870.18	0.36 ± 0.00	4326784	0.71 ± 0.00
$M_{ll} > 12/10\text{ GeV}$	879.85 ± 0.89	8933.66 ± 9.82	172771.33 ± 134.49	20041.83 ± 38.48	1661368.39 ± 282.25	621964.49 ± 1160.06	3614488.72 ± 3678.68	6099268.41 ± 3870.11	0.36 ± 0.00	4326642	0.71 ± 0.00
SF: Z Veto	879.85 ± 0.89	8933.66 ± 9.82	172771.33 ± 134.49	20041.83 ± 38.48	1661368.39 ± 282.25	621964.49 ± 1160.06	3614488.72 ± 3678.68	6099268.41 ± 3870.11	0.36 ± 0.00	4326642	0.71 ± 0.00
Leptons ID, singleFakes 1 anti-ID, doubleFake 2 anti-ID	596.53 ± 0.74	5630.33 ± 7.80	126996.63 ± 115.72	11152.41 ± 25.92	1165160.56 ± 237.51	257221.77 ± 444.61	1150777.06 ± 1622.64	2716938.76 ± 1703.28	0.36 ± 0.00	1587474	0.58 ± 0.00
Apply QCD FF weight	596.53 ± 0.74	5630.33 ± 7.80	126996.63 ± 115.72	11152.41 ± 25.92	1165160.56 ± 237.51	257221.77 ± 444.61	36980.51 ± 255.27	1603142.22 ± 577.39	0.47 ± 0.00	1587474	0.99 ± 0.00
Apply fake factor	596.53 ± 0.74	5630.33 ± 7.80	126996.63 ± 115.72	11152.41 ± 25.92	1165160.56 ± 237.51	257221.77 ± 444.61	36980.51 ± 255.27	1603142.22 ± 577.39	0.47 ± 0.00	1587474	0.99 ± 0.00

Table 5.5: Cutflow in the pre-selection region.

The following plots show kinematic distributions after all preselection cuts with fake factors applied directly after a 2-jet cut. These show good modelling with data over a variety of kinematic variables including those used in the Top+ WW vs. VBF BDT. We are not able to directly examine modelling of input variables to this BDT because we require blinding in the signal region. Modelling in the pre-selection region is studied in lieu of this and shows no evidence of bias.

CHAPTER 5. EVENT SELECTION




Figure 5.3: Distributions of ΔY_{jj} , $\Delta Y_{\ell\ell}$, $\Delta \Phi_{\ell\ell}$, m_{jj} , $m_{\ell\ell}, m_T$, jet p_T^{lead} , jet p_T^{sublead} , and $\Delta \Phi_{jj}$ in the preselection region before blinding is applied. Distributions show good MC modelling of variables used in the signal region BDT.

5.4.2 Signal region selection

In addition to the pre-selection cuts described, a number of cuts are applied to the VBF signal region which differ from those used in the coupling ggF analysis. These cuts include a requirement for at least 2 jets ($n_{jets} \geq 2$) and a b -veto using the DL1r b -tagging algorithm. A central-jet-veto (CJV) and an outside-lepton-veto (OLV) are also applied. These cuts remove first events with $p_T > 20\text{GeV}$ which lie between the tagging jets in pseudo-rapidity and next any where the two charged leptons are not within the tag jets' rapidity gap. Two additional cuts that differ from the VBF HWW couplings analysis are also added. These are cuts on mass of the two jets ($m_{jj} > 200\text{GeV}$) and on the rapidity difference between the two jets ($DY_{jj} > 2.1$). These further purify the signal region against a range of backgrounds, notably top. The plots below show signal and background yields as well as signal significance at a variety of m_{jj} and ΔY_{jj} values. The values used here are chosen for their effects on signal significance while retaining high signal statistics.




Figure 5.4: Signal and background yields for VBF and total background (aside from fakes) at various m_{jj} and ΔY_{jj} cut values. MC simulation for mc16a campaign only. Choice of cut at $m_{jj} > 200\text{GeV}$ and $DY_{jj} > 2.1$ nearly halves background yields while reducing signal by $< 5\%$

Signal region cuts are listed and described in the table below.

Yields for all samples as well as data for signal region cuts are shown in the following table.

The following plots show kinematic distributions after all signal region cuts. Here only

CHAPTER 5. EVENT SELECTION

Signal region cut	Description
2-jet (30,30)	Require at least 2 jets with $p_T \geq 30\text{GeV}$
b-veto	Use DL1 efficiency b-tag reject events with b-jets/apply b-tag weight
Blinding (VBF)	Blinding condition on data (cut on VBF vs. Top+WW BDT)
CJV (20GeV)	Cut events with a third central rapidity jet $p_T > 20\text{GeV}$
OLV bool	Leading lepton η required to be between two η 2 leading jets
$Z \rightarrow \tau\tau$ veto	$m_{\tau\tau} < m_Z - 25\text{GeV}$
m_{jj} cut	$m_{jj} > 200\text{GeV}$
ΔY_{jj} cut	$\Delta Y_{jj} > 2.1$

Table 5.6: Table describing VBF signal region cuts

$\sqrt{s} = 13\text{TeV}$, $\mathcal{L} = 139\text{fb}^{-1}$ (Full Run 2)	H_{VBF}	H_{ggF}	WW	Other VV	Top	Zjets	Mis-Id	Total Bkg	Significance	Data	Data/MC
Scale factors					NF = 0.99 ± 0.01	NF = 1.03 ± 0.05					
2-jet (30,30) DJVT	366.47 ± 0.58	1061.66 ± 3.08	24353.25 ± 30.46	3459.16 ± 10.99	910264.96 ± 201.55	26791.10 ± 110.14	12382.41 ± 170.01	978312.54 ± 287.60	0.37 ± 0.00	976091	1.00 ± 0.00
Scale factors					NF = 0.99 ± 0.01	NF = 1.03 ± 0.05					
b-veto	323.67 ± 0.54	914.43 ± 2.86	21075.69 ± 28.85	2701.86 ± 10.03	63729.30 ± 57.17	22477.17 ± 104.35	4407.46 ± 70.33	115305.91 ± 141.58	0.95 ± 0.00	109677	0.95 ± 0.00
Scale factors					NF = 0.99 ± 0.01	NF = 1.03 ± 0.05					
blinding (2-jet)	323.67 ± 0.54	914.43 ± 2.86	21075.69 ± 28.85	2701.86 ± 10.03	63729.30 ± 57.17	22477.17 ± 104.35	4407.46 ± 70.33	115305.91 ± 141.58	0.95 ± 0.00	86168	0.75 ± 0.00
Scale factors					NF = 0.99 ± 0.01	NF = 1.03 ± 0.05					
CJV (20GeV)	256.58 ± 0.48	668.26 ± 2.45	15050.73 ± 24.95	1919.48 ± 8.97	43095.38 ± 47.58	16399.61 ± 90.71	3078.82 ± 57.99	80212.29 ± 120.68	0.91 ± 0.00	61566	0.77 ± 0.00
Scale factors					NF = 0.99 ± 0.01	NF = 1.03 ± 0.05					
OLV bool	199.59 ± 0.43	182.05 ± 1.29	2741.46 ± 11.72	410.62 ± 4.92	9409.54 ± 22.24	3719.07 ± 45.98	539.20 ± 25.93	17001.94 ± 58.69	1.53 ± 0.00	10412	0.61 ± 0.01
Scale factors					NF = 0.99 ± 0.01	NF = 1.03 ± 0.05					
$Z \rightarrow \tau\tau$ veto	172.20 ± 0.40	161.62 ± 1.22	1701.95 ± 9.30	239.95 ± 4.33	6051.92 ± 17.81	1351.94 ± 36.47	358.93 ± 19.82	9866.31 ± 46.34	1.73 ± 0.01	5448	0.54 ± 0.01
Scale factors					NF = 0.99 ± 0.01	NF = 1.03 ± 0.05					
$M_{jj}, 200$	165.51 ± 0.39	110.73 ± 1.00	1160.80 ± 8.03	156.65 ± 4.00	3565.45 ± 13.71	870.82 ± 31.48	233.45 ± 15.88	6097.90 ± 38.90	2.11 ± 0.01	3021	0.48 ± 0.01
Scale factors					NF = 0.99 ± 0.01	NF = 1.03 ± 0.05					
$DY_{jj}, 2,1$	163.07 ± 0.39	101.90 ± 0.97	1027.28 ± 7.90	133.53 ± 3.94	3003.00 ± 12.73	804.53 ± 31.16	206.57 ± 14.59	5276.81 ± 37.75	2.23 ± 0.01	2513	0.46 ± 0.01
Scale factors					NF = 0.99 ± 0.01	NF = 1.03 ± 0.05					
BDT Zjets 0.5	153.15 ± 0.37	92.91 ± 0.92	950.37 ± 7.52	116.48 ± 3.84	2845.05 ± 12.39	337.73 ± 20.62	173.00 ± 13.09	4515.55 ± 28.68	2.27 ± 0.01	2334	0.50 ± 0.01

Table 5.7: Cutflow in the signal region.

MC predictions are shown since data is blinded in the signal region.

VBF signal boosted decision tree discriminant

Finally, this analysis uses a number of boosted decision trees (BDTs) to both amplify our VBF signal and discriminate and reject a number of specific backgrounds. In the next chapter each major background will be described along with their control or validation regions and the BDTs discriminants trained and used in the overall fit. This next section will focus on the BDT trained and used in the signal region to isolate the VBF signal from dominant backgrounds (top and WW events). This discriminant is the results of numerous studies as to the best training parameters, input variables, and multivariate analysis techniques to increase discrimination. Appendix A shows some results from studies on using a multidimensional

CHAPTER 5. EVENT SELECTION




Figure 5.5: Distributions of $\Delta Y_{\ell\ell}$, $\Delta\Phi_{\ell\ell}$, $m_{\ell\ell}$, m_T , p_{tot}^T , lep p_{lead}^T , lep p_{sublead}^T , jet p_{lead}^T , jet p_{sublead}^T , $\Delta\Phi_{jj}$, and $E_{\text{T},\text{rel}}^{\text{miss}}$ in the differential VBF signal region, many used as input to the BDT discriminating VBF from top + WW backgrounds.

CHAPTER 5. EVENT SELECTION

dimensional BDT to simultaneously discriminate VBF, ggF, and WW background samples. While initially this showed promising results, estimation of ggF backgrounds through use of multiple control regions (summarized in the next chapter) showed better determination of the ggF background than the 3D BDT and so the one-dimensional method here was adopted.

A decision tree is a collection of cuts designed to classify events as signal-like or background-like. A given signal event is correctly identified if it is placed in a signal-dominated leaf and vice-versa for background events. After the initial tree is built another tree is grown to better separate the signal and background events misidentified by the first tree. This proceeds iteratively until there is a collection of a specified number of trees, in a process known as boosting. A weighted average is taken from all these trees to form a BDT output discriminant with values ranging from -1 to 1.

This BDT is trained using $e\mu + \mu e$ events after the VBF selection and the signal regions cuts including that on n_{jets} , b -veto, OLV, CJV, M_{jj} and DY_{jj} . In this way, the phase space in which we train the BDT is exactly the same as the one where we apply it. The training includes only the top and WW backgrounds and the VBF signal. The MC statistics used in the training are half those available after all signal region cuts (as the other half are later used to test the training). This corresponds to $\approx 90,000$ un-weighted WW and top events and $\approx 100,000$ raw VBF events. This training includes MC weights on events to best account for overall event distributions. There are ≈ 2000 total weighted top and WW events used in the training and ≈ 80 weighted VBF events.

The TMVA BDTG interface is used to train and test the BDT. The optimal parameters were found through a scan of reasonable values and the final set is summarized in Table 5.8.

This BDT utilizes a wide range of lepton and jet kinematic variables (12) to distinguish between signal and background events. These include ΔY_{jj} , $\Delta Y_{\ell\ell}$, $\Delta\Phi_{\ell\ell}$, m_{jj} , $m_{\ell\ell}$, m_T , η_{j0} ,

CHAPTER 5. EVENT SELECTION

Parameter	Value
Boosting algorithm	Gradient
Maximum tree depth	22
Number of trees	400
Minimum number of events requires per mode	5%
Number of cuts	7

Table 5.8: BDT parameters used for the VBF vs. top + WW training.

η_{j1} , p_{j0}^T , p_{j1}^T , $\Delta\Phi_{jj}$, and \sum centralities (L). While a larger variety of variables have been tested, these demonstrated the highest discrimination between VBF and top/ WW background. Plots shown in 5.6 and 5.7 demonstrate the input distributions used to train the BDT and their correlations. The BDT training successfully separates VBF signal and top/ WW




Figure 5.6: Distributions of input variables to VBF vs. top+ WW BDT. Samples are weighted and normalized to even numbers of background and signal events. Signal represents VBF and background top+ WW .

background. These backgrounds are considered together both because they have very similar signatures compared to the VBF signal and because our overall fit uses one parameter to estimate both WW and top backgrounds together. In order to quantify the discrimination we use the integrated-ROC calculated through TMVA for weighted normalized samples and find an optimal value of 0.960. Comparisons between the test and training show that the BDT is un-biased- differences between testing and training samples would imply overtraining, or the BDT using too many parameters on too few events. Visually, once can see that the

CHAPTER 5. EVENT SELECTION




Figure 5.7: Correlations of input variables to VBF vs. top +WW BDT. Signal represents VBF and background top+WW.

testing and trainings samples are quite similar. Additionally, a Kolmogorov-Smirnov test is performed to measure if the two test and training distributions differ significantly. If the two distributions are random samples of the same parent distribution, the KS-test would give a uniformly distributed value between zero and one (or an average value of 0.5). The closer to 0.5 the KS-test, the greater likelihood the curves come from the same parent, however this calculation is heavily skewed toward lower values so any value above zero (or not very close to zero, on order 10^{-4}) can be considered not indicative of overtraining. For signal and background we find KS-test values of 0.107 and 0.154, and so no evidence of over-training. We can visualize the BDT output variable both on un-weighted normalized samples and on samples with all event weights applied. The following plot shows BDT results applied to normalized samples of VBF signal and top/WW backgrounds.

We aim to fit this distribution in the signal region with high significance in uppermost bins of the distribution. Since this BDT is trained and applied in the signal region we cannot directly test the modelling of input variables. However, modelling at the pre-selection level for each of these variables, shown earlier in the chapter show no evidence of mis-modelling.

CHAPTER 5. EVENT SELECTION




Figure 5.8: Unweighted, normalized samples of VBF (signal) and top+WW samples (background) plotted over BDT output distribution on left, overlaid testing and training samples shown. Right, full weighted samples of VBF signal and all backgrounds plotted over BDT output distribution after signal region selection. Data is blinded here

The binning for this discriminant used in the statistical fit and its result (using Asimov data) are shown in the final chapter.

Chapter 6

Backgrounds and Systematics

6.1 Backgrounds

While the VBF $H \rightarrow WW \rightarrow \ell\nu\ell\nu$ channel gives a fairly clear signal with E_T^{miss} , two leptons, and 2 jets, there are substantial backgrounds with similar final states. These include Drell-Yan processes (in which a Z boson is formed from quark anti-quark annihilation of the two colliding protons), top quark final states (predominantly $t\bar{t}$), diboson events (led by SM WW events), Higgs decays from the other production modes (mainly ggF), and background from mis-identified leptons, called fakes. In this section I will detail each background and describe our methods for estimating their role in overall results. I will describe how this analysis estimates and minimizes contributions from each background process. Finally, I demonstrate purity and modelling of these backgrounds in validation and control regions. I made key contributions in this analysis in optimizing and testing control region definitions and their effects on our overall results.




Figure 6.1: Feynman diagram for Drell-Yan process in which two quarks produce a photon or Z -boson that then decays leptonically [43].

6.1.1 Drell-Yan background ($Z \rightarrow \tau\tau$)

The Drell-Yan background, alternatively called $Z \rightarrow \tau\tau$ or $Z+\text{jets}$ is produced in the initial collision when two quarks produce a Z boson which then decays to two leptons (in the largest background, τ leptons which then decay to electron/muons and neutrinos). Thus there are two jets and two leptons in its final state as well as missing energy from neutrinos, all of which are indicative of signal VBF Higgs events as well. A typical Drell-Yan feynman diagram is shown in Figure 6.1. The Drell-Yan background can be discriminated from our signal as well as the other backgrounds in two main ways- first in terms of the mass of the overall event aligning with the mass of the Z boson. To select for Drell-Yan events we can select those which $m_{\tau\tau}$ is near the Z mass peak. $m_{\tau\tau}$ is defined in the previous section and cutting on this in a 25 GeV window about the Z -mass peak drastically improves our $Z+\text{jets}$ purity from only 19% of total events to $\approx 52\%$.

We have also tested and trained a secondary discriminant for $Z+\text{jets}$ in our analysis, a BDT trained to discriminate $Z+\text{jets}$ and VBF signal events. This BDT significantly decreases overall $Z+\text{jets}$ background in our signal region but we found that the subsequent decrease in statistics from using this cut leads to very similar overall results. The results

CHAPTER 6. BACKGROUNDS AND SYSTEMATICS

from this comparison are shown in the Appendix.

Drell-Yan Control Region

The Z +jets control region definition is quite similar to the VBF signal region except that the Z +jets veto cut is inverted. Thus instead of removing events near the Z -mass window we select for them by applying a cut on $m_{ll} < 80$ GeV, $66.2 \text{ GeV} < m_{\tau\tau} < 116.2$ GeV, and the same OLV and CJV cuts as in the VBF signal region. The Z +jets control region has a purity of $\approx 82\%$ and yields in this region are shown in the table below.

$\sqrt{s} = 13\text{TeV}$, $\mathcal{L} = 139\text{fb}^{-1}$ (Full Run 2)	H_{VBF}	H_{qgF}	WW	Other VV	Top	Zjets	Mis-Id	Total Bkg	Significance	Data	Data/MC
Scale factors					NF = 0.99 ± 0.01	NF = 1.03 ± 0.05		NFs Applied			
$Z \rightarrow \tau\tau$ CR: $ m_{\tau\tau} - m_Z < 25$, bVeto	27.15 ± 0.16	67.22 ± 0.76	1809.83 ± 8.26	394.27 ± 4.91	5630.06 ± 17.10	9040.34 ± 45.91	507.91 ± 26.88	17449.63 ± 56.70	0.21 ± 0.00	16400	0.94 ± 0.01
Scale factors					NF = 0.99 ± 0.01	NF = 1.03 ± 0.05		NFs Applied			
$Z \rightarrow \tau\tau$ CR: $M_{ll} < 80$ GeV	26.48 ± 0.16	64.96 ± 0.75	589.77 ± 4.57	221.55 ± 4.43	1702.27 ± 9.24	8802.21 ± 41.65	284.22 ± 21.72	11664.97 ± 48.30	0.25 ± 0.00	10805	0.92 ± 0.01
Scale factors					NF = 0.99 ± 0.01	NF = 1.03 ± 0.05		NFs Applied			
$Z \rightarrow \tau\tau$ CR: CJV< 20 GeV	20.86 ± 0.14	46.58 ± 0.63	419.89 ± 3.94	161.68 ± 4.14	1149.53 ± 7.66	6569.13 ± 36.52	224.08 ± 18.68	8570.90 ± 42.12	0.23 ± 0.00	7931	0.92 ± 0.01
Scale factors					NF = 0.99 ± 0.01	NF = 1.03 ± 0.05		NFs Applied			
$Z \rightarrow \tau\tau$ CR: OLV	16.04 ± 0.12	11.63 ± 0.32	86.60 ± 1.87	34.00 ± 1.35	291.98 ± 3.89	1413.62 ± 17.49	30.62 ± 8.22	1868.45 ± 19.85	0.37 ± 0.00	1832	0.97 ± 0.02
Scale factors					NF = 0.99 ± 0.01	NF = 1.03 ± 0.05		NFs Applied			
BDT Zjets < 0.5	3.52 ± 0.06	4.42 ± 0.20	34.38 ± 1.27	13.76 ± 1.03	106.25 ± 2.36	809.42 ± 14.54	9.33 ± 6.18	977.56 ± 16.06	0.11 ± 0.00	979	1.00 ± 0.04

Table 6.1: Cutflow in the Z +jets control region.

Data/MC shows decent agreement over various variable distributions as seen in the below.

Some of these variables are those used in the BDT described in Appendix A.

Normalization factors (NF) are derived in the Z +jets control region to correct for Data/MC mis-modelling. These factors are applied to the Z +jets sample in the signal region. The NF factors are 1.03 ± 0.05 .

6.1.2 Top background

The top background consists of two main components, Wt and $t\bar{t}$ events where the W decays leptonically and the top quarks decay to jets (notably b -jets). The top background is dominated by $t\bar{t}$ and is the largest background in our signal region, composing about 60% of the total background. Though top backgrounds are numerous, discrimination between top and signal Higgs events is possible through training a BDT on variables that have very

CHAPTER 6. BACKGROUNDS AND SYSTEMATICS




Figure 6.2: Distributions of $m_{\tau\tau}$, p_T^{tot} , m_T , in E_T^{miss} , $E_T^{\text{miss}}, \text{significance}$, $E_T^{\text{miss}, \text{track}}$, $E_T^{\text{miss}, \text{track}}$, $E_{\text{T,rel}}^{\text{miss}}$, $\Delta\phi_{\ell\ell, E_T^{\text{miss}, \text{track}}}$ in the $Z + \text{jets}$ control region.

different distributions between these two types of events like $m_{\ell\ell}$ and ΔY_{jj} . As demonstrated in Chapter 4, this BDT discriminates between signal and top background quite well. The top

CHAPTER 6. BACKGROUNDS AND SYSTEMATICS

background is therefore able to be defined in the signal region and separated using dedicated discriminants. The final BDT used in the final statistical fit is discussed in the following chapter but in this section I will describe an additional BDT used to isolate top background events- that splitting top+ WW and all other backgrounds. This is added to the overall fit as the main discriminant for top backgrounds and is considered in combination with WW due to their similar signatures and their estimation as a combined parameter in the final fit.

We also define a top validation region to test top Monte Carlo modelling as well as to calculate a normalization factor that is used to correct top mis-modelling in the signal region. The top validation region is described similarly to the signal region with one major difference, a b -tag applied in the signal region requiring all events with a b -jet to be removed is (almost) reversed. Instead we require exactly one b -tagged jet. We require exactly one in order to define the validation region as similarly to the signal region as possible while increasing top purity. The result is a highly pure top validation region where the flavor composition of the tagged jets is close to the signal region. Purity of the top control region is $\approx 97\%$ and yields in this region are shown in the table below.

$\sqrt{s} = 13 TeV, \mathcal{L} = 139 fb^{-1}$ (Full Run 2)	H_{VBF}	H_{ggF}	WW	Other VV	Top	Zjets	Mis-Id	Total Bkg	Significance	Data	Data/MC
Scale factors					$NF = 0.99 \pm 0.01$	$NF = 1.03 \pm 0.05$		NFs Applied			
$n_{b-jets} = 1$	39.79 ± 0.19	129.20 ± 1.08	2993.57 ± 9.53	690.94 ± 4.40	349498.67 ± 128.68	3739.16 ± 34.69	4824.17 ± 101.77	361875.73 ± 168.02	0.07 ± 0.00	359758	0.99 ± 0.00
Scale factors					$NF = 0.99 \pm 0.01$	$NF = 1.03 \pm 0.05$		NFs Applied			
$Z \rightarrow \tau\tau$ veto	18.05 ± 0.13	18.70 ± 0.42	175.96 ± 2.68	43.24 ± 1.19	29996.21 ± 38.10	183.57 ± 8.79	287.05 ± 30.06	30704.73 ± 49.41	0.10 ± 0.00	30709	1.00 ± 0.01
Scale factors					$NF = 0.99 \pm 0.01$	$NF = 1.03 \pm 0.05$		NFs Applied			
CJV < 20 GeV	29.69 ± 0.17	84.86 ± 0.88	1905.77 ± 7.80	454.48 ± 3.81	238439.98 ± 107.23	2524.42 ± 30.90	3083.10 ± 83.79	246492.61 ± 139.83	0.06 ± 0.00	244811	0.99 ± 0.00
Scale factors					$NF = 0.99 \pm 0.01$	$NF = 1.03 \pm 0.05$		NFs Applied			
OLV	20.85 ± 0.14	20.90 ± 0.44	294.56 ± 3.34	76.17 ± 1.52	46224.53 ± 47.42	507.30 ± 13.10	427.86 ± 36.88	47551.32 ± 61.60	0.10 ± 0.00	47182	0.99 ± 0.00
Scale factors					$NF = 0.99 \pm 0.01$	$NF = 1.03 \pm 0.05$		NFs Applied			
$Z \rightarrow \tau\tau$ veto	18.05 ± 0.13	18.70 ± 0.42	175.96 ± 2.68	43.24 ± 1.19	29996.21 ± 38.10	183.57 ± 8.79	287.05 ± 30.06	30704.73 ± 49.41	0.10 ± 0.00	30709	1.00 ± 0.01

Table 6.2: Cutflow in the top control region.

Data/MC in the top validation region shows good agreement over various variable distributions as seen in the below. Many of these distributions of used as inputs to the top+ WW BDT.

CHAPTER 6. BACKGROUNDS AND SYSTEMATICS




Figure 6.3: Distributions of $m_{\ell\ell}$, $\Delta Y_{\ell\ell}$, ΔY_{jj} , m_{jj} , $\Delta\Phi_{jj}$, p_{tot}^T , m_T , and $E_{T,\text{rel}}^{\text{miss}}$ in the top validation region.

Top and WW BDT discriminant (against all other samples)

This BDT is trained using $e\mu + \mu e$ events after the VBF selection and all signal regions cuts so that the phase space in which we train the BDT is exactly the same as the one where 103

CHAPTER 6. BACKGROUNDS AND SYSTEMATICS

we apply it in the final fit. The training includes top and WW trained against weighted samples of VBF, ggF, $Z+jets$, and $V\gamma$ events. The MC statistics used in the training are half those available after all signal region cuts and the other half are later used to test the training. This corresponds to $\approx 90,000$ un-weighted WW and top events and $\approx 115,000$ raw VBF, ggF, $Z+jets$ and $V\gamma$ events. This training includes MC weights on events to best account for overall event distributions and there are ≈ 2000 total weighted top and WW events used in the training and ≈ 550 weighted other events.

The TMVA BDTG interface is used to train and test the BDT. The optimal parameters were found through a scan of reasonable values and the final set is summarized in Table 6.3.

Parameter	Value
Boosting algorithm	Gradient
Maximum tree depth	22
Number of trees	200
Minimum number of events requires per mode	5%
Number of cuts	7

Table 6.3: BDT parameters used for the top + WW vs. other backgrounds training.

This BDT utilizes a range of lepton and jet kinematic variables (8) to distinguish between signal and background events. These include ΔY_{jj} , the combination of lepton/jet masses M_{l0j0} for leading and subleading leptons and jets, $\Delta\Phi_{\ell\ell}$, m_T , η_{j0} , η_{j1} , $\Delta\Phi_{jj}$, and \sum centralities (L). While a larger variety of variables have been tested, these demonstrated the highest discrimination between top/ WW events and other samples. Plots shown in 6.4 and 6.5 demonstrate the input distributions used to train the BDT and their correlations. The BDT training successfully separates top/ WW background and other samples (ggF, VBF, $Z+jets$, and $V\gamma$). In order to quantify the discrimination we use the integrated-ROC calculated through TMVA for weighted normalized samples and find an optimal value of 0.920.

CHAPTER 6. BACKGROUNDS AND SYSTEMATICS




Figure 6.4: Distributions of input variables to top+ WW vs. other samples BDT. Samples are weighted and normalized to even numbers of background and signal events. Signal represents top+ WW and background all other samples.




Figure 6.5: Correlations of input variables to top + WW vs other samples BDT. Signal represents top+ WW and background other samples.

CHAPTER 6. BACKGROUNDS AND SYSTEMATICS

Comparisons between the test and training show that the BDT is un-biased- differences between testing and training samples would imply overtraining, or the BDT using to many parameters on too few events. For signal and background we find KS-test values of 0.107 and 0.154, and so no evidence of over-training. We can visualize the BDT output variable both on un-weighted normalized samples and on samples with all event weights applied. The following plot shows BDT results applied to normalized samples of VBF signal and top/WW backgrounds.




Figure 6.6: Unweighted, normalized samples of top+WW samples (signal) and other samples (background) plotted over BDT output distribution on left, overlaid testing and training samples shown. Right, full weighted samples of all signal and background events plotted over BDT output distribution after signal region selection. Data is blinded here

We aim to fit this distribution in the signal region with high significance for top/WW events in uppermost bins of the distribution. Since this BDT is trained and applied in the signal region we cannot directly test the modelling of input variables. However, modelling at the pre-selection level for each of these variables as well as in the top validation region described earlier show no evidence of mis-modelling. The binning for this discriminant used in the statistical fit and its result (using Asimov data) are shown in the final chapter. We can visualize this BDT discriminant in the top validation region to gain an idea of overall

CHAPTER 6. BACKGROUNDS AND SYSTEMATICS

modelling (particularly of the reasonably pure top events).




Figure 6.7: Full weighted samples of all signal and background plotted over Top + WW vs. VBF ggF + $Z+jets+V\gamma$ BDT output distributions in top validation region

The BDT to discriminate top+ WW background from VBF signal events is trained and applied in the signal region and so described in the previous chapter, but its distribution in the top validation region is also shown.

A designated BDT discriminates between top and WW backgrounds and is described further in the WW background section (next). However, this distribution in the top validation region with all weighted samples is shown below. to demonstrate its effect on primarily predominantly top-like events.

Normalization factors (NF) are derived in the top validation region to correct for data/MC mis-modelling. These factors are applied to the top sample in the signal region. The NF factors are 0.99 ± 0.01 .




Figure 6.8: Full weighted samples of all signal and background plotted over Top + WW vs. VBF BDT output distributions in top validation region




Figure 6.9: Full weighted samples of all signal and background plotted over Top vs. WW BDT output distributions in top validation region

CHAPTER 6. BACKGROUNDS AND SYSTEMATICS

6.1.3 Diboson background

The WW background consists primarily of QCD WW +jets events (highly dominating electroweak vertices). This background is estimated along with the top background using a joint parameter due to their similarities in signature as well as the difficulty in defining a pure WW control region without top contamination. A WW validation region is defined to demonstrate WW MC modelling in a targeted WW region. The WW validation region is defined with at least 2 jets, a b -veto $N_{b-jet} < 1$ and a central-jet-veto of below 20GeV as in the signal region. Two additional cuts differ from the signal region - $m_T > 130\text{GeV}$ and $m_{T2} > 160\text{GeV}$ where m_{T2} is defined as

$$m_{T2} = \min_{p_T^1 + p_T^2 = p_T} (\max(m_T^2(p_T^1, p_T^a), m_T^2(p_T^2, p_T^b))) \quad (6.1)$$

This represents a lower bound on the parent particle's mass, so using a large m_{T2} cut eliminates contamination from many $t\bar{t}$ decays which have an upper limit near the top mass. The purity of the region is $\approx 37\%$ and the cutflow for this region is shown below.

$\sqrt{s} = 13\text{TeV}, \mathcal{L} = 139\text{fb}^{-1}$ (Full Run 2)	H_{VBF}	H_{ggF}	WW	Other VV	Top	Zjets	Mis-Id	Total Bkg	Significance	Data	Data/MC
Scale factors					$\text{NF} = 0.99 \pm 0.01$	$\text{NF} = 1.03 \pm 0.05$					
b-veto	323.67 ± 0.54	914.43 ± 2.86	21075.69 ± 28.85	2701.86 ± 10.03	63729.30 ± 57.17	22477.17 ± 104.35	4407.46 ± 70.33	115305.91 ± 141.58	0.95 ± 0.00	109677	0.95 ± 0.00
Scale factors					$\text{NF} = 0.99 \pm 0.01$	$\text{NF} = 1.03 \pm 0.05$					
$M_T > 130\text{ GeV}$	32.37 ± 0.17	130.66 ± 1.09	16127.61 ± 24.93	1573.71 ± 6.42	50042.33 ± 50.78	885.51 ± 34.16	2078.63 ± 49.84	70838.44 ± 83.03	0.12 ± 0.00	68255	0.96 ± 0.00
Scale factors					$\text{NF} = 0.99 \pm 0.01$	$\text{NF} = 1.03 \pm 0.05$					
$M_{T2} > 160\text{ GeV}$	18.87 ± 0.13	50.22 ± 0.67	6528.07 ± 14.33	591.47 ± 3.59	11542.51 ± 24.79	306.95 ± 14.38	602.98 ± 26.34	19622.20 ± 41.64	0.13 ± 0.00	18672	0.95 ± 0.01
Scale factors					$\text{NF} = 0.99 \pm 0.01$	$\text{NF} = 1.03 \pm 0.05$					
CJV (20GeV)	13.61 ± 0.11	32.51 ± 0.54	4363.49 ± 11.96	385.58 ± 2.86	6410.03 ± 18.98	187.58 ± 13.07	392.37 ± 20.45	11771.56 ± 33.18	0.13 ± 0.00	11245	0.95 ± 0.01

Table 6.4: Cutflow in the WW validation region.

Data/MC shows good agreement over various variable distributions as seen in the below.

Top vs. WW BDT discriminant

This BDT is trained using $e\mu + \mu e$ events after the VBF selection and all signal regions cuts so that the phase space in which we train the BDT is exactly the same as the one where

CHAPTER 6. BACKGROUNDS AND SYSTEMATICS




Figure 6.10: Distributions of $m_{\ell\ell}$, $\Delta Y_{\ell\ell}$, ΔY_{jj} , m_{jj} , $\Delta\Phi_{jj}$, p_{tot}^T , m_T , and $E_{T,\text{rel}}^{\text{miss}}$ in the WW validation region.

we apply it in the final fit. The training includes weighted samples of top and WW events trained against one another. The MC statistics used in the training are half those available

CHAPTER 6. BACKGROUNDS AND SYSTEMATICS

after all signal region cuts where the other half are later used to test the training. This corresponds to $\approx 35,000$ un-weighted top and $\approx 55,000$ raw WW events. This training includes MC weights on events to best account for overall event distributions and there are ≈ 1500 total weighted top used in the training and ≈ 500 weighted WW events.

The TMVA BDTG interface is used to train and test the BDT. The optimal parameters were found through a scan of reasonable values and the final set is summarized in Table 6.5.

Parameter	Value
Boosting algorithm	Gradient
Maximum tree depth	22
Number of trees	200
Minimum number of events requires per mode	5%
Number of cuts	7

Table 6.5: BDT parameters used for the top vs. WW training.

This BDT utilizes a collection of lepton and jet kinematic variables (14) to distinguish between top and WW events. These include ΔY_{jj} , $\Delta Y_{\ell\ell}$, $\Delta \Phi_{\ell\ell}$, m_T , η_{l0} , η_{l1} , p_{j0}^T , p_{j1}^T , $\Delta \Phi_{jj}$, \sum centralities (L), and combined masses M_{l0j0} , M_{l0j1} , M_{l1j0} , M_{l1j1} . While a larger variety of variables have been tested, these demonstrated the highest discrimination between top and WW events. Plots shown in 6.11 and 6.12 demonstrate the input distributions used to train the BDT and their correlations. The BDT training separates top and WW backgrounds somewhat. These signals have similar kinematic distributions that make high discrimination difficult. In order to quantify the discrimination we use the integrated-ROC calculated through TMVA for weighted normalized samples and find an optimal value of 0.653. While relatively low, this shows some minimal amount of discrimination. This is useful in the fit to differentiate top and WW backgrounds, though high levels of separation are not necessary because top and WW are estimated together with a shared parameter in the simultaneous fit.

CHAPTER 6. BACKGROUNDS AND SYSTEMATICS




Figure 6.11: Distributions of input variables to top vs. WW BDT. Samples are weighted and normalized to even numbers of background and signal events. Signal represents top and background WW .




Figure 6.12: Correlations of input variables to top vs. WW BDT. Signal represents top and background WW .

CHAPTER 6. BACKGROUNDS AND SYSTEMATICS

Comparisons between the test and training show that the BDT is un-biased. For signal and background we find KS-test values of 0.067 and 0.037, and so no evidence of over-training. We can visualize the BDT output variable both on un-weighted normalized samples and on samples with all event weights applied. The following plot shows BDT results applied to normalized samples of top and WW backgrounds.




Figure 6.13: Unweighted, normalized samples of top samples (signal) and WW samples (background) plotted over BDT output distribution on left, overlaid testing and training samples shown. Right, full weighted samples of all signal and background events plotted over the BDT output distribution after signal region selection. Data is blinded here

We aim to fit this distribution in the signal region with high significance for top events in uppermost bins of the distribution (and WW in the lowermost). Since this BDT is trained and applied in the signal region we cannot directly test the modelling of input variables. However, modelling at the pre-selection level for each of these variables as well as in the WW validation region described earlier show no evidence of mis-modelling. The binning for this discriminant used in the statistical fit and its result (using Asimov data) are shown in the final chapter. We can visualize this BDT discriminant in the WW validation region to gain an idea of overall modelling particularly of the WW events.

The BDT to discriminate top+ WW background from VBF, ggF, $Z+jets$ and $V\gamma$ events




Figure 6.14: Full weighted samples of all signal and background plotted over BDT output distributions in WW validation region

CHAPTER 6. BACKGROUNDS AND SYSTEMATICS

is trained and applied in the signal region and described in the previously, but its distribution in the WW validation region is also shown.




Figure 6.15: Full weighted samples of all signal and background plotted over Top + WW vs. VBF ggF + $Z+jets+V\gamma$ BDT output distributions in WW validation region

The BDT to discriminate WW background from VBF signal events is trained and applied in the signal region and so described in the previous chapter, but its distribution in the WW validation region is shown below. This BDT discriminates both WW and top (combined) against the VBF signal as they have similar kinematic distributions and are treated together in the final simultaneous fit.

6.1.4 ggF background

Other Higgs production modes are considered backgrounds in this analysis- and while vector mediated Higgs production (VH) and top produced Higgs (tth) are small in our

CHAPTER 6. BACKGROUNDS AND SYSTEMATICS




Figure 6.16: Full weighted samples of all signal and background plotted over Top + WW vs. VBF BDT output distributions in WW validation region

CHAPTER 6. BACKGROUNDS AND SYSTEMATICS

signal region so don't play a large role in the analysis - gluon fusion Higgs production (ggF) represents a significant background. Signal region cuts requiring at least 2-jets and the central jet veto significantly reduce background yields, but because it is kinematically very similar to VBF signal, it has a large effect on the VBF differential coupling measurement. To mitigate these effects, ggF events are simultaneously estimated from three control regions and the signal region. The control regions are chosen to minimize both the statistical and the modelling uncertainties, in particular those originating from the modelling of higher order QCD corrections. In particular, the regions inspired from a subdivision based on the jet multiplicity. The first two regions, ggF CR1 and ggF CR2, are built from all preselection cuts and cuts requiring at least two jets, a b -veto, $Z \rightarrow \tau\tau$ -veto, and orthogonal cuts on the central jet veto (CJV) and opposite lepton veto (OLV). The third ggF control region, ggF CR3, uses all preselection cuts, a b veto, $Z \rightarrow \tau\tau$ veto, and requirement for less than 2 jets. The full definitions are described:

- **GGF-CR1** Preselection criteria and $N_{\text{jet}} \geq 2$, b veto, $Z \rightarrow \tau\tau$ veto, CJV < 1 and OLV > 1 or CJV > 1 and OLV < 1 .
- **GGF-CR2** Preselection criteria and $N_{\text{jet}} \geq 2$, b veto, $Z \rightarrow \tau\tau$ veto, CJV > 1 and OLV > 1 .
- **GGF-CR3** Preselection criteria and $N_{\text{jet}} \geq 2$, b veto, and $Z \rightarrow \tau\tau$ veto.

The full ggF expectation in the signal region is built from the following ratio (inspired by the ABCD method)

$$\mu_{\text{GGF-SR}} = \frac{\mu_3^{\text{GGF-CR}} \cdot \mu_1^{\text{GGF-CR}}}{\mu_2^{\text{GGF-CR}}}. \quad (6.2)$$

where $\mu_{\text{ggF-X}}$ represents the yield modifier in each ggF control region. The final event yield in the signal region $\mu_{\text{ggF-SR}}$ is determined in the simultaneous fit as described in the next

CHAPTER 6. BACKGROUNDS AND SYSTEMATICS

chapter. In each ggF category the ggF yield is extracted from simulation-extracted template fits based on dedicated discriminants. For each control region a dedicated multivariate discriminant is trained and applied to discriminate between ggF and all the backgrounds combined. Results and training parameters from each of these discriminants is summarized next.

Discriminant in ggF CR1

The multivariate discriminant used for the ggF CR1 is a boosted decision tree (BDT) trained using $e\mu + \mu e$ events that pass all ggF CR1 cuts. The training includes ggF events trained against VBF signal and all backgrounds. The MC statistics used in the training are half those available after the ggF CR1 cuts (as the other half are later used to test the training). This corresponds to $\approx 8,000$ ggF events and $\approx 250,000$ other signal and background events. Events are trained with MC weights applied to accurately represent relative importance of each event and the training uses ≈ 130 weighted ggF samples and $\approx 10,000$ weighted other background samples.

The TMVA BDTG interface is used to train and test the BDT. The optimal parameters were found through a scan of reasonable values and the final set is summarized in Table 6.6. For this BDT various distributions (8) are used to take advantage of differences

Parameter	Value
Boosting algorithm	Gradient
Maximum tree depth	10
Number of trees	600
Minimum number of events requires per mode	5%
Number of cuts	7

Table 6.6: BDT parameters used for the ggF CR1 training.

in distributions between ggF events and other sample types. These variables include $\Delta Y_{\ell\ell}$,

CHAPTER 6. BACKGROUNDS AND SYSTEMATICS

$\Delta\Phi_{\ell\ell, m_{\ell\ell}, m_T}$, jet p_{lead}^T , jet p_{sublead}^T , $\Delta\Phi_{jj}$, and $E_{\text{T}}^{\text{miss}}$. Distributions for these variables in the ggF CR1 region where the BDT is trained are shown below demonstrating data/MC modelling for each.

Plots shown in 6.18 and 6.19 demonstrate the input distributions used to train the BDT and their correlations where each distribution is normalized to equal number of background and signal events.

In order to quantify the discrimination we use the integrated-ROC calculated through TMVA for weighted normalized samples and find an optimal value of 0.898. Comparisons between the test and training show that the BDT is un-biased- differences between testing and training samples would imply overtraining, or the BDT using too many parameters on too few events. Visually, we can see that the testing and trainings samples are quite similar. Additionally, a Kolmogorov-Smirnov test is performed to measure if the two test and training distributions differ significantly and no evidence of over-training is present (values 0.191, 0.049 for signal, background). We can visualize the BDT output variable both on weighted normalized samples and on samples without normalization. The following plots show BDT results applied to normalized and full weighted samples.

Discriminant in ggF CR2

The multivariate discriminant used for ggF CR2 is a boosted decision tree trained using $e\mu + \mu e$ events that pass all ggF CR2 cuts. The training includes ggF events trained against VBF signal and all backgrounds. The MC statistics used in the training are half those available after ggF CR2 cuts (as the other half are later used to test the training). This corresponds to $\approx 22,200$ ggF events and $\approx 90,000$ other signal and background events.

The TMVA BDTG interface is used to train and test the BDT. The optimal parameters were found through a scan of reasonable values and the final set is summarized in

CHAPTER 6. BACKGROUNDS AND SYSTEMATICS




Figure 6.17: Distributions of $\Delta Y_{\ell\ell}$, $\Delta\Phi_{\ell\ell}$, $m_{\ell\ell}$, m_T , jet p_T^T , jet p_{sublead}^T , $\Delta\Phi_{jj}$, and E_T^{miss} in the ggF CR1 used as input to the BDT discriminating ggF from all other samples.

Table 6.7. For this BDT various distributions (6) are used to take advantage of differences in distributions between ggF events and other sample types. These variables include $\Delta Y_{\ell\ell}$,

CHAPTER 6. BACKGROUNDS AND SYSTEMATICS




Figure 6.18: Distributions of input variables to ggF CR1 BDT. Samples are unweighted and normalized to even numbers of background and signal events. Signal represents ggF and background all other samples.




Figure 6.19: Correlations of input variables to ggF CR1 BDT. Signal represents ggF and background all other samples.

CHAPTER 6. BACKGROUNDS AND SYSTEMATICS




Figure 6.20: Normalized samples of ggF (signal) and all other samples (background) plotted over BDT output distribution on left, overlaid testing and training samples shown. Right, full weighted samples of ggF signal and all other backgrounds plotted over BDT output distribution after ggF CR 1.

Parameter	Value
Boosting algorithm	Gradient
Maximum tree depth	10
Number of trees	50
Minimum number of events requires per mode	5%
Number of cuts	7

Table 6.7: BDT parameters used for the ggF CR2 training.

$\Delta\Phi_{\ell\ell, m_{\ell\ell}, m_T}$, $\Delta\Phi_{jj}$, and E_T^{miss} . Distributions for these variables in the ggF CR2 region where the BDT is trained are shown below demonstrating data/MC modelling for each.

Plots shown in 6.22 and 6.23 demonstrate the input distributions used to train the BDT and their correlations where each distribution is weighted and normalized to equal number of background and signal events.

In order to quantify the discrimination we use the integrated-ROC calculated through TMVA for weighted normalized samples and find an optimal value of 0.902. Comparisons between the test and training show that the BDT is un-biased- visually, once can see that the testing and trainings samples are quite similar. Additionally, a Kolmogorov-Smirnov test is

CHAPTER 6. BACKGROUNDS AND SYSTEMATICS




Figure 6.21: Distributions of $\Delta Y_{\ell\ell}$, $\Delta \Phi_{\ell\ell}$, $m_{\ell\ell}$, m_T , $\Delta \Phi_{jj}$, and E_T^{miss} in the ggF CR2 used as input to the BDT discriminating ggF from all other samples.

performed to measure if the two test and training distributions differ significantly and with values of 0.012 (0.303) for signal (background) no evidence of over-training is present. We can visualize the BDT output variable both on weighted normalized samples and on samples with full event weights applied. The following plots show BDT results applied to normalized and full weighted samples.

Discriminant in ggF-CR3

The multivariate discriminant used for the ggF CR3 is a boosted decision tree trained using $e\mu + \mu e$ events that pass all ggF CR3 cuts. The training includes only ggF events

CHAPTER 6. BACKGROUNDS AND SYSTEMATICS




Figure 6.22: Distributions of input variables to ggF CR2 BDT. Samples are weighted and normalized to even numbers of background and signal events. Signal represents ggF and background all other samples.




Figure 6.23: Correlations of input variables to ggF CR2 BDT. Signal represents ggF and background all other samples.

CHAPTER 6. BACKGROUNDS AND SYSTEMATICS




Figure 6.24: Weighted, normalized samples of ggF (signal) and all other samples (background) plotted over BDT output distribution on left, overlaid testing and training samples shown. Right, full weighted samples of ggF signal and all other backgrounds plotted over BDT output distribution after ggF CR2 cuts.

trained against VBF signal and all backgrounds. The MC statistics used in the training are half those available after the ggF CR3 cuts (as the other half are later used to test the training). This corresponds to $\approx 200,000$ ggF events and $\approx 2,800,000$ other signal and background events.

The TMVA BDTG interface is used to train and test the BDT. The optimal parameters were found through a scan of reasonable values and the final set is summarized in Table 6.8. For this BDT various distributions (4) are used to take advantage of differences

Parameter	Value
Boosting algorithm	Gradient
Maximum tree depth	10
Number of trees	1000
Minimum number of events requires per mode	5%
Number of cuts	7

Table 6.8: BDT parameters used for the ggF CR3 training.

in distributions between ggF events and other samples types. These variables include $\Delta Y_{\ell\ell}$, $\Delta\Phi_{\ell\ell, m_{\ell\ell}}$, and m_T . Distributions for these variables in the ggF CR3 region where the BDT

CHAPTER 6. BACKGROUNDS AND SYSTEMATICS

is trained are shown below demonstrating data/MC modelling for each.




Figure 6.25: Distributions of $\Delta Y_{\ell\ell}$, $\Delta \Phi_{\ell\ell}$, $m_{\ell\ell}$, and m_T in the ggF CR3 used as input to the BDT discriminating ggF from all other samples.

Plots shown in 6.26 and 6.27 demonstrate the input distributions used to train the BDT and their correlations where each distribution is weighted and normalized to equal number of background and signal events.

In order to quantify the discrimination we use the integrated-ROC calculated through TMVA for weighted normalized samples and find an optimal value of 0.888. Comparisons between the test and training show that the BDT is un-biased, we can see that the testing and training samples are quite similar. Additionally, a Kolmogorov-Smirnov test is performed and shows no sign of bias with values of 0.276 (0.673) for signal (background). We can

CHAPTER 6. BACKGROUNDS AND SYSTEMATICS




Figure 6.26: Distributions of input variables to the ggF CR3 BDT. Samples are weighted and normalized to even numbers of background and signal events. Signal represents ggF and background all other samples.




Figure 6.27: Correlations of input variables to the ggF CR3 BDT. Signal represents ggF and background all other samples.

CHAPTER 6. BACKGROUNDS AND SYSTEMATICS

visualize the BDT output variable both on weighted normalized samples and on samples with full event weights applied. The following plots show BDT results applied to normalized and fully weighted samples.




Figure 6.28: Normalized samples of ggF (signal) and all other samples (background) plotted over BDT output distribution on left, overlaid testing and training samples shown. Right, full weighted samples of ggF signal and all other backgrounds plotted over BDT output distribution after ggF CR3 cuts.

6.1.5 Fake backgrounds

The final substantial background in the VBF HWW differential coupling analysis comes from mis-identified leptons. These are jets that are mistakenly identified as leptons in reconstruction and in this analysis these predominantly come from W +jet events. In these events a W boson decays leptonically leading to one true lepton and one mistaken lepton (a jet from the primary vertex is mistaken for a lepton). This event mimics our desired two lepton signature and so creates an additional HWW background sample. This background is estimated as in the HWW coupling analysis using the fake factor method, similarly to in the 2016 HWW coupling analysis. Further detail on the overall fake factor method and its mathematical formulation can be found here [?] while further details on HWW-specific

CHAPTER 6. BACKGROUNDS AND SYSTEMATICS

fake background studies are here [?]. In this section I will give a brief summary of fake estimations along with the slightly augmented fake factor control region definition used in the HWW differential coupling analysis.

The fake background is estimated using data which is measured in a region defined by signal region cuts with one important distinction- one or both of the leptons used are “anti-identified” meaning they pass some looser lepton identification criteria but not the one used in the analysis signal region. These fakes are then extrapolated to the true signal region (with two “identified” electrons fake factors. Fake factors are measured as functions of p_T and η in jet-enriched $Z+jets$ samples and are defined as the ratio of identified to anti-identified leptons. The fake backgrounds can be considered split between “single-fake” (with one “anti-ID” lepton) from the predominant $W+jets$ background and “double-fake” (with two “anti-ID” leptons from QCD processes. The total signal sample can be defined as:

$$N_{id+id} = N_{id+id}^{EW} + N_{id+id}^{W+jets} + N_{id+id}^{QCD} \quad (6.3)$$

so that the total events include all electroweak processes (N_{id+id}^{EW}) as well as fake backgrounds from $W+jets$ and QCD events. In order to estimate the total fake background in the signal region we need to estimate the number of $W+jets$ and QCD events in “id+anti-id” events and then apply the fake factor to extrapolate into the “id+id” region. The $N_{id+anti-id}$ for fake backgrounds is calculated after subtraction of electroweak backgrounds (two true leptons that contaminate the “id+anti-id region”) from Monte Carlo simulations as follows:

$$N_{id+anti-id}^{W+jets} + N_{id+anti-id}^{QCD} = N_{id+anti-id} - N_{id+anti-id}^{EWMC} \quad (6.4)$$

Fake factors are derived from $Z+jets$ samples and then applied to $W+jets$ regions so the differences between these two samples are important to understand fully. The fake factor is

CHAPTER 6. BACKGROUNDS AND SYSTEMATICS

Table 6.9: Requirements for “identified” and “anti-identified” electrons and muons.

“id” electron	“anti-id” electron	“id” muon	“anti-id” muon
$p_T > 15 \text{ GeV}$ $ \eta < 2.47$, excluding $1.37 < \eta < 1.52$ $ z_0 \sin\theta < 0.5 \text{ mm}$ Pass <i>Tight</i> if $p_T < 25 \text{ GeV}$ Pass <i>Medium</i> if $p_T > 25 \text{ GeV}$ $ d_0 \sigma(d_0) < 5$ Pass FixedCutTrackCone40 if $p_T < 25 \text{ GeV}$ Pass IsoGradient if $p_T < 25 \text{ GeV}$	Pass <i>Loose</i> Veto against identified electron	$p_T > 15 \text{ GeV}$ $ \eta < 2.45$ $ z_0 \sin\theta < 0.5 \text{ mm}$ Pass <i>Tight</i> $ d_0 \sigma(d_0) < 3$ $E_T^{\text{cone}20}/p_T < 0.09$ $p_T^{\text{varcone}30}/p_T < 0.06$	Pass <i>Medium</i> $ d_0 \sigma(d_0) < 15$ Veto against identified muon

defined

$$F.F. = \frac{N_{id}}{N_{anti-id}} \quad (6.5)$$

and is measured separately for electron and muons and measured in bins of η and p_T . The following table summarizes to “id” and “anti-id” requirements.

$Z+jets$ event are selected to contain exactly three loosely identified leptons with $p_T > 15 \text{ GeV}$. Further requirements are for an opposite sign ee or $\mu\mu$ lepton pair with $7 \text{ GeV} < m_{\ell\ell} < 110 \text{ GeV}$. Both Z candidate leptons must be “identified” so that the third is the fake candidate. An additional WZ veto is applied using $m_T > 50 \text{ GeV}$ to mitigate electroweak background in the $Z+jets$ sample. Dedicated MC model electroweak backgrounds in the $Z+jets$ region and include $V + \gamma$, diboson (WW , WZ , and ZZ), single top and $t\bar{t}$. Finally, WZ backgrounds are normalized ot their measured cross-sections. Fake yields for the $Z+jets$ samples are finally calculated by subtracting data from all MC electroweak backgrounds. The $Z+jets$ cutflow for fake estimation in the $ee + \mu\mu$ channel is shown next.

Add $Z+jets$ cutflow and plots?

The fake factor is computed as a binned ratio in p_T , $[15, 20, 25, 35, \inf]$ and for muons also in η for $[0, 1.5, 2.5]$ exluding the electromagnetic calorimeter crack region. The following table summarized fake factors used in this analysis.

Further studies on fake factors include comparisons to their MC simulation expectations

CHAPTER 6. BACKGROUNDS AND SYSTEMATICS

Table 6.10: Fake factors binned for muons and electrons in p_T and η with their statistical uncertainties

$p_T(\text{GeV})$	Muon FF ($ \eta < 1.5$)	Muon FF ($1.5 < \eta < 2.5$)	Electron FF
15-25	0.038 ± 0.004	0.057 ± 0.005	0.079 ± 0.005
20-25	0.021 ± 0.007	0.0357 ± 0.0077	0.097 ± 0.001
25-35	0.029 ± 0.011	0.064 ± 0.014	0.157 ± 0.017
35-inf	0.049 ± 0.023	0.11 ± 0.04	0.19 ± 0.026

and studies on differences between the $Z+\text{jets}$ control sample used to calculate fake factors and the $W+\text{jets}$ sample on which they are applied. These studies are beyond the scope of this thesis but support the use of the fake factor method in our analysis. Specifically, jet kinematics and heavy flavor fraction in both samples have been studied by the HWW coupling analysis and factor into a correction factor applied to transfer between $Z+\text{jets}$ and $W+\text{jets}$ samples. More information on these studies and correction factors can be found in [?].

The $W+\text{jets}$ control region is defined as in the signal region, with cuts specifying at least 2 jets, a b -veto, $m_{\tau\tau}$ based Z-veto, an opposite lepton veto, central jet veto, and cuts on m_{jj} and ΔY_{jj} . Unlike the signal region, this region also specified one “id” and one “anti-id” lepton instead of two identified leptons. The cutflow for this region is shown below where fakes are defined from the subtraction of total electroweak background from data. Fake purity in this control region is 74% after all signal region-like cuts are applied.

Distributions below show the electroweak background processes (blue) in the fake control region along with data (black). The $W+\text{jets}$ or fake background is taken as the difference between data and EW background and shown in the plots as light blue points. High fake purity is shown and these backgrounds are used along with fake factors to estimate fakes in contaminating our signal region.

The EW background subtracted from data in the $W+\text{jets}$ control region consists of $V\gamma$,

CHAPTER 6. BACKGROUNDS AND SYSTEMATICS

$\sqrt{s} = 13\text{TeV}$, $\mathcal{L} = 139\text{fb}^{-1}$ (Full Run 2)	Total Bkg	data	fakes	fake purity(%)
Channel Selection	8116768.84 \pm 3042.09	13124937	5008168.16 \pm 4730.67	38.16 \pm 0.04
Trigger Selection	8116768.84 \pm 3042.09	13124937	5008168.16 \pm 4730.67	38.16 \pm 0.04
Trigger Matching	7906126.65 \pm 2917.00	13057932	5151805.35 \pm 4644.01	39.45 \pm 0.04
W+jets flavour split muon	5896841.78 \pm 2703.75	10572340	4675498.22 \pm 4228.78	44.22 \pm 0.04
W+jets flavour split electron	3791849.84 \pm 2494.76	7417426	3625576.16 \pm 3693.41	48.88 \pm 0.05
Jet Cleaning	3791849.84 \pm 2494.76	7417426	3625576.16 \pm 3693.41	48.88 \pm 0.05
Overlap: Vgamma/Vjets	3791849.84 \pm 2494.76	7417426	3625576.16 \pm 3693.41	48.88 \pm 0.05
Only two Leptons	3754720.02 \pm 2485.95	7376514	3621793.98 \pm 3681.91	49.10 \pm 0.05
$p_t^{\text{lead}} > 22\text{ GeV}$	3754720.02 \pm 2485.95	7376514	3621793.98 \pm 3681.91	49.10 \pm 0.05
$p_t^{\text{sublead}} > 15$	3753322.61 \pm 2485.49	7373346	3620023.39 \pm 3681.17	49.10 \pm 0.05
OS Leptons	3751223.46 \pm 2483.45	7365693	3614469.54 \pm 3678.75	49.07 \pm 0.05
$M_{\ell\ell} > 12/10\text{ GeV}$	3751147.28 \pm 2483.41	7365336	3614188.72 \pm 3678.68	49.07 \pm 0.05
SF: Z Veto	3751147.28 \pm 2483.41	7365336	3614188.72 \pm 3678.68	49.07 \pm 0.05
Leptons ID, singleFakes 1 anti-ID,1 ID, doubleFakes 2*anti-ID	780838.94 \pm 837.46	1931616	1150777.06 \pm 1622.64	59.58 \pm 0.09
2-jet (30,30) fJVT	384692.76 \pm 242.92	657536	272843.24 \pm 846.49	41.49 \pm 0.14
b-veto	62361.10 \pm 199.23	198158	135796.90 \pm 487.70	68.53 \pm 0.29
CJV (20GeV)	43093.51 \pm 171.93	137885	94791.49 \pm 409.20	68.75 \pm 0.35
OLV bool	9162.00 \pm 78.28	28068	18906.00 \pm 184.92	67.36 \pm 0.77
$Z \rightarrow \tau\tau$ veto	5142.69 \pm 60.99	18069	12926.31 \pm 147.61	71.54 \pm 0.97
$M_{jj} < 200$	3233.48 \pm 51.55	11911	8677.52 \pm 120.70	72.85 \pm 1.21
DYjj;2.1	2838.02 \pm 50.56	10904	8065.98 \pm 116.02	73.97 \pm 1.28
BDT Zjets ; 0.5	2136.60 \pm 41.35	7338	5201.40 \pm 95.12	70.88 \pm 1.54

Table 6.11: Cutflow in the fakes control region.

diboson, top, and $Z+jets$ events.

Add fake factor closure test

6.2 Systematic uncertainties

The experimental, theoretical, and statistical uncertainties in this analysis all impact the final measurement substantially. In this section I describe first the experimental systematic uncertainties then the theoretical in detail. Statistical uncertainties are derived from the limited number of MC events used to model each of signal and background samples per bin in the differential measurement. A single MC-stat nuisance parameter is implemented based on Poisson statistics and used in our overall statistical fit.

CHAPTER 6. BACKGROUNDS AND SYSTEMATICS




Figure 6.29: Distributions of $p_T^{\ell\ell}$, $m_{\ell\ell}$, m_T , p_T^{tot} , lep p_{lead}^T , lep p_{sublead}^T , m_{jj} , and E_T^{miss} in the differential VBF $W + \text{jets}$ control region.

CHAPTER 6. BACKGROUNDS AND SYSTEMATICS

6.2.1 Experimental uncertainties

Several types of experimental systematics uncertainties affect this analysis. Each of these are calculated for this analysis using the current recommendations for each of the following Combined Performance groups: muons, electrons, Jet/ E_T^{miss} , Trigger, flavor tagging (?), and pile-up. The current set of experimental systematics uses 102 nuisance parameters, each of which will be briefly described next.

There are two different ways systematic uncertainties are calculated and applied. First, for systematic uncertainties on the energy of jets and electrons and momentum of muons, the scale and resolutions is calculated. These test the number of events in the final state affected by shifting energy/momentum by a scale factor. This “smearing” procedure is done for a nominal scaling value and values with $\pm 1\sigma$. Next, systematic uncertainty on the scale factors is calculated by comparing nominal event numbers to those after particular weights are applied. These weights are derived from data-MC agreement and their uncertainty is the relative difference between nominal and modified weight sums.

At the end of this section there is a table which describes all key experimental uncertainties and then tables of experimental uncertainties in the signal region, $Z \rightarrow \tau\tau$ and top control regions are shown. Not included yet are shape systematics on the BDT distribution are also considered for all the experimental systematic uncertainties, maybe necessary, and acceptance in the signal region, $Z \rightarrow \tau\tau$ and top control regions.

Muon reconstruction and identification There are four nuisance parameters associated with muon reconstruction and identification, these are reconstruction efficiency statistic and systematic uncertainties ($MUON_EFF_RECO_STAT$, $MUON_EFF_RECO_SYS$). There are additional separate uncertainties for low p_T muons, or those with $3 < p_T < 20$ GeV ($MUON_EFF_RECO_STAT_LOWPT$, $MUON_EFF_RECO_SYS_LOWPT$).

CHAPTER 6. BACKGROUNDS AND SYSTEMATICS

Muon momentum scale and resolution Muon momentum resolution uncertainties are divided into those from the Inner Detector (*MUON_ID*) and those from the muon spectrometer (*MUON_MS*). Muon momentum scale uncertainties are contained in *MUON_SCALE*. Finally, two additional systematic uncertainties *MUON_SAGITTA_RESBIAS* and *MUON_SAGITTA_* come from a correction applied to muons to account for residual ID/MS misalignments which create a charge dependent bias.

Electron reconstruction and identification lectrone reconstruction and identification uncertainties come from nuisance parameters *EL_EFF_Reco_TOTAL_1NPCOR_PLUSUNCOR*, *EL_EF* 0 to 15, and *EL_EFF_ID_SIMPLIFIED_UncorrUncertaintyNP* 0 to 17.

Electron energy scale and resolution The standard systematics included for electron energy scale and resolution are *EG_SCALE_AF2* and *EG_SCALE_ALL*.
Missing resolution NP?

Jet energy scale and resolution There are a number of jet energy scale and resolution systematics included in the analysis: *JET_BJES_Response*, *JET_EffectiveNP_Detector* 1 to 2, *JET_EffectiveNP_Mixed* 1 to 3, *JET_EffectiveNP_Modelling* 1 to 4, *JET_EffectiveNP_Stat* 1 to 6, *JET_EtaIntercalibration_Modelling*, *JET_EtaIntercalibration_NonClosure_highE*, *JET_EtaIntercalibration_NonClosure_negEta*, *JET_EtaIntercalibration_NonClosure_posEta*, *JET_EtaIntercalibration_TotalStat*, *JET_Flavor_Composition*, *JET_Flavor_Response*, *JET_JER_DataVsMC*, *JET_JER_EffectiveNP* 1 to 7, *JET_JvtEfficiency*, *JET_PunchThrough_M*, *JET_SingleParticle_HighPt*.

Isolation One isolation systematic uncertainty included is for electrons (*EL_EFF_Iso_TOTAL_1NPC*
and another two are included for muons:*MUON_ISO_STAT* and *MUON_ISO_SYS* .
These aren't included currently?

Trigger efficiency Three nuisance parameters are included for trigger efficiency- one for electrons (*EL_EFF_Trigger_TOTAL_1NPCOR_PLUSUNCOR*) and two for muons

CHAPTER 6. BACKGROUNDS AND SYSTEMATICS

(*MUON_EFF_TrigStatUncertainty* and *MUON_EFF_TrigSystUncertainty*) Muon Syst isn't included....

Flavor tagging efficiency Flavor tagging efficiency nuisance parameters considered are *FT_EFF_Eigen_B_* 0 to 2, *FT_EFF_Eigen_C_* 0 to 2, *FT_EFF_Eigen_Light_* 0 to 3, *FT_EFF_extrapolation*, and *FT_EFF_extrapolation_from_charm*.

Pileup reweighting and integrated luminosity The data scale factor for the pileup $\langle \mu \rangle$ value is 1.0/1.09 [Update this value?](#). Systematics of the pileup $\langle \mu \rangle$ value are accounted for with the nuisance parameter *PRW_DATASF* and evaluated by varying the data scale factor upward and downward. Jet pileup nuisance parameters are also included: *JET_Pileup_OffsetMu*, *JET_Pileup_OffsetNPV*, *JET_Pileup_PtTerm*, and *JET_Pileup_RhoTopology*.

Integrated luminosity Uncertainty on the integrated luminosity is $\pm 2.1\%$ for the 2015+2016 dataset, $\pm 2.4\%$ for the 2017 dataset and $\pm 2.0\%$ for the 2018 dataset. This results in an overall integrated luminosity uncertainty of $\pm 1.7\%$ [?].

6.2.2 Theoretical uncertainties

Signal theory uncertainties

ggF background theory uncertainties

Top background theory uncertainties

WW background theory uncertainties

$Z \rightarrow \tau\tau$ background theory uncertainties

CHAPTER 6. BACKGROUNDS AND SYSTEMATICS

Systematic uncertainty	Short description
	Event
Luminosity PRW_DATASF	uncertainty on total integrated luminosity uncertainty on pileup reweighting
	Electrons
EL_EFF_Trigger_Total_1NPCOR_PLUS_UNCOR EL_EFF_Reco_Total_1NPCOR_PLUS_UNCOR EL_EFF_ID_CorrUncertaintyNP (0 to 15) EL_EFF_ID_SIMPLIFIED_UncorrUncertaintyNP (0 to 17) EL_EFF_Iso_Total_1NPCOR_PLUS_UNCOR EG_SCALE_ALL EG_SCALE_AF2	trigger efficiency uncertainty reconstruction efficiency uncertainty ID efficiency uncertainty splits in 16 components ID efficiency uncertainty splits in 18 components isolation efficiency uncertainty energy scale uncertainty
	Muons
MUON_EFF_TrigStatUncertainty MUON_EFF_STAT MUON_EFF_SYS MUON_EFF_STAT_LOWPT MUON_EFF_SYST_LOWPT MUON_ID MUON_MS MUON_SCALE MUON_SAGITTA_RHO MUON_SAGITTA_RESBIAS	trigger efficiency uncertainty reconstruction and ID efficiency uncertainty for muons with $p_T > 20$ GeV reconstruction and ID efficiency uncertainty for muons with $p_T \leq 20$ GeV momentum resolution uncertainty from inner detector momentum resolution uncertainty from muon system momentum scale uncertainty charge dependent momentum scale uncertainty
	Jets
JET_EffectiveNP_Detector (1 to 2) JET_JER_EffectiveNP_ (1 to 7) JET_JER_DataVsMC JET_BJES_Response JET_EffectiveNP_Mixed (1 to 3) JET_EffectiveNP_Modelling (1 to 4) JET_EffectiveNP_Statistical (1 to 6) JET_EtaIntercalibration_NonClosure_negEta JET_EtaIntercalibration_NonClosure_posEta JET_EtaIntercalibration_TotalStat JET_Pileup_OffsetMu JET_Pileup_OffsetNPV JET_Pileup_PtTerm JET_Pileup_RhoTopology JET_Flavor_Composition JET_Flavor_Response JET_PunchThrough_MC16 JET_SingleParticle_HighPt JET_JvtEfficiency FT_EFF_Eigen_B (0 to 2) FT_EFF_Eigen_C (0 to 2) FT_EFF_Eigen_Light (0 to 3) FT_EFF_Eigen_extrapolation FT_EFF_Eigen_extrapolation_from_charm	detector related energy scale uncertainty energy resolution uncertainty energy resolution modelling uncertainty energy scale uncertainty on b-jets energy resolution uncertainty energy scale uncertainty on eta-intercalibration (modeling) statistical resolution uncertainty energy scale uncertainty on eta-intercalibrations (non-closure) energy scale uncertainty on eta-intercalibrations (non-closure) energy scale uncertainty on eta-intercalibrations (statistics/method) energy scale uncertainty on pile-up (mu dependent) energy scale uncertainty on pile-up (NPV dependent) energy scale uncertainty on pile-up (pt term) energy scale uncertainty on pile-up (density ρ) energy scale uncertainty on flavour composition energy scale uncertainty on samples' flavour response energy scale uncertainty for punch-through jets energy scale uncertainty from the behaviour of high- p_T jets JVT efficiency uncertainty <i>b</i> -tagging efficiency uncertainties (“BTAG_MEDIUM”): 3 components for b jets, 3 for c jets and 4 for light jets <i>b</i> -tagging efficiency uncertainty on the extrapolation to high- p_T jets <i>b</i> -tagging efficiency uncertainty on tau jets

Table 6.12: Summary of the experimental systematic uncertainties considered.

Chapter 7

Results

Thus far I have outlined all reconstruction-level inputs to our analysis including event selection, all backgrounds and their estimation methods, and all systematic uncertainties. Each of these components is critical to our final result and this section will focus on the methods used to extract differential cross-section measurements and their uncertainties from the signal and control regions previously defined.

7.1 Statistical analysis

7.1.1 Likelihood functions

This analysis rests on the estimation of a parameter μ which describes the statistical significance of our signal yield relative to its standard model prediction and in this particular analysis defines the signal cross-section. We build a likelihood function $\mathcal{L}(\mu, \Theta)$ where the signal strength μ is a parameter of interest (POI) and nuisance parameters (NPs) $\Theta = \Theta_a, \Theta_b, \dots$ represent all relevant uncertainties. None of these values are known a priori so the likelihood is built to represent the probability of particular values for the POI and NPs.

CHAPTER 7. RESULTS

The analysis uses a maximum likelihood estimator to find the inputs which maximize the likelihood (or equivalently and more mathematically tractable, minimize the negative log of the likelihood). Here I will briefly outline how a likelihood function can incorporate regions of interest, this discussion uses [44] as a guide. First, the Poisson distribution $\mathcal{P}(n|\lambda)$ describes the probability of n events with a true unknown yield λ :

$$\mathcal{P}(n|\lambda) = \lambda^n \frac{e^{-\lambda}}{n!} \quad (7.1)$$

Next we can define a variable observable x with probability density $f(x)$ hence with n events the probability density of each is multiplied. The likelihood of λ can now be written

$$\mathcal{L}(\lambda) = \mathcal{P}(n|\lambda) \prod_{\text{event}}^n f(x) \quad (7.2)$$

Our likelihood also must take into account multiple regions (the signal region as well as the control regions) and so these likelihoods are multiplied together with their own distinct Poisson distributions

$$\mathcal{L}(\lambda) = \prod_r^{\text{regions}} (\mathcal{P}(n_r|\lambda_r) \prod_{\text{event}}^n f(x)) \quad (7.3)$$

A simultaneous fit maximizes this likelihood function and so produces signal strengths λ for each free parameter simultaneously. The λ values here represent predicted yields where $\lambda_{r,b} = \mu\lambda_{\text{sig}} + \lambda_{\text{bkg}}$. Maximizing the overall likelihood is made simpler by applying the natural logarithm (as the products between Poisson distributions become summations) and negating the likelihood, so as to take advantage of minimizing software.

Particle physics defines discovery with rigorous standards using hypothesis testing. The null hypothesis is considered $\mu = 0$ and is considered “background-only” while the alternative hypothesis is that there is a signal above the background. The probability that the null

CHAPTER 7. RESULTS

hypothesis is rejected (or that the signal is discovered) is defined as a p -value. The p -value can be converted to a number of Gaussian standard deviations and in high energy physics three σ significance (or a p -value of 1.35×10^{-3}) shows evidence while a five σ result (p -value 2.87×10^{-7}) is considered a discovery.

7.2 Unfolding

7.3 Results and future measurements

Chapter 8

Conclusions

Bibliography

- [1] M. Tanabashi, K. Hagiwara, K. Hikasa, K. Nakamura, Y. Sumino, F. Takahashi, J. Tanaka, K. Agashe, G. Aielli, C. Amsler, M. Antonelli, D. M. Asner, H. Baer, Sw. Banerjee, R. M. Barnett, T. Basaglia, C. W. Bauer, J. J. Beatty, V. I. Belousov, J. Beringer, S. Bethke, A. Bettini, H. Bichsel, O. Biebel, K. M. Black, E. Blucher, O. Buchmuller, V. Burkert, M. A. Bychkov, R. N. Cahn, M. Carena, A. Cecucci, A. Cerri, D. Chakraborty, M.-C. Chen, R. S. Chivukula, G. Cowan, O. Dahl, G. D'Ambrosio, T. Damour, D. de Florian, A. de Gouv  a, T. DeGrand, P. de Jong, G. Dissertori, B. A. Dobrescu, M. D'Onofrio, M. Doser, M. Drees, H. K. Dreiner, D. A. Dwyer, P. Eerola, S. Eidelman, J. Ellis, J. Erler, V. V. Ezhela, W. Fettscher, B. D. Fields, R. Firestone, B. Foster, A. Freitas, H. Gallagher, L. Garren, H.-J. Gerber, G. Gerbier, T. Gershon, Y. Gershtein, T. Gherghetta, A. A. Godizov, M. Goodman, C. Grab, A. V. Gritsan, C. Grojean, D. E. Groom, M. Grunewald, A. Gurtu, T. Gutsche, H. E. Haber, C. Hanhart, S. Hashimoto, Y. Hayato, K. G. Hayes, A. Hebecker, S. Heinemeyer, B. Heltsley, J. J. Hern  ndez-Rey, J. Hisano, A. H  cker, J. Holder, A. Holtkamp, T. Hyodo, K. D. Irwin, K. F. Johnson, M. Kado, M. Karliner, U. F. Katz, S. R. Klein, E. Klemp, R. V. Kowalewski, F. Krauss, M. Kreps, B. Krusche, Yu. V. Kuyanov, Y. Kwon, O. Lahav, J. Laiho, J. Lesgourgues, A. Liddle, Z. Ligeti, C.-J. Lin, C. Lippmann, T. M. Liss, L. Littenberg, K. S. Lugovsky, S. B. Lugovsky, A. Lusiani, Y. Makida,

BIBLIOGRAPHY

F. Maltoni, T. Mannel, A. V. Manohar, W. J. Marciano, A. D. Martin, A. Massoni, J. Matthews, U.-G. Meißner, D. Milstead, R. E. Mitchell, K. Mönig, P. Molaro, F. Moortgat, M. Moskovic, H. Murayama, M. Narain, P. Nason, S. Navas, M. Neubert, P. Nevski, Y. Nir, K. A. Olive, S. Pagan Griso, J. Parsons, C. Patrignani, J. A. Peacock, M. Pennington, S. T. Petcov, V. A. Petrov, E. Pianori, A. Piepke, A. Pomarol, A. Quadt, J. Rademacker, G. Raffelt, B. N. Ratcliff, P. Richardson, A. Ringwald, S. Roesler, S. Rolli, A. Romanouk, L. J. Rosenberg, J. L. Rosner, G. Rybka, R. A. Ryutin, C. T. Sachrajda, Y. Sakai, G. P. Salam, S. Sarkar, F. Sauli, O. Schneider, K. Scholberg, A. J. Schwartz, D. Scott, V. Sharma, S. R. Sharpe, T. Shutt, M. Silari, T. Sjöstrand, P. Skands, T. Skwarnicki, J. G. Smith, G. F. Smoot, S. Spanier, H. Spieler, C. Spiering, A. Stahl, S. L. Stone, T. Sumiyoshi, M. J. Syphers, K. Terashi, J. Terning, U. Thoma, R. S. Thorne, L. Tiator, M. Titov, N. P. Tkachenko, N. A. Tornqvist, D. R. Tovey, G. Valencia, R. Van de Water, N. Varelas, G. Venanzoni, L. Verde, M. G. Vincenter, P. Vogel, A. Vogt, S. P. Wakely, W. Walkowiak, C. W. Walter, D. Wands, D. R. Ward, M. O. Wascko, G. Weiglein, D. H. Weinberg, E. J. Weinberg, M. White, L. R. Wiencke, S. Willocq, C. G. Wohl, J. Womersley, C. L. Woody, R. L. Workman, W.-M. Yao, G. P. Zeller, O. V. Zenin, R.-Y. Zhu, S.-L. Zhu, F. Zimmermann, P. A. Zyla, J. Anderson, L. Fuller, V. S. Lugovsky, and P. Schaffner. Review of particle physics. *Phys. Rev. D*, 98:030001, Aug 2018.

- [2] F. Halzen and A. Martin. *Quarks and Leptons: An Introductory Course in Modern Particle Physics*. 1984.
- [3] John Ellis. Higgs Physics. (KCL-PH-TH-2013-49. KCL-PH-TH-2013-49. LCTS-2013-36. CERN-PH-TH-2013-315):117–168. 52 p, Dec 2013. 52 pages, 45 figures, Lectures

BIBLIOGRAPHY

presented at the ESHEP 2013 School of High-Energy Physics, to appear as part of the proceedings in a CERN Yellow Report.

- [4] Jonathan M. Butterworth, Guenther Dissertori, and Gavin P. Salam. Hard Processes in Proton-Proton Collisions at the Large Hadron Collider. *Ann. Rev. Nucl. Part. Sci.*, 62:387–405, 2012.
- [5] John M. Campbell, J. W. Huston, and W. J. Stirling. Hard Interactions of Quarks and Gluons: A Primer for LHC Physics. *Rept. Prog. Phys.*, 70:89, 2007.
- [6] W.J. Stirling. Private Communication... 2018.
- [7] Abdelhak Djouadi. The Anatomy of electro-weak symmetry breaking. I: The Higgs boson in the standard model. *Phys. Rept.*, 457:1–216, 2008.
- [8]
- [9] Jon Butterworth et al. PDF4LHC recommendations for LHC Run II. *J. Phys.*, G43:023001, 2016.
- [10] Georges Aad et al. Observation of a new particle in the search for the Standard Model Higgs boson with the ATLAS detector at the LHC. *Phys. Lett.*, B716:1–29, 2012.
- [11] Tom W. B. Kibble. The Standard Model of Particle Physics. 2014.
- [12] N.R. Council, D.E.P. Sciences, B.P. Astronomy, and C.E.P. Physics. *Elementary-Particle Physics: Revealing the Secrets of Energy and Matter*. Physics in a New Era: A Series. National Academies Press, 1998.

BIBLIOGRAPHY

- [13] Georges Aad et al. Combined measurements of Higgs boson production and decay using up to 80 fb^{-1} of proton-proton collision data at $\sqrt{s} = 13 \text{ TeV}$ collected with the ATLAS experiment. *Phys. Rev.*, D101(1):012002, 2020.
- [14] Georges Aad et al. Measurement of fiducial differential cross sections of gluon-fusion production of Higgs bosons decaying to $WW^* \rightarrow e\nu\mu\nu$ with the ATLAS detector at $\sqrt{s} = 8 \text{ TeV}$. *JHEP*, 08:104, 2016.
- [15] L Rossi. Superconductivity: Its Role, Its Success and Its Setbacks in the Large Hadron Collider of CERN. *Supercond. Sci. Technol.*, 23(CERN-ATS-2010-006):034001. 27 p, Jan 2010.
- [16] Rende Steerenberg. Lhc report: Protons: mission accomplished, 2018.
- [17] M. Aaboud, G. Aad, B. Abbott, O. Abdinov, B. Abelos, D.K. Abhayasinghe, S.H. Abidi, O.S. AbouZeid, N.L. Abraham, H. Abramowicz, and et al. Measurements of gluon-gluon fusion and vector-boson fusion higgs boson production cross-sections in the $h \rightarrow ww^* \rightarrow e\nu\mu\nu$ decay channel in pp collisions at $\sqrt{(s)}=13\text{tev}$ with the atlas detector. *Physics Letters B*, 789:508–529, Feb 2019.
- [18] M. Aaboud, G. Aad, B. Abbott, O. Abdinov, B. Abelos, S. H. Abidi, O. S. AbouZeid, N. L. Abraham, H. Abramowicz, and et al. Measurement of the higgs boson coupling properties in the $h \rightarrow zz^* \rightarrow 4\ell$ decay channel at $\sqrt{s} = 13 \text{ tev}$ with the atlas detector. *Journal of High Energy Physics*, 2018(3), Mar 2018.
- [19] M. Aaboud, G. Aad, B. Abbott, O. Abdinov, B. Abelos, S. H. Abidi, O. S. AbouZeid, N. L. Abraham, H. Abramowicz, H. Abreu, and et al. Measurements of higgs boson properties in the diphoton decay channel with 36 fb^{-1} of pp collision data at $\sqrt{s} = 13 \text{ tev}$ with the atlas detector. *Physical Review D*, 98(5), Sep 2018.

BIBLIOGRAPHY

- [20] M. Aaboud, G. Aad, B. Abbott, O. Abdinov, B. Abelos, D. K. Abhayasinghe, S. H. Abidi, O. S. AbouZeid, N. L. Abraham, H. Abramowicz, and et al. Cross-section measurements of the higgs boson decaying into a pair of τ -leptons in proton-proton collisions at $\sqrt{s}=13$ tev with the atlas detector. *Physical Review D*, 99(7), Apr 2019.
- [21] Search for squarks and gluinos in final states with jets and missing transverse momentum using 139 fb^{-1} of $\sqrt{s} = 13 \text{ TeV}$ pp collision data with the ATLAS detector. Technical Report ATLAS-CONF-2019-040, CERN, Geneva, Aug 2019.
- [22] CERN. A new schedule for the lhc and its successor, 2019.
- [23] M. Benedikt, P. Collier, V. Mertens, J. Poole, and K. Schindl. LHC Design Report. 3. The LHC injector chain. 2004.
- [24] Radiofrequency cavities. Sep 2012.
- [25] Lyndon Evans. The large hadron collider. *New Journal of Physics*, 9(9):335–335, sep 2007.
- [26] G Aad et al. The ATLAS Experiment at the CERN Large Hadron Collider. *JINST*, 3:S08003. 437 p, 2008. Also published by CERN Geneva in 2010.
- [27] ATLAS Collaboration. Technical Design Report for the ATLAS Inner Tracker Strip Detector. Technical Report CERN-LHCC-2017-005. ATLAS-TDR-025, CERN, Geneva, Apr 2017.
- [28] Jet global sequential corrections with the ATLAS detector in proton-proton collisions at $\text{sqrt}(s) = 8 \text{ TeV}$. Technical Report ATLAS-CONF-2015-002, CERN, Geneva, Mar 2015.

BIBLIOGRAPHY

- [29] Morad Aaboud et al. Jet reconstruction and performance using particle flow with the ATLAS Detector. *Eur. Phys. J.*, C77(7):466, 2017.
- [30] JET ETmiss group. Public plots, 2019.
- [31] Matteo Cacciari, Gavin P. Salam, and Gregory Soyez. The anti- k_t jet clustering algorithm. *JHEP*, 04:063, 2008.
- [32] Morad Aaboud et al. Determination of jet calibration and energy resolution in proton-proton collisions at $\sqrt{s} = 8$ TeV using the ATLAS detector. 2019.
- [33] Object-based missing transverse momentum significance in the ATLAS detector. Technical Report ATLAS-CONF-2018-038, CERN, Geneva, Jul 2018.
- [34] Morad Aaboud et al. Performance of missing transverse momentum reconstruction with the ATLAS detector using proton-proton collisions at $\sqrt{s} = 13$ TeV. *Eur. Phys. J.*, C78(11):903, 2018.
- [35] ATLAS Collaboration. 14(12):P12006–P12006, dec 2019.
- [36] J. Illingworth and J. Kittler. A survey of the hough transform. *Computer Vision, Graphics, and Image Processing*, 44(1):87 – 116, 1988.
- [37] Performance of the ATLAS Silicon Pattern Recognition Algorithm in Data and Simulation at $\sqrt{s} = 7$ TeV. Technical Report ATLAS-CONF-2010-072, CERN, Geneva, Jul 2010.
- [38] G. Aad, B. Abbott, J. Abdallah, O. Abdinov, B. Abelos, R. Aben, M. Abolins, O. S. AbouZeid, N. L. Abraham, and et al. Muon reconstruction performance of the atlas

BIBLIOGRAPHY

detector in proton–proton collision data at

$$\sqrt{s}$$

$s = 13$ tev. *The European Physical Journal C*, 76(5), May 2016.

- [39] Georges Aad et al. Measurement of the muon reconstruction performance of the ATLAS detector using 2011 and 2012 LHC proton–proton collision data. Measurement of the muon reconstruction performance of the ATLAS detector using 2011 and 2012 LHC proton-proton collision data. *Eur. Phys. J. C*, 74(CERN-PH-EP-2014-151. CERN-PH-EP-2014-151):3130. 34 p, Jul 2014. Comments: 21 pages plus author list + cover pages (34 pages total), 21 figures, 2 tables, submitted to EPJC, All figures including auxiliary figures are available at <http://atlas.web.cern.ch/Atlas/GROUPS/PHYSICS/PAPERS/PERF-2014-05/>.
- [40] Johannes Josef Junggeburth. Muon identification and reconstruction efficiencies in full Run-2 dataset. Technical Report ATL-COM-PHYS-2019-1001, CERN, Geneva, Aug 2019.
- [41] G. Aad, B. Abbott, J. Abdallah, A. A. Abdelalim, A. Abdesselam, O. Abdinov, B. Abi, M. Abolins, H. Abramowicz, and et al. Jet energy measurement with the atlas detector in proton-proton collisions at $\sqrt{s} = 7$ TeV. *The European Physical Journal C*, 73(3), Mar 2013.
- [42] Miha Zgubič, Johannes Josef Junggeburth, and K. Technical report.
- [43] Travis Martin and W Martin. Analyzing new neutral gauge bosons at the lhc using third generation fermions. 05 2020.

BIBLIOGRAPHY

- [44] Kyle Cranmer. Practical statistics for the lhc, 2015.