

**Vector boson fusion Higgs boson  
fiducial cross-section measured in the  
 $WW^* \rightarrow \ell\nu\ell\nu$  decay channel with the  
ATLAS detector**

A Dissertation

Presented to

The Faculty of the Graduate School of Arts and Sciences

Brandeis University

Department of Physics

Professor Gabriella Sciolla, Advisor

In Partial Fulfillment  
of the Requirements for the Degree  
Doctor of Philosophy

by

Laura Bergsten

June, 2020

This dissertation, directed and approved by Laura Bergsten's committee, has been accepted and approved by the Graduate Faculty of Brandeis University in partial fulfillment of the requirements for the degree of:

**DOCTOR OF PHILOSOPHY**

Eric Chasalow, Dean of Arts and Sciences

Dissertation Committee:

Professor Gabriella Sciolla, Chair

Craig Blocker

Alessandro Tricoli

©Copyright by

Laura Bergsten

2020

# Abstract

## **Vector boson fusion Higgs boson fiducial cross-section measured in the $WW^* \rightarrow \ell\nu\ell\nu$ decay channel with the ATLAS detector**

A dissertation presented to the Faculty of  
the Graduate School of Arts and Sciences of  
Brandeis University, Waltham, Massachusetts

by Laura Bergsten

The Standard Model vector boson fusion Higgs boson fiducial cross-section is measured in the  $H \rightarrow WW^* \rightarrow \ell\nu\ell\nu$  decay channel using  $139 \text{ fb}^{-1}$  of proton-proton collisions. Collisions from the Large Hadron Collider at  $\sqrt{s} = 13 \text{ TeV}$  are recorded by the ATLAS detector. The fiducial cross-section measured to be  $\text{---}$ .

# Contents

<b>Abstract</b>	v
<b>1 Introduction</b>	1
<b>2 Theory</b>	3
2.1 Standard Model . . . . .	3
2.2 LHC Physics/Phenomenology . . . . .	19
2.3 Brief history of SM tests . . . . .	26
2.4 Measurement motivation . . . . .	31
<b>3 The LHC and the ATLAS detector</b>	33
3.1 ATLAS, A Toroidal LHC ApparatuS . . . . .	36
3.2 The High-Luminosity LHC and Inner Tracker (ITk) . . . . .	38
<b>4 Tracking and Isolation in ATLAS</b>	49
4.1 Tracking . . . . .	50
4.2 Electrons . . . . .	51
4.3 Muons . . . . .	55
4.4 Jets . . . . .	65
4.5 Missing transverse energy . . . . .	70
<b>5 Event Selection</b>	74
5.1 Data and Monte Carlo samples . . . . .	74
5.2 Object definitions . . . . .	77
5.3 Common Observables . . . . .	82
5.4 Event selection . . . . .	84
<b>6 Backgrounds and Systematics</b>	96
6.1 Backgrounds . . . . .	96
6.2 Systematic uncertainties . . . . .	131

*CONTENTS*

<b>7 Results</b>	<b>145</b>
7.1 Statistical analysis . . . . .	145
7.2 Fiducial and inclusive cross sections . . . . .	154
7.3 Differential cross-section measurements . . . . .	157
7.4 Future results and measurements . . . . .	160
<b>8 Conclusions</b>	<b>163</b>

# Chapter 1

## Introduction

The Standard Model describes particles and their interactions at fundamental scales and provides the theoretical tools to test the limits of that knowledge. No significant deviation has thus far been observed between Standard Model predictions and experimental results, but the search continues. Our observable universe shows evidence of physics beyond the current theory and the search for evidence of what this new physics might be motivates continued measurements of Standard Model parameters. Chapter 2 describes the theoretical motivations for this thesis and the key experimental results that precede it.

The ATLAS detector is a complex experiment tailor-made to precisely measure a wide-range of Standard Model parameters. The design and performance of the ATLAS detector is summarized in Chapter 3. This thesis uses prescriptions for reconstruction, identification, isolation and measurements of efficiency, scale, and resolution from a number of dedicated ATLAS performance groups. Chapter 4 discusses methodology for these recommendations and their particular use in the final state of our measured decay mode.

This thesis focuses on the fiducial cross-section of vector boson fusion Higgs boson production in the  $WW^*$  decay channel. This decay channel has a final state that requires

## *CHAPTER 1. INTRODUCTION*

accurate reconstruction of leptons and jets within the detector and greatly benefits from increased statistics available from the latest data. Chapter 5 summarizes the dataset, simulations and observables used in the cross-section measurement as well as the signal event selection. Chapter 6 describes each major background and uncertainty that affects the final measurement and Chapter 7 details the statistical analysis and final results.

The  $H \rightarrow WW^* \rightarrow \ell\nu\ell\nu$  cross-section measurement is the cornerstone of this thesis and has been my primary focus for the past two years. I also had the opportunity to work on detector upgrades and muon performance studies during my Ph.D. studies so these topics are described with particular attention in Chapters 3 and 4.

All measurements and studies included here were made possible through my analysis team and the advice and software from countless others. All figures contained in this thesis are cited from either publications or individuals unless I was there direct author.

# Chapter 2

## Theory

The measurement at the heart of this thesis can only be understood within the context of a vast amount of preceding theoretical and experimental work. I have tried to condense and summarize key concepts that will motivate the central measurement's strategy and results. This chapter begins with a brief summary of the Standard Model itself, first describing fundamental particles and their forces before delving into a succinct mathematical formulation. Next, the chapter discusses the history of the Standard Model and crucial tests of the theory up until current work at the Large Hadron Collider (LHC). The next section outlines some of the recent physics analyses at the LHC with a focus on Higgs boson measurements. Finally, I introduce my thesis' main focus, fiducial cross-section measurements of Vector-Boson-Fusion (VBF) Higgs decaying into two W-bosons.

### 2.1 Standard Model

The Standard Model (SM) is one of the most successful scientific theories to date. Its predictions encompass all of the visible universe and continue to undergo careful testing.

## *CHAPTER 2. THEORY*

The SM combines three forces- electromagnetic, weak, and strong - but it is not complete. One known force, gravity, is not included in the Standard Model and further questions, like an explanation for dark matter and dark energy, remain.

### **2.1.1 Particles and forces**

The particles we define in high energy physics are the most minute portions of observable matter. They are generally considered point-like, have no internal structure, and cannot be further split. Each particle we can define has a unique set of quantum numbers and its own anti-particle (with the same mass and spin, but opposite electrical charge and quantum numbers).

Particles can be sorted into two distinct groups: bosons, with integer spin, and fermions, with half-integer spin. Bosons are ‘force carriers’ meaning they are exchanged any time particles interact. Fermions are at the heart of all conventional matter and can be split further into two categories- leptons and quarks. Quarks have fractional integer charge and interact strongly, while leptons have integer charge and interact solely through the weak or electromagnetic forces. Both quarks and leptons are made of three generations of particles, each heavier and more unstable than the previous. Charts showing quark/lepton families and their key quantum numbers are shown below in Figure 2.1. Each generation of quarks and leptons contains a particle doublet. Each lepton doublet contains a charged lepton and a neutrino while each quark doublet contains one  $+2/3$  charged particle and one with a  $-1/3$  charge. Each lepton and quark also has an anti-particle. All conventional, stable matter is made from the first generation of quarks and leptons.

There are four gauge bosons and one scalar boson predicted through the Standard Model. These correspond to three fundamental forces in nature (the fourth, gravity, is so small on the

## CHAPTER 2. THEORY

scale of particle interactions as to not be considered). The strongest force on the subatomic scale is the strong force - this is mediated by the gluon - and works primarily to bind quarks together to form composite particles like protons or neutrons. The electromagnetic force is about 60 times weaker than the strong force and is mediated by the photon. This force accounts for all electric interactions like that between an electron and an atomic nucleus. Finally, the weak force ( $10^4$  times weaker than the EM) mediates  $\beta$ -decay with massive  $W$  and  $Z$  bosons. The final boson predicted by the Standard Model is the Higgs boson. The only scalar boson, it has no charge or intrinsic spin. The Higgs gives mass to all other particles through Spontaneous Symmetry Breaking, which will be described in later sections.

### Standard Model of Elementary Particles




Figure 2.1: Three generations of quarks and leptons are shown along with all SM bosons [1]

## CHAPTER 2. THEORY

Photons are massless, spin-1 particles and mediate all electromagnetic interactions. They couple directly to any particle with electric charge- so quarks, leptons, and  $W/Z$  bosons but not neutrinos. Since the photon is massless, the electromagnetic force can operate on infinitely long scales but its force decreases with  $1/r^2$ .

Gluons are massless particles with no charge and spin-1. They couple to color charges, which are a property only of quarks and gluons. Each quark has one of three colors (RGB). Colors are conserved ‘charges’ similar to electric charge. Quarks are never found alone as they couple so strongly to one another as to be confined in groups of two or three. These groups are “color-confined” meaning the quarks contain colors which add up to a color neutral sum. For instance, a two quark meson  $u\bar{u}$  may have colors R and anti-R while a three quark hadron  $uud$  (proton) may have colors R, G, and B. Gluons differ from photons in that they are not neutral in the charge they couple to. Gluons have two colors (8 total combinations) and can thus couple to each other. This makes the strong force distinct from the electromagnetic and has implications for long-distance interactions.

$W$  and  $Z$  bosons, unlike gluons and photons, are massive. However, like their other gauge boson counterparts, they have spin-1 and mediate a charge (weak).  $W^\pm$  mediates charged-current interactions which can violate flavor conservation between quarks and/or leptons and their neutrinos.  $Z^0$  mediates neutral-current interactions which conserve flavor.  $W^\pm$  bosons contain electric charge so can interact through EM as well. In addition,  $W$  and  $Z$  bosons contain weak charge (as do all fermions) so can self-couple as well as couple with all fermions.

The Higgs boson will be further motivated and described in later sections but suffice to say it is a massive spin-0 particle which couples to all particles with mass (including itself). It does not mediate any force but it is still an integral part of the Standard Model.

## CHAPTER 2. THEORY

### 2.1.2 Gauge Invariance

According to Noether's theorem, for every continuous transformation of a field that leaves the Lagrangian invariant, there is a conserved current. In other words, symmetries found in physical theories lead to conservation laws (and vice-versa). The Standard Model is a gauge theory built on symmetries; all interactions between particles result from requiring the theory to be invariant under local gauge transformations. Each part of the Standard Model- from quantum electrodynamics (QED) to quantum chromodynamics (QCD) - is a gauge theory on its own. Each part has gauge invariance symmetries. In this section I step through the basic mathematic formalism for QED, QCD, and the combined electro-weak theory to illustrate the physical ramifications of gauge invariance and set the stage for the Higgs mechanism. The following sections are written with guidance from text [2].

### Quantum Electrodynamics

Quantum electrodynamics (QED) is the first, and simplest, physical gauge theory, describing how light and matter interact even under relativistic conditions. The theory produces extremely good agreement with experiment due to the success of perturbative QED calculations. Entire textbooks are dedicated to QED formalism and predictions. Here I will highlight only the effects of its local gauge invariance symmetry and generating the full QED Lagrangian beginning with the Dirac Lagrangian of a free fermion.

The Dirac Lagrangian describes a free fermion of mass  $m$

$$\mathcal{L} = i\psi\gamma^\mu\partial_\mu\psi - m\bar{\psi}\psi, \quad (2.1)$$

where  $\psi$  is a Dirac spinor and  $\gamma^\mu$  represent the Dirac matrices. To demonstrate local gauge

## CHAPTER 2. THEORY

invariance we need to transform

$$\psi(x) \rightarrow e^{i\alpha(x)}\psi(x), \quad (2.2)$$

where  $\alpha(x)$  depends on space and time arbitrarily. Directly substituting this into our Lagrangian shows that  $\mathcal{L}$  is not invariant, and the  $\partial_\mu$  term causes this

$$\partial_\mu\psi \rightarrow e^{i\alpha(x)}\partial_\mu\psi + ie^{i\alpha(x)}\partial_\mu\alpha. \quad (2.3)$$

In order to mandate the theory is invariant we need to change this term to the “covariant derivative”  $D_\mu$  which transforms

$$D_\mu\psi \rightarrow e^{i\alpha(x)}D_\mu\psi. \quad (2.4)$$

The “covariant derivative” must contain a vector field  $A_\mu$  and this field must transform so as to cancel with the unwanted part of the transformed  $D_\mu$  in order to transform as required by gauge invariance.

$$D_\mu \equiv \partial_\mu - ieA_\mu \quad (2.5)$$

where

$$A_\mu \rightarrow A_\mu + \frac{1}{e}\partial_\mu\alpha. \quad (2.6)$$

Now the original Dirac equation is replaced with the following:

$$\mathcal{L} = \bar{\psi}(i\gamma^\mu\partial_\mu - m)\psi + e\bar{\psi}\gamma^\mu\psi A_\mu. \quad (2.7)$$

By requiring local gauge invariance we have introduced a gauge field  $A_\mu$  which couples to the Dirac particle just as the photon. In fact, if we take this as the photon gauge field and so add a kinetic energy term (which is also local gauge invariant!) we find the Lagrangian

## CHAPTER 2. THEORY

of QED:

$$\mathcal{L} = \bar{\psi}(i\gamma^\mu\partial_\mu - m)\psi + e\bar{\psi}\gamma^\mu\psi A_\mu - \frac{1}{4}F_{\mu\nu}F^{\mu\nu}. \quad (2.8)$$

One can also see that adding a mass term to the Lagrangian for the new field ( $\frac{1}{2}m^2A_\mu A^\mu$ ) would break gauge invariance, indicating the photon must be massless. From the free fermion Lagrangian, imposing local gauge invariance leads to the full interacting field theory of QED. This is not a curiosity but an essential component of the theory, and the use of local gauge symmetry in deriving particle interactions does not end here.

### Quantum chromodynamics

Quantum chromodynamics differs from QED in a few crucial ways. Since quark color fields exist, the QED  $U(1)$  gauge group is replaced with  $SU(3)$  and the free Lagrangian contains indices  $j$  to denote the three color fields:

$$\mathcal{L} = \bar{q}_j(i\gamma^\mu\partial_\mu - m)q_j. \quad (2.9)$$

QCD also carries three quark flavors, which will be ignored here for simplicity. The QCD group is non-Abelian since not all generators of the group commute with each other. These generators will be defined as  $T_a$  where  $a = 1, \dots, 8$  and are linearly independent traceless  $3 \times 3$  matrices (the Gell-Mann matrices  $\lambda_a$  are conventional). The local color phase transformation required is thus

$$q(x) \rightarrow e^{i\alpha_a(x)T_a}q(x). \quad (2.10)$$

We can consider an infinitesimal phase transformation as

$$\begin{aligned} q(x) &\rightarrow [1 + i\alpha_a(x)T_a]q(x), \\ \partial_u q &\rightarrow (1 + i\alpha_a T_a)\partial_\mu q + iT_a q \partial_\mu \alpha_a. \end{aligned} \quad (2.11)$$

## CHAPTER 2. THEORY

Just as in the QED example, the last line breaks the invariance of  $\mathcal{L}$  and we can proceed similarly by introducing a new gauge field (or in this case eight) called  $G_\mu^a$  which transform

$$G_\mu^a \rightarrow G_\mu^a - \frac{1}{g} \partial_\mu \alpha_a - f_{abc} \alpha_b G_\mu^c. \quad (2.12)$$

The last term added here is to cope with the non-Abelian nature of QCD <sup>1</sup>. Just as in QED this invariance forms a covariant derivative:

$$D_\mu = \partial_\mu u + igT_a G_\mu^a. \quad (2.13)$$

Replacing the derivative in our Lagrangian and adding a gauge invariant energy term for each of the  $G_m u^a$  fields ( $\frac{1}{4} G_{\mu\nu}^a G_{\mu\nu}^a$ ) yields the final gauge invariant QCD Lagrangian:

$$\mathcal{L} = \bar{q}(i\gamma^\mu \partial_\mu - m)q - g(\bar{q}\gamma^\mu T_a q)G_\mu^a - \frac{1}{4} G_{\mu\nu}^a G_{\mu\nu}^a. \quad (2.14)$$

Just as in the QED case, imposing local color phase invariance produced 8 new interacting fields with coupling  $g$ . These are the gluon fields and just like photons, local gauge invariance requires them to be massless. Unlike the QED case, this Lagrangian's new kinetic term includes self-interaction between the gauge bosons - another key feature of QCD that is mandated by local color phase invariance. Gluons themselves must carry color charge and therefore they self-couple. The structure of these self coupling terms and their single coupling strength  $g$  are uniquely determined by gauge invariance.

---

<sup>1</sup>Non-Abelian here means all the generators  $T_a$  commute with each other.

## CHAPTER 2. THEORY

### Electroweak unification

Thus far, I have summarized the theoretical backgrounds for symmetries (and so conserved quantities) in both quantum electrodynamics and chromodynamics. The weak force is the final Standard Model force and weak interactions are mediated by  $Z$  and  $W$  bosons. Unlike the gluons and photons of QCD and QED, these gauge bosons are massive. This is explained through spontaneous symmetry breaking of the electroweak force, which is outlined in the following section. Assuming that  $W/Z$  bosons are massive, the weak force can be combined with QED and a central electroweak force (with its associated symmetries) can be described.

The weak neutral current  $J_\mu^{NC}$  and weak charged currents ( $J_\mu$ ,  $J_\mu^\dagger$ ) can form a symmetry group for weak interactions. The charged currents correspond to the charged weak interaction through  $W^\pm$  bosons while the neutral current is associated with the  $Z^0$  boson:

$$\begin{aligned} J_\mu &= \bar{\nu}_L \gamma_\mu \nu_L, \\ J_\mu^\dagger &= \bar{e}_L \gamma_\mu \nu_L J_\mu^3 = \frac{1}{2} \bar{\nu}_L \gamma_\mu \nu_L - \frac{1}{2} \bar{e}_L \gamma_\mu e_L, \end{aligned} \tag{2.15}$$

where  $L$  denotes that these are left-handed spinors. The charged currents can be written as a doublet using the Pauli spin matrices  $\tau_i$  where  $\tau_\pm = \frac{1}{2}(\tau_1 \pm i\tau_2)$  and

$$\chi_L = \begin{bmatrix} \nu \\ e^- \end{bmatrix} \tag{2.16}$$

as

$$\begin{aligned} J_\mu^+(x) &= \bar{\chi}_L \gamma_\mu \tau_+ \chi_L, \\ J_\mu^-(x) &= \bar{\chi}_L \gamma_\mu \tau_- \chi_L, \\ J_\mu^3(x) &= \bar{\chi}_L \gamma_\mu \frac{1}{2} \tau_i \chi_L \text{ with } i = 1, 2, 3 \end{aligned} \tag{2.17}$$

## CHAPTER 2. THEORY

The corresponding charge  $T^i = \int J_0^i(x)d^3x$  can be introduced so we have an  $SU(2)_L$  algebra

$$[T^i, T^j] = i\epsilon_{ijk}T^k. \quad (2.18)$$

Unfortunately while these currents create an  $SU(2)$  group, they do not correspond with the weak neutral current symmetry in an obvious way. Unlike the charged currents, the neutral current has a right-handed component. One way to attain the expected symmetry is to add in the electromagnetic current, as it is a neutral current with left and right-handed components

$$j_\mu^{em}(x) = -\bar{e}_R \gamma_\mu e_R - \bar{e}_L \gamma_\mu e_L \quad (2.19)$$

so the the electromagnetic current  $j_\mu$  can be written using the coupling  $e$

$$j_\mu = ej_\mu^{em} = e\bar{\psi}\gamma_\mu Q\psi \quad (2.20)$$

with  $Q$  the charge operator and generator of the  $U(1)$  symmetry group of EM. In order to “save” the symmetry of the weak neutral current, we can define an electromagnetic current  $j_\mu^Y$ , or the weak hypercharge current, that is unchanged by  $SU(2)_L$  transformations. We define a weak hypercharge  $Y$  and its current  $j_\mu^Y$

$$\begin{aligned} Q &= T^3 + \frac{Y}{2}, \\ j_\mu^Y &= \bar{\psi}\gamma_\mu Y\psi. \end{aligned} \quad (2.21)$$

The combined current

$$j_\mu^{em} = J_\mu^3 + \frac{1}{2}j_\mu^Y \quad (2.22)$$

now generates the symmetry group  $U(1)_Y$  and so the electromagnetic interaction and weak interaction are combined into one  $SU(2)_L \times U(1)_Y$  group. While unified in this way, the two

## CHAPTER 2. THEORY

forces still have independent coupling strengths. This brief introduction into electroweak unification is not the complete picture- EM and weak *interactions* still have to be unified. In the Standard Model framework electroweak currents just have to be coupled to vector bosons to complete unification. In the electroweak  $SU(2)_L \times U(1)_Y$  group there is an isotriplet of vector fields  $W_\mu^i$  coupled with strength  $g$  to the weak isospin current  $J_\mu^i$ , while a single vector field  $B_\mu$  is coupled to the weak hypercharge current  $j_\mu^Y$  with strength  $g'/2$ . The electroweak Lagrangian interaction term can thus be defined:

$$-ig(J^i)^\mu W_\mu^i - i\frac{g'}{2}(j^Y)^\mu B_\mu. \quad (2.23)$$

This summary of the unified electroweak force will be the starting point for a derivation of the Higgs boson and an explanation for mass of the weak force's vector bosons (and all the fermions). The electroweak theory is unique in its calculability even at higher order scales. Consequently, theoretical uncertainties are relatively low and many deviations from theory could potentially be observed at current energy scales. The measurement central to my thesis probes for such discrepancies in electroweak theory. The mechanisms for this will be explained in the last section in this chapter.

### Spontaneous Symmetry Breaking

Unlike QED and QCD, the weak force is mediated by massive gauge bosons. We therefore can not apply the same gauge invariance prescription that we did in the last sections. If a mass term is added to the Lagrangian we break the gauge invariance we aimed to find. If we instead ignore the gauge invariance and add a mass term to the Lagrangian, all predictive power of the theory is lost due to unrenormalizable divergences. With “spontaneous symmetry breaking” we can gain massive gauge bosons while maintaining the integrity of the

## CHAPTER 2. THEORY

theory. In this section, I first describe the "spontaneous symmetry breaking" mechanism in terms of an Abelian theory composed of complex scalar fields to illustrate the overall strategy. This mechanism is then applied to the non-Abelian electroweak theory to gain massive weak gauge bosons,  $W^{+/-}$  and  $Z$ , with the Higgs field appearing as a 'spontaneous' result.

The Lagrangian for a  $U(1)$  gauge symmetry can be given as

$$\phi \rightarrow e^{i\alpha(x)}\phi. \quad (2.24)$$

As in the QED case, we introduce a gauge field  $A_\mu$  and covariant derivative  $D_\mu = \partial_\mu - ieA_\mu$  to obtain the gauge invariant Lagrangian,

$$\mathcal{L} = (\partial^\mu + ieA^\mu)\phi^*(\partial_\mu u - ieA_\mu)\phi - \mu^2\phi^*\phi - \lambda(\phi^*\phi)^2 - \frac{1}{4}F_{\mu\nu}F^{\mu\nu}. \quad (2.25)$$

In this example if  $\mu^2 > 0$  we gain back the QED Lagrangian for a charged scalar particle of mass  $\mu$  with an additional self-interaction term. However, if we take  $\mu^2 < 0$  the potential  $V(\phi^*\phi) = \mu^2\phi^*\phi - \lambda(\phi^*\phi)^2$  acquires a non-zero vacuum expectation value (v.e.v.) and there is a set of equivalent minima shown in Figure 2.2. Choosing one of these minima spontaneously breaks the potential's rotational symmetry. We can perturbatively expand the field about a minimum through

## CHAPTER 2. THEORY




Figure 2.2: Higgs potential when  $\mu^2 < 0$ , choosing a minima spontaneously breaks the  $U(1)$  rotational symmetry [3]

$$\phi(x) = \sqrt{\frac{1}{2}[\nu + \eta(x) + i\xi(x)]}. \quad (2.26)$$

We substitute this perturbation into the Lagrangian and gain

$$\mathcal{L}' = \frac{1}{2}(\partial_\mu \xi)^2 + \frac{1}{2}(\partial_\mu \eta)^2 - \nu^2 \lambda \eta^2 + \frac{1}{2}e^2 \nu_\mu^A A^\mu - e\nu A_\mu \partial^\mu \xi - \frac{1}{4}F_{\mu\nu} F^{\mu\nu} + \text{interaction terms}. \quad (2.27)$$

Three particles emerge here: a massless Goldstone boson  $\xi$ , a massive vector  $A_\mu$  with  $m_A = e\nu$ , and a massive scalar  $\eta$  with  $m_\eta = \sqrt{2\lambda\nu^2}$ . However, the number of particles does not correspond to the expected polarization degrees of freedom expected. A longitudinal polarization was added, creating an unphysical field, and injecting extra degrees of freedom.

To eliminate the unphysical field we can substitute a new set of fields:

$$\phi \rightarrow \sqrt{\frac{1}{2}(\nu + h(x))e^{i\theta(x)/\nu}} \quad (2.28)$$

and

$$A_\mu \rightarrow A_\mu + \frac{1}{e\nu} \partial_\mu \theta. \quad (2.29)$$

## CHAPTER 2. THEORY

Introducing these substitutions, the Goldstone boson field disappears and the new Lagrangian becomes

$$\mathcal{L} = \frac{1}{2}(\partial_\mu h)^2 - \lambda\nu^2 h^2 + \frac{1}{2}e^2\nu^2 A_\mu^2 - \lambda\nu h^3 - \frac{1}{4}\lambda h^4 + \frac{1}{2}e^2 A_\mu^2 h^2 + \nu e^2 A_\mu^2 h - \frac{1}{4}F_{\mu\nu}F^{\mu\nu}. \quad (2.30)$$

Here the degrees of freedom before our substitutions remains the same and a massive boson  $A_\mu$  is preserved along with a massive scalar  $h$ . The “Higgs mechanism” applied to a scalar field succeeded in creating a massive boson and determined the existence of a massive scalar boson. This same mechanism can be applied in the more complicated Standard Model electroweak field. Through electroweak symmetry breaking we not only gain massive gauge bosons and a massive scalar boson (the Higgs) but we also gain a way to calculate testable Standard Model predictions for many other quantities. We start with the  $SU(2)\times U(1)$  gauge symmetry of electroweak interactions derived in the previous section. In order to gain masses for three gauge bosons and keep the photon massless we need at least 3 degrees of freedom added and a simple choice is to use the  $SU(2)$  doublet of scalar fields  $\phi$ , with four fields in an isospin doublet of weak hypercharge  $Y = 1$ :

$$\mathcal{L} = (D^\mu\phi)^\dagger(D_\mu\phi) - \mu^2\phi^\dagger\phi - \lambda(\phi^\dagger\phi)^2, \quad (2.31)$$

where  $\phi$  is a  $SU(2)$  doublet of complex scalar fields

$$\phi = \sqrt{\frac{1}{2}} \begin{bmatrix} \phi_1 + i\phi_2 \\ \phi_3 + i\phi_4 \end{bmatrix}. \quad (2.32)$$

Local gauge invariance can be achieved just as in the  $U(1)$  case with the covariant derivative,

## CHAPTER 2. THEORY

though now a bit more complex:

$$D_\mu = \partial_\mu + ig\frac{\tau_a}{2}W_\mu^a \quad (2.33)$$

with three gauge fields  $W_\mu^a(x)$  and  $a = 1, 2, 3$ . An infinitesimal transformation is defined as

$$\phi(x) \rightarrow \phi'(x) = (1 + i\alpha(x) \cdot \tau/2)\phi(x) \quad (2.34)$$

and so the Lagrangian potential becomes

$$V(\phi) = \mu^2\phi^\dagger\phi + \lambda(\phi^\dagger\phi)^2. \quad (2.35)$$

Once again if we choose the conditions  $\mu^2 < 0$  and  $\lambda > 0$  there is rotational symmetry in our choice of vacuum expectation value. In this case, the choice of v.e.v. is limited. For the photon to remain massless, the vacuum must be invariant under  $U(1)$  (or electromagnetic) transformations, and not be charged in either direction (charge conservation). Thus the chosen minima to spontaneous break electroweak symmetry is

$$\phi_0 = \sqrt{\frac{1}{2}} \begin{bmatrix} 0 \\ \nu \end{bmatrix}. \quad (2.36)$$

Next, substituting the vacuum expectation value  $\phi_0$  for  $\phi(x)$  and expanding perturbatively yields

$$\phi(x) \rightarrow \begin{bmatrix} 0 \\ \sqrt{\frac{1}{2}(\nu + H(x))} \end{bmatrix}. \quad (2.37)$$

Fully expanding this term in the Lagrangian gives a complex and illuminating result, the Goldstone bosons have been consumed and there is only a Higgs field ( $H(x)$ ) remaining. Next, masses for the vector bosons are found from expanding one key parameter in the

## CHAPTER 2. THEORY

Lagrangian

$$|(-ig\frac{\tau}{2} \cdot W_\mu - i\frac{g'}{2}B_\mu)\phi|^2 = \frac{1}{8} \left| \begin{bmatrix} gW_\mu^3 + g'B_\mu & g(W_\mu^1 - iW_\mu^2) \\ g(W_\mu^1 + iW_\mu^2) & igW_\mu^3 + g'B_\mu \end{bmatrix} \begin{bmatrix} 0 \\ \nu \end{bmatrix} \right|^2. \quad (2.38)$$

Expanding further and substituting  $W^\pm = (W^1 \pm iW^2)/\sqrt{2}$  gives the result

$$|(-ig\frac{\tau}{2} \cdot W_\mu - i\frac{g'}{2}B_\mu)\phi|^2 = (\frac{1}{2}\nu g)^2 W_\mu^+ W^{-\mu} + \frac{1}{8}(W_\mu^3, B_\mu) \begin{bmatrix} g^2 & -gg' \\ -gg' & g'^2 \end{bmatrix} \begin{bmatrix} W^{\mu 3} \\ B^\mu \end{bmatrix}. \quad (2.39)$$

It is immediately clear that there is a mass-term for the  $W^\pm$ ,  $M_W = \frac{1}{2}\nu g$ . Masses for the photon and  $Z$ -boson are also apparent after expanding the last final term

$$\frac{1}{8}\nu^2(g^2(W_3^\mu)^2 - 2gg'W_\mu^3B^\mu + g'^2B_\mu^2) = \frac{1}{8}\nu^2(gW_\mu^3 - g'B_\mu)^2, \quad (2.40)$$

and using the substitutions

$$\begin{aligned} A_\mu &= \frac{g'W_\mu^3 + gB_\mu}{\sqrt{(g^2 + g'^2)}} \text{ with } M_A = 0, \text{ and} \\ Z_\mu &= \frac{gW_\mu^3 - g'B_\mu}{\sqrt{(g^2 + g'^2)}} \text{ with } M_Z = \frac{1}{2}\nu\sqrt{(g^2 + g'^2)}. \end{aligned} \quad (2.41)$$

Now the Higgs field exists just as in the previous example and the theory contains a massive scalar boson and three massive vector gauge fields - one for each of the  $W^\pm$  and  $Z$  bosons. The Goldstone bosons degrees of freedom were used to give mass to the vector bosons. Choosing a ground state and so breaking the gauge symmetry does not eliminate this symmetry altogether, since the theory is still renormalizable. Fermion masses can also be derived from their interactions with the Higgs boson using this Lagrangian. These derivations can be used to predict masses of bosons and fermions and couplings to the Higgs boson. It

## *CHAPTER 2. THEORY*

is important to note that the Higgs mechanism gives mass to all fermions and massive gauge bosons, it does not determine what the Higgs mass ought to be. This is left as an empirical input to the theory that can then be used to calculate other observables.

The Standard Model has been proven over decades to be an incredibly robust theory. Since 2010, the Large Hadron Collider (LHC) has become its key testing ground.

## **2.2 LHC Physics/Phenomenology**

The Large Hadron Collider (LHC) is the foremost Standard Model testing ground and the proton-proton collisions recorded through ATLAS, CMS, ALICE, and LHCb have demonstrated the breadth and accuracy of the theory. Fermion and gauge boson masses and couplings, including the mass of the Higgs boson, have been measured with high precision. In the next chapter the mechanics of the LHC and ATLAS detector will be discussed. Here I will introduce the motivations and observations of LHC physics. This section will begin with the mechanics of proton-proton collisions and their decay products, then discuss the concept of decay cross-sections, and finally focus more closely on the Higgs boson and its properties.

One of the LHC's central goal was to discover the missing Standard Model Higgs boson. The protons in the LHC collide at a center-of-mass energy of 13 TeV, but began at half that in 2010. The electroweak symmetry breaking scale was theoretically known to be between 100-1000 GeV and so probing at 7 TeV provided near certainty of finding either the Higgs or an inconsistency in the Standard Model. The motivation for a proton collider was multifaceted. Foremost, using the tunnels built for the electron-positron detector LEP with protons allows the collider to reach higher energies, as protons do not lose as much energy to synchrotron radiation as electrons. However, proton collisions have added complexity from

## CHAPTER 2. THEORY

their component quarks. Each parton carries some fraction of the momentum of the proton described by parton distribution functions.

Figure 2.3 shows a proton-proton collision schematic. In this example the hard process comes from the up quark in each proton [4]. “Hardness” refers to the fraction of proton momentum involved in the collision. In contrast, “soft” collisions are those from remaining partons in each proton and usually involve low momentum transfer. These soft collisions are considered the underlying event shown in the figure 2.3. Parton scatter is the most common hard process at the LHC by far due to the high density of gluons in the proton and the scale of QCD couplings above electroweak coupling strength.




Figure 2.3: Example proton-proton collision with quark-gluon scattering and final state jet and Z-boson [4]

Quarks and gluons emitted from the high energy hard scatter do not appear in the detector directly. QCD, in one of its key differences to QED, becomes stronger with larger distances. As a parton reaches high enough energy, it will begin to radiate low-energy gluons until resultant partons are able to bind into color-neutral hadrons. These hadrons are seen collimated in groups in the detector as “jets”. The energy and momentum of jets

## CHAPTER 2. THEORY

are considered reflections of the initial scattered partons. Various “jet algorithms” can be used to determine initial parton properties as reproducibly and accurately as possible. The jet algorithm used in this analysis will be described in detail in Chapter 4. However, it is important to note that the algorithm used by all LHC experiments- anti- $k_t$ - is collinear and infrared safe, or unaffected by small angle and soft scatterings that occur in a parton shower. Without these qualities, perturbation theory applied to the parton shower would find infinities at high orders.

Cross-sections (denoted  $\sigma$ ) measure the probability that a certain process will occur in the collision of two particles, in our case protons. In high-energy physics, cross-sections are measured in inverse femtobarns ( $\text{fb}^{-1}$ ). A barn is the cross-sectional area of a typical nucleus,  $10^{-28} \text{ m}^2$ , and was named to describe the large target area needed in order to have direct strikes on a nucleus. The name was inspired by the expression “could not hit the broad side of a barn”. Inverse femtobarns ( $10^{15}$  inverse barns) are used to measure the number of particle collision events per femtobarn area of a target and quantifies time-integrated luminosity.

Hard scattering cross-sections in hadron-hadron collisions can be calculated using the QCD factorization theorem, and to leading-order these calculations are relatively simple. In the factorization theorem, developed by Drell and Yan, deep inelastic scattering parton model processes could apply to hadron-hadron collisions. The Drell-Yan process is the production of a massive lepton pair by quark-antiquark annihilation. According to the factorization theorem, a hadronic cross-section  $\sigma(AB \rightarrow \mu^+ \mu^- + X)$  could be calculated by rescaling the Drell-Yan sub-process cross-section  $\hat{\sigma}$  for  $\bar{q}q \rightarrow \mu^+ \mu^-$  with parton distribution functions  $f_{q/A}(x)$  which come from deep inelastic scattering [5]:

$$\sigma_{AB} = \int dx_a dx_b f_{a/A}(x_a) f_{b/B}(x_b) \hat{\sigma}_{ab \rightarrow X}, \quad (2.42)$$

## CHAPTER 2. THEORY

where  $X$  represents the two resulting leptons and  $ab$  the two annihilated quarks. This parton model provides good agreement with measured cross sections and so allows understanding of particular hard scattering processes. Predictions for some key Standard Model processes are shown in Figure 2.4. Noting the logarithmic scales it is clear that the Higgs boson of mass 125 GeV is orders of magnitude more numerous at the LHC than the Tevatron and that certain high mass particles like the  $b$  quark and  $W/Z$  bosons are abundantly produced at the LHC [5]. In addition, the plot shows cross-sections of particular Higgs decay modes. These will be discussed next.




Figure 2.4: Predicted Standard Model cross-sections for the Tevatron and LHC [6].

Higgs production at the LHC occurs via four main processes: gluon-gluon fusion, vector-boson fusion, associated production with  $W/Z$  bosons, and associated production with top

## CHAPTER 2. THEORY

or bottom quarks. The Feynman diagrams for these processes are shown in Figure 2.5.




Figure 2.5: Feynman diagrams for the leading Higgs boson production modes at the LHC [7].

The LHC Cross-section Working Group produces predictions on cross-sections, branching ratios, and pseudo-observables for the Higgs boson. Four CERN reports bring together Higgs prescriptions for current and planned LHC efforts [8]. Figure 2.6 shows the Higgs cross-section for the main production modes as a function of the Higgs mass and the collision center-of-mass energy. The Standard Model does not predict the Higgs mass, but experimentally we know that  $m_H \approx 125$  GeV. Examining cross-section as a function of center-of-mass demonstrates the increase in statistics for events of interest that running the detector at higher energy levels can accomplish. Figure 2.6 also shows that gluon-gluon fusion Higgs production is the leading production mechanism by far. The Higgs production cross-section is currently known at NNLO in QCD with NLO EW corrections.

The second highest production cross-section is from vector-boson-fusion. As seen in the Feynman diagram, two outgoing quarks are produced in the interaction. These quarks produce two hard jets back to back in the forward region with the Higgs boson appearing between them. To leading order, VBF Higgs production is solely electroweak, and QCD corrections (calculated at NLO) have a smaller impact than in ggF. NLO EW corrections

## CHAPTER 2. THEORY

are also applied. As a result, VBF theoretical uncertainties are smaller than those on ggF. Vector boson associated Higgs production through a  $W/Z$  boson are less common than VBF but also dominated by electroweak processes with a small QCD correction (NNLO). Finally, associated production with top and bottom quarks is shown, though these are quite rare and have high NLO and NNLO QCD corrections.




Figure 2.6: Higgs production cross-sections over Higgs mass at center-of-mass energy 14 TeV (left) and over center-of-mass energy for a Higgs mass of 125 GeV (right)[8]

Theoretical uncertainties shown as colored bars in 2.6 are calculated from choice of PDFs and renormalization and factorization scales. Parton distribution functions (PDFs) are described in more detail by the PDF4LHC working group [9]. This group performs studies of PDFs and their predictions at the LHC and makes recommendations for methods of estimating PDF uncertainties.

## CHAPTER 2. THEORY




Figure 2.7: Predicted branching ratios for the Higgs boson at the LHC as a function of Higgs mass [8].

Since the Higgs boson couplings are directly proportional to the masses of each particle, the Higgs decays most readily into the heaviest particles possible. Figure 2.7 shows key decay mode branching ratios near the experimentally known Higgs mass. While the branching ratios demonstrate the relative abundance of each Higgs decay, these do not translate directly into their ease of discovery or measurement. The current status of Higgs boson coupling and cross-section measurements for each decay mode will be detailed in the next section. The Higgs boson discovery was made through a combination of searches in many channels though dominated by  $H \rightarrow ZZ^*$ ,  $H \rightarrow \gamma\gamma$ , and  $H \rightarrow WW$  [10]. This is because though other decay branching ratios are higher, like  $H \rightarrow b\bar{b}$ , the backgrounds associated with this decay are much higher. As previously mentioned, proton-proton collisions create large amounts of QCD jets that are difficult to discern from target hard QCD processes. Because of this, Higgs decays to quarks and gluons are particularly difficult and those with decays to heavy and light bosons (ZZ decays to 4 leptons, decays directly to photons, and WW decays to two leptons and two neutrinos) are easier to reconstruct. As energy and integrated luminosity

## CHAPTER 2. THEORY

increased during LHC runs 1 and 2, measurements of even rare and background-heavy Higgs decay channels could be made. This thesis focuses on the decay of  $H \rightarrow W^+W^- \rightarrow \ell\nu\ell\nu$  through VBF production. In the last section of this chapter I will motivate the choice of these conditions for probing new physics beyond the SM.

### 2.3 Brief history of SM tests

Here I summarize key moments in the history of the development of the Standard Model and its experimental tests. I use both [11] and [12] as a guide.

The history of the Standard Model could start with any number of physicists well before the formalism of the theory itself. As far back as the fifth century B.C. philosophers posited that matter is composed of discrete “particles” in its most fundamental state. This idea was only tested beginning in the nineteenth century, when scientists were able to detect physical evidence of atoms and their structure. The first of the gauge theories, QED, was invented in the 1930s but only calculated to first order. Renormalization theory was invented simultaneously by Feynman, Schwinger, and Tomonaga in the 1940s. This made calculations of higher order QED results possible. Following this advancement, QED was verified experimentally with high precision.

Physicists next attempted to understand and formalize the other fundamental forces- strong and weak- in the same way. Symmetries for these theories were not as easy to find as that of QED and in 1954 the first new gauge theory for QCD was proposed by Shaw, Yang, and Mills. Though ultimately incorrect, it led to proposals for a gauge theory of weak interactions by Schwinger in 1956 (unifying weak and electromagnetic interactions with photons and massive  $W^\pm$  bosons). Glashow added a fourth boson,  $Z$ , to the theory, but the problem of a broken symmetry necessary to give mass to the  $W/Z$  bosons remained.

## *CHAPTER 2. THEORY*

Spontaneous symmetry breaking was a known and tested concept, but its use in the theory led to unwanted massless “Nambu-Goldstone” bosons. Some thought this problem was inevitable in using SSB in the gauge theory. In 1964 Englert and Brout, then Higgs, and a few months later Guralnik, Hagen, and Tribble all published papers with the same conclusion - the Goldstone theorem can be applied to gauge theories and massive bosons can “eat” the Nambu-Goldstone bosons to gain mass. This then creates a new scalar field whose particles are now termed Higgs bosons. A few years later Weinberg unified these ideas into the electroweak theory we recognize today. The theory was confirmed many times over at experiments over the next decades.

While electroweak symmetry breaking and its implications were being understood in the 1960s, QCD was gradually being assembled. Experimental discoveries of a host of new particles led to Gell-Man and Zweig’s development of a theory that these were all composed of the same three base particles, “quarks”. Han and Nambu understood that an octet of colored gauge bosons, “gluons”, mediated the strong force through color interactions. In 1973 Gross, Wilczek, and Politzer demonstrated the asymptotic freedom of QCD, its weakness at short distances which allows perturbation theory to calculate high-energy interactions. The Standard Model, as it is composed of electroweak dynamics and QCD, has been remarkably predictive but there is much it does not explain, such as the existence of dark matter particles. While theoretical physicists work to expand the Standard Model (or replace it entirely), experimentalists search for deviations from Standard Model predictions, which may be the next hint of entirely new physics.

Large amounts of experimental evidence for the predictive power of QED had amassed over time, and by the 1970s high-energy accelerators at CERN, Fermilab, Brookhaven and SLAC began making first measurements of predicted electroweak and QCD observables. In 1969 physicists at SLAC collided electrons and protons and found that electron scattering

## *CHAPTER 2. THEORY*

behaved as though the proton was made up of point-like particles, quarks. Electroweak theory's predicted neutral weak current interaction between quarks and leptons was discovered at CERN and then Fermilab in 1973, giving evidence to electroweak unification. The  $W$  and  $Z$  bosons predicted by the theory with masses  $\approx 80$  GeV were discovered at the SPS at CERN. This was the first proton-antiproton collider and with a center-of-mass energy of 540 GeV, it was the highest energy collider ever built. The collider was built for the main purpose of finding the predicted  $W$  and  $Z$  bosons and it succeeded in discovering both in 1983. Colliders like LEP at CERN and SLC at SLAC were able to produce millions of  $W$  and  $Z$  bosons and so test electroweak predictions with high precision. The theory continued to prove extremely accurate. QCD remains more difficult to test precisely, as calculating QCD parameters theoretically is a computationally difficult task. However, evidence of QCD's accuracy has accumulated. In 1978, SPEAR at DESY was able to indirectly detect the first gluon. The electron-positron collider produced two quarks that formed two jets of particles with equal energy in opposite directions. QCD predicts that there would also be 3-jet events, where a gluon would be radiated from one of the scattered quarks and form a third quark. The strength of the strong interaction was first measured in 1978 and thereafter with more and more precision and has always followed QCD predictions. Quarks beyond the first generation (up, down) were discovered in order of increasing mass - strange first, followed by charm, then bottom, and finally in 1995 the top. Similarly, the muon was discovered well before the heavier tau-lepton. After the discovery of the top quark, the last Standard Model particle left undiscovered was the Higgs boson. Its mass is an input rather than a prediction to the Standard Model, but with the large amount of data taken at increasingly high energy colliders, experimentalists and theorists were confident that if it existed, its mass would be between 100 GeV and 1 TeV. In order to search for the Higgs boson, the Large Hadron Collider was planned and built at CERN. The collider and its largest all-purpose detector,

## CHAPTER 2. THEORY

ATLAS, are discussed in the next chapter.

In 2012, the Higgs boson was discovered at the LHC in both the ATLAS and CMS detectors. After just one year of data-taking at a center-of-mass energy of 7-8 TeV, combined searches in the  $H \rightarrow ZZ^* \rightarrow \ell\ell\ell\ell$ ,  $H \rightarrow \gamma\gamma$  and  $H \rightarrow WW \rightarrow \ell\nu\ell\nu$  channels found a particle compatible with the Standard Model Higgs with a significance of 5.9 standard deviations at a mass close to 125 GeV. The LHC continued data-taking after the discovery with a new goal: measuring this new particle's properties with accuracy and precision. In 2015, the center-of-mass energy increased to 13 TeV and more than  $20\times$  the data used in the first discovery has now been recorded.

Measurements of the Higgs boson are now numerous and quite precise, but no deviations were observed from the theoretical expectation. While the Standard Model has proven a successful model of known interactions, there are many phenomena that it does not predict - from dark matter to neutrino masses. There must be physics beyond this theory. In continuing to probe new aspects of the model, the LHC may find deviations from the known forces or physics beyond the Standard Model.

The mass of the Higgs boson is measured through a combination of decay modes and in combination with CMS to be  $125.10 \pm 0.14$  GeV [1]. The latest combined Higgs cross-section measurement from ATLAS uses data from 2015-2017 and finds the production cross-sections (normalized to their Standard Model predictions) as shown in Figure 2.8. Branching ratios of relevant decay modes are also measured and are shown in Figure 2.9 multiplied by their cross-sections in each relevant production mode. The differences in theoretical and systematic uncertainty for certain decays (QCD heavy VBF  $H \rightarrow b\bar{b}$  in comparison to the leptonically decaying VBF  $H \rightarrow WW^*$ ) show how discernable backgrounds play a major role in the viability of a measurement, even when the branching ratio is high [13].

## CHAPTER 2. THEORY




Figure 2.8: Production cross-sectios for ggF, VBF, VH, and  $t\bar{t}H + tH$  normalized to their SM predictions. Total, systematic, and statistical uncertainties are shown [13]




Figure 2.9: Branching ratios for measured Higgs decays normalized to their SM predictions. Total, systematic, and statistical uncertainties are shown [13]

## CHAPTER 2. THEORY

This thesis details my work on the fiducial cross-section measurement of VBF  $H \rightarrow WW^* \rightarrow \ell\nu\ell\nu$  decays. The clear sign of VBF Higgs production (2 jets recoiling in opposite high  $\eta$  directions, combined with the leptonically decaying  $W$  bosons) gives the decay a distinctive signature. This channel can be measured with high accuracy which may allow small deviations from theory to be visible. In addition, the measurement is made with the goal of eventual differential cross-section measurements, which can probe higher order perturbative contributions to QCD or EW. Details on the particular motivations for this thesis' analysis will be outlined in the next section. The last ATLAS  $H \rightarrow WW$  differential measurement was published in 2016 with  $20.3\text{fb}^{-1}$  at center-of-mass energy 8 TeV and measured only ggF Higgs boson production differential cross-sections [14].

## 2.4 Measurement motivation

There are many reasons why the VBF  $H \rightarrow \ell\nu\ell\nu$  channel is particularly interesting to investigate. Since the  $W$  bosons decay leptonically, they create a clear signal in the detector which allows for a precise reconstruction of the hard processes. Further, the VBF production mode has a similarly clear signal with the two jets accompanying the decaying  $W$  bosons with a high  $\eta$  separation. The pseudorapidity of these jets and the rapidity between them are particularly sensitive to the electroweak symmetry breaking mechanism, so a differential cross-section measurement over these variables (along with other detailed and motivated in further sections) tests the Higgs mechanism. In Run-1 VBF  $H \rightarrow WW^* \rightarrow \ell\nu\ell\nu$  cross-sections were measured with an experimental uncertainty of  $\approx 50\%$  [15]. This new measurement uses the entirety of Run-2 data, with a factor of  $3.9\times$  higher statistics. This VBF  $H \rightarrow WW$  fiducial cross-section measurement uses new analysis techniques and improvements in theoretical calculations as well as higher statistics to make an accurate

## CHAPTER 2. THEORY

probe of new physics within the electroweak symmetry breaking mechanism.

This thesis focuses on the VBF  $H \rightarrow WW$  fiducial cross-section measurement, but this measurement is the first piece to a larger goal beyond the scope of my Ph.D. Our group plans to make a simultaneous measurement of VBF  $H \rightarrow WW$  and VBF  $W + \text{jets}$  fiducial and differential cross-sections. The VBF  $W + jj$  production cross section is sensitive to the  $WWV$  vertex. In vector boson scattering, we expect (with minimal assumptions) that new physics ought to be at a scale above the current experiment. Effects from this type of new physics would simultaneously affect  $HWW$  and  $WWV$  vertices. In particular, dimension-6 operators affect VBF Higgs mechanisms and VBF  $W$  production, while the interplay of VBF and VBS mechanisms allows for the probe of dimension-6 and dimension-8 operators. Measuring fiducial and differential cross-section ratios (of VBF Higgs and  $Wjj$ ) enhances sensitivity to new physics because any new phenomena that simultaneously affect both vertices are suppressed and so sensitivity to the phenomena affecting just the  $WWV$  vertex (for example, anomalous quadruple-gauge couplings) is enhanced. VBF Higgs and  $Wjj$  productions are also characterized by final-states with two jets with similar kinematics. Because of this, the ratio of cross sections measurement allows for the cancellations of correlated systematic uncertainties. As both process are characterized by large (10% – 20%) jet-related uncertainties, a reduction via the ratio cancellation would greatly improve the sensitivity to the VBF Higgs process and to the search for new phenomena. This VBF  $HWW$  measurement is the first part of the ratio of fiducial differential cross sections for VBF  $HWW$  and  $Wjj$ , which will set limits on EFT parameters, benefiting from enhanced sensitivity through the cancellation of correlated uncertainties.

# Chapter 3

## The LHC and the ATLAS detector

The Large Hadron Collider (LHC) is a proton-proton storage ring operating at CERN and for its 9 years of operation, it has been the world's highest energy particle collider. During LHC operation thus far, protons have collided with increased center-of-mass approaching the design energy of 14 TeV. Instantaneous luminosity has also successively increased, surpassing design instantaneous luminosity of  $1 \times 10^{34} \text{cm}^{-2}\text{s}^{-1}$  in 2018 to reach  $2 \times 10^{34} \text{cm}^{-2}\text{s}^{-1}$ [16]. The overall data recorded in the ATLAS detector totals more than  $10^{16}$  collisions. Operation of the LHC has led to the discovery of the Higgs boson and some of the most precise measurements of its properties including the coupling of the Higgs boson to bottom quarks [17],  $W$  and  $Z$  bosons [15], [18], photons[19] and tau leptons [20]. The LHC has also facilitated searches for new physics over a wide parameter space, setting confidence level exclusion limits on masses of supersymmetric particles like squarks, gluinos and neutralinos [21].

The LHC can run continuously for a few years before detector components need to be repaired and replaced. The schedule of data-taking consists of long periods of data accumulation (Run 1 from 2015-2016, and Run 2 from 2017-2018) paired with long shutdown periods. The LHC is set to begin Run 3, in which design center-of-mass energy should

### CHAPTER 3. THE LHC AND THE ATLAS DETECTOR

be reached, in 2021. Following Run 3, detector upgrades will be installed during a long shutdown. Then the High-Luminosity LHC (HL-LHC) will begin colliding protons with unprecedented ( $10\times$ ) luminosity in 2027 [22]. The HL-LHC and its goals will be explained further later in this chapter. Suffice to say LHC-physics is progressing quickly and promises exciting developments in the near future.

In a brief explanation of the LHC operation, one could begin with the small volume of  $\approx 10^{11}$  protons that are accelerated per bunch. Linac-2 is the primary accelerator for CERN colliders and has been since the early 1990s [23]. This injects protons at 50 MeV into the Proton Synchrotron Booster (PSB) where they are further accelerated to 1.4 GeV. In the the Proton Synchrotron (PS), the protons are separated into bunches with a spacing of 25 ns and are futher accelerated to 25 GeV before being extracted to the Super Proton Synchrotron (SPS), where they reach 450 GeV. Finally the bunches of protons enters the LHC, where they are accelerated to their final energy of 6.5 TeV. Linac 2, PSB, PS, and SPS were all operational accelerators before the LHC era though each had to be majorly upgraded to handle the energy and beam intensity required for LHC collisions [23].

The LHC layout mimics that of the Large Electron Positron collider (LEP) that was housed in the same tunnels. Figure 3.1 shows the positioning of each experiment at the LHC as well as injection systems and other features. Once proton bunches enter the LHC in two opposing beams, they are accelerated with radio frequency (RF) systems. Located at Point 4 in the LHC schematic, the system consists of 16 RF cavities operating at twice the frequency of the SPS injector. RF cavities are metallic chambers containing oscillating electromagnetic fields; in the LHC this oscillation frequency is 400 Mz. The tuning of this frequency ensures that protons of the ideal energy are not accelerated further and simply maintain their momentum while particles arriving in an RF cavity slightly before or after will be decelerated or accelerated toward the ideal proton energy. This acceleration process

### CHAPTER 3. THE LHC AND THE ATLAS DETECTOR

can also be used to split beams of protons into discrete bunches, and this is first done with RF cavities in the PS. After proton bunches have circled the LHC approximately 1 million times (15 minutes), peak energy is reached and collisions can commence [24].




Figure 3.1: LHC layout [25]

Superconducting magnets in the LHC main dipoles create a magnetic field of  $\approx 8$  T to bend the proton beams into the circular path of the collider. Figure 3.2 shows the flux in a dipole cross-section. The opposing direction beamlines are shown centered and the flux is shown to be high (and directionally opposed) in the center of each beam. To maintain these fields, the magnets operate at below 1.9K. Pressurized superfluid helium chosen for its low viscosity and high specific heat cools the dipole magnets. Once the two LHC rings are filled from the SPS the center-of-mass energy of the beams increases until it reaches peak energy after about 28 minutes. Finally, proton bunches separated by 25ns collide simultaneously in each detector.




Figure 3.2: Flux within an LHC dipole cross-section [25]

### 3.1 ATLAS, A Toroidal LHC ApparatuS

The LHC creates proton-proton collisions at the rate and energy necessary for pushing the boundaries of particle physics, but identifying and reconstructing the tracks of such energetic particles is no mean feat. A Toroidal LHC ApparatuS (ATLAS) and the Compact Muon Solenoid (CMS) are multi-purpose detectors built to search for a wide range of particle interactions and the measurement of their properties. Both experiments measured a particle consistent with the Higgs boson in 2012 and their agreement was a key verification of the discovery. The following sections describe each major component of the ATLAS detector so to highlight their role in the measurement of  $H \rightarrow WW \rightarrow \ell\nu\ell\nu$ .

ATLAS utilizes a coordinate system with its origin at the center of the detector (the “interaction point”) and has a z-axis along the beam pipe. The x-axis points from the interaction point to the center of the LHC ring, and the y-axis points upward. The experiment uses cylindrical coordinates  $(r, \phi)$  where  $\phi$  is the azimuthal angle around the beam pipe. The pseudorapidity and the transverse momentum are defined in terms of the polar angle  $\theta$  as

### CHAPTER 3. THE LHC AND THE ATLAS DETECTOR

$\eta = -\ln(\tan(\theta/2))$  and  $p_T = p \sin \theta$ .




Figure 3.3: Computer-simulated ATLAS detector schematic [26]

The Inner Detector (ID) detects charged particles with  $|\eta| < 2.5$  operating in a 2 T solenoidal field. It consists of 3 layers of pixel sensors, 4 layers of silicon strips, and 72 straw layers of transition radiation tracker modules. The ID describes particles closest to the interaction point and locates track parameters with great resolution due to its high granularity [26].

The ATLAS detector contains 3 superconducting magnet systems- the central solenoid, barrel toroid, and 2 end-cap toroids. The central solenoid provides a magnetic field for the inner detector while the toroids create a strong magnetic field for the muon detector. These magnets were built to create the largest possible uniform magnetic field to maximize the momentum resolution on particle tracks. They also need to use as little material as possible so as to not unduly influence particles in the detector. The toroids in the barrel and endcap each have 8 coils and create a 4 T magnetic field while the central solenoid creates

### CHAPTER 3. THE LHC AND THE ATLAS DETECTOR

a 2 T magnetic field in the inner detector. Combined the magnet systems contain  $>100$  km of superconducting wire which are cooled to working temperatures below 5K [26].

The Muon Spectrometer precision chambers provide muon momentum measurements at a high resolution over a wide range of  $p_T$ . The MS consists of 3 layers of Monitored Drift Tube chambers covering  $|\eta| < 2.7$  and an inner layer of Cathode Strip Chambers with  $|\eta| > 2.0$ . In addition, it includes trigger chambers that contain 3 layers of Resistive Plate Chambers ( $|\eta| < 1.05$ ) and 3 layers of Thin Gap Chambers ( $1.05 < |\eta| < 2.4$ ). As the outermost subdetector, the MS provides precise muon momentum measurements along the muon trajectory and the muon chambers are located with a precision of under  $60\ \mu m$ . The MS also contains a system of three superconducting toroidal magnets each with eight coils providing a magnetic field with a bending integral of up to 6 Tm [26].

Calorimeters provide detailed information about the energy deposited as particles pass through. Electromagnetic calorimeters detect and halt the motion of electrons and photons while the hadronic calorimeter does the same for hadrons. Muons and neutrinos are able to pass through the calorimeters to the MS. The electromagnetic and hadronic calorimeters, made of liquid Argon and scintillating tiles respectively, are able to pass information from the location of energy deposits to the various idenfication and reconstruction algorithms [26].

## 3.2 The High-Luminosity LHC and Inner Tracker (ITk)

The LHC succeeded in its paramount goal of discovering the Higgs boson in 2012. Its continuous operation at higher energy and luminosity has led to more rigorous measurements of Higgs boson properties as well as searches for new physics beyond the Standard Model. While more data collection is planned in Run-3 starting in 2021, new colliders and detectors take decades to design, develop and build, so the plans for upgrading the current detectors

### *CHAPTER 3. THE LHC AND THE ATLAS DETECTOR*

are well underway. The High Luminosity LHC will operate at 14 TeV starting in 2027. The HL-LHC will begin with  $5 - 7 \times$  the luminosity of the LHC and peak at the design instantaneous luminosity of  $10 \times$  the LHC, or  $12.6 \times 10^{-34} \text{cm}^{-2}\text{s}^{-1}$ . This huge increase in number of collisions requires substantial upgrades to the LHC including new 11-12 T superconducting magnet systems, compact superconducting cavities for beam rotation and phase control, and new technology beam collimation [27].

Just as the LHC had to be re-designed, so too do all the experiments to be able to handle the much higher luminosity. The detectors must be built to withstand more radiation, as the increased collision rate also means a high radiation rate, especially closest to the beamline. They also have to provide greater granularity to be able to reconstruct tracks with good enough resolution that individual tracks can be discriminated. Finally, they have to faster to be able to deal with increased pile-up. Pile-up is caused by high numbers of collisions occurring at each bunch crossing. When there is a large amount of pile-up it becomes difficult to trace which particle tracks come from the same interaction point. Finally, the increased data in and of itself creates a complex problem for the detectors to solve. The trigger system must quickly select and store the events that may hold interesting information.

Detectors for high energy colliders are not built often - expensive and time-consuming to design and test, they are made to last at least a decade. I was lucky to have the opportunity to work on ATLAS detector research and development during the 1.5 years I worked at Brookhaven National Laboratory during my Ph.D. Though my thesis is not directly related to this work, it was formative and extremely interesting, so I touch on this in the next section. Because I worked on the new ATLAS Inner Detector for the HL-LHC (termed Inner Tracker or ITk) I will discuss solely this sub-detector and the particular role I played in its assembly.

### 3.2.1 Inner Tracker (ITk)

The Inner Tracker is planned to be an all-silicon detector that will completely replace the current Inner Detector. While the current ID has been extremely successful during Runs 1 and 2 (and will certainly continue to be in Run 3), it does not have the capacity to withstand the radiation and pile-up conditions of the HL-LHC. The ITk is designed to operate for 10 years under an instantaneous luminosity of  $7.5 \times 10^{-34} \text{ cm}^{-2}\text{s}^{-1}$  with 25 ns between bunch crossing. This will result in  $1,000 \text{ fb}^{-1}$  and average pile-up up to  $\langle \mu \rangle = 200$  [28]. The current solenoid magnet will remain in the detector with a 2 T magnetic field. The ITk will consist of an innermost section with silicon pixels and an outermost section of silicon strips. The pixel detector will contain four barrel layers and six forward region disks, while the strip detector will contain five barrel layers and seven disks. The rapidity range matches the coverage of the Muon Spectrometer with  $|\eta| < 2.7$ . This layout is shown in 3.4.




Figure 3.4: ITk layout as defined in ITk Technical Design Report [28]

## *CHAPTER 3. THE LHC AND THE ATLAS DETECTOR*

### **Building ITk Strip barrel staves**

At Brookhaven National Laboratory, I made key contributions to the ITk Strip barrel stave assembly effort. The goal of stave assembly is to glue silicon modules to carbon fiber stave cores within a  $25\ \mu\text{m}$  tolerance. Brookhaven is responsible for assembly of 200 ITk staves and their accurate assembly is necessary for the ITk to reduce uncertainty on track positions as well as to ensure a symmetric detector. I was tasked with co-creating a stave assembly software system through LABView to automatically calibrate required module positions, apply a layer of adhesive gel, and guide a user in accurately placing a module into its specified location. This project was highly collaborative and evolved further after I left the laboratory, but the overall process remains unchanged.

The basic design of the Inner Tracker for both barrel and endcap components is the same - a carbon fiber core (containing titanium cooling pipes) is covered on each side with co-cured kapton service tapes. The carbon fiber core is designed to reduce material inside the detector and the similar design in the barrel and endcap adds to simplicity. Silicon modules are glued to stave cores. Similar silicon strip detectors have been used previously in both ATLAS and CMS, but have never covered so much fiducial area. The modules consist of one silicon sensor and one or two low-mass PCB's (hybrid) which host ASICs. Module design has optimized producibility and low cost while maintaining readout goals. Overall module design is the same in barrel and endcap regions, while strip lengths and geometries vary. Components of a short-strip barrel module are shown in [3.5](#).

Each barrel stave core needs to be “loaded” with 14 modules, as shown in the assembled electrical prototype in [3.6](#).

Brookhaven National Laboratory is one of two sites responsible for assembling barrel staves. Assembly procedures have been tested with the production of numerous prototypes

### CHAPTER 3. THE LHC AND THE ATLAS DETECTOR




Figure 3.5: Short-strip barrel module components [28]




Figure 3.6: Electrical stave prototype at Brookhaven National Laboratory (G. van Nieuwenhuizen)

including a thermomechanical double-sided stave and a fully operational electrical stave. The thermomechanical prototype was later used for various thermal tests, including IR imaging. The electrical stave was used for testing the full electrical read-out. Stave assembly is composed of three main parts: system calibration, module placement, and survey of results.

Staves at BNL are assembled on a granite table housing an Aerotech XYZ Stage accurate to the micron level. The stage is equipped with a 10-megapixel camera that gives real-time feedback to a nearby computer and a glue dispenser. The stave assembly software system is implemented by a user who interacts with a LabView GUI and monitors progress. The stave is fixed to optical rails drilled into the granite table. In order to accurately place modules in their correct positions a series of calibrations need to be completed including camera calibrations to test the optimal working point, focus, and pixel-to-micron conversion. Next, the position of the stave with relation to the XYZ stage needs to be calibrated. Transforming

### *CHAPTER 3. THE LHC AND THE ATLAS DETECTOR*

coordinates of the XYZ stage to that of the stave requires locating a fixed point on the stave core as well as the angle of the stave relative to the XYZ stage. Pattern matching algorithms find the exact locations of particular features on the stave core and allow calculation of required positions for all modules based on specifications. Once specified, module positions are calculated, calibrations are completed and it is time to apply glue and afix modules.

Next, an epoxy (SE4445) is loaded into the glue dispenser on the XYZ-stage which is connected to a vaccuum controlled by the LABView software system. The epoxy is automatically dispensed in lines to cover  $\approx 60\%$  of area under the module. Then the module is lifted with a custom-made “pick-up” tool which uses vacuum applied to module corners to hold the module in place and move it to the needed position along the optical rails. Using real-time feedback from the software system and its pattern matching algorithm, the user is directed on how to finetune module position using knobs on the “pick-up” tool. Markings etched in the silicon sensor at each corner are used to position the module accurately. The output of the module alignment GUI is shown in [3.7](#). When the module is within specifications it is lowered into position above the epoxy and held in place for 24 hours until the glue has completely dried.

After the glue has set, a final survey of module positions is recorded using pattern matching to find positions of etched markings on each module corner. These results are saved into an ITk database and checked for any biases. After module placement on the stave is complete, the loaded stave is moved to another station in the lab for wirebonding so that all data from the modules can be read-out to stave-wide electronics. The module assembly system has been successful at placing modules accurately for all prototypes, achieving specification requirements for almost all modules. Results of the first prototype stave’s module placement are shown in [3.8](#). While a few module corners are slightly out of the ideal range, the majority are well within specifications. Throughout prototype assembly issues and inefficiencies

### CHAPTER 3. THE LHC AND THE ATLAS DETECTOR



Figure 3.7: GUI interface showing etched marking on module corner located in real-time to guide user on how to adjust module position (H. Herde)

were found and corrected. New hardware, like an improved glue system and temperature monitoring, were also added. The methods described continue to be in use now and will be utilized for the production of 200 ITk staves at BNL starting in 2021.




Figure 3.8: Histograms show difference between ideal and final position of each module corner. Left shows difference from specification in X and right in Y (P. Bhattacharai)

### IR Testing of ITk Strip barrel staves

The first full US stave prototype was the thermo-mechanical stave built in the summer of 2017. Building this stave was the first test of stave assembly procedures and the results

### *CHAPTER 3. THE LHC AND THE ATLAS DETECTOR*

proved that the module placement algorithm we developed worked. This stave was also used to test the thermal and mechanical properties of a fully loaded barrel stave. Multiple studies were conducted, including thermal measurements using thermistors and IR imaging, thermal cycling and thermal shock tests, mechanical studies, and bending tests. I will give a short summary of IR imaging tests, as these were another focus of my time at BNL.

The thermo-mechanical stave consists of 13 modules mounted on each side. The modules used are thermo-mechanical, which means that instead of the usual readout chips their hybrids employed copper resistors to mimic the power dissipation and location of the chips. The powerboard can vary the TM hybrid power dissipation of each module individually. Three thermistors were mounted on each TM module- one on the DC-DC converter and one on each of the two hybrids. A custom End-of-Substructure (EoS) card is attached to a RaspberryPi and Arduino to power on or off each module.

Thermal testing of barrel staves had a few main goals: to validate Finite Element Analysis (FEA) simulations by testing that all temperature trends are as we expect, to make sure that individual modules do not exhibit abnormal thermal behavior, and finally to check that loaded staves can cope with large changes in temperature they might face during operation. I will highlight a few key results which demonstrate that each of these goals have been accomplished.

Thermal measurements were taken both through the mounted thermistors on each module and through IR imaging. IR imaging provides information about the entirety of the loaded stave, rather than at just a few module positions so provides a more complete picture. The loaded stave was spray-painted black with a high emissivity, low conductivity black paint since silicon is transparent to the IR camera spectrum ( $8\text{-}14\mu\text{m}$ ). In order to image the entire stave core, the IR camera was attached to rails above and pulled at a constant speed with an external motor as it recorded a video. The frames were then stitched together into one

### CHAPTER 3. THE LHC AND THE ATLAS DETECTOR

image. A section of the painted stave is shown in 3.9.




Figure 3.9: Portion of the thermo-mechanical stave after spray-painted for increased emissivity

FEA simulations for the thermal performance of a stave were completed by Prof. Graham Beck at Queen Mary University of London. These calculations quickly became intractable if convection was included so conditions of the stave and coolant were adjusted to minimize convective contributions or make sure that the total electrical power and power absorbed into the coolant were identical. At BNL we adjusted the coolant temperature until we obtained the convective power minimization and then recorded module temperatures under these conditions. These results were compared to the FEA simulations by averaging hybrid temperatures recorded through IR imaging and recording NTC thermistor readings. These comparisons are shown in 3.10. The measurements show very good agreement with FEA calculations, within 5% of the expected values.

During module assembly some slight variations were tested, including varying glue thickness below modules, glue curing time, and FEAST versions. Modules with and without these variations were compared at varying coolant temperatures and output power settings. Overall, no significant difference in module temperature change was observed for any of these assembly modifications. Stave thermal properties are thus robust to such assembly modifi-

CHAPTER 3. THE LHC AND THE ATLAS DETECTOR




Figure 3.10: IR measurements, NTC thermistor measurements, and FEA simulations for the TM stave are compared. Agreement with FEA simulation within 5%.

cations. Figure 3.11 shows a full IR image of the fully loaded stave. It is clear that there are no obvious module-to-module variations in silicon, hybrid, or FEAST temperature. The module sensors increase in temperature as they get closer to the EoS, which is expected since it dissipates power to the stave.




Figure 3.11: IR image of fully loaded thermo-mechanical stave

The thermo-mechanical stave was pushed to limits beyond what we would expect loaded staves to encounter during operation and never exhibited unexpected behavior. Thermal cycles, thermal shocks and bend tests showed the loaded stave to be robust against temperature variation and that the carbon fiber core is as stiff as it was prior to loading. Another test was how neighboring modules would perform if one module malfunctioned and was powered

### CHAPTER 3. THE LHC AND THE ATLAS DETECTOR

off. Figure 3.12(top) shows the temperature of each module when one of them (fourth to the left) is turned off. The rest of the modules continue to operate normally and temperature changes from the unpowered module do not propagate very far. The bottom image in the figure shows the reverse of the stave when a module is powered off (fourth from the right). The temperature effects are greater on the module directly below than those adjacent to, which is expected due to material differences.




Figure 3.12: IR image of fully loaded thermo-mechanical stave

My experiences on ATLAS detector upgrades for the HL-LHC provided context for the bulk of my thesis research. This project provided me with in depth and hands-on knowledge of the ATLAS detector and its component parts, as well as the scale of effort required to build new detector components. I have an abiding appreciation for the people and technology necessary for data-taking at the LHC, both of which make measurements like the Higgs cross-section detailed in this thesis possible.

# Chapter 4

## Tracking and Isolation in ATLAS

The measurement of  $H \rightarrow WW^* \rightarrow \ell\nu\ell\nu$  cross-sections relies on the accurate determination of a number of “physics objects” within the detector, primarily jets, missing transverse energy ( $E_T^{miss}$ ), electrons, and muons. The final state particles of this interaction consist of one electron, one muon, and two neutrinos, which appear as missing transverse energy in the detector since they cannot be directly detected. The different flavored leptons are required in our final state to eliminate large contamination from Drell-Yan ( $Z/\gamma^* \rightarrow ee/\mu\mu$ ) backgrounds. Drell-Yan backgrounds still play a role in our different flavor selection analysis but primarily come from far less numerous  $Z/\gamma \rightarrow \tau\tau$  events which then decay to an electron and muon. Because our search focuses on VBF production Higgs, we also require two jets in the final state. Each physics object has a dedicated performance group which is tasked with providing recommendations for reconstruction, identification, isolation and measurements of efficiency, scale, and resolution. Biases or omissions in any of these would severely impact the precision of our analysis. Understanding the uncertainties associated with each reconstructed object is critical. In this chapter I briefly outline the algorithms used for tracking and isolation to accurately determine the kinematics of electrons, muons,

## CHAPTER 4. TRACKING AND ISOLATION IN ATLAS

jets and missing transverse energy. Particular attention will be paid to muons, as I spent a significant amount of time working in the Muon Performance Group applying and validating corrections to muon momentum scale and resolution.

### 4.1 Tracking

Track reconstruction is fundamental to accurately identifying and describing electrons, muons and jets and primarily utilizes information from the ID. Charged particles entering the ID and solenoid field follow a circular trajectory in the transverse plane. They can be fully explained with a set of five parameters:

- $q/p_T$ , the charged curvature where  $q$  is electric charge and  $p_T$  transverse momentum
- $\phi$ , the azimuthal angle
- $\theta$ , the polar angle
- $d_0$ , the transverse impact parameter or distance of closest approach of the beamspot position in the  $x - y$  plane
- $z_0$ , the longitudinal impact parameter, or  $z$  coordinate of the track at the point of closest approach.

Track reconstruction calculates these descriptive parameters for each located track through an initial *inside-out* procedure and second *outside-in* method [29]. First, track seeds are built from three hits in the silicon detectors, required to be in different silicon detector layers. Hits are then added to seeds moving away from the interaction point. If the final number of hits exceeds a determined threshold a track candidate is created and if not, the hits are discarded. Next for the *outside-in* approach, segments are reconstructed in the TRT

## CHAPTER 4. TRACKING AND ISOLATION IN ATLAS

and silicon hits are added through back-tracking. This approach finds tracks from secondary interactions which do not begin at the primary interaction point. Requirements on hits, holes (points where measurement is expected but not found), and outliers (hits that reduce track fit quality) reduce probability of tracks which can not be matched to the primary vertex or secondary particles [30]. The ATLAS Tracking Performance group determines reconstruction performance for tracks and demonstrate their robustness to potential fake track candidates and pile-up. Reconstructed tracks are used as input for the identification and measurement of kinematic variables for each of the physics objects that are described in this chapter.

## 4.2 Electrons

Accurate reconstruction and calibration of electrons within the ATLAS detector is integral to precision measurements, including the  $H \rightarrow WW$  measurements in which an electron is required in the final state. Reconstruction, identification, and energy measurements of electrons and photons are the goals of the Electron and Photon Performance group. This section will summarize each of these calibration processes and their performances with a focus on electrons.

Electrons are defined through energy deposits in the calorimeter, the superclusters, each with a matching track from the ID. Photons are defined strictly through a calorimeter cluster. Figure 4.1 shows the procedure for electron and photon reconstruction. First, topo-clusters in the EM calorimeter and tracks in the ID are selected and matched together. Topo-clusters are defined based on signal to noise significance in calorimeter cells and calorimeter cell proximity. Standard reconstruction takes place in the ID and potential tracks are assigned to topo-clusters if their positions are within a region-of-interest compatible with that topo-clusters EM shower. Next, super-clusters are built from track-matched topo-clusters. Topo-

## CHAPTER 4. TRACKING AND ISOLATION IN ATLAS

clusters are tested for use as seed cluster candidates which begin super-clusters. Remaining topo-clusters are tested for compatibility as satellite clusters to each seed candidate. The resultant combination of seed and satellite clusters form super-clusters which are defined independently for photons and electrons. Finally, tracks are added to super-clusters, energy calibration and position corrections are applied, and analysis-level electrons and photons are created. Reconstruction efficiency for electrons is quite high, approaching the tracking efficiency at high  $p_T$ . Photon reconstruction efficiency is significantly lower due to their dependence solely on calorimeter clusters [31].




Figure 4.1: Electron and photon reconstruction algorithm [31]

Electron and photon energy resolution is determined from EM calorimeter showers and is optimized with multivariate regression algorithms. Energy scale is also corrected using calibration from  $Z \rightarrow ee$  decays and verified with other  $Z$ -boson decays. Similarly, these calibrations are calculated for photons using  $Z \rightarrow \ell\ell\gamma$ . Systematic uncertainties that effect these calibrations include passive material between the interaction point and the EM calorimeter and pile-up [31].

## CHAPTER 4. TRACKING AND ISOLATION IN ATLAS

While electron and photon objects are already identified and reconstructed, further ‘identification’ selections are used to improve electron and photon purity. Prompt electrons are identified with a likelihood discriminant which takes into account track parameters and electromagnetic shower properties. The primary electron track must stretch into the two inner pixel layers and hit multiple points in the silicon-strip detector. The likelihood discriminant is calculated from probability density functions formed by smoothing histograms of 13 discriminating variables with an adaptive kernel density estimator. These are calculated separately for both the likelihood that an event reconstructs a prompt electron (signal) or that it does not (background). These likelihoods are derived from  $Z \rightarrow ee$  and  $J/\Psi \rightarrow ee$  events recorded in Run-2. A tag-and-probe method is used to evaluate this likelihood - one electron in each decay must satisfy strict Run-1 likelihood discriminant requirements while the other serves as a probe of the new likelihood discriminant. Three electron working points are derived- Loose, Medium, and Tight, each with lower efficiencies and higher purity than the previous. Physics analyses use the working points optimized for their analysis. For the  $H \rightarrow WW$  measurement we use ‘Medium’ reconstructed electrons. Figure 4.2 shows electron identification efficiency for each of the working points in a sample of  $Z \rightarrow ee$  events where efficiency is calculated through comparisons to MC simulated  $Z \rightarrow ee$  and  $J/\Psi \rightarrow ee$  events. The average efficiencies for electroweak processes are 93%, 88% and 80%, respectively, for the Loose, Medium, and Tight operating points and gradually increase at high  $E_T$  [31].

Track hits and calorimeter deposits near reconstructed electrons and muons can bias energy, momentum, and position measurements. Isolation performance is defined for calorimeter clusters by  $E_T^{cone}$ , the sum of transverse energy within a cone  $\Delta R$  near a photon or electron cluster after correcting for leakage and pile-up effects. Track isolation  $p_T^{cone}$  is defined as a sum of the transverse momentum of tracks within a cone about the electron track or interpolated photon track. For electrons the distance between nearby decay products is

## CHAPTER 4. TRACKING AND ISOLATION IN ATLAS




Figure 4.2: Electron reconstruction efficiency as a function of  $E_T$  (left) and  $\eta$  (right) in  $Z \rightarrow ee$  events for Losse, Medium, and Tight electrons [31]

directly related to electron  $p_T$  so a variable cone size can be used such that

$$\Delta R = \min\left(\frac{10}{p_T}, \Delta R_{max}\right), \quad (4.1)$$

where  $\Delta R_{max}$  is typically 0.2. Isolation working points strike a balance between efficiency and rejection of misidentified prompt electrons. Electron isolation for the  $H \rightarrow WW^*$  analysis uses two different selections based on electron track  $p_T$ . The *IsoGradient* working point is used for  $p_T > 25 \text{ GeV}$ . This requires calorimeter and track isolation about cones of  $\Delta R = 0.2$  (*topoetcone20*, *ptvarcone20*). Electron isolation is changed to the fixed cut track cone isolation for  $p_T < 25 \text{ GeV}$  to further eliminate fake background contributions. High efficiency at each working point requires cuts on  $E_T^{cone20}$  and  $p_T^{varcone20}$  derived from  $J\Psi \rightarrow ee$  and  $Z \rightarrow ee$  MC simulations. Isolation efficiency for electrons are shown in Figure 4.3 for Medium identified electrons in  $Z \rightarrow ee$  events. The Gradient working point delivers efficiency that is stable across  $\eta$  and is coupled with high background rejection of misidentified electrons [31].

Photon isolation is solely calorimeter-based.  $Z \rightarrow \ell\ell\gamma$  events are used for photon isolation

## CHAPTER 4. TRACKING AND ISOLATION IN ATLAS




Figure 4.3: Electron isolation efficiency shown for four working points in  $Z \rightarrow ee$  events as a function of  $E_T$  and  $\eta$  [31]

efficiency measurements and three working points are available which balance efficiency and background rejection just as in the electron case.

Electron reconstruction, calibration, identification, and isolation are all integral for the  $H \rightarrow WW$  analysis in order to accurately identify electrons in the final state of candidate Higgs events. Systematic uncertainties from electron identification and isolation efficiency will be discussed further in later chapters and play a significant role in  $H \rightarrow WW$  cross-section measurements.

## 4.3 Muons

Muons help lead to some of the most interesting physics analyses produced by the ATLAS experiment including  $H \rightarrow WW \rightarrow \ell\nu\ell\nu$  measurements, which require a muon in the final state. The Muon Combined Performance group (MCP) is tasked with producing the most accurate muon calibrations for physics analyses. This includes defining muon identification and isolation criteria and measuring efficiency as well as muon momentum scales and resolutions. The group’s goal is to create a number of “working points” tailored to different

## CHAPTER 4. TRACKING AND ISOLATION IN ATLAS

types of physics analyses. The working points are continuously updated and improved before being tested and implemented on different analyses. My work with the MCP group has focused on applying corrections necessary for muon momentum scale at the per mille level and resolution at the percent level in simulation to match the data.

Muon reconstruction is performed independently using tracks reconstructed in the ID and MS and for combined tracks. This section will focus on reconstruction in the MS, which is particular to muons and the combined reconstruction, which uses information from both the ID and MS.

Muon reconstruction begins with a search for hit patterns in each MS subdetector, which are called segments. The middle of the MS typically exhibits the largest number of trigger hits, therefore tracks are built by working out from the center of the MS and connecting segments layer-by-layer. Criteria such as hit multiplicity and fit quality determine track acceptance. At least two segments are needed to build a track. Hits associated with each track candidate are fitted using a global  $\chi^2$  fit. A track candidate is accepted if it passes the selection criteria [30].

The combined ID-MS reconstruction uses different algorithms to find different *muon types*. There are four main types outlined below. When the same muon is reconstructed in more than one category, preference is given to Combined (CB), then Segment-tagged (ST), and finally Calorimeter Tagged (CT) muons. These algorithms have been continuously improved to increase precision, speed, and robustness against misidentification [32].

- **Combined muons (CB):** They combine tracks from the ID and MS detectors using a global refit on all hits (some may be removed or added to improve quality). Most muons are reconstructed using an outside-in method.
- **Segment-tagged muons (ST):** ST muons are assigned an ID track that is associated

## CHAPTER 4. TRACKING AND ISOLATION IN ATLAS

with at least one local MDT or CSC track after extrapolation. These are used when muons cross only one layer of the MS because of low  $p_T$  or regions out of most MS layer boundaries.

- **Calorimeter-tagged muons (CT):** These muons are identified by an ID track that can be matched to a minimum ionizing particle energy deposit in the calorimeter. They have the lowest purity but are optimized for  $|\eta| < 0.1$  and  $1.5 < p_T < 100$  GeV where the MS is only partially instrumented.
- **Extrapolated muons (ME):** They are reconstructed in the MS with the addition of silicon points and with a loose requirement that the muon track originated at the IP. In general, these muons are required to traverse 2–3 layers of MS chambers. These are mainly used to extend acceptance in the region  $2.5 < |\eta| < 2.7$ , which is not covered by the ID.

In order to distinguish muons from other particles (like backgrounds from pion and kaon decays) strict quality requirements must be set to select prompt muons with high efficiency. Reconstruction targets  $W$  and  $Z$  decays (as opposed to light-hadron decays) which originate from the interaction point. We use the following variables to identify such muons:

- $q/p$  significance, the absolute value of the difference between the ratio of the charge and momentum of muons in the ID and MS divided by the sum in quadrature of their corresponding uncertainties
- $\rho'$ , the absolute value of the difference between the  $p_T$  measurements in the ID and MS divided by the  $p_T$  of the combined track
- $\chi^2$ , the normalized fit parameter of the combined track

## CHAPTER 4. TRACKING AND ISOLATION IN ATLAS

Specific requirements on the number of hits in the ID and MS assure that inefficiencies are expected and momentum measurements are robust. There are four muon identification selections that each addresses specific needs of physics analyses [32].

- **Loose Muons:** The *Loose* criteria maximizes the reconstruction efficiency, losing very few potential muons, while providing satisfactory tracks. All muon types are used in this criteria.
- **Medium Muons:** *Medium* is the default selection for muons in ATLAS because it minimizes systematic uncertainties associated with reconstruction and calibration. Only CB and ME tracks are used with requirements for over 3 hits in at least two MDT layers in most regions. All *Medium* muons are included in the *Loose* criteria.
- **Tight Muons:** *Tight* selects muons with the highest purity, but sacrifices efficiency. All *Tight* muons are included in the *Medium* selection, but only CB muons with at least two hits in the MS are considered, and the  $\chi^2$  value must be less than 8.
- **High- $p_T$  Muons:** *High- $p_T$*  muons have good momentum resolution for tracks with  $p_T > 100$  GeV. This is beneficial to searches for high-mass  $Z'$  and  $W'$  resonances. CB muons in the *Medium* selection with at least 3 hits in 3 MS stations are included.

The  $H \rightarrow WW$  analysis uses Tight muons with the added condition that  $p_T > 15$  GeV and  $\eta < 2.5$  to gain the highest purity possible and eliminate background from misidentified leptons, which constitute a significant background even with this selection.

We measure the muon reconstruction efficiency in two different ways in the regions  $|\eta| < 2.5$  and  $2.5 < |\eta| < 2.7$ . First, in the barrel region, we use the **Tag-and-Probe** method. In this method we select an almost-pure sample of  $J/\psi$  and  $Z$  decays and require the leading

## CHAPTER 4. TRACKING AND ISOLATION IN ATLAS

muon to be a *Medium* muon labeled the **tag**. The subleading muon, the **probe**, must be reconstructed independently. There are three types of probes:

- **ID track:** Allows measurement of MS efficiency and of tracks not accessible to CT muons.
- **CT tracks:** Allows measurement of MS efficiency and has powerful rejection of background (especially at low  $p_T$ ). This is the most commonly used probe.
- **MS tracks:** Allows measurement of ID and CT efficiency.

To find the overall efficiency of *Medium*, *Tight*, or *High- $p_T$*  muons, we multiply the efficiencies associated with each type of probe. The efficiency  $\epsilon(X|CT)$  ( $X = Medium / Tight / High-p_T$ ) of reconstructing these muons assuming a reconstructed ID track is measured using a CT muon as probe. This result is corrected by the efficiency  $\epsilon(ID|MS)$  of the ID track reconstruction measured using MS probes.

$$\epsilon(X|ID) \cdot \epsilon(ID) = \epsilon(X|CT) \cdot \epsilon(ID|MS) \quad (X = Medium/Tight/High-p_T) \quad (4.2)$$

The ID track reconstruction efficiency must be independent from the muon spectrometer track reconstruction ( $\epsilon(ID) = \epsilon(ID|MS)$ ). In addition, the use of a CT muon as a probe instead of an ID track must not affect the probability for *Medium*, *Tight*, or *High- $p_T$*  reconstruction ( $\epsilon(X|ID) = \epsilon(X|CT)$ ). These assumptions are largely true with simulations showing some small deviations. These deviations are taken into account when calculating systematic errors.

The reconstruction efficiency of *Loose* muons is measured separately for CT muons within  $|\eta| < 0.1$  and all other *Loose* types. The CT muon efficiency is measured using MS probe tracks, and the efficiency of other muons is evaluated similarly to the *Medium*, *Tight*, and

## CHAPTER 4. TRACKING AND ISOLATION IN ATLAS

*High- $p_T$*  muons using CT probe muons [32]. For  $|\eta| > 2.5$ , the efficiency is calculated using the ME muons in the **Loose** and **Medium** selections. The number of muons observed in this region is normalized to the number of muons observed in the region  $2.2 < |\eta| < 2.5$ . A more detailed discussion of the efficiency measurement in this region can be found in Ref [33].

**Scale factors** are defined as the ratios between the efficiency of data and the efficiency of Monte Carlo simulations. They are used to describe the deviation between simulated and real detector behavior and are used in physics analyses to correct simulations.




Figure 4.4: On the left, reconstruction efficiency for *Tight* muons from  $Z \rightarrow \mu\mu$  events is displayed as a function of the  $p_T$  of the muon in the region  $0.1 < |\eta| < 2.5$  with systematic and statistical uncertainties. On the right, muon reconstruction efficiency is shown as a function of  $\eta$  in  $Z \rightarrow \mu\mu$  events for muons with  $p_T > 15$  GeV for *Medium*, *Loose* and *Tight* muons. In both plots the full  $139\text{fb}^{-1}$  Run-2 dataset is used [34].

Figure 4.4 displays reconstruction efficiency for *Medium* muons over a range of  $p_T$  and all other working plotted over  $\eta$ . While *Medium* muons have a higher efficiency than the *Tight* selection used in this analysis, both have an efficiency above 95% for a large range of  $\eta$  and  $p_T$ .  $J/\psi$  decays probe low  $p_T$  muons while  $Z$  decays probe muons of a higher  $p_T$  allowing a large range to be defined. MC simulations match data within 1 – 2%. The only significant loss of efficiency is seen at extremely low  $\eta$  due to criteria excluding ID muons. Overall, the

## CHAPTER 4. TRACKING AND ISOLATION IN ATLAS

default *Tight* muon selection demonstrates reconstruction efficiency around 95% for muons in our selection  $p_T$  range.

Isolation distinguishes muons from  $W/Z$  decays from those produced in the decay of  $b$  and  $c$  mesons. When heavy particles like  $W$ ,  $Z$ , and Higgs bosons decay they often produce muons in isolation. Semileptonic decays of  $b$  and  $c$  hadrons, on the other hand, typically produce muons embedded in jets.

The MCP group uses two muon isolation variables: a track-based variable ( $p_T^{varcone30}$ ) and a calorimeter-based variable ( $E_T^{topocone20}$ ).  $p_T^{varcone30}$  is defined as the scalar sum of the transverse momenta of tracks with  $p_T > 1$  GeV in a cone around the muon of transverse momentum  $p_T$  excluding the muon track itself. The cone size is  $p_T$ -dependent to improve the performance for muons produced in decays with a large transverse momentum.  $E_T^{topocone20}$  is defined as the sum of the transverse energy of topological clusters in a cone around the muon after subtracting the contribution from the energy deposit of the muon itself and correcting for pile-up effects [35].

Table 4.1 defines seven isolation selection criteria - called “isolation working points” - that optimize different physics analyses. The *LooseTrackOnly* and *FixedCutTightTrackOnly* working points are defined by cuts on the relative track-based isolation variable. All other working points are defined by cuts applied separately on both relative isolation variables. All cuts are tuned as a function of the  $\eta$  and  $p_T$  of the muon to obtain a uniform performance. The target efficiencies of the different working points are described in Table ???. The efficiencies for the seven isolation working points are measured in data and simulation using the **Tag-and-Probe** method described previously. Figure 4.5 shows the isolation efficiency measured for *Medium* muons in data and simulation as a function of the muon  $p_T$  for two different working points. In both the *GradientLoose* and *FixedCutTightTrackOnly* working points, efficiency is above 90% and matches simulation well within errors for muons for higher

## CHAPTER 4. TRACKING AND ISOLATION IN ATLAS

$p_T$  muons. In the  $HWW$  analysis, fixed isolation cuts are optimized independent of Muon Performance working points. Only muons with  $p_T > 15$  GeV are considered so the efficiency is optimized for muons which pass this and the *Tight* selection cuts.

Isolation WP	Discriminating variable(s)	Definition
<i>LooseTrackOnly</i>	$p_T^{\text{varcone}30}/p_T^\mu$	99% efficiency constant in $\eta$ and $p_T$
<i>Loose</i>	$p_T^{\text{varcone}30}/p_T^\mu, E_T^{\text{topocone}20}/p_T^\mu$	99% efficiency constant in $\eta$ and $p_T$
<i>Tight</i>	$p_T^{\text{varcone}30}/p_T^\mu, E_T^{\text{topocone}20}/p_T^\mu$	96% efficiency constant in $\eta$ and $p_T$
<i>Gradient</i>	$p_T^{\text{varcone}30}/p_T^\mu, E_T^{\text{topocone}20}/p_T^\mu$	$\geq 90(99)\%$ efficiency at 25 (60) GeV
<i>GradientLoose</i>	$p_T^{\text{varcone}30}/p_T^\mu, E_T^{\text{topocone}20}/p_T^\mu$	$\geq 95(99)\%$ efficiency at 25 (60) GeV
<i>FixedCutTightTrackOnly</i>	$p_T^{\text{varcone}30}/p_T^\mu$	$p_T^{\text{varcone}30}/p_T^\mu < 0.06$
<i>FixedCutLoose</i>	$p_T^{\text{varcone}30}/p_T^\mu, E_T^{\text{topocone}20}/p_T^\mu$	$p_T^{\text{varcone}30}/p_T^\mu < 0.15, E_T^{\text{topocone}20}/p_T^\mu < 0.30$

Table 4.1: The seven isolation working points are described by their discriminating variables and defining criteria [32].




Figure 4.5: Isolation efficiency for the GradientLoose (left) and FixedCutTrackOnly (right) muon isolation working points. The efficiency is displayed as a function of  $p_T$  in  $Z \rightarrow \mu\mu$  events. The black markers show efficiency measured in data samples while the red show MC simulations. The bottom panel shows the ratio of the efficiency between the two as well as both statistical and systematic uncertainties [36].

The isolation cuts used by the  $H \rightarrow WW$  analysis are  $p_T^{\text{varcone}30}/p_T < 0.06$  and  $p_T^{\text{topocone}20}/p_T <$

## CHAPTER 4. TRACKING AND ISOLATION IN ATLAS

0.09 which are relatively tight selections and designed to reduce misidentified lepton backgrounds.

The muon momentum scale and resolution are studied using  $Z$  and  $J/\psi$  decays. In order to obtain agreement between simulation and data in muon momentum scale to the per mille level and in resolution to the percent level, we need to apply a set of corrections to the simulated muon momentum. After applying the corrections we validate them by comparing the muon momentum scale and resolution between simulation and data over  $\eta$ ,  $\phi$ , and  $p_T$ .

We extract the calibration parameters with the transverse momentum of the ID and MS components of a CB track. The corrected transverse momentum is described by the following equation:

$$p_T^{\text{Cor,Det}} = \frac{p_T^{\text{MC,Det}} + \sum_{n=0}^1 s_n^{\text{Det}}(\eta, \phi)(p_T^{\text{MC,Det}})^n}{1 + \sum_{m=0}^2 \Delta r_m^{\text{Det}}(\eta, \phi)(p_T^{\text{MC,Det}})^{m-1} g_m}. \quad (4.3)$$

Here the  $g_m$  terms are normally distributed random variables with zero mean and unit width. The  $\Delta r$  and  $s$  terms describe momentum resolution smearing and scale corrections applied in specific detector regions, respectively. Both the ID and MS are divided into 18 pseudorapidity regions and the MS is divided into two  $\phi$  bins separating the large and small sectors. Each of these bins leverages different alignment techniques and has different material distributions.

There are two  $s$  terms that represent different types of corrections.  $s_1$  corrects for inaccuracy in the description of the magnetic field integral and the detector in the direction perpendicular to the magnetic field.  $s_0$  corrects for the inaccuracy in the simulation of energy loss in the calorimeter and other materials. Since this loss is negligible in the ID, it is only nonzero in the MS [32].

The denominator introduces momentum smearing which broadens the  $p_T$  resolution in

## CHAPTER 4. TRACKING AND ISOLATION IN ATLAS

simulation. The parametrization of the smearing is defined as

$$\frac{\sigma(p_T)}{p_T} = r_0/p_T \oplus r_1 \oplus r_2 \cdot p_T. \quad (4.4)$$

In this equation  $r_0$  is related to the fluctuations in energy loss in the traversed material,  $r_1$  accounts for multiple scattering, local magnetic field inhomogeneities, and local radial displacements of hits, and  $r_2$  describes intrinsic resolution effects caused by the spatial resolution of the hit measurements and by residual misalignment of the MS [32]. Correction parameters are extracted from data using a binned maximum-likelihood fit with templates derived from simulation which compares the invariant mass distributions for  $J/\psi$  and  $Z$  decay candidates in data and simulation. The muons are carefully selected to be compatible with tracks that start at the interaction point and penetrate both the ID and the MS. Muons are also selected to pass specific momentum and isolation criteria. The dimuon mass distribution of these tracks in data is fitted using a Crystal Ball function convoluted with an exponential background distribution in the ID and MS fits. The background model and its normalization are then used in the template fit. The fits are performed in  $\eta - \phi$  regions of fit (ROFs) which compromise regions with uniform features in the ID and MS [32].

From these fits, we can find the smearing terms across all  $\eta$  regions. Once the corrections are applied, we can validate that the agreement between data and MC is excellent. This is shown in Figure 4.6.  $r_0$  is set to zero across all  $\eta$  regions since energy loss is negligible in the ID.  $r_1$  and  $r_2$  increase as  $\eta$  increases since spatial resolution decreases and inhomogeneities increase as we move from the barrel to end-cap regions of both the ID and MS. Muon momentum corrections are continuously studied during ATLAS runs to validate muon calibration performance and account for discrepancies.




Figure 4.6: The  $r$ - values from each of 10 fits of resolution to  $p_T$  for ID muon simulations are shown. Each value corresponds to a particular ROF or  $\eta$  region. These plots show  $r_1$  (left) and  $r_2$  (right) as functions of leading muon  $\eta$ .

## 4.4 Jets

Quarks and gluons emitted from high-energy hard scattering do not appear in the detector directly. Quarks and gluons with high enough energy radiate low energy gluons until partons are able to bind into color-neutral hadrons. These hadrons are collimated in groups as “jets”. Jet energy and momentum are used in physics analyses as proxies for initial scattered partons. Pile-up presents the main difficulty in jet calibration as multiple interactions occurring in the detector at once create significant, often hadronic, background. The hard interaction of interest must be separated from pile-up background, which is most often soft. During Run-1, the ATLAS experiment reconstructed jets using either only the calorimeter or the tracker, though most often the calorimeter. Topological clusters of calorimeter cells (topo-clusters) were used to trace jet tracks. At the end of Run-1, the jet energy scale (JES) correction factor used to calibrate jets to the particle level was re-calculated using additional track information from the Inner Detector and Muon Spectrometer, which greatly improved jet resolution [37].

## CHAPTER 4. TRACKING AND ISOLATION IN ATLAS

In Run-2, a new algorithm for jet reconstruction took advantage of the improvements shown in Run-1 by including information from the tracker. ‘Particle flow’ uses the tracker’s higher momentum resolution for low-energy charged particles and its greater angular resolution of single charged particles. This is complemented by the calorimeter’s ability to reconstruct both charged and neutral particles and the calorimeter’s higher energy resolution for high energy physics objects. The calorimeter also has an extended acceptance so in the forward region only calorimeter topo-clusters are used. One potential difficulty with the ‘particle flow’ algorithm is the possibility for double counting particles if the jets reconstructed by the tracker are not properly matched with the corresponding calorimeter signals. This is avoided in the algorithm through the condition that if a particle’s track measurement is used, its corresponding energy must be subtracted from the calorimeter measurement. The success of the algorithm in removing only energy deposits from the tracked jet represents a key criteria for its overall performance [38].




Figure 4.7: Flow chart of the particle flow algorithm beginning with track selection and ending with charged particles and changed/unchanged topo-clusters [38].

Topo-clusters are described by two main properties -  $\epsilon$ , which represents the fraction of true energy deposited in the cluster out of the total true energy deposited in all topo-clusters for that object, and  $\rho$ , the fraction of a particle’s true energy which lies within the topo-cluster. High  $\rho$ , high  $\epsilon$  topo-clusters allow contributions from different particles to be distinguished and so are easier to apply hadronic shower subtraction [38].

Jet tracks need to meet strict criteria. In this analysis, tracks included in jets are required

## CHAPTER 4. TRACKING AND ISOLATION IN ATLAS

to satisfy the “tight” selection criteria, which includes requirements for at least nine hits in silicon detectors, no missing Pixel hits,  $\eta < 2.5$ , and  $40 > p_T > 0.5$  GeV. High  $p_T$  tracks are excluded because of their poor isolation. In addition, tracks which are identified as electrons or muons are excluded [38].

With topo-clusters and tracks assembled, the algorithm matches each to one another. Topo-clusters are ranked through the distance metric

$$\Delta R' = \sqrt{\left(\frac{\Delta\phi}{\sigma_\phi}\right)^2 + \left(\frac{\Delta\eta}{\sigma_\eta}\right)^2}, \quad (4.5)$$

where  $\sigma_\phi$  and  $\sigma_\eta$  denote angular topo-cluster widths. A requirement that  $E^{clus}/p^{trk} > 0.1$  is applied so that the energy of the topo-cluster must contain a significant portion of the energy of the track. This requirement rejects about 30 – 40% of the incorrect topo-clusters at  $p_T > 5$  GeV. The closest topo-cluster to each track in  $\Delta R'$  is taken to be the correct match. If no topo-cluster is within a cone of  $\Delta R' = 1.64$  it is assumed that the particle did not form a topo-cluster in the calorimeter [38].

Topo-clusters are thus matched to particle tracks and the energy deposited by the particle is subtracted from the calorimeter. The average energy deposited by a particle with momentum  $p^{trk}$  is  $\langle E_{dep} \rangle = p^{trk} \langle E_{ref}^{clus}/p_{ref}^{trk} \rangle$ , where  $\langle E_{ref}^{clus}/p_{ref}^{trk} \rangle$  is calculated using single-particle samples without pile-up by summing topo-cluster energies within  $\Delta R = 0.4$  about the track position. These are calculated at varying  $p_T$  and  $\eta$  values to capture effects from detector geometry and shower development. Particles often split their energy between multiple topo-clusters and this split can be determined through the significance of the difference between expected energy and that of the matched topo-cluster. This new full set of matched clusters is considered for energy subtraction [38].

Energy subtraction is performed cell-by-cell unless  $\langle E_{dep} \rangle$  is greater than the energy

## CHAPTER 4. TRACKING AND ISOLATION IN ATLAS

of the total matched topo-clusters, in which case they are all removed. Rings are formed in  $(\eta, \phi)$  about the extrapolated track and are one calorimeter cell wide. The average energy density in each ring is computed and the ring with the highest energy density is subtracted first. This continues to lower density rings until  $\langle E_{dep} \rangle$  is reached. Finally, energy clusters from shower fluctuations are removed. Ideally now the sum of selected tracks and remaining topo-clusters together represent the reconstructed event without double counting. An example display of particle flow events is shown in Figure 4.8




Figure 4.8: Sum of transverse momenta of neutral and charged particle flow objects in an area  $\Delta\eta \times \Delta\phi = 0.2 \times 0.2$  from a 2017 event with mean number of interactions per beam crossing  $\mu = 38$  [39]

Jet-finding algorithms aim to approximate the hadrons that generate calorimeter and track-based jets in the detector. They bridge the divide between observable jet objects and theoretical predictions from QCD. There are several jet algorithms in use currently and all are collinear and infrared safe. This means that neither splitting a jet collinearly nor soft emissions should change jet structure. Without these properties perturbation theory diverges at high orders.

Anti- $k_t$  is the jet algorithm used by ATLAS and is infrared collinear-safe by construction

## CHAPTER 4. TRACKING AND ISOLATION IN ATLAS

and resilient to soft radiation [40].

Jet reconstruction and isolation does not end with particle flow and the anti- $k_t$  algorithm. Further corrections have to be applied to improve the agreement between data and MC simulations. MC simulation are calibrated to better model pile-up and to improve jet angular resolution. Global sequential calibrations are calculated with MC using calorimeter, track, and muon chamber information to improve jet resolution. Data is used to constrain uncertainties with through known dijet samples. These calibrations lead to final measurements of jet energy scale (JES) and jet energy resolution (JER) and provide reconstructed jet events in MC and data as well as a number recommended uncertainties for physics analyses to use. Figures 4.9 show jet energy scale and resolution as a function of  $p_T$  for PFlow jets modeling with the anti- $k_t$  algorithm after JES corrections are applied. Fully combined systematic uncertainties are also shown [41].




Figure 4.9: Left: data-to-simulation ratio for average jet  $p_T$  as a function of jet  $p_T$ . Three techniques shown as well as their their combination (black) and the combination total uncertainty. PFlow Anti- $k_t$  jets with  $R=0.4$  and JES correction. Right: relative jet energy resolution as a function of  $p_T$  for anti- $k_t$  PFlow jets with  $R = 0.4$  in 2017. JES calibrations applied and compared to MC with full systematic uncertainties [39].

Jet reconstruction and calibration are crucial to this analysis and later chapters will detail

how impactful jet energy scale and resolution uncertainties are to the overall precision of the  $H \rightarrow WW$  cross-section measurement.

## 4.5 Missing transverse energy

Colliding protons in collider experiments like ATLAS have momentum solely in the plane of the beam. Conservation of momentum implies that in the plane transverse to the beam ( $x - y$ ) the sum of momentum from all interaction by-products is zero. This is most often not the case and the non-zero transverse momentum from any interaction is termed  $E_T^{miss}$ . Missing transverse energy is a sign of final state neutrinos.  $E_T^{miss}$  could also point to new particles that cannot be directly detected, like neutralinos or dark matter particles.  $E_T^{miss}$  could also signify interacting particles which evade detection in ATLAS due to detector acceptance or poor reconstruction [42].

Missing transverse energy is determined using a combination of all reconstructed particles in an event. This is challenging because it involves all detector components and final particle types. The Jet/ $E_T^{miss}$  performance group delivers calibrations for multiple  $E_T^{miss}$  variables that we use in the  $H \rightarrow WW$ . This section will discuss  $E_T^{miss}$  reconstruction and performance followed by definitions of a few additional  $E_T^{miss}$  observables. Reconstructed  $E_T^{miss}$  calculations take into account both *hard* and *soft* event signals. Hard-events are composed of fully reconstructed and calibrated particles like electrons, muons, photons,  $\tau$ -leptons and jets. All other objects are considered *soft*. Reconstruction for all particle types happens independently, which means that the same signal may be used to identify two distinct particles. This double-counting is taken into account in  $E_T^{miss}$  resolution. At its most basic,  $E_T^{miss}$  is defined:

$$E_{x(y)}^{miss} = - \sum_{i \in \text{hard objects}} p_{x(y),i} - \sum_{j \in \text{soft objects}} p_{x(y),j} \quad (4.6)$$

## CHAPTER 4. TRACKING AND ISOLATION IN ATLAS

where overall  $E_T^{miss}$  is a vector composed of  $x, y$  components. In order to avoid double-counting the same detector signal in multiple particle reconstruction algorithms, hard objects are considered in order: electrons, photons, hadronically decaying  $\tau$ -leptons, and then jets. Muons have little overlap since they are reconstructed using MS and ID tracks (and muons alone leave tracks in the MS). As particles are reconstructed in this sequence, signals used more than once are rejected to avoid overlap. Another key variable is  $\sum E_T$  which is defined

$$\sum E_T = \sum_{\text{electrons}} p_T^e + \sum_{\text{photons}} p_T^\gamma + \sum_{\tau\text{-leptons}} p_T^\tau + \sum_{\text{muons}} p_T^\mu + \sum_{\text{jets}} p_T^{jet} - \sum_{\text{unused tracks}} p_T^{track} \quad (4.7)$$

The first five terms show the hard term while the last represents the soft-term. Selections are applied to reconstructed jets and particles to achieve optimal  $E_T^{miss}$  for a particular analysis [43]. In the context of  $H \rightarrow WW$  we use a “tight” configuration for  $E_T^{miss}$  classified by strict conditions on accepted jets. This working point has the greatest pile-up rejection which is integral to our analysis.

$E_T^{miss}$  reconstruction contains the complexity of each of its component parts and their  $p_T$  resolutions all affect total  $E_T^{miss}$  resolution. Pile-up and total event activity also play a large role in  $E_T^{miss}$  performance. Validations for  $E_T^{miss}$  are performed on a variety of observables and MC modelling is compared to reconstructed data. Systematic uncertainties are derived from comparing the reproducibility of these observables and their successful modelling of data. Resolution for reconstructed jets and leptons are also propagated to overall  $E_T^{miss}$  uncertainty.  $E_T^{miss}$  performance is evaluated using  $Z \rightarrow \mu\mu$ ,  $Z \rightarrow e^-e^+$ , and  $W \rightarrow e\nu$  events, the first two with no genuine  $E_T^{miss}$  and the third with significant  $E_T^{miss}$  from neutrinos. After specific event selection and  $E_T^{miss}$  reconstruction,  $E_T^{miss}$  for each of these samples is studied and data and MC are demonstrated to match within uncertainties. Selected performance observables are shown for the full Run-2 dataset in 4.10. Here  $Z \rightarrow e^+e^-$  reconstructed

## CHAPTER 4. TRACKING AND ISOLATION IN ATLAS

track-based soft term  $E_T^{miss}$  and  $E_T^{miss}$  significance distributions are shown. Data and MC show good agreement.




Figure 4.10: Left: Track-based Soft Term  $E_T^{miss}$  shown for the complete Run-2 dataset with an integrated luminosity of  $140 \text{ fb}^{-1}$ . Monte Carlo simulations are compared to data for the tight  $E_T^{miss}$  working point. Right: Track-based Soft Term  $E_T^{miss}$  significance modelling shown in the tight  $E_T^{miss}$  working point [39]

In this analysis, a number of  $E_T^{miss}$  variables are used. These variables are used to reject backgrounds particularly from  $Z \rightarrow \tau\tau$  events. Some of the variables used for these selections include track-based  $E_T^{miss}$ , track-based soft term  $E_T^{miss}$ ,  $m_{\tau\tau}$ ,  $E_T^{miss}$  significance and  $p_T^{tot}$ . The  $E_T^{miss}$  soft term is composed of a track-based and calo-based component. The calorimeter-based component is highly dependent on pile-up and so using a track-based soft term reduces overall pile-up dependence. Track-based  $E_T^{miss}$  uses only reconstructed ID tracks from the primary vertex and in our analysis this has been just as powerful discriminant for isolating  $Z \rightarrow \tau\tau$  background events from VBF signal as overall  $E_T^{miss}$  without the same pile-up dependence and correlation with other kinematic variables.  $E_T^{miss}$  significance is a newly defined and recommended variable from the Jet/ $E_T^{miss}$  group. Described further in [42],

## CHAPTER 4. TRACKING AND ISOLATION IN ATLAS

it discriminates real missing energy from momentum resolution effects using a calculated likelihood. Shown in 4.10,  $E_T^{miss}$  significance peaks at low values if  $E_T^{miss}$  likely comes from resolution effects and not from a real invisible particle in the event. Our analysis defines two additional variables based on  $E_T^{miss}$ :  $p_T^{tot}$  describes the total transverse momentum from all hard objects in the event, and  $m_{\tau\tau}$ , defined as

$$m_{\tau\tau} = \frac{m_{\ell\ell}}{\sqrt{x_1 * x_2}}, \quad (4.8)$$

where

$$\begin{aligned} x_1 &= \frac{p_x^{\ell 0} * p_y^{\ell 1} - p_y^{\ell 0} * p_x^{\ell 1}}{p_y^{\ell 1} * E_{Tx}^{miss} - p_x^{\ell 1} * E_{Ty}^{miss} + p_x^{\ell 0} * p_y^{\ell 1} - p_y^{\ell 0} * p_x^{\ell 1}}, \text{ and} \\ x_2 &= \frac{p_x^{\ell 0} * p_y^{\ell 1} - p_y^{\ell 0} * p_x^{\ell 1}}{p_x^{\ell 1} * E_{Ty}^{miss} - p_y^{\ell 0} * E_{Tx}^{miss} + p_x^{\ell 0} * p_y^{\ell 1} - p_y^{\ell 0} * p_x^{\ell 1}} \end{aligned} \quad (4.9)$$

Each of these variables contributes to our signal region selection and elimination of background. A number of systematic uncertainties from reconstructed  $E_T^{miss}$  are defined and used in our analysis, though these are small compared to the uncertainties from jets and other final state particles.

In this chapter, I have outlined the procedures for building all the physics objects we use in the  $H \rightarrow WW$  analysis. The next chapters will use these physics objects and their kinematic variables to understand our background and signal events.

# Chapter 5

## Event Selection

This chapter steps through some of the object definitions and inputs used in the analysis. First, the data and Monte Carlo simulations used for signal and background modelling are summarized. Then, the kinematic variables and observables used in event selection and later for the fiducial cross-section measurement are defined. Finally, the signal selection is outlined and its results shown. I played a primary role in testing and optimizing the analysis selection using reconstructed events.

### 5.1 Data and Monte Carlo samples

#### 5.1.1 Data samples

The full Run-2 dataset containing all proton-proton collision data collected from 2015-2018 at  $\sqrt{s} = 13$  TeV with a 25ns bunch spacing configuration are used.

In 2015, 2016, 2017, and 2018  $3.86\text{ fb}^{-1}$ ,  $35.6\text{ fb}^{-1}$ ,  $46.9\text{ fb}^{-1}$ , and  $62.2\text{ fb}^{-1}$  of luminosity were recorded, respectively. Peak instantaneous luminosity increased from  $5.0 \times 10^{33}\text{ cm}^{-2}\text{s}^{-1}$  in 2015 to  $21.4 \times 10^{33}\text{ cm}^{-2}\text{s}^{-1}$  in 2018. Average (peak) pile-up also increased from  $\langle\mu\rangle = 13.6$

(40.5) in 2015 to  $\langle \mu \rangle = 37.0$  (90) in 2018. These pile-up distributions datasets are shown in 5.1.




Figure 5.1: The luminosity-weighted distribution of the mean number of interactions per crossing is shown for Run-2 pp collision data [44].

Events are only used if all relevant detector components are operating normally. These events are part of the standard “All Good” Good Run List comprising a total integrated luminosity of  $139 \text{ fb}^{-1}$  and data quality efficiency of 91.5%.

Throughout Run-2, the instantaneous luminosity and pile-up changed over time. This is accounted for in our MC modelling, as described in the next section.

### 5.1.2 Monte Carlo samples

Monte Carlo samples are generated to compare to data in order to test Standard Model predictions and search for discrepancies. These simulations are also used to optimize the analysis selection criteria. Monte Carlo events are fully simulated using the ATLAS detector simulation in the GEANT4 framework [45] and reconstructed with standard ATLAS reconstruction software. Pile-up is simulated as additional  $pp$  interations in a separate simulation

## CHAPTER 5. EVENT SELECTION

step during digitization where minimum bias events are superimposed on the simulated signal events. These additional events are added based on that year’s recorded pile-up to account for dataset differences.

Separate programs are used to generate the hard scattering process and to model the parton showering (PS), hadronization, and the underlying event (UE). The next sections summarize the simulation techniques for our signal VBF samples, other Higgs production modes, and relevant backgrounds.

The data is split into three time-based categories. Data-taking conditions in 2015 and 2016 are averaged and described together as MC16a. 2017 and 2018 data-taking conditions are considered separately as MC16d and MC16e respectively.

### Vector boson fusion Higgs samples

VBF Higgs events are generated through POWHEG [46] interfaced with PYTHIA8 with the PDF4LHC15 parton distribution function (PDF) set [9]. Cross sections are calculated with full NLO QCD and EW corrections [47, 48] with an approximate NNLO QCD correction applied [49]. These cross-sections as well as associated branching ratios are calculated by the LHC Higgs Cross Section Working Group Yellow Report 4 [8]. Generated events are normalized to calculated cross-sections.

### Other production mode Higgs samples

Other Higgs production modes are considered in the analysis including gluon fusion (ggF), associated Higgs boson production ( $VH$ ,  $V = W, Z$ ), and Higgs boson production in association with a heavy quark pair ( $t\bar{t}H$ ). Though only ggF has an appreciable yield in the signal region, each of these are studied and considered in the analysis. As in VBF Higgs sample, ggF,  $VH$ , and  $t\bar{t}H$  are produced with PYTHIA8[50, 51] for decay, parton shower,

hadronisation and multiple parton interactions.

### Background samples

Main Standard Model backgrounds include events from production of dibosons, top-quark,  $Z$ +jets,  $W$ +jets and multijets. The  $WW$  samples are generated using SHERPA 2.2.2 interfaced with NNPDF3.0 NNLO PDFs.  $Z$ +jets (or Drell-Yan) production is simulated with SHERPA 2.2.1 using the NNPDF3.0 NNLO PDFs with dedicated parton shower tuning developed by Sherpa authors. Top-quark pair production ( $t\bar{t}$ ) is simulated using POWHEG with the POWHEG-Box framework using the NNPDF 3.0 PDFs and interfaced with PYTHIA 8 using NNPDF 2.3 PDFs for parton showering. Single top, mainly  $Wt$ , production is generated with POWHEG-Box 2.0 interfaced to PYTHIA 6.428 for parton showering.  $Z\gamma$  and  $W\gamma$  productions are modeled using SHERPA 2.2.2 at the NLO accuracy for 0- and 1-jet. The  $W$ +jets process modeling is based on a data-driven method described in the next chapter. MC samples (with the same MC generators for both  $W$ +jets and  $Z$ +jets samples) are used to validate the fake estimation and estimate sample composition uncertainties. These processes are generated with POWHEG MiNLO interfaced to PYTHIA 8 with the AZNLO tune.

## 5.2 Object definitions

This analysis utilizes a number of calibrated physics objects and the accurate use of each determines the precision of our  $H \rightarrow WW$  cross-section measurement. Lepton, jet, and missing transverse energy reconstruction, isolation, and calibration were described in some detail in the Chapter 4. This chapter focuses on the particular parameters applied for these physics objects in this analysis. The object definitions are used in accordance with

## CHAPTER 5. EVENT SELECTION

the  $H \rightarrow WW$  coupling analysis for consistency where optimization focussed on  $H \rightarrow WW$  measurements at large. Finally, further observables used in event selection are defined and described here as well.

### 5.2.1 Lepton

Our choice of lepton identification algorithm impacts both the rejection of fake lepton backgrounds ( $W+jets$ ) and QCD background. Tighter requirements on lepton identification decrease signal efficiency so the optimal criteria balance high signal efficiency and high background rejection.

Overall, all events must have at least two leptons tracks and each lepton track must have  $p_T > 400$  MeV. Further, leptons are all required to originate at the hard-scatter primary vertex, which is defined as the primary vertex with the largest track  $\sum p_T$ . Leptons must pass two impact parameter selections to ensure they originate at the primary vertex. The longitudinal impact parameter of each lepton track is defined as  $|z_0 \sin \theta|$  where  $z_0$  is the impact parameter and  $\theta$  the track angle. Each lepton longitudinal impact parameter must be less than 0.5 mm. The significance of the transverse impact parameter is calculated with respect to the beam line ( $|d_0|/\sigma_{d_0}$ ) and has a requirement of three (five) for muons (electrons) as recommended by the Muon and E/ $\gamma$  combined performance groups.

#### Electron

This analysis uses both the *Medium* and *Tight* identification selections which have 94% and 88% identification efficiency respectively for an electron with  $E_T = 100$  GeV. Electrons with  $E_T > 25$  GeV must pass the *Medium* identification while those with  $15 < E_T < 25$  GeV are required to pass the *Tight* selection. This selection is the same used in the  $H \rightarrow WW$

## CHAPTER 5. EVENT SELECTION

coupling analysis where optimization studies demonstrated its impact on overall significance.

Electrons are identified in the range  $|\eta| < 2.47$ , where the transition region between barrel and endcaps in the LAr calorimeter ( $1.37 < |\eta| < 1.52$ ) is excluded.

Electron isolation uses two different selections based on electron track  $p_T$ . The *IsoGradient* working point is used for  $p_T > 25$  GeV. This requires calorimeter and track isolation about cones of  $\Delta R = 0.2$  (*topoetcone20*, *ptvarcone20*). Electron isolation is changed to the fixed cut track cone isolation for  $p_T < 25$  GeV to further eliminate fake background contributions. These require both track and calorimeter variables to fall below the  $p_T$  dependent  $0.1143 \times p_T + 92.14$ .

Table 5.1 summarizes electron selection criteria.

Table 5.1: Electron selections

$p_T$ range	$ \eta $ range	Electron ID	Isolation	Impact parameter
< 25GeV	0 – 1.37, 1.52 – 2.47	Tight	FixedCutTrackCone40	$ z_0 \sin\theta  < 0.5$ , $ d_0 /\sigma_{d_0} < 5$
> 25GeV		Medium	IsoGradient	

## Muon

As described previously, muons can be reconstructed using inner detector tracks, calorimeter deposits, and muon spectrometer tracks. Muons in this analysis must fit the *Tight* selection as well as pass cuts  $p_T > 15$  GeV and  $|\eta| < 2.5$ . Muon identification and isolation criteria match those used in the  $H \rightarrow WW$  coupling analysis and optimize background rejection and signal efficiency. Muon isolation uses calorimeter and track isolation with a cone of  $\Delta R = 0.2$  (*topoetcone20*, *ptvarcone20*). Muon selection criteria are summarized in table 5.2.

## CHAPTER 5. EVENT SELECTION

Table 5.2: Muon selections

$p_T$ range	$ \eta $ range	Muon ID	Calo Isolation	Track Isolation	Impact parameter
$> 15\text{GeV}$	$< 2.5$	“Tight”	$E_T^{\text{cone}20}/p_T < 0.09$	$p_T^{\text{parcone}30}/p_T < 0.06$	$ z_0 \sin \theta  < 0.5,  d_0 /\sigma_{d_0} < 3$

### 5.2.2 Jets

Jets constitute an important part of the analysis both in their number and characteristics. Our signal region selection and most control regions require at least two jets. In order to estimate and reject  $ggF$  Higgs background events, regions with less than two jets are also studied. “Tagging” jets refer to those considered in our jet requirement, though others may be defined. As detailed in Section 4.4, jets reconstruction uses the anti- $k_t$  algorithm to create jet tracks from calorimeter energy deposits within a cone of  $R = 0.4$  and this analysis uses particle flow jet reconstruction.

Jets are required to have  $p_T > 30$  GeV to eliminate high potential for pile-up jets in the full  $\eta$  range,  $|\eta| < 4.5$ . “Jet vertex tagger” variables suppress pile-up events through combinations of multiple variables into a single jet tagger. For jets with  $p_T < 60$  GeV and  $|\eta| < 2.4$ , the JVT variable is required to be larger than 0.59. In the forward region, a designated tagger called “fJVT” ought to be applied. However, the current analysis leaves out this requirement leading to more jet pile-up than expected.

The VBF signal region contains an additional jet observable, the central jet veto (CJV), which rejects events with additional jets (other than the two “tagged” jets) with  $p_T > 20$  GeV within the rapidity gap between two leading jets. This veto increases VBF sensitivity through removal of hadronic backgrounds.

***b*-tagged jet**

The Jet/ $E_T^{miss}$  group provides tools to identify jets formed from bottom quarks. While the previous HWW analysis [15] used the MV2C10 jet tagging algorithm, the current analysis utilizes the DL1 tagger. Both tools use jet kinematics, impact parameters, and secondary vertex variables as input to machine learning classifiers. The DL1 tagger uses a neural network training method (the MV2C10 a boosted decision tree) trained with  $t\bar{t}$  signal to discriminate  $b$ -quarks from light and  $c$ -quarks. The DL1 tagger shows greater b-veto efficiency in isolating top background events and so is used in this analysis.

**5.2.3 Missing transverse energy**

Missing transverse energy is used to both suppress background and build other variables used in signal selection like  $m_{tt}$  and  $p_T^{tot}$ .  $E_T^{miss}$  definitions and calculations are explained in Section 4.5. This analysis uses the “Tight”  $E_T^{miss}$  working point which has proven the most robust against increasing pile-up.

**5.2.4 Overlap removal**

Overlap removal is applied to electrons, muons, and jets following the latest algorithm recommended by the Analysis Software Group (ASG). Current removal steps can be summarized as followed:

- If a muon and electron share an ID track the electron is removed and if a calo-tagged muon shares an ID track with an electron, the muon is removed.
- If  $\Delta R(\text{jet}, e) < 0.2$  the jet is removed to eliminate overlap with the nearby electron. The electron is removed if  $\Delta R(\text{jet}, e) < \min(0.4, 0.04 + 10 \text{ GeV}/p_T^e)$ .

## CHAPTER 5. EVENT SELECTION

- If  $\Delta R(\text{jet}, \mu) < 0.2$  and the jet has less than three associated tracks with  $p_T > 500$  MeV the jet is removed. The jet is also removed if the  $p_T$  ratio of the muon and jet is larger than 0.5 and the ratio of the muon  $p_T$  and the sum of the  $p_T$  of all jet tracks with  $p_T > 500$  GeV is larger than 0.7. The muon is removed if  $\Delta R(\text{jet}, \mu) < \min(0.4, 0.04 + 10 \text{ GeV}/p_T^\mu)$  for any surviving jets.

### 5.3 Common Observables

Dedicated observables are used in this analysis to reject and isolate backgrounds as well as to enhance VBF signal strength. This section lists and describe key observables used in the analysis.

- $m_{\tau\tau}$ : This can be defined as in Section 4.5, if charged leptons are the products of a pair of  $\tau$  leptons, the neutrinos are assumed to be collinear with the charged leptons, and the neutrinos are the only source of the observed  $E_T^{miss}$  in the event. We cut on this variable to suppress  $Z \rightarrow \tau\tau$  events.
- $N_{b-jet}$ : Defines the number of jets with  $p_T > 20$  GeV identified as  $b$ -jets from the  $b$ -tagging algorithm (DL1) and is used to reject  $t\bar{t}$  background.
- $\Delta\phi_{ll}$ : The two leptons from HWW decays tend to be collimated especially compared to non-resonant WW backgrounds. This is due to the spin-zero initial state of the resonant process.
- $m_{ll}$ : The invariant mass of the two leptons from the hard scattering interaction, this targets Drell Yan events and requires  $m_{ll} > 70$  GeV in the control region.

## CHAPTER 5. EVENT SELECTION

- $m_T$ : Transverse mass is used as discriminant variable though not directly cut. It is defined as

$$m_T = \sqrt{(E_{ll} + E_T^{miss})^2 - |p_{ll} + E_T^{miss}|^2}, \quad (5.1)$$

where  $E_{ll} = \sqrt{|p_{ll}|^2 + m_{ll}^2}$ .

- $p_T^{\text{tot}}$ : Total transverse momentum  $p_T^{\text{tot}}$  is defined in Section 4.5. This variable isolates events with significant soft gluon radiation but not high  $p_T$  jets.
- $\Delta Y_{jj}$ : VBF Higgs signal events are characterized by a large separation of the two tagging jets in rapidity. This variable is used as a signal region cut.
- $m_{jj}$ : VBF Higgs events tend to have a high jet invariant mass ( $m_{jj}$ ), defined by combining mass of tagged jets. This analysis applies a cut on  $m_{jj}$  in our signal region.
- $\eta_{lep}$  centrality: This analysis uses an outside lepton veto defined to reject events with leptons outside the rapidity gap between the two tagged jets. The sum of the centralities of each lepton is used as a discriminant in various BDTs in the analysis. The OLV variable is defined using pseudorapidity of the tagged jets and leptons as follows:

$$\begin{aligned} \text{OLV}_{l_0} &= 2 \cdot \left| \frac{\eta_{l_0} - \bar{\eta}}{\eta_{j_0} - \eta_{j_1}} \right|, \\ \text{OLV}_{l_1} &= 2 \cdot \left| \frac{\eta_{l_1} - \bar{\eta}}{\eta_{j_0} - \eta_{j_1}} \right|, \\ \eta_{\text{lep}} \text{ centrality} &= \text{OLV}_{l_0} + \text{OLV}_{l_1}, \end{aligned} \quad (5.2)$$

where  $\bar{\eta} = (\eta_{j_0} + \eta_{j_1})/2$  and so for each lepton:

$$\text{OLV}_l \left\{ \begin{array}{ll} = 0 & \rightarrow \text{The lepton is right in the middle of the rapidity gap between the two tagged jets.} \\ < 1 & \rightarrow \text{The lepton lies within the rapidity gap between the two tagged jets.} \\ > 1 & \rightarrow \text{The lepton is outside the rapidity gap between the two tagged jets.} \end{array} \right. \quad (5.3)$$

- $\sum_{l,j} M_{lj}$ : The sum of the invariant masses of all possible lepton-jet pairs are used to train our signal discriminant as VBF Higgs signal peaks at a higher value than the dominant backgrounds. VBF signal jets tend to be very forward while lepton central so a large separation between lepton and jets is expected as opposed to typical background topologies.

## 5.4 Event selection

Figure 5.2 shows the Feynman diagram for the vector boson fusion Higgs production mode. Two energetic jets with large separation in rapidity accompany the Higgs and its decay products. In addition, since the Higgs is created through the fusion of two electro-weak bosons, there is no QCD interaction between the jets and the Higgs boson or its decay products. The main backgrounds this analysis seeks to mitigate include top quark production, Drell-Yan/ $Z$ , di-bosons ( $WW$ ,  $WZ$ ,  $ZZ$ , and  $W\gamma$ ), ggF Higgs production, and  $W+jets$ .

This analysis controls and estimates these backgrounds with a variety of methods depending on the characteristics of each background. Major backgrounds are estimated in CRs and minor backgrounds are estimated using MC simulation. Top quark production is suppressed in the SR by a  $b$ -jet veto requirement, and can be studied in a very pure orthogonal set of bins in the signal region defined by a boosted decision tree discriminant. A pure validation




Figure 5.2: Feynman diagram for VBF Higgs production [7]

region is also defined to cross-check top MC modelling. The Drell-Yan/ $Z$  background is suppressed by kinematic requirements and is constrained in a dedicated and orthogonal CR and ultimately normalized in the signal phase space. The  $ggF$  Higgs production mode is particularly difficult to separate from the VBF production mode, therefore several CRs are defined to constrain it in data, and ultimately its normalization is determined in the signal phase space. Other Higgs production modes ( $VH$  and  $t\bar{t}H$ ) are analyzed and included as backgrounds, but play a very small role in our overall result. The  $W$ +jets background is estimated with data-driven techniques, as described in the Chapter 6, and then fixed in the simultaneous fit of SR and CRs.

The analysis employs multivariate analysis (MVA), more specifically gradient boosted decision trees trained to discriminate between samples. There are seven trained BDTs used to separate:  $WW$ +top and  $VBF$  events,  $VBF$  and  $ggF$  events,  $ggF$  events from all other samples in each of their three control regions,  $WW$ +top events and all other samples, and  $WW$  and top events from one another. The use of these BDTs separates the backgrounds from our signal in the signal region and maintains discrimination between the  $WW$  and top background samples.

The  $WW$ +top vs. VB BDT is discussed in this chapter while the others are explained with their associated backgrounds in Chapter 6. In addition, two other types of BDT have

## CHAPTER 5. EVENT SELECTION

been tested in this analysis: a  $Z$ +jets vs. VBF BDT and a 3D BDT to separate  $ggF$ ,  $WW$  and VBF events. Results from these tests showed that though these led to minimization of background (in the case of  $Z$ +jets vs. VBF) and good discrimination between key backgrounds in the signal region (in the case of the 3D BDT), when used in the final fit with systematic uncertainties they showed no significant decrease in calculated error in the VBF coupling value. Results from these studies are shown in Appendix I and II.

Background	Relative size Pre-selection	Relative size Signal Region	Estimation	Generator
$WW$	$\sim 8\%$	$\sim 20\%$	MC only + BDT	SHERPA 2.2.2 ( $gg \rightarrow WW$ : SHERPA 2.1)
Top	$\sim 73\%$	$\sim 61\%$	MC only + BDT	POWHEG+PYTHIA 8
$ggF$ Higgs	$< 1\%$	$\sim 2\%$	MC only + BDT	POWHEG+PYTHIA 8 NNLOPS
$W$ +jets	$\sim 2.5\%$	$\sim 4\%$	Data-driven	-
$Z \rightarrow \tau\tau$	$\sim 16\%$	$\sim 7\%$	Data+MC ( $Z \rightarrow \tau\tau$ CR)	SHERPA 2.2.1
$W\gamma, W\gamma^*$	$\sim 1\%$	$\sim 2.5\%$	MC only	SHERPA 2.1/SHERPA 2.2.2

Table 5.3: Each background and its relative size of in SR (after all cuts), estimation method, and generator.

### 5.4.1 Pre-selection

The VBF differential analysis shares object definitions with the VBF and ggF coupling analysis and pre-selection cuts are based off of those used in the 2016 ggF and VBF  $H \rightarrow WW$   $36.1\text{fb}^{-1}$  cross-section paper [15]. The VBF selection utilizes observables defined in the previous section. Cuts applied in the preselection region are listed and described in the table 5.4.

Yields for all samples as well as data for pre-selection cuts are shown in the table 5.5

The plots 5.3 show kinematic distributions after all preselection cuts with fake factors applied directly after a 2-jet cut. These show good modelling with data over a variety of kinematic variables including those used in the Top+ $WW$  vs. VBF BDT. We are not able to directly examine modelling of input variables to this BDT because we require blinding in

## CHAPTER 5. EVENT SELECTION

Pre-sel cut	Description
Channel Selection	Fits $e\mu/\mu e$ channel/event weight applied
Trigger Selection	Trigger selected
Trigger Matching	Trigger weight applied
Data driven muon/electron	W+jets split into electron and muon fake flavors
Cut Jet Cleaning	Pass loose jet cleaning
Cut $V\gamma/Z$ +jets overlap	Eliminate any $Z$ +jets events that overlap with $V\gamma$
Only two leptons	Require exactly 2 leptons (in case of fakes make sure 1 ID/1 Anti-ID)
Lead lepton $p_T$	$p_T^{lead} > 22\text{GeV}$
Sublead lepton $p_T$	$p_T^{sublead} > 15\text{GeV}$
Opposite sign leptons	Require zero dilepton charge
Fake Factor	W+jets weight applied

Table 5.4: Table describing pre-selection cuts applied in common with VBF and  $ggF$  coupling analyses

$\mathcal{L} = 139\text{fb}^{-1}$	$H_{VBF}$	$H_{ggF}$	$\tau\tau H$	$WW$	Top	Zjets	Mis-Id	Total Bkg	Data	Data/MC
Channel Selection	904.81 ± 0.92	9452.28 ± 10.11	2989.84 ± 3.62	175857.28 ± 140.42	1709307.70 ± 287.98	655027.79 ± 1233.45	5121458.35 ± 4457.28	7802403.00 ± 4702.65	4374979	0.56 ± 0.00
Trigger	882.26 ± 0.90	9154.83 ± 9.82	2901.46 ± 3.52	173094.52 ± 138.41	1678206.02 ± 283.57	624251.33 ± 1168.16	5275683.79 ± 4374.34	7882480.56 ± 4600.06	4352644	0.55 ± 0.00
Jet Cleaning	882.26 ± 0.90	9154.83 ± 9.82	2901.46 ± 3.52	173094.52 ± 138.41	1678206.02 ± 283.57	624251.33 ± 1168.16	3905166.04 ± 3213.25	6511962.80 ± 3514.36	4352644	0.67 ± 0.00
Only two Leptons	880.17 ± 0.89	9147.93 ± 9.81	2895.82 ± 3.52	172870.71 ± 138.36	1662432.27 ± 282.33	622191.46 ± 1160.36	3903952.62 ± 3197.39	6490421.92 ± 3496.01	4331979	0.67 ± 0.00
$p_T^{lead} > 22\text{ GeV}$	880.17 ± 0.89	9147.93 ± 9.81	2895.82 ± 3.52	172870.71 ± 138.36	1662432.27 ± 282.33	622191.46 ± 1160.36	3903952.62 ± 3197.39	6490421.92 ± 3496.01	4331979	0.67 ± 0.00
$p_T^{sublead} > 15$	879.98 ± 0.89	9147.27 ± 9.81	2895.32 ± 3.52	172854.38 ± 138.35	1661877.68 ± 282.29	622068.28 ± 1160.19	3902218.38 ± 3196.64	6487838.74 ± 3495.19	4330240	0.67 ± 0.00
OS Leptons	879.86 ± 0.89	9147.00 ± 9.81	2894.70 ± 3.52	172771.82 ± 134.49	1661388.47 ± 282.25	622135.47 ± 1154.02	3896428.83 ± 3190.85	6481093.73 ± 3487.00	4326784	0.67 ± 0.00
$M_{\ell\ell} > 12/10\text{ GeV}$	879.85 ± 0.89	9146.97 ± 9.81	2894.69 ± 3.52	172771.33 ± 134.49	1661368.39 ± 282.25	622133.63 ± 1154.02	3896151.69 ± 3190.77	6480785.01 ± 3486.93	4326642	0.67 ± 0.00
ID leptons	596.53 ± 0.74	5800.47 ± 7.84	1354.56 ± 2.41	126996.63 ± 115.72	1165160.56 ± 237.51	257302.75 ± 431.92	1144546.96 ± 1611.02	2721373.46 ± 1701.42	1587474	0.58 ± 0.00
Apply fake factor	596.53 ± 0.74	5800.47 ± 7.84	1354.56 ± 2.41	126996.63 ± 115.72	1165160.56 ± 237.51	257302.75 ± 431.92	32107.94 ± 241.33	1608934.44 ± 598.08	1587474	0.99 ± 0.00

Table 5.5: Cutflow in the pre-selection region with statistical errors shown.

the signal region. Modelling in the pre-selection region is studied in lieu of this and shows no evidence of bias. In addition, normalization factors (NF) are calculated by data/MC comparisons in top ( $Z \rightarrow \tau\tau$ ) validation (control) regions. These scale factors are added to  $Z \rightarrow \tau\tau$  background events in order to correct for MC mis-modelling and are calculated to be  $0.99 \pm 0.01$  and  $1.01 \pm 0.04$  respectively.

### 5.4.2 Signal region selection

In addition to the pre-selection cuts which are in common with the  $ggF$  coupling analysis, additional selection criteria are applied to the VBF signal region. These selections include a requirement for at least 2 jets ( $n_{jets} \geq 2$ ) and a  $b$ -veto using the DL1r  $b$ -tagging algorithm.

## CHAPTER 5. EVENT SELECTION




Figure 5.3: Distributions of  $\Delta Y_{jj}$ ,  $\Delta Y_{\ell\ell}$ ,  $\Delta \Phi_{\ell\ell}$ ,  $m_{jj}$ ,  $m_{\ell\ell}$ ,  $m_T$ , jet  $p_T^{\text{lead}}$ , jet  $p_T^{\text{sublead}}$ , and  $\Delta \Phi_{jj}$  in the preselection region before blinding is applied. Distributions show good MC modelling of variables used in the signal region BDT.

## CHAPTER 5. EVENT SELECTION

A central-jet-veto (CJV) and an outside-lepton-veto (OLV) are also applied. These remove first events with  $p_T > 20$  GeV which lie between the tagging jets in pseudo-rapidity and next any event where the two charged leptons are not within the tagging jets' rapidity gap. Two additional cuts that differ from the VBF HWW couplings analysis are also added. These require that the mass of the two jets be  $> 200$  GeV and that the rapidity difference between the two jets be  $> 2.1$ . These further purify the signal region against a range of backgrounds, notably top. The plots 5.4 show signal and background yields as well as signal significance at a variety of  $m_{jj}$  and  $\Delta Y_{jj}$  values. The values used here are chosen for their effects on signal significance while retaining high signal statistics.




Figure 5.4: Signal and background yields for VBF and total background (aside from fakes) at various  $m_{jj}$  and  $\Delta Y_{jj}$  cut values. Simulated events for MC16a campaign only. Choice of cut at  $m_{jj} > 200$ GeV and  $DY_{jj} > 2.1$  nearly halves background yields while reducing signal by  $< 5\%$ .

Signal region cuts are listed in table 5.6.

Signal region cut	Description
2-jet (30,30)	Require at least 2 jets with $p_T \geq 30$ GeV
b-veto	Use DL1 efficiency b-tag reject events with b-jets/apply b-tag weight
Blinding (VBF)	Blinding condition on data (cut on VBF vs. Top+WW BDT)
CJV (20GeV)	Cut events with a third central rapidity jet $p_T > 20$ GeV
OLV bool	Leading lepton $\eta$ required to be between two $\eta$ 2 leading jets
$Z \rightarrow \tau\tau$ veto	$m_{\tau\tau} < m_Z - 25$ GeV
$m_{jj}$ cut	$m_{jj} > 200$ GeV
$\Delta Y_{jj}$ cut	$\Delta Y_{jj} > 2.1$

Table 5.6: Table describing VBF signal region cuts

## CHAPTER 5. EVENT SELECTION

Yields for all samples, as well as data for signal region cuts are shown in table 5.7.

$\mathcal{L} = 139 fb^{-1}$	$H_{VBF}$	$H_{ggF}$	$H \rightarrow \tau\tau$	$WW$	Top	Zjets	Mis-Id	Total Bkg	Significance	Data	Data/MC
2-jet (30,30) fJVT	$366.47 \pm 0.58$	$1295.72 \pm 3.71$	$341.06 \pm 1.18$	$24353.25 \pm 30.46$	$911102.17 \pm 201.73$	$26401.33 \pm 108.44$	$11519.17 \pm 164.08$	$980281.52 \pm 300.00$	$0.37 \pm 0.00$	976091	$1.00 \pm 0.00$
b-veto	$323.67 \pm 0.54$	$1109.94 \pm 3.42$	$287.53 \pm 1.07$	$21075.69 \pm 28.85$	$63787.91 \pm 57.22$	$22151.24 \pm 102.73$	$3793.55 \pm 70.69$	$116387.21 \pm 168.10$	$0.95 \pm 0.00$	109677	$0.94 \pm 0.00$
blinding (2-jet)	$323.67 \pm 0.54$	$1109.94 \pm 3.42$	$287.53 \pm 1.07$	$21075.69 \pm 28.85$	$63787.91 \pm 57.22$	$22151.24 \pm 102.73$	$3793.55 \pm 70.69$	$116387.21 \pm 168.10$	$0.95 \pm 0.00$	107456	$0.92 \pm 0.00$
CJV (20GeV)	$256.58 \pm 0.48$	$807.85 \pm 2.92$	$214.70 \pm 0.92$	$15050.73 \pm 24.95$	$43135.02 \pm 47.62$	$16139.77 \pm 89.34$	$2629.36 \pm 59.14$	$81067.63 \pm 144.34$	$0.90 \pm 0.00$	75589	$0.93 \pm 0.00$
OLV bool	$199.59 \pm 0.43$	$218.02 \pm 1.52$	$75.81 \pm 0.47$	$2741.46 \pm 11.72$	$9418.19 \pm 22.26$	$3664.17 \pm 45.30$	$460.46 \pm 26.24$	$17212.14 \pm 68.78$	$1.52 \pm 0.00$	15644	$0.90 \pm 0.01$
$Z \rightarrow \tau\tau$ veto	$172.20 \pm 0.40$	$192.86 \pm 1.43$	$12.62 \pm 0.22$	$1701.95 \pm 9.30$	$6057.49 \pm 17.83$	$1331.98 \pm 35.93$	$311.20 \pm 20.27$	$9979.97 \pm 54.56$	$1.72 \pm 0.01$	8971	$0.88 \pm 0.01$
$M_{jj} > 200$	$165.51 \pm 0.39$	$133.74 \pm 1.19$	$9.99 \pm 0.19$	$1160.80 \pm 8.03$	$3568.73 \pm 13.73$	$857.97 \pm 31.02$	$200.52 \pm 15.37$	$6164.63 \pm 41.76$	$2.10 \pm 0.01$	5401	$0.85 \pm 0.01$
$DY_{jj} > 2.1$	$163.07 \pm 0.39$	$121.34 \pm 1.13$	$9.59 \pm 0.18$	$1027.28 \pm 7.90$	$3005.76 \pm 12.74$	$792.65 \pm 30.70$	$180.37 \pm 14.10$	$5332.00 \pm 40.16$	$2.22 \pm 0.01$	4572	$0.83 \pm 0.01$

Table 5.7: Cutflow in the signal region.

The following plots 5.5 show kinematic distributions after the described selection. Here only MC predictions are shown since data is blinded in the signal region. **Unblinded distributions and cutflows will be added.**

### VBF signal boosted decision tree discriminant

This analysis uses a number of boosted decision trees (BDTs) to both amplify our VBF signal and discriminate and reject a number of specific backgrounds. A decision tree is a collection of cuts designed to classify events as signal-like or background-like. A given signal event is correctly identified if it is placed in a signal-dominated leaf and vice-versa for background events. After the initial tree is built another tree is grown to better separate the signal and background events misidentified by the first tree. This proceeds iteratively until there is a collection of a specified number of trees, in a process known as boosting. A weighted average is taken from all these trees to form a BDT output discriminant with values ranging from -1 to 1.

This section will focus on the BDT trained and used in the signal region to isolate the VBF signal from dominant backgrounds (top and  $WW$  events). The discriminant used in this analysis is the result of numerous studies on training parameters, input variables, and multivariate analysis techniques. Appendix II shows results from studies on using a multidimensional BDT to simultaneously discriminate VBF,  $ggF$ , and  $WW$  background sam-

## CHAPTER 5. EVENT SELECTION




Figure 5.5: Distributions of  $\Delta Y_{\ell\ell}$ ,  $\Delta \Phi_{\ell\ell}$ ,  $m_{\ell\ell}$ ,  $m_T$ ,  $p_{tot}^T$ , lep  $p_{\text{lead}}^T$ , jet  $p_{\text{lead}}^T$  and  $E_{\text{T},\text{rel}}^{\text{miss}}$  in the differential VBF signal region, many used as input to the BDT discriminating VBF from top +  $WW$  backgrounds.

plexes. While initially this 3D BDT showed promising results, estimation of ggF backgrounds through use of multiple control regions (summarized in the next chapter) showed better determination of the ggF background, so the one-dimensional method here was adopted.

This BDT is trained using  $e\mu + \mu e$  events after the VBF selection and the signal regions cuts including that on  $n_{\text{jets}}$ ,  $b$ -veto, OLV, CJV,  $M_{jj}$  and  $\Delta Y_{jj}$ . In this way, the phase

## CHAPTER 5. EVENT SELECTION

space in which we train the BDT is exactly the same as the one where we apply it. The training includes only the top and  $WW$  backgrounds and the VBF signal. Half of the available events are used for training and the other half are used to test the training. This corresponds to about 90,000 un-weighted  $WW$  and top events and 100,000 raw VBF events. This training includes MC weights on events to best account for overall event distributions. There are approximately 2000 total weighted top and  $WW$  events used in the training and 80 weighted VBF events. The TMVA BDTG interface is used to train and test the BDT [52]. The optimal parameters were found through a scan of reasonable values and the final set is summarized in Table 5.8. This BDT utilizes 12 lepton and jet kinematic variables to

Parameter	Value
Boosting algorithm	Gradient
Maximum tree depth	22
Number of trees	400
Minimum number of events requires per mode	5%
Number of cuts	7

Table 5.8: BDT parameters used for the VBF vs. top +  $WW$  training.

distinguish between signal and background events. These include  $\Delta Y_{jj}$ ,  $\Delta Y_{\ell\ell}$ ,  $\Delta \Phi_{\ell\ell}$ ,  $m_{jj}$ ,  $m_{\ell\ell}$ ,  $m_T$ ,  $\eta_{j0}$ ,  $\eta_{j1}$ ,  $p_{j0}^T$ ,  $p_{j1}^T$ ,  $\Delta \Phi_{jj}$ , and  $\sum$  centralities (L). While a larger variety of variables have been tested, these demonstrated the highest discrimination between VBF and top/ $WW$  background. Plots shown in 5.6 and 5.7 demonstrate the input distributions used to train the BDT and their correlations.

The BDT training successfully separates VBF signal and top/ $WW$  background. These backgrounds are considered together because our overall fit uses one parameter to estimate both  $WW$  and top backgrounds due to their similar signatures. One useful metric to test the success of a classifier is a receiver operating characteristic curve (ROC curve). This curve measures signal acceptance over background rejection and its integral for a perfect classifier

## CHAPTER 5. EVENT SELECTION




Figure 5.6: Distributions of input variables to VBF vs. top+WW BDT. Samples are weighted and normalized to even numbers of background and signal events. Signal represents VBF and background top+WW (S. Addepalli).




Figure 5.7: Correlations of input variables to VBF vs. top +WW BDT. Signal represents VBF and background top+WW (S. Addepalli).

## CHAPTER 5. EVENT SELECTION

would be 1. In order to quantify the discrimination we use the integrated-ROC calculated through TMVA for weighted normalized samples and find an optimal value of 0.960, showing very good classification. Comparisons between the test and training show that the BDT is un-biased as one can see that the testing and trainings samples are quite similar. Differences between testing and training samples would imply overtraining, or the BDT using to many parameters on too few events. A Kolmogorov-Smirnov test is performed to measure if the two test and training distributions differ significantly. If the two distributions are random samples of the same parent distribution, the KS-test would give a uniformly distributed value between zero and one (or an average value of 0.5). The closer to 0.5 the KS-test, the greater likelihood the curves come from the same parent, however this calculation is heavily skewed toward lower values so any value above zero (or not very close to zero, on order  $10^{-4}$ ) can be considered not indicative of overtraining. For signal and background we find KS-test values of 0.107 and 0.154, and so no evidence of over-training. We can visualize the BDT output variable both on weighted normalized samples and on samples with all event weights applied. The plot 5.8 shows BDT results applied to normalized samples of VBF signal and top/ $WW$  backgrounds and applied to all fully weighted samples in the signal region.

We aim to fit this distribution in the signal region with high significance in uppermost bins of the distribution. Since this BDT is trained and applied in the signal region, we cannot directly test the modelling of input variables. However, modelling at the pre-selection level for each of these variables, shown earlier in the chapter show no evidence of mis-modelling. The binning for this discriminant used in the statistical fit and its result are shown in the final chapter.

## CHAPTER 5. EVENT SELECTION




Figure 5.8: Weighted, normalized samples of VBF (signal) and top+WW samples (background) plotted over BDT output distribution on left, overlaid testing and training samples shown. Right, full weighted samples of VBF signal and all backgrounds plotted over BDT output distribution after signal region selection. **Data is blinded here** (S. Addepalli).

# Chapter 6

## Backgrounds and Systematics

### 6.1 Backgrounds

While the VBF  $H \rightarrow WW \rightarrow \ell\nu\ell\nu$  channel gives a fairly clear signal with  $E_T^{\text{miss}}$ , two leptons, and 2 jets, there are substantial backgrounds with similar final states. These include Drell-Yan processes (in which a  $Z$  boson is formed from quark anti-quark annihilation of the two colliding protons), top quark final states (predominantly  $t\bar{t}$ ), diboson events (led by SM  $WW$  events), Higgs decays from the other production modes (mainly ggF), and background from mis-identified leptons, called fakes. In this section I will detail each background and describe our methods for estimating their role in overall results. I will describe how this analysis estimates and minimizes contributions from each background process. Finally, I demonstrate purity and modelling of these backgrounds in validation and control regions. I made key contributions in this analysis in optimizing and testing control region definitions and their effects on our overall results.




Figure 6.1: Feynman diagram for Drell-Yan process in which two quarks produce a photon or  $Z$ -boson that then decays leptonically [53].

### 6.1.1 Drell-Yan background ( $Z \rightarrow \tau\tau$ )

The Drell-Yan background, alternatively called  $Z \rightarrow \tau\tau$  or  $Z+\text{jets}$  is produced in the initial collision when two quarks produce a  $Z$  boson which then decays to two leptons (in the largest background,  $\tau$  leptons which then decay to electron/muons and neutrinos). Thus there are two jets and two leptons in its final state as well as missing energy from neutrinos, all of which are indicative of signal VBF Higgs events as well. A typical Drell-Yan Feynman diagram is shown in Figure 6.1. The Drell-Yan background can be discriminated from our signal as well as the other backgrounds in two main ways- first in terms of the mass of the overall event aligning with the mass of the  $Z$  boson. To select for Drell-Yan events we can select those which  $m_{\tau\tau}$  is near the  $Z$  mass peak.  $m_{\tau\tau}$  is defined in the previous section and cutting on this in a 25 GeV window about the  $Z$ -mass peak drastically improves our  $Z+\text{jets}$  purity from only 19% of total events to  $\approx 52\%$ .

We have also tested and trained a secondary discriminant for  $Z+\text{jets}$  in our analysis, a BDT trained to discriminate  $Z+\text{jets}$  and VBF signal events. This BDT significantly decreases overall  $Z+\text{jets}$  background in our signal region but we found that the subsequent decrease in statistics from using this cut leads to very similar overall results. The results from this comparison are shown in the Appendix.

## CHAPTER 6. BACKGROUNDS AND SYSTEMATICS

### Drell-Yan Control Region

The  $Z$ +jets control region definition is quite similar to the VBF signal region except that the  $Z$ +jets veto cut is inverted. Thus instead of removing events near the  $Z$ -mass window we select for them by applying a cut on  $m_{ll} < 80$  GeV,  $66.2$  GeV  $< m_{\tau\tau} < 116.2$  GeV, and the same OLV and CJV cuts as in the VBF signal region. The  $Z$ +jets control region has a purity of  $\approx 82\%$  and yields in this region are shown in the table below.

$\sqrt{s} = 13TeV, \mathcal{L} = 139fb^{-1}$ (Full Run 2)	$H_{VBF}$	$H_{ggF}$	$H_{VH}$	$H \rightarrow \tau\tau$	$WW$	Other VV	Top	Zjets	Mis-Id	Total Blk	Significance	Data	Data/MC
$Z \rightarrow \tau\tau$ CR: $ m_{\tau\tau} - m_Z  < 25$ , bVeto	$27.15 \pm 0.16$	$84.85 \pm 0.95$	$13.02 \pm 0.17$	$88.88 \pm 0.58$	$1809.83 \pm 8.26$	$730.35 \pm 31.87$	$NF = 0.99 \pm 0.01$ $5635.24 \pm 17.11$	$NF = 1.01 \pm 0.04$ $8909.14 \pm 45.18$	$417.01 \pm 26.17$	$17688.32 \pm 64.07$	$0.20 \pm 0.00$	$16400$	$0.93 \pm 0.01$
$Z \rightarrow \tau\tau$ CR: $M_H < 80$ GeV	$26.48 \pm 0.16$	$82.16 \pm 0.93$	$10.92 \pm 0.16$	$73.07 \pm 0.52$	$589.77 \pm 4.57$	$491.77 \pm 26.73$	$NF = 0.99 \pm 0.01$ $1703.83 \pm 9.25$	$NF = 1.01 \pm 0.04$ $8674.53 \pm 40.97$	$232.18 \pm 20.92$	$11858.22 \pm 54.20$	$0.24 \pm 0.00$	$10805$	$0.91 \pm 0.01$
$Z \rightarrow \tau\tau$ CR: CJV < 20 GeV	$20.86 \pm 0.14$	$58.69 \pm 0.79$	$8.29 \pm 0.15$	$55.57 \pm 0.45$	$419.89 \pm 3.94$	$370.64 \pm 23.57$	$NF = 0.99 \pm 0.01$ $1150.59 \pm 7.67$	$NF = 1.01 \pm 0.04$ $6474.42 \pm 35.91$	$185.09 \pm 18.00$	$8723.18 \pm 47.38$	$0.22 \pm 0.00$	$7931$	$0.91 \pm 0.01$
$Z \rightarrow \tau\tau$ CR: OLV	$16.04 \pm 0.12$	$14.49 \pm 0.39$	$1.40 \pm 0.06$	$19.60 \pm 0.23$	$86.60 \pm 1.87$	$70.67 \pm 12.02$	$NF = 0.99 \pm 0.01$ $292.25 \pm 3.90$	$NF = 1.01 \pm 0.04$ $1392.75 \pm 17.23$	$15.06 \pm 8.64$	$1892.82 \pm 23.12$	$0.37 \pm 0.00$	$1832$	$0.96 \pm 0.03$

Table 6.1: Cutflow in the  $Z$ +jets control region.

Data/MC shows decent agreement over various variable distributions as seen in the below. Some of these variables are those used in the BDT described in Appendix A.

Normalization factors (NF) are derived in the  $Z$ +jets control region to correct for Data/MC mis-modelling. These factors are applied to the  $Z$ +jets sample in the signal region. The NF factors are  $1.01 \pm 0.04$ .

#### 6.1.2 Top background

The top background consists of two main components,  $Wt$  and  $t\bar{t}$  events where the  $W$  decays leptonically and the top quarks decay to jets (notably  $b$ -jets). The top background is dominated by  $t\bar{t}$  and is the largest background in our signal region, composing about 60% of the total background. Though top backgrounds are numerous, discrimination between top and signal Higgs events is possible through training a BDT on variables that have very different distributions between these two types of events like  $m_{\ell\ell}$  and  $\Delta Y_{jj}$ . As demonstrated in Chapter 4, this BDT discriminates between signal and top background quite well. The top

## CHAPTER 6. BACKGROUNDS AND SYSTEMATICS




Figure 6.2: Distributions of  $m_{\tau\tau}$ ,  $p_{\text{tot}}^T$ ,  $m_T$ ,  $E_{\text{T}}^{\text{miss}}$ ,  $E_{\text{T},\text{rel}}^{\text{miss}}$ ,  $E_{\text{T}}^{\text{miss}}, \text{significance}$ ,  $E_{\text{T}}^{\text{miss}, \text{track}}$ , and  $E_{\text{T},\text{rel}}^{\text{miss}}$  in the  $Z + \text{jets}$  control region.

background is therefore able to be defined in the signal region and separated using dedicated discriminants. The final statistical fit is discussed in the following chapter but in this section

## CHAPTER 6. BACKGROUNDS AND SYSTEMATICS

I will describe the BDT used to isolate top background events- that splitting top+ $WW$  and all other backgrounds. This is added to the overall fit as the main discriminant for top backgrounds and is considered in combination with  $WW$  due to their similar signatures and their estimation as a combined parameter in the final fit.

We also define a top validation region to test top Monte Carlo modelling as well as to calculate a normalization factor that is used to correct top mis-modelling in the signal region. The top validation region is described similarly to the signal region with one major difference, a  $b$ -tag applied in the signal region requiring all events with a  $b$ -jet to be removed is (almost) reversed. Instead we require exactly one  $b$ -tagged jet. We require exactly one in order to define the validation region as similarly to the signal region as possible while increasing top purity. The result is a highly pure top validation region where the flavor composition of the tagged jets is close to the signal region. Purity of the top control region is  $\approx 97\%$  and yields in this region are shown in the table below.

$\sqrt{s} = 13\text{TeV}, \mathcal{L} = 139\text{fb}^{-1}$ (Full Run 2)	$H_{VBF}$	$H_{e\eta F}$	$H_{VH}$	$H \rightarrow \tau\tau$	$WW$	Other VV	Top	Zjets	Mis-Id	Total Bkg	Significance	Data	Data/MC
Scale factors							NF = $0.99 \pm 0.01$	NF = $1.01 \pm 0.03$		NFs Applied			
<b>1</b>	$39.79 \pm 0.19$	$162.38 \pm 1.34$	$42.78 \pm 0.32$	$46.63 \pm 0.47$	$2993.57 \pm 9.53$	$928.64 \pm 30.21$	$13820.12 \pm 128.80$	$3683.96 \pm 34.18$	$4601.97 \pm 97.54$	$362289.00 \pm 108.16$	$0.07 \pm 0.00$	359758	$0.99 \pm 0.00$
Scale factors							NF = $0.99 \pm 0.01$	NF = $1.01 \pm 0.04$		NFs Applied			
Scale factors							NF = $0.99 \pm 0.01$	NF = $1.01 \pm 0.04$		NFs Applied			
CJV < 20 GeV	$29.69 \pm 0.17$	$106.25 \pm 1.08$	$30.48 \pm 0.27$	$31.94 \pm 0.39$	$1905.77 \pm 7.80$	$612.51 \pm 25.84$	$238659.28 \pm 107.33$	$24871.15 \pm 30.44$	$2941.00 \pm 80.13$	$246771.20 \pm 139.99$	$0.06 \pm 0.00$	244811	$0.99 \pm 0.00$
Scale factors							NF = $0.99 \pm 0.01$	NF = $1.01 \pm 0.04$		NFs Applied			
OLV	$20.85 \pm 0.14$	$25.71 \pm 0.53$	$5.12 \pm 0.11$	$8.72 \pm 0.17$	$294.56 \pm 3.34$	$116.45 \pm 13.55$	$46267.04 \pm 47.47$	$499.81 \pm 12.91$	$415.70 \pm 35.23$	$47633.11 \pm 62.10$	$0.10 \pm 0.00$	47182	$0.99 \pm 0.00$
Scale factors							NF = $0.99 \pm 0.01$	NF = $1.01 \pm 0.04$		NFs Applied			
$Z \rightarrow \tau\tau$ veto	$18.05 \pm 0.13$	$22.85 \pm 0.50$	$4.22 \pm 0.10$	$1.44 \pm 0.08$	$175.96 \pm 2.68$	$69.39 \pm 12.54$	$30023.80 \pm 38.13$	$180.86 \pm 8.66$	$280.21 \pm 28.51$	$30758.74 \pm 50.03$	$0.10 \pm 0.00$	30709	$1.00 \pm 0.01$

Table 6.2: Cutflow in the top control region.

Data/MC in the top validation region shows good agreement over various variable distributions as seen in the below. Many of these distributions of used as inputs to the top+ $WW$  BDT.

### Top and $WW$ BDT discriminant (against all other samples)

This BDT is trained using  $e\mu + \mu e$  events after the VBF selection and all signal regions cuts so that the phase space in which we train the BDT is exactly the same as the one where we apply it in the final fit. The training includes top and  $WW$  trained against weighted

## CHAPTER 6. BACKGROUNDS AND SYSTEMATICS




Figure 6.3: Distributions of  $m_{\ell\ell}$ ,  $\Delta Y_{\ell\ell}$ ,  $\Delta Y_{jj}$ ,  $\sum$  centralities,  $M_{l0j0}$ ,  $M_{l1j1}$ ,  $\Delta\Phi_{jj}$ ,  $p_T^{tot}$ , and  $m_T$  in the top validation region.

samples of VBF, ggF,  $Z+jets$ , and  $V\gamma$  events. The MC statistics used in the training are half those available after all signal region cuts and the other half are later used to test the

## CHAPTER 6. BACKGROUNDS AND SYSTEMATICS

training. This corresponds to  $\approx 90,000$  un-weighted  $WW$  and top events and  $\approx 115,000$  raw VBF, ggF,  $Z+jets$  and  $V\gamma$  events. This training includes MC weights on events to best account for overall event distributions and there are  $\approx 2000$  total weighted top and  $WW$  events used in the training and  $\approx 550$  weighted other events.

The TMVA BDTG interface is used to train and test the BDT. The optimal parameters were found through a scan of reasonable values and the final set is summarized in Table 6.3.

Parameter	Value
Boosting algorithm	Gradient
Maximum tree depth	22
Number of trees	200
Minimum number of events requires per mode	5%
Number of cuts	7

Table 6.3: BDT parameters used for the top +  $WW$  vs. other backgrounds training.

This BDT utilizes a range of lepton and jet kinematic variables (8) to distinguish between signal and background events. These include  $\Delta Y_{jj}$ , the combination of lepton/jet masses  $M_{l0j0}$  for leading and subleading leptons and jets,  $\Delta\Phi_{\ell\ell}$ ,  $m_T$ ,  $\eta_{j0}$ ,  $\eta_{j1}$ ,  $\Delta\Phi_{jj}$ , and  $\sum$  centralities (L). While a larger variety of variables have been tested, these demonstrated the highest discrimination between top/ $WW$  events and other samples. Plots shown in 6.4 and 6.5 demonstrate the input distributions used to train the BDT and their correlations. The BDT training successfully separates top/ $WW$  background and other samples (ggF, VBF,  $Z+jets$ , and  $V\gamma$ ). In order to quantify the discrimination we use the integrated-ROC calculated through TMVA for weighted normalized samples and find an optimal value of 0.920. Comparisons between the test and training show that the BDT is un-biased- differences between testing and training samples would imply overtraining, or the BDT using to many parameters on too few events. For signal and background we find KS-test values of 0.107

## CHAPTER 6. BACKGROUNDS AND SYSTEMATICS




Figure 6.4: Distributions of input variables to top+ $WW$  vs. other samples BDT. Samples are weighted and normalized to even numbers of background and signal events. Signal represents top+ $WW$  and background all other samples.\*




Figure 6.5: Correlations of input variables to top + $WW$  vs other samples BDT. Signal represents top+ $WW$  and background other samples.\*

## CHAPTER 6. BACKGROUNDS AND SYSTEMATICS

and 0.154, and so no evidence of over-training. We can visualize the BDT output variable both on weighted normalized samples and on samples with full event weights applied. The following plot shows BDT results applied to all samples.




Figure 6.6: Weighted, normalized samples of top+ $WW$  samples (signal) and other samples (background) plotted over BDT output distribution on left, overlaid testing and training samples shown. Right, full weighted samples of all signal and background events plotted over BDT output distribution after signal region selection. Data is blinded here\*

We aim to fit this distribution in the signal region with high significance for top/ $WW$  events in uppermost bins of the distribution. Since this BDT is trained and applied in the signal region we cannot directly test the modelling of input variables. However, modelling at the pre-selection level for each of these variables as well as in the top validation region described earlier show no evidence of mis-modelling. The binning for this discriminant used in the statistical fit and its result (using Asimov data) are shown in the final chapter. We can visualize this BDT discriminant in the top validation region to gain an idea of overall modelling (particularly of the reasonably pure top events).

The BDT to discriminate top+ $WW$  background from VBF signal events is trained and applied in the signal region and so described in the previous chapter, but its distribution in the top validation region is also shown.

A designated BDT discriminates between top and  $WW$  backgrounds and is described

## CHAPTER 6. BACKGROUNDS AND SYSTEMATICS




Figure 6.7: Full weighted samples of all signal and background plotted over Top +  $WW$  vs. VBF  $ggF + Z+jets + V\gamma$  BDT output distributions in top validation region




Figure 6.8: Full weighted samples of all signal and background plotted over Top +  $WW$  vs. VBF BDT output distributions in top validation region

## CHAPTER 6. BACKGROUNDS AND SYSTEMATICS

further in the  $WW$  background section (next). However, this distribution in the top validation region with all weighted samples is shown below. to demonstrate its effect on primarily predominantly top-like events.




Figure 6.9: Full weighted samples of all signal and background plotted over Top vs.  $WW$  BDT output distributions in top validation region

Normalization factors (NF) are derived in the top validation region to correct for data/MC mis-modelling. These factors are applied to the top sample in the signal region. The NF factors are  $0.99 \pm 0.01$ .

### 6.1.3 Diboson background

The  $WW$  background consists primarily of QCD  $WW$ +jets events (highly dominating electroweak vertices). This background is estimated along with the top background using a joint parameter due to their similarities in signature as well as the difficulty in defining a pure  $WW$  validation region without top contamination. A  $WW$  validation region is defined to

## CHAPTER 6. BACKGROUNDS AND SYSTEMATICS

demonstrate  $WW$  MC modelling in a targeted  $WW$  region. The  $WW$  validation region is defined with at least 2 jets, a  $b$ -veto ( $N_{b-jet} < 1$ ) and a central-jet-veto of below 20GeV as in the signal region. Two additional cuts differ from the signal region:  $m_T > 130\text{GeV}$  and  $m_{T2} > 160\text{GeV}$  where  $m_{T2}$  is defined as

$$m_{T2} = \min_{p_T^1 + p_T^2 = p_T} (\max(m_T^2(p_T^1, p_T^a), m_T^2(p_T^2, p_T^b))) \quad (6.1)$$

This represents a lower bound on the parent particle's mass, so using a large  $m_{T2}$  cut eliminates contamination from many  $t\bar{t}$  decays which have an upper limit near the top mass. The purity of the region is  $\approx 37\%$  and the cutflow for this region is shown below.

$\sqrt{s} = 13\text{TeV}, \mathcal{L} = 139\text{fb}^{-1}$ (Full Run 2)	$H_{VBF}$	$H_{ggF}$	$H_{VH}$	$H \rightarrow \tau\tau$	$WW$	Other VV	Top	Zjets	Mis-Id	Total Bkg	Significance	Data	Data/MC
Scale factors							$\text{NF} = 0.99 \pm 0.01$	$\text{NF} = 1.01 \pm 0.01$		116387.21 $\pm$ 168.10	0.95 $\pm$ 0.00	109677	0.94 $\pm$ 0.00
b-veto	323.67 $\pm$ 0.54	1109.94 $\pm$ 3.42	150.32 $\pm$ 0.59	287.53 $\pm$ 1.07	21075.69 $\pm$ 28.85	4031.02 $\pm$ 92.67	63787.91 $\pm$ 57.22	22151.24 $\pm$ 102.73	3793.55 $\pm$ 70.69	<b>NFs Applied</b>			
Scale factors							$\text{NF} = 0.99 \pm 0.01$	$\text{NF} = 1.01 \pm 0.01$		<b>NFs Applied</b>			
$M_T > 130\text{ GeV}$	32.37 $\pm$ 0.17	155.17 $\pm$ 1.28	39.90 $\pm$ 0.27	20.08 $\pm$ 0.35	16127.61 $\pm$ 24.93	1893.54 $\pm$ 44.79	50088.89 $\pm$ 50.83	872.43 $\pm$ 33.66	1784.17 $\pm$ 49.41	70981.25 $\pm$ 93.74	0.12 $\pm$ 0.00	68255	0.96 $\pm$ 0.00
Scale factors							$\text{NF} = 0.99 \pm 0.01$	$\text{NF} = 1.01 \pm 0.01$		<b>NFs Applied</b>			
$M_{T2} > 160\text{ GeV}$	18.87 $\pm$ 0.13	62.44 $\pm$ 0.82	10.50 $\pm$ 0.15	6.79 $\pm$ 0.19	6528.07 $\pm$ 14.33	743.72 $\pm$ 32.98	11553.13 $\pm$ 24.82	302.42 $\pm$ 14.17	515.71 $\pm$ 26.44	19722.79 $\pm$ 53.01	0.13 $\pm$ 0.00	18672	0.95 $\pm$ 0.01
Scale factors							$\text{NF} = 0.99 \pm 0.01$	$\text{NF} = 1.01 \pm 0.04$		<b>NFs Applied</b>			
CJV (20GeV)	13.61 $\pm$ 0.11	40.50 $\pm$ 0.66	7.02 $\pm$ 0.12	4.28 $\pm$ 0.16	4363.49 $\pm$ 11.96	497.52 $\pm$ 30.27	6415.93 $\pm$ 19.00	184.81 $\pm$ 12.88	327.54 $\pm$ 20.73	11841.08 $\pm$ 44.91	0.13 $\pm$ 0.00	11245	0.95 $\pm$ 0.01

Table 6.4: Cutflow in the  $WW$  validation region.

Data/MC shows good agreement over various variable distributions as seen below.

### Top vs. $WW$ BDT discriminant

This BDT is trained using  $e\mu + \mu e$  events after the VBF selection and all signal regions cuts so that the phase space in which we train the BDT is exactly the same as the one where we apply it in the final fit. The training includes weighted samples of top and  $WW$  events trained against one another. The MC statistics used in the training are half those available after all signal region cuts where the other half are later used to test the training. This corresponds to  $\approx 35,000$  un-weighted top and  $\approx 55,000$  raw  $WW$  events. This training includes MC weights on events to best account for overall event distributions and there are  $\approx 1500$  total weighted top used in the training and  $\approx 500$  weighted  $WW$  events.

## CHAPTER 6. BACKGROUNDS AND SYSTEMATICS




Figure 6.10: Distributions of  $\Delta Y_{\ell\ell}$ ,  $\Delta Y_{jj}$ ,  $\sum$  centralities,  $M_{l0j0}$ ,  $M_{l1j1}$ ,  $p_T^{j0}$ ,  $\Delta\Phi_{jj}$ ,  $p_T^T$ , and  $m_T$  in the  $WW$  validation region.

The TMVA BDTG interface is used to train and test the BDT. The optimal parameters were found through a scan of reasonable values and the final set is summarized in Table 6.5.

## CHAPTER 6. BACKGROUNDS AND SYSTEMATICS

Parameter	Value
Boosting algorithm	Gradient
Maximum tree depth	22
Number of trees	200
Minimum number of events requires per mode	5%
Number of cuts	7

Table 6.5: BDT parameters used for the top vs.  $WW$  training.

This BDT utilizes a collection of lepton and jet kinematic variables (14) to distinguish between top and  $WW$  events. These include  $\Delta Y_{jj}$ ,  $\Delta Y_{\ell\ell}$ ,  $\Delta\Phi_{\ell\ell}$ ,  $m_T$ ,  $\eta_{l0}$ ,  $\eta_{l1}$ ,  $p_{j0}^T$ ,  $p_{j1}^T$ ,  $\Delta\Phi_{jj}$ ,  $\sum$  centralities (L), and combined masses  $M_{l0j0}$ ,  $M_{l0j1}$ ,  $M_{l1j0}$ ,  $M_{l1j1}$ . While a larger variety of variables have been tested, these demonstrated the highest discrimination between top and  $WW$  events. Plots shown in 6.11 and 6.12 demonstrate the input distributions used to train the BDT and their correlations. The BDT training separates top and  $WW$  backgrounds




Figure 6.11: Distributions of input variables to top vs.  $WW$  BDT. Samples are weighted and normalized to even numbers of background and signal events. Signal represents top and background  $WW$ .\*

somewhat. These signals have similar kinematic distributions that make high discrimination difficult. In order to quantify the discrimination we use the integrated-ROC calculated through TMVA for weighted normalized samples and find an optimal value of 0.653. While relatively low, this shows some minimal amount of discrimination. This is useful in the fit to

## CHAPTER 6. BACKGROUNDS AND SYSTEMATICS




Figure 6.12: Correlations of input variables to top vs.  $WW$  BDT. Signal represents top and background  $WW^*$

differentiate top and  $WW$  backgrounds, though high levels of separation are not necessary because top and  $WW$  are estimated together with a shared parameter in the simultaneous fit. Comparisons between the test and training show that the BDT is un-biased. For signal and background we find KS-test values of 0.067 and 0.037, and so no evidence of over-training. We can visualize the BDT output variable both on weighted normalized samples and on samples with all event weights applied. The following plot shows BDT results applied to all samples.

We aim to fit this distribution in the signal region with high significance for top events in uppermost bins of the distribution (and  $WW$  in the lowermost). Since this BDT is trained and applied in the signal region we cannot directly test the modelling of input variables. However, modelling at the pre-selection level for each of these variables as well as in the  $WW$  validation region described earlier show no evidence of mis-modelling. The binning for this discriminant used in the statistical fit and its result (using Asimov data) are shown in the final chapter. We can visualize this BDT discriminant in the  $WW$  validation region to gain an idea of overall modelling particularly of the  $WW$  events.

The BDT to discriminate top+ $WW$  background from VBF, ggF,  $Z+jets$  and  $V\gamma$  events

## CHAPTER 6. BACKGROUNDS AND SYSTEMATICS




Figure 6.13: Weighted, normalized samples of top samples (signal) and WW samples (background) plotted over BDT output distribution on left, overlaid testing and training samples shown. Right, full weighted samples of all signal and background events plotted over the BDT output distribution after signal region selection. Data is blinded here\*




Figure 6.14: Full weighted samples of all signal and background plotted over BDT output distributions in WW validation region

## CHAPTER 6. BACKGROUNDS AND SYSTEMATICS

is trained and applied in the signal region and described in the previously, but its distribution in the  $WW$  validation region is also shown.




Figure 6.15: Full weighted samples of all signal and background plotted over  $Top + WW$  vs.  $VBF\ ggF + Z+jets + V\gamma$  BDT output distributions in  $WW$  validation region

The BDT to discriminate  $WW$  background from VBF signal events is trained and applied in the signal region and so described in the previous chapter, but its distribution in the  $WW$  validation region is shown below. This BDT discriminates both  $WW$  and top (combined) against the VBF signal as they have similar kinematic distributions and are treated together in the final simultaneous fit.

### 6.1.4 ggF background

Other Higgs production modes are considered backgrounds in this analysis- and while vector mediated Higgs production ( $VH$ ) and top produced Higgs ( $tth$ ) are small in our signal region so don't play a large role in the analysis - gluon fusion Higgs production (ggF) represents a significant background. Signal region cuts requiring at least 2-jets and the central jet veto significantly reduce background yields, but because it is kinematically very

## CHAPTER 6. BACKGROUNDS AND SYSTEMATICS




Figure 6.16: Full weighted samples of all signal and background plotted over Top +  $WW$  vs. VBF BDT output distributions in  $WW$  validation region

similar to VBF signal, it has a large effect on the VBF differential coupling measurement. To mitigate these effects, ggF events are simultaneously estimated from three control regions and the signal region. The control regions are chosen to minimize both the statistical and the modelling uncertainties, in particular those originating from the modelling of higher order QCD corrections. The regions begin with a subdivision based on the jet multiplicity. The first two regions, ggF CR1 and ggF CR2, are built from all preselection cuts and cuts requiring at least two jets, a  $b$ -veto,  $Z \rightarrow \tau\tau$ -veto, and orthogonal cuts on the central jet veto (CJV) and opposite lepton veto (OLV). The third ggF control region, ggF CR3, uses all preselection cuts, a  $b$  veto,  $Z \rightarrow \tau\tau$  veto, and requirement for less than 2 jets. The full definitions are described:

- **GGF-CR1** Preselection criteria and  $N_{\text{jet}} \geq 2$ ,  $b$  veto,  $Z \rightarrow \tau\tau$  veto, CJV  $< 1$  and OLV  $> 1$  or CJV  $> 1$  and OLV  $< 1$ .
- **GGF-CR2** Preselection criteria and  $N_{\text{jet}} \geq 2$ ,  $b$  veto,  $Z \rightarrow \tau\tau$  veto, CJV  $> 1$  and OLV  $> 1$ .

## CHAPTER 6. BACKGROUNDS AND SYSTEMATICS

- **GGF-CR3** Preselection criteria and  $N_{\text{jet}} \geq 2$ ,  $b$  veto, and  $Z \rightarrow \tau\tau$  veto.

The full ggF expectation in the signal region is built from the following ratio (inspired by the ABCD method)

$$\mu_{\text{GGF-SR}} = \frac{\mu_3^{\text{GGF-CR}} \cdot \mu_1^{\text{GGF-CR}}}{\mu_2^{\text{GGF-CR}}}. \quad (6.2)$$

where  $\mu_{\text{ggF-X}}$  represents the yield modifier in each ggF control region. The final event yield in the signal region  $\mu_{\text{ggF-SR}}$  is determined in the simultaneous fit as described in the next chapter. In each ggF category the ggF yield is extracted from simulation-extracted template fits based on dedicated discriminants. For each control region a dedicated multivariate discriminant is trained and applied to discriminate between ggF and all the backgrounds combined. Results and training parameters from each of these discriminants is summarized next.

### Discriminant in ggF CR1

The multivariate discriminant used for the ggF CR1 is a boosted decision tree (BDT) trained using  $e\mu + \mu e$  events that pass all ggF CR1 cuts. The training includes ggF events trained against VBF signal and all backgrounds. The MC statistics used in the training are half those available after the ggF CR1 cuts (as the other half are later used to test the training). This corresponds to  $\approx 8,000$  ggF events and  $\approx 250,000$  other signal and background events. Events are trained with MC weights applied to accurately represent relative importance of each event and the training uses  $\approx 130$  weighted ggF samples and  $\approx 10,000$  weighted other background samples.

The TMVA BDTG interface is used to train and test the BDT. The optimal parameters were found through a scan of reasonable values and the final set is summarized in Table 6.6. For this BDT various distributions (8) are used to take advantage of differences

## CHAPTER 6. BACKGROUNDS AND SYSTEMATICS

Parameter	Value
Boosting algorithm	Gradient
Maximum tree depth	10
Number of trees	600
Minimum number of events requires per mode	5%
Number of cuts	7

Table 6.6: BDT parameters used for the ggF CR1 training.

in distributions between ggF events and other sample types. These variables include  $\Delta Y_{\ell\ell}$ ,  $\Delta\Phi_{\ell\ell}, m_{\ell\ell}, m_T$ , jet  $p_{\text{lead}}^T$ , jet  $p_{\text{sublead}}^T$ ,  $\Delta\Phi_{jj}$ , and  $E_T^{\text{miss}}$ . Distributions for these variables in the ggF CR1 region where the BDT is trained are shown below demonstrating data/MC modelling for each.

Plots shown in [6.18](#) and [6.19](#) demonstrate the input distributions used to train the BDT and their correlations where each distribution is normalized to equal number of background and signal events.

In order to quantify the discrimination we use the integrated-ROC calculated through TMVA for weighted normalized samples and find an optimal value of 0.898. Comparisons between the test and training show that the BDT is un-biased- differences between testing and training samples would imply overtraining, or the BDT using too many parameters on too few events. Visually, we can see that the testing and trainings samples are quite similar. Additionally, a Kolmogorov-Smirnov test is performed to measure if the two test and training distributions differ significantly and no evidence of over-training is present (values 0.191, 0.049 for signal, background). We can visualize the BDT output variable both on weighted normalized samples and on samples without normalization. The following plots show BDT results applied to normalized and full weighted samples.

## CHAPTER 6. BACKGROUNDS AND SYSTEMATICS




Figure 6.17: Distributions of  $\Delta Y_{\ell\ell}$ ,  $\Delta\Phi_{\ell\ell}$ ,  $m_{\ell\ell}$ ,  $m_T$ , jet  $p_T^T$ , jet  $p_{\text{sublead}}^T$ ,  $\Delta\Phi_{jj}$ , and  $E_T^{\text{miss}}$  in the ggF CR1 used as input to the BDT discriminating ggF from all other samples.

### Discriminant in ggF CR2

The multivariate discriminant used for ggF CR2 is a boosted decision tree trained using  $e\mu + \mu e$  events that pass all ggF CR2 cuts. The training includes ggF events trained against

## CHAPTER 6. BACKGROUNDS AND SYSTEMATICS




Figure 6.18: Distributions of input variables to ggF CR1 BDT. Samples are weighted and normalized to even numbers of background and signal events. Signal represents ggF and background all other samples.\*




Figure 6.19: Correlations of input variables to ggF CR1 BDT. Signal represents ggF and background all other samples.\*

## CHAPTER 6. BACKGROUNDS AND SYSTEMATICS




Figure 6.20: Normalized samples of ggF (signal) and all other samples (background) plotted over BDT output distribution on left, overlaid testing and training samples shown. Right, full weighted samples of ggF signal and all other backgrounds plotted over BDT output distribution after ggF CR 1.\*

VBF signal and all backgrounds. The MC statistics used in the training are half those available after ggF CR2 cuts (as the other half are later used to test the training). This corresponds to  $\approx 22,200$  ggF events and  $\approx 90,000$  other signal and background events.

The TMVA BDTG interface is used to train and test the BDT. The optimal parameters were found through a scan of reasonable values and the final set is summarized in Table 6.7. For this BDT various distributions (6) are used to take advantage of differences

Parameter	Value
Boosting algorithm	Gradient
Maximum tree depth	10
Number of trees	50
Minimum number of events requires per mode	5%
Number of cuts	7

Table 6.7: BDT parameters used for the ggF CR2 training.

in distributions between ggF events and other sample types. These variables include  $\Delta Y_{\ell\ell}$ ,

## CHAPTER 6. BACKGROUNDS AND SYSTEMATICS

$\Delta\Phi_{\ell\ell, m_{\ell\ell}, m_T}$ ,  $\Delta\Phi_{jj}$ , and  $E_T^{\text{miss}}$ . Distributions for these variables in the ggF CR2 region where the BDT is trained are shown below demonstrating data/MC modelling for each.




Figure 6.21: Distributions of  $\Delta Y_{\ell\ell}$ ,  $\Delta\Phi_{\ell\ell}$ ,  $m_{\ell\ell}$ ,  $m_T$ ,  $\Delta\Phi_{jj}$ , and  $E_T^{\text{miss}}$  in the ggF CR2 used as input to the BDT discriminating ggF from all other samples.

Plots shown in 6.22 and 6.23 demonstrate the input distributions used to train the BDT and their correlations where each distribution is weighted and normalized to equal number of background and signal events.

In order to quantify the discrimination we use the integrated-ROC calculated through TMVA for weighted normalized samples and find an optimal value of 0.902. Comparisons between the test and training show that the BDT is un-biased- visually, once can see that the testing and trainings samples are quite similar. Additionally, a Kolmogorov-Smirnov test is

## CHAPTER 6. BACKGROUNDS AND SYSTEMATICS




Figure 6.22: Distributions of input variables to ggF CR2 BDT. Samples are weighted and normalized to even numbers of background and signal events. Signal represents ggF and background all other samples.\*




Figure 6.23: Correlations of input variables to ggF CR2 BDT. Signal represents ggF and background all other samples.\*

## CHAPTER 6. BACKGROUNDS AND SYSTEMATICS

performed to measure if the two test and training distributions differ significantly and with values of 0.012 (0.303) for signal (background) no evidence of over-training is present. We can visualize the BDT output variable both on weighted normalized samples and on samples with full event weights applied. The following plots show BDT results applied to normalized and full weighted samples.




Figure 6.24: Weighted, normalized samples of ggF (signal) and all other samples (background) plotted over BDT output distribution on left, overlaid testing and training samples shown. Right, full weighted samples of ggF signal and all other backgrounds plotted over BDT output distribution after ggF CR2 cuts.\*

### Discriminant in ggF-CR3

The multivariate discriminant used for the ggF CR3 is a boosted decision tree trained using  $e\mu + \mu e$  events that pass all ggF CR3 cuts. The training includes only ggF events trained against VBF signal and all backgrounds. The MC statistics used in the training are half those available after the ggF CR3 cuts (as the other half are later used to test the training). This corresponds to  $\approx 200,000$  ggF events and  $\approx 2,800,000$  other signal and

## CHAPTER 6. BACKGROUNDS AND SYSTEMATICS

background events.

The TMVA BDTG interface is used to train and test the BDT. The optimal parameters were found through a scan of reasonable values and the final set is summarized in Table 6.8. For this BDT various distributions (4) are used to take advantage of differences

Parameter	Value
Boosting algorithm	Gradient
Maximum tree depth	10
Number of trees	1000
Minimum number of events requires per mode	5%
Number of cuts	7

Table 6.8: BDT parameters used for the ggF CR3 training.

in distributions between ggF events and other samples types. These variables include  $\Delta Y_{\ell\ell}$ ,  $\Delta\Phi_{\ell\ell}, m_{\ell\ell}$ , and  $m_T$ . Distributions for these variables in the ggF CR3 region where the BDT is trained are shown below demonstrating data/MC modelling for each.

Plots shown in 6.26 and 6.27 demonstrate the input distributions used to train the BDT and their correlations where each distribution is weighted and normalized to equal number of background and signal events.

In order to quantify the discrimination we use the integrated-ROC calculated through TMVA for weighted normalized samples and find an optimal value of 0.888. Comparisons between the test and training show that the BDT is un-biased, we can see that the testing and training samples are quite similar. Additionally, a Kolmogorov-Smirnov test is performed and shows no sign of bias with values of 0.276 (0.673) for signal (background). We can visualize the BDT output variable both on weighted normalized samples and on samples with full event weights applied. The following plots show BDT results applied to normalized and fully weighted samples.

## CHAPTER 6. BACKGROUNDS AND SYSTEMATICS




Figure 6.25: Distributions of  $\Delta Y_{\ell\ell}$ ,  $\Delta\phi_{\ell\ell}$ ,  $m_{\ell\ell}$ , and  $m_T$  in the ggF CR3 used as input to the BDT discriminating ggF from all other samples.

### Discriminant for ggF and VBF in the signal region

There is one final multivariate discriminant used to discriminate ggF backgrounds. This focuses on discriminating ggF from VBF events in the signal region. The boosted decision tree uses  $e\mu + \mu e$  events that pass all signal region cuts. The training includes only ggF events trained against VBF signal. The MC statistics used in the training are half those available after the SR cuts (as the other half are later used to test the training). This corresponds to  $\approx 6,000$  ggF events and  $\approx 95,000$  signal events.

The TMVA BDTG interface is used to train and test the BDT. The optimal parameters

## CHAPTER 6. BACKGROUNDS AND SYSTEMATICS




Figure 6.26: Distributions of input variables to the ggF CR3 BDT. Samples are weighted and normalized to even numbers of background and signal events. Signal represents ggF and background all other samples.\*




Figure 6.27: Correlations of input variables to the ggF CR3 BDT. Signal represents ggF and background all other samples.\*

## CHAPTER 6. BACKGROUNDS AND SYSTEMATICS




Figure 6.28: Normalized samples of ggF (signal) and all other samples (background) plotted over BDT output distribution on left, overlaid testing and training samples shown. Right, full weighted samples of ggF signal and all other backgrounds plotted over BDT output distribution after ggF CR3 cuts.\*

were found through a scan of reasonable values and the final set is summarized in Table 6.9.

For this BDT various distributions (7) are used to take advantage of differences in distribu-

Parameter	Value
Boosting algorithm	Gradient
Maximum tree depth	10
Number of trees	300
Minimum number of events requires per mode	5%
Number of cuts	7

Table 6.9: BDT parameters used for the ggF vs. VBF training.

tions between ggF and VBF events. These variables include  $\Delta Y_{jj}$ ,  $\Delta \Phi_{jj}$ ,  $m_T$ ,  $p_T^{j0}$ ,  $p_T^{j1}$ ,  $\sum M_{lj}$ .

Plots shown in 6.29 and 6.30 demonstrate the input distributions used to train the BDT and their correlations where each distribution is weighted and normalized to equal number of background and signal events.

## CHAPTER 6. BACKGROUNDS AND SYSTEMATICS




Figure 6.29: Distributions of input variables to the ggF vs. VBF BDT. Samples are weighted and normalized to even numbers of background and signal events. Signal represents VBF and background ggF samples.




Figure 6.30: Correlations of input variables to the ggF vs. VBF BDT. Signal represents VBF and background ggF samples.

## CHAPTER 6. BACKGROUNDS AND SYSTEMATICS

In order to quantify the discrimination we use the integrated-ROC calculated through TMVA for weighted normalized samples and find an optimal value of 0.785. Comparisons between the test and training show that the BDT is un-biased, since the testing and training samples are similar. Additionally, a Kolmogorov-Smirnov test is performed and shows no sign of bias with values of 0.008 (0.047) for signal (background). We can visualize the BDT output variable both on weighted normalized samples and on samples with full event weights applied. The following plots show BDT results applied to normalized and fully weighted samples.




Figure 6.31: Normalized samples of VBF (signal) and ggF (background) plotted over BDT output distribution on left, overlaid testing and training samples shown. Right, full weighted samples of VBF signal and ggF background plotted over BDT output distribution after SR cuts.

### 6.1.5 Fake backgrounds

The final substantial background in the VBF HWW differential coupling analysis comes from mis-identified leptons. These are jets that are mistakenly identified as leptons in reconstruction and in this analysis these predominantly come from  $W$ +jet events. In these events a  $W$  boson decays leptonically leading to one true lepton and one mistaken lepton

## CHAPTER 6. BACKGROUNDS AND SYSTEMATICS

(a jet from the primary vertex is mistaken for a lepton). This event mimics our desired two lepton signature and so creates an additional HWW background sample. This background is estimated as in the HWW coupling analysis using the fake factor method, similarly to in the 2016 HWW coupling analysis. Further detail on the overall fake factor method and its mathematical formulation can be found here [54] while further details on HWW-specific fake background studies are here [? ]. In this section I will give a brief summary of fake estimations along with the slightly augmented fake factor control region definition used in the HWW differential coupling analysis.

The fake background is estimated using data which is measured in a region defined by signal region cuts with one important distinction- one or both of the leptons used are “anti-identified” meaning they pass some looser lepton identification criteria but not the one used in the analysis signal region. These fakes are then extrapolated to the true signal region (with two “identified” leptons) using fake factors. Fake factors are measured as functions of  $p_T$  and  $\eta$  in jet-enriched  $Z+jets$  samples and are defined as the ratio of identified to anti-identified leptons. The fake backgrounds can be considered split between “single-fake” (with one “anti-ID” lepton) from the predominant  $W+jets$  background and “double-fake” (with two “anti-ID” leptons) from QCD processes. The total signal sample can be defined as:

$$N_{id+id} = N_{id+id}^{EW} + N_{id+id}^{W+jets} + N_{id+id}^{QCD} \quad (6.3)$$

so that the total events include all electroweak processes ( $N_{id+id}^{EW}$ ) as well as fake backgrounds from  $W+jets$  and QCD events. In order to estimate the total fake background in the signal region we need to estimate the number of  $W+jets$  and QCD events in “id+anti-id” events and then apply the fake factor to extrapolate into the “id+id” region. The  $N_{id+anti-id}$  for fake backgrounds is calculated after subtraction of electroweak backgrounds (two true leptons

## CHAPTER 6. BACKGROUNDS AND SYSTEMATICS

Table 6.10: Requirements for “identified” and “anti-identified” electrons and muons.

“id” electron	“anti-id” electron	“id” muon	“anti-id” muon
$p_T > 15 \text{ GeV}$ $ \eta  < 2.47$ , excluding $1.37 <  \eta  < 1.52$ $ z_0 \sin \theta  < 0.5 \text{ mm}$ Pass <i>Tight</i> if $p_T < 25 \text{ GeV}$ Pass <i>Medium</i> if $p_T > 25 \text{ GeV}$ $ d_0  \sigma(d_0) < 5$ Pass FixedCutTrackCone40 if $p_T < 25 \text{ GeV}$ Pass IsoGradient if $p_T < 25 \text{ GeV}$	Pass <i>Loose</i> Veto against identified electron	$p_T > 15 \text{ GeV}$ $ \eta  < 2.45$ $ z_0 \sin \theta  < 0.5 \text{ mm}$ Pass <i>Tight</i> $ d_0  \sigma(d_0) < 3$ $E_T^{\text{cone}20}/p_T < 0.09$ $p_T^{\text{varcone}30}/p_T < 0.06$	Pass <i>Medium</i> $ d_0  \sigma(d_0) < 15$ Veto against identified muon

that contaminate the “id+anti-id region”) from Monte Carlo simulations as follows:

$$N_{\text{id+anti-id}}^{W+\text{jets}} + N_{\text{id+anti-id}}^{\text{QCD}} = N_{\text{id+anti-id}} - N_{\text{id+anti-id}}^{\text{EWMC}} \quad (6.4)$$

Fake factors are derived from  $Z+\text{jets}$  samples and then applied to  $W+\text{jets}$  regions so the differences between these two samples are important to understand fully. The fake factor is defined

$$F.F. = \frac{N_{\text{id}}}{N_{\text{anti-id}}} \quad (6.5)$$

and is measured separately for electron and muons and measured in bins of  $\eta$  and  $p_T$ . The following table summarizes to “id” and “anti-id” requirements.

$Z+\text{jets}$  event are selected to contain exactly three loosely identified leptons with  $p_T > 15 \text{ GeV}$ . Further requirements are for an opposite sign  $ee$  or  $\mu\mu$  lepton pair with  $7 \text{ GeV} < m_{\ell\ell} < 110 \text{ GeV}$ . Both  $Z$  candidate leptons must be “identified” so that the third is the fake candidate. An additional  $WZ$  veto is applied using  $m_T > 50 \text{ GeV}$  to mitigate electroweak background in the  $Z+\text{jets}$  sample. Dedicated MC simulations model electroweak backgrounds in the  $Z+\text{jets}$  region and include  $V + \gamma$ , diboson ( $WW$ ,  $WZ$ , and  $ZZ$ ), single top and  $t\bar{t}$ .  $WZ$  backgrounds are normalized to their measured cross-sections. Fake yields for the  $Z+\text{jets}$  samples are finally calculated by subtracting data from all MC electroweak backgrounds.

## CHAPTER 6. BACKGROUNDS AND SYSTEMATICS

Table 6.11: Fake factors binned for muons and electrons in  $p_T$  and  $\eta$  with their statistical uncertainties

$p_T(\text{GeV})$	Muon FF ( $ \eta  < 1.5$ )	Muon FF ( $1.5 <  \eta  < 2.5$ )	Electron FF
15-25	$0.038 \pm 0.004$	$0.057 \pm 0.005$	$0.079 \pm 0.005$
20-25	$0.021 \pm 0.007$	$0.0357 \pm 0.0077$	$0.097 \pm 0.001$
25-35	$0.029 \pm 0.011$	$0.064 \pm 0.014$	$0.157 \pm 0.017$
35-inf	$0.049 \pm 0.023$	$0.11 \pm 0.04$	$0.19 \pm 0.026$

The fake factor is computed as a binned ratio in  $p_T$ , [15, 20, 25, 35, inf] and for muons also in  $\eta$  for [0, 1.5, 2.5] excluding the electromagnetic calorimeter crack region. The following table summarized fake factors used in this analysis.

Further studies on fake factors include comparisons to their MC simulation expectations and studies on differences between the  $Z+\text{jets}$  control sample used to calculate fake factors and the  $W+\text{jets}$  sample on which they are applied. These studies are beyond the scope of this thesis but support the use of the fake factor method in our analysis. Specifically, jet kinematics and heavy flavor fraction in both samples have been studied by the HWW coupling analysis and factored into a correction factor applied to transfer between  $Z+\text{jets}$  and  $W+\text{jets}$  samples. More information on these studies and correction factors can be found in [? ]. Comparisons between data-driven fake yields in the signal region and predicted fake events from various Monte Carlo generators (Powheg, MadGraph) are included in Appendix B.

The  $W+\text{jets}$  control region is defined as in the signal region, with cuts specifying at least 2 jets, a  $b$ -veto,  $m_{\tau\tau}$  based Z-veto, an opposite lepton veto, central jet veto, and cuts on  $m_{jj}$  and  $\Delta Y_{jj}$ . Unlike the signal region, this region also specified one “id” and one “anti-id” lepton instead of two identified leptons. The cutflow for this region is shown below where fakes are defined from the subtraction of total electroweak background from data. Once fake factors are applied, these events constitute the fake background estimate in the signal region.

## CHAPTER 6. BACKGROUNDS AND SYSTEMATICS

Fake purity in this control region is 74% after all signal region-like cuts are applied.

$\sqrt{s} = 13\text{TeV}$ , $\mathcal{L} = 139\text{fb}^{-1}$ (Full Run 2)	Total Bkg	data	fakes	fake purity(%)
Channel Selection	$8116768.84 \pm 3042.09$	13124937	$5008168.16 \pm 4730.67$	$38.16 \pm 0.04$
Trigger Selection	$8116768.84 \pm 3042.09$	13124937	$5008168.16 \pm 4730.67$	$38.16 \pm 0.04$
Trigger Matching	$7906126.65 \pm 2917.00$	13057932	$5151805.35 \pm 4644.01$	$39.45 \pm 0.04$
W+jets flavour split muon	$5896841.78 \pm 2703.75$	10572340	$4675498.22 \pm 4228.78$	$44.22 \pm 0.04$
W+jets flavour split electron	$3791849.84 \pm 2494.76$	7417426	$3625576.16 \pm 3693.41$	$48.88 \pm 0.05$
Jet Cleaning	$3791849.84 \pm 2494.76$	7417426	$3625576.16 \pm 3693.41$	$48.88 \pm 0.05$
Overlap: Vgamma/Vjets	$3791849.84 \pm 2494.76$	7417426	$3625576.16 \pm 3693.41$	$48.88 \pm 0.05$
Only two Leptons	$3754720.02 \pm 2485.95$	7376514	$3621793.98 \pm 3681.91$	$49.10 \pm 0.05$
$p_t^{\text{lead}} > 22\text{ GeV}$	$3754720.02 \pm 2485.95$	7376514	$3621793.98 \pm 3681.91$	$49.10 \pm 0.05$
$p_t^{\text{sublead}} > 15$	$3753322.61 \pm 2485.49$	7373346	$3620023.39 \pm 3681.17$	$49.10 \pm 0.05$
OS Leptons	$3751223.46 \pm 2483.45$	7365693	$3614469.54 \pm 3678.75$	$49.07 \pm 0.05$
$M_{\ell\ell} > 12/10\text{ GeV}$	$3751147.28 \pm 2483.41$	7365336	$3614188.72 \pm 3678.68$	$49.07 \pm 0.05$
SF: Z Veto	$3751147.28 \pm 2483.41$	7365336	$3614188.72 \pm 3678.68$	$49.07 \pm 0.05$
Leptons ID, singleFakes 1 anti-ID,1 ID, doubleFakes 2*anti-ID	$780838.94 \pm 837.46$	1931616	$1150777.06 \pm 1622.64$	$59.58 \pm 0.09$
2-jet (30,30) fJVT	$384692.76 \pm 242.92$	657536	$272843.24 \pm 846.49$	$41.49 \pm 0.14$
b-veto	$62361.10 \pm 199.23$	198158	$135796.90 \pm 487.70$	$68.53 \pm 0.29$
CJV (20GeV)	$43093.51 \pm 171.93$	137885	$94791.49 \pm 409.20$	$68.75 \pm 0.35$
OLV bool	$9162.00 \pm 78.28$	28068	$18906.00 \pm 184.92$	$67.36 \pm 0.77$
$Z \rightarrow \tau\tau$ veto	$5142.69 \pm 60.99$	18069	$12926.31 \pm 147.61$	$71.54 \pm 0.97$
$M_{jj} > 200$	$3233.48 \pm 51.55$	11911	$8677.52 \pm 120.70$	$72.85 \pm 1.21$
$DY_{jj} > 2.1$	$2838.02 \pm 50.56$	10904	$8065.98 \pm 116.02$	$73.97 \pm 1.28$

Table 6.12: Cutflow in the fakes control region.

Distributions below show the electroweak background processes (blue) in the fake control region along with data (black). The  $W+\text{jets}$  or fake background is taken as the difference between data and EW background and shown in the plots as light blue points. High fake purity is shown and these backgrounds are used along with fake factors to estimate fakes contaminating our signal region.

The EW background subtracted from data in the  $W+\text{jets}$  control region consists of  $V\gamma$ , diboson, top, and  $Z+\text{jets}$  events.

## 6.2 Systematic uncertainties

The experimental, theoretical, and statistical uncertainties in this analysis all impact the final measurement substantially. In this section I describe first the experimental systematic uncertainties then the theoretical in detail. Statistical uncertainties are derived from the limited number of MC events used to model each of signal and background samples per bin

## CHAPTER 6. BACKGROUNDS AND SYSTEMATICS




Figure 6.32: Distributions of  $p_T^{\ell\ell}$ ,  $m_{\ell\ell}$ ,  $m_T$ ,  $p_{tot}^T$ , lep  $p_{lead}^T$ , lep  $p_{sublead}^T$ ,  $m_{jj}$ , and  $E_T^{\text{miss}}$  in the differential VBF  $W+jets$  control region.

in the differential measurement. A single MC-stat nuisance parameter is implemented based on Poisson statistics and used in our overall statistical fit.

### 6.2.1 Experimental uncertainties

Several types of experimental systematics uncertainties affect this analysis. Each of these are calculated for this analysis using the current recommendations for each of their respective

## CHAPTER 6. BACKGROUNDS AND SYSTEMATICS

Combined Performance groups and can be split into those derived from muons, electrons, Jet/ $E_T^{miss}$ , Trigger, flavor tagging and pile-up. The current set of experimental systematics uses 102 nuisance parameters, each of are briefly described with their associated names in the ???. The impact of these systematics on our overall result as well as rankings and interpretation of the most impactful systematics is in the next chapter.

There are two different ways systematic uncertainties are calculated and applied. First, for systematic uncertainties on the energy of jets and electrons and momentum of muons, the scale and resolutions is calculated. These test the number of events in the final state affected by shifting energy/momentum by a scale factor. This “smearing” procedure is done for a nominal scaling value and values with  $\pm 1\sigma$ . Next, systematic uncertainty on the scale factors is calculated by comparing nominal event numbers to those after particular weights are applied. These weights are derived from data-MC agreement and their uncertainty is the relative difference between nominal and modified weight sums.

There are four nuisance parameters associated with muon reconstruction and identification, these are the reconstruction efficiency statistical and systematic uncertainties along with additional separate uncertainties for low  $p_T$  muons, or those with  $3 < p_T < 20$  GeV. Muon momentum resolution uncertainties are divided into those from the Inner Detector and those from the muon spectrometer. Muon momentum scale uncertainties are contained in one parameter and two additional systematic uncertainties come from a correction applied to muons to account for residual ID/MS misalignments which create a charge dependent bias. Two nuisance parameters account for muon isolation uncertainties and an additional two for muon trigger uncertainties.

Electron reconstruction and identification uncertainties are contained in 35 recommended nuisance parameters. There are three additional standard systematics included for electron energy scale and resolution and a final two are included for electron isolation and trigger

## CHAPTER 6. BACKGROUNDS AND SYSTEMATICS

efficiency.

There are a number of jet energy scale and resolution systematics included in the analysis including parameters to estimate energy scale and resolution uncertainties,  $b$ -jet scale and efficiency uncertainties, flavor composition, response, energy scale dependence on pile-up, and  $\eta$  calibrations.

Systematics of the pileup  $\langle \mu \rangle$  value are determined by varying the data scale factor and accounted for with one overall nuisance parameter (though pile-up effects on jets are considered independently). Overall integrated luminosity uncertainty is  $\pm 1.7\%$  as calculated for the complete 2015-2018 dataset [55].

### 6.2.2 Theoretical uncertainties

Theoretical uncertainties are determined and evaluated per sample rather than on reconstructed physics objects and determine the accuracy of each of our Monte Carlo generated signal and background samples. In this analysis theoretical uncertainties are calculated and included in our overall fit for VBF signal, and ggF, top,  $WW$  and  $Z \rightarrow \tau\tau$  background events. In this section I will briefly discuss the uncertainties associated with each of these samples and demonstrate the scale of their effects. Their impacts on the overall results is further discussed in the next chapter.

#### Signal theory uncertainties

The VBF theoretical uncertainties come from four main sources: shower, PDFs,  $\alpha_s$ , and QCD. Parton shower uncertainties are derived from comparison between different parton shower generators (Powheg+Pythia8 (nominal) and Powheg+Herwig7) and constitute the largest theoretical uncertainties in the signal region. PDF uncertainties are derived

## CHAPTER 6. BACKGROUNDS AND SYSTEMATICS

Systematic uncertainty	Short description
	Event
Luminosity PRW_DATASF	uncertainty on total integrated luminosity uncertainty on pileup reweighting
	Electrons
EL_EFF_Trigger_Total_1NPCOR_PLUS_UNCOR EL_EFF_Reco_Total_1NPCOR_PLUS_UNCOR EL_EFF_ID_CorrUncertaintyNP (0 to 15) EL_EFF_ID_SIMPLIFIED_UncorrUncertaintyNP (0 to 17) EL_EFF_Iso_Total_1NPCOR_PLUS_UNCOR EG_SCALE_ALL EG_SCALE_AF2	trigger efficiency uncertainty reconstruction efficiency uncertainty ID efficiency uncertainty splits in 16 components ID efficiency uncertainty splits in 18 components isolation efficiency uncertainty energy scale uncertainty
	Muons
MUON_EFF_TrigStatUncertainty MUON_EFF_STAT MUON_EFF_SYS MUON_EFF_STAT_LOWPT MUON_EFF_SYST_LOWPT MUON_ID MUON_MS MUON_SCALE MUON_SAGITTA_RHO MUON_SAGITTA_RESBIAS	trigger efficiency uncertainty reconstruction and ID efficiency uncertainty for muons with $p_T > 20$ GeV reconstruction and ID efficiency uncertainty for muons with $p_T \leq 20$ GeV momentum resolution uncertainty from inner detector momentum resolution uncertainty from muon system momentum scale uncertainty charge dependent momentum scale uncertainty
	Jets
JET_EffectiveNP_Detector (1 to 2) JET_JER_EffectiveNP_ (1 to 7) JET_JER_DataVsMC JET_BJES_Response JET_EffectiveNP_Mixed (1 to 3) JET_EffectiveNP_Modelling (1 to 4) JET_EffectiveNP_Statistical (1 to 6) JET_EtaIntercalibration_NonClosure_negEta JET_EtaIntercalibration_NonClosure_posEta JET_EtaIntercalibration_TotalStat JET_Pileup_OffsetMu JET_Pileup_OffsetNPV JET_Pileup_PtTerm JET_Pileup_RhoTopology JET_Flavor_Composition JET_Flavor_Response JET_PunchThrough_MC16 JET_SingleParticle_HighPt JET_JvtEfficiency FT_EFF_Eigen_B (0 to 2) FT_EFF_Eigen_C (0 to 2) FT_EFF_Eigen_Light (0 to 3) FT_EFF_Eigen_extrapolation FT_EFF_Eigen_extrapolation_from_charm	detector related energy scale uncertainty energy resolution uncertainty energy resolution modelling uncertainty energy scale uncertainty on b-jets energy resolution uncertainty energy scale uncertainty on eta-intercalibration (modeling) statistical resolution uncertainty energy scale uncertainty on eta-intercalibrations (non-closure) energy scale uncertainty on eta-intercalibrations (non-closure) energy scale uncertainty on eta-intercalibrations (statistics/method) energy scale uncertainty on pile-up (mu dependent) energy scale uncertainty on pile-up (NPV dependent) energy scale uncertainty on pile-up (pt term) energy scale uncertainty on pile-up (density $\rho$ ) energy scale uncertainty on flavour composition energy scale uncertainty on samples' flavour response energy scale uncertainty for punch-through jets energy scale uncertainty from the behaviour of high- $p_T$ jets JVT efficiency uncertainty <i>b</i> -tagging efficiency uncertainties (“BTAG_MEDIUM”): 3 components for $b$ jets, 3 for $c$ jets and 4 for light jets <i>b</i> -tagging efficiency uncertainty on the extrapolation to high- $p_T$ jets <i>b</i> -tagging efficiency uncertainty on tau jets

Table 6.13: Summary of the experimental systematic uncertainties considered.

## CHAPTER 6. BACKGROUNDS AND SYSTEMATICS

from internal re-weighting within the nominal Powheg+Pythia8 samples, QCD uncertainties include factorization and renormalization scales and are found with an 8-point envelope scheme. VBF signal theory systematics within the signal region, control/validation regions is shown in 6.14.

Fit regions	Impact high / low (+/- stat) [%]			
	SR	CR.Ztt	VR.top	VR.WW
theo_VBF_shower (VBF)	-8.8 ± 0.5	-8.0 ± 1.7	-15.3 ± 1.5	0.0 ± 2.0
theo_VBF_qcd_muRmuF (VBF)	0.9 ± 0.0 / -0.8 ± 0.0	1.3 ± 0.0 / -1.1 ± 0.0	1.9 ± 0.0 / -1.3 ± 0.0	1.2 ± 0.0 / -0.3 ± 0.0
theo_VBF_pdf4lhc (VBF)	1.7 ± 0.0	1.7 ± 0.0	3.4 ± 0.0	1.7 ± 0.0
theo_VBF_alphas (VBF)	0.5 ± 0.3	0.4 ± 1.1	1.0 ± 1.0	0.4 ± 1.2

Table 6.14: VBF theory uncertainties- NP breakdown

We examine these uncertainties for any shape effects on the BDT output used in the final fit. Plots 6.33 show very little shape effects.

### ggF background theory uncertainties

The ggF theoretical uncertainties come from four main sources: shower, PDFs,  $\alpha_s$ , and QCD. Parton shower uncertainties use the same prescription described for the VBF samples and similarly constitute the largest theoretical uncertainties from ggF events. PDF uncertainties are derived from internal re-weighting as for VBF and QCD uncertainties include factorization and renormalization scales and are derived using the WG1 scheme [56]. ggF theory uncertainties derived in the signal and validation regions are show in the table 6.15.

Fit regions	Impact high / low (+/- stat) [%]				
	SR	CR.Ztt	CR_ggF1	CR_ggF2	VR_top
PDF uncert (ggF)	1.7 ± 0.0	1.8 ± 0.0	1.9 ± 0.0	2.0 ± 0.0	2.1 ± 0.0
$\alpha_s$ uncert (ggF)	3.2 ± 1.3	3.3 ± 3.7	3.2 ± 0.6	3.2 ± 1.2	3.4 ± 2.8
Shower uncert (ggF)	-14.4 ± 1.2	0.7 ± 4.0	3.5 ± 0.6	18.6 ± 1.5	-2.7 ± 3.1
QCD mu uncert (ggF)	5.1 ± 1.4	6.4 ± 4.1	7.2 ± 0.6	7.3 ± 1.4	6.8 ± 3.2
QCD res uncert (ggF)	4.7 ± 1.4	6.1 ± 4.1	6.8 ± 0.6	7.0 ± 1.4	6.5 ± 3.2
QCD mig01 uncert (ggF)	3.7 ± 1.4	4.3 ± 4.0	4.4 ± 0.6	4.4 ± 1.3	4.3 ± 3.1
QCD mig12 uncert (ggF)	4.8 ± 1.4	8.4 ± 4.2	10.2 ± 0.6	11.0 ± 1.4	10.0 ± 3.3
QCD qm1 uncert (ggF)	1.0 ± 1.4	2.4 ± 4.0	1.3 ± 0.6	1.9 ± 1.3	1.7 ± 3.0
QCD $p_T^H$ uncert (ggF)	1.3 ± 1.4	4.9 ± 4.1	2.4 ± 0.6	3.2 ± 1.3	2.5 ± 3.0
QCD vbf2j uncert (ggF)	5.3 ± 1.4	3.3 ± 4.0	1.3 ± 0.6	1.1 ± 1.3	2.7 ± 3.1
QCD vbf3j uncert (ggF)	-2.1 ± 1.3	-1.4 ± 3.8	0.5 ± 0.6	0.6 ± 1.3	-0.4 ± 3.0

Table 6.15: ggF theory uncertainties- NP breakdown

## CHAPTER 6. BACKGROUNDS AND SYSTEMATICS




Figure 6.33: Up and down variations shown for four theoretical uncertainties from the VBF signal displayed against the VBF vs. top +WW BDT. The nominal sample is shown in black and slopes are calculated for up and down variations to display any potential linear shape effects. Distributions are shown after all signal region cuts.

We examine these uncertainties for any shape effects on the BDT output used in the final fit. Plots 6.34 show very little shape effects.

### Top background theory uncertainties

Top theoretical uncertainties include those from a variety of sources including hard scatter generation, parton shower, factorization and renormalization scales, initial state radiation (ISR), final state radiation (FSR), and PDFs. For parton shower uncertainties, the same

## CHAPTER 6. BACKGROUNDS AND SYSTEMATICS




Figure 6.34: Up and down variations shown for four theoretical uncertainties from the ggF background displayed against the VBF vs. top +WW BDT. The nominal sample is shown in black and slopes are calculated for up and down variations to display any potential linear shape effects. Distributions are shown after all signal region cuts.

scheme used in ggF and VBF samples is used here (comparisons between Powheg+Pythia8 (nominal) and Powheg+Herwig7 are made). Hard scatter generation uncertainties are similarly derived from comparing the nominal generation method with aMC@Nlo+Pythia8. QCD scale uncertainties from renormalization and factorization are added with a 6-point envelope scheme [? ]. ISR and FSR uncertainties are determined through varying internal weights and the PDF uncertainty is completed with internal reweighting and errors produced within the nominal Powheg+Pythia8 samples.

## CHAPTER 6. BACKGROUNDS AND SYSTEMATICS

The top background considered in this analysis includes  $t\bar{t}$  and  $Wt$  where  $t\bar{t}$  is the dominant sample by far. Theoretical uncertainties are produced for both single top and  $t\bar{t}$  independently and both include those from a variety of sources including hard scatter generation, parton shower, factorization and renormalization scales, initial state radiation (ISR), final state radiation (FSR), and PDFs. For parton shower uncertainties, the same scheme used in ggF and VBF samples is used here (comparisons between Powheg+Pythia8 (nominal) and Powheg+Herwig7 are made). Hard scatter generation uncertainties are similarly derived from comparing the nominal generation method with aMCatNlo+Pythia8. QCD scale uncertainties from renormalization and factorization are added with a 6-point envelope scheme. ISR and FSR uncertainties are determined through varying internal weights and the PDF uncertainty is completed with internal reweighting and errors produced within the nominal Powheg+Pythia8 samples.  $Wt$  events contain one additional systematic uncertainty - interference. This is derived from removing  $t\bar{t}$ -like contributions at the level of amplitude (DR) or matrix elements (DS) [57]. Table 6.16 shows theory uncertainty from  $t\bar{t}$  events in the signal, control and validation regions while table 6.17 shows the same for  $Wt$ . While  $Wt$  uncertainties are large, the background has a minimal contribution to the analysis so these do not have a large effect on overall results.

Fit regions	Impact high / low (+/- stat) [%]					
	SR	CR_Ztt	VR_top	VR_WW	CR_ggF1	CR_ggF2
theo_ttbar_shower (ttbar_vbf)	4.1 ± 0.6	3.6 ± 1.8	-0.5 ± 0.1	11.2 ± 0.4	10.6 ± 0.2	17.6 ± 0.3
theo_ttbar_generator (ttbar_vbf)	18.3 ± 1.2	17.7 ± 3.7	5.6 ± 0.2	7.3 ± 0.7	18.4 ± 0.4	16.0 ± 0.7
theo_ttbar_scale (ttbar_vbf)	13.6 / -12.9	13.8 / -12.6	11.6 / -11.5	18.1 / -15.0	13.5 / -12.6	14.4 / -13.1
theo_ttbar_isr (ttbar_vbf)	-0.6 / 0.6	-1.0 / 0.8	-0.6 / 0.5	-0.4 / 0.3	0.3 / -0.3	1.3 / -1.2
theo_ttbar_fsr (ttbar_vbf)	-1.3 / 2.2	-2.0 / 7.4	1.2 / -1.7	-1.6 / 3.1	-3.4 / 5.2	-4.2 / 5.8
theo_ttbar_pdf (ttbar_vbf)	1.9	1.7	1.5	1.6	1.7	1.6

Table 6.16:  $t\bar{t}$  theory uncertainties- NP breakdown

We examine these uncertainties for any shape effects on the BDT output used in the final fit. Plots ?? show too little statistics to discern any shape effects.

Plots ?? show very little shape effects from theoretical uncertainties on far more numerous

## CHAPTER 6. BACKGROUNDS AND SYSTEMATICS

Fit regions	Impact high / low (+/- stat) [%]					
	SR	CR_Ztt	VR_top	VR_WW	CR_ggF1	CR_ggF2
theo_wt_shower (Wt_vbf)	18.9 ± 3.3	11.5 ± 10.0	4.6 ± 1.0	24.5 ± 2.0	30.9 ± 1.2	45.7 ± 2.8
theo_wt_generator (Wt_vbf)	21.4 ± 4.5	18.5 ± 13.6	17.7 ± 1.4	19.4 ± 2.5	20.6 ± 1.6	21.1 ± 3.3
theo_wt_scale (Wt_vbf)	5.1 / -5.0	5.5 / -5.2	5.2 / -5.0	7.3 / -6.5	5.3 / -5.1	5.4 / -5.1
theo_wt_isr (Wt_vbf)	-0.1 / 0.1	-0.9 / -0.1	-0.6 / 0.5	-0.6 / 0.3	0.3 / -0.2	0.5 / -0.7
theo_wt_fsr (Wt_vbf)	-0.2 / 3.9	-1.9 / 3.5	2.0 / -4.1	-0.8 / 0.6	-2.3 / 4.4	-2.3 / 4.9
theo_wt_interference (Wt_vbf)	4.4 ± 2.5	14.6 ± 7.7	1.2 ± 0.9	12.3 ± 1.4	-3.1 ± 0.9	7.6 ± 1.7
theo_wt_pdf (Wt_vbf)	2.0	1.9	1.9	2.0	1.9	1.9

Table 6.17:  $Wt$  theory uncertainties- NP breakdown

$t\bar{t}$  backgrounds.

### WW background theory uncertainties

The  $WW$  background theoretical uncertainties come from three dominant sources: PDFs,  $\alpha_s$ , and QCD scale. **Add CKKM uncertainties and generator?**. PDF uncertainties are derived from internal re-weighting within the nominal Powheg+Pythia8 samples, QCD uncertainties include factorization and renormalization scales, and generator uncertainties are derived from a comparison with  $MG5_aMC@NLO$  interfaced with Pythia8 (and so include shower uncertainty).  $WW$  theory systematics within the signal region, control/validation regions are shown in 6.18.

Fit regions	Impact high / low (+/- stat) [%]					
	SR	CR_Ztt	VR_top	VR_WW	CR_ggF1	CR_ggF2
theo_ww_scale (WW_vbf)	20.0 / -14.9	24.6 / -17.8	26.4 / -19.1	26.8 / -19.0	25.2 / -18.1	29.2 / -20.1
theo_ww_pdf (WW_vbf)	1.0	1.0	1.0	0.9	1.0	0.9
theo_ww_alphas (WW_vbf)	2.0	2.1	2.3	2.3	2.2	2.4

Table 6.18:  $WW$  theory uncertainties- NP breakdown

We examine these uncertainties for any shape effects on the BDT output used in the final fit. Plots ?? show no shape effects.

### 6.2.3 $Z \rightarrow \tau\tau$ background theory uncertainties

The  $Z \rightarrow \tau\tau$  theoretical uncertainties come from four main sources: PDFs,  $\alpha_s$ , QCD scale, and generator. PDF uncertainties are derived from internal re-weighting within the

## CHAPTER 6. BACKGROUNDS AND SYSTEMATICS




Figure 6.35: Up and down variations shown for four theoretical uncertainties from the  $Wt$  background displayed against the VBF vs. top +WW BDT. The nominal sample is shown in black and slopes are calculated for up and down variations to display any potential linear shape effects. Distributions are shown after all signal region cuts.

## CHAPTER 6. BACKGROUNDS AND SYSTEMATICS




Figure 6.36: Up and down variations shown for four theoretical uncertainties from the  $t\bar{t}$  background displayed against the VBF vs. top +WW BDT. The nominal sample is shown in black and slopes are calculated for up and down variations to display any potential linear shape effects. Distributions are shown after all signal region cuts.

nominal Powheg+Pythia8 samples, QCD uncertainties include factorization and renormalization scales and use a seven-point envelope,  $\alpha_s$  uncertainties use an up-down comparison with two *NNPDF3.0* NNLO  $\alpha_s$  variations, and generator uncertainties are derived from a comparison with *MG5<sub>a</sub>MC@NLO* interfaced with Pythia8.  $Z \rightarrow \tau\tau$  theory systematics within the signal region, control/validation regions is shown in 6.19.

We examine these uncertainties for any shape effects on the BDT output used in the final fit. Plots 6.38 show very little shape effects.

## CHAPTER 6. BACKGROUNDS AND SYSTEMATICS




Figure 6.37: Up and down variations shown for four theoretical uncertainties from the  $Z \rightarrow \tau\tau$  background displayed against the VBF vs. top +WW BDT. The nominal sample is shown in black and slopes are calculated for up and down variations to display any potential linear shape effects. Distributions are shown after all signal region cuts.

Fit regions	Impact high / low (+/- stat) [%]					
	SR	CR_Ztt	VR_top	VR_WW	CR_ggF1	CR_ggF2
theo_ztautau_scale (ztautau_VBF)	6.6 / -5.7	12.1 / -9.5	7.8 / -11.2	4.8 / -5.7	11.7 / -9.3	21.4 / -13.1
theo_ztautau_pdf (ztautau_VBF)	1.6	1.2	1.7	1.2	1.4	1.2
theo_ztautau_alphas (ztautau_VBF)	1.4	1.7	2.6	1.6	1.8	2.1
theo_ztautau_generator (ztautau_VBF)	-17.2 $\pm$ 5.9	32.9 $\pm$ 4.9	33.0 $\pm$ 14.7	-3.9 $\pm$ 14.2	-2.4 $\pm$ 2.6	-19.2 $\pm$ 4.7

Table 6.19:  $Z \rightarrow \tau\tau$  theory uncertainties- NP breakdown

## CHAPTER 6. BACKGROUNDS AND SYSTEMATICS




Figure 6.38: Up and down variations shown for four theoretical uncertainties from the  $Z \rightarrow \tau\tau$  background displayed against the VBF vs. top + $WW$  BDT. The nominal sample is shown in black and slopes are calculated for up and down variations to display any potential linear shape effects. Distributions are shown after all signal region cuts.

# Chapter 7

## Results

Thus far I have outlined all reconstruction-level inputs to our analysis including event selection, all backgrounds and their estimation methods, and all systematic uncertainties. Each of these components is critical to our final result and this section will focus on the methods used to extract differential cross-section measurements from the signal and control regions previously defined.

### 7.1 Statistical analysis

#### 7.1.1 Likelihood functions

This analysis rests on the estimation of a parameter  $\mu$  which describes the statistical significance of our signal yield relative to its Standard Model prediction and in this particular analysis defines the signal cross-section and signal and background yields. We build a likelihood function  $\mathcal{L}(\mu, \Theta)$  where the signal strength  $\mu$  is a parameter of interest (POI) and nuisance parameters (NPs)  $\Theta = \Theta_a, \Theta_b, \dots$  represent all relevant uncertainties. None of these values are known a priori so the likelihood is built to represent the probability of particular

## CHAPTER 7. RESULTS

values for the POIs and NPs. The analysis uses a maximum likelihood estimator to find the inputs which maximize the likelihood (or equivalently and more mathematically tractable, minimize the negative log of the likelihood). Here I will briefly outline how a likelihood function can incorporate regions of interest. This discussion uses [58] as a guide. First, the Poisson distribution  $\mathcal{P}(n|\lambda)$  describes the probability of  $n$  events with a true unknown yield  $\lambda$ :

$$\mathcal{P}(n|\lambda) = \lambda^n \frac{e^{-\lambda}}{n!} \quad (7.1)$$

Next we can define a variable observable  $x$  with probability density  $f(x)$  hence with  $n$  events the probability density of each is multiplied. The likelihood of  $\lambda$  can now be written

$$\mathcal{L}(\lambda) = \mathcal{P}(n|\lambda) \prod_{\text{event}}^n f(x) \quad (7.2)$$

Our likelihood also must take into account multiple regions (the signal region as well as the control regions) and so these likelihoods are multiplied together with their own distinct Poisson distributions

$$\mathcal{L}(\lambda) = \prod_r^{\text{regions}} (\mathcal{P}(n_r|\lambda_r) \prod_{\text{event}}^n f(x)) \quad (7.3)$$

A simultaneous fit maximizes this likelihood function and so produces yield  $\lambda$  for each free parameter simultaneously. The  $\lambda$  values here represent predicted yields where  $\lambda_{r,b} = \mu\lambda_{\text{sig}} + \lambda_{\text{bkg}}$ . Maximizing the overall likelihood is made simpler by applying the natural logarithm (as the products between Poisson distributions become summations) and negating the likelihood, so as to take advantage of minimizing software.

Particle physics defines discovery with rigorous standards using hypothesis testing. The null hypothesis is considered  $\mu = 0$  and is considered “background-only” while the alternative hypothesis is that there is a signal above this background. The probability that the null

## CHAPTER 7. RESULTS

hypothesis is rejected (or that the signal is discovered) is defined using a  $p$ -value. The  $p$ -value can be converted to a number of Gaussian standard deviations and in high energy physics three  $\sigma$  significance (or a  $p$ -value of  $1.35 \times 10^{-3}$ ) shows evidence while a five  $\sigma$  result ( $p$ -value  $2.87 \times 10^{-7}$ ) is considered a discovery.

In this analysis we use HistFitter software which is compiled with RooStats, RooFit, and HistFactory to complete our likelihood fit. We calculate the signal VBF Higgs cross-section in the fiducial space defined with signal region cuts (described in Chapter 4) with a simultaneous fit of the signal and control regions. Each region has a different distribution used in the fit to maximize that samples' discrimination against all other events. Overall we include six floating parameters: signal strength for  $\mu_{VBF}$ ,  $\mu_{TopWW}$ ,  $\mu_{Z+jets}$ , and  $\mu_{ggF}$ ,  $\mu_{ggF1}$  and  $\mu_{ggF2}$ . Other backgrounds (fakes,  $V/\gamma$ ,  $VH$ , and  $tth$ ) are included as fixed parameters in the fit.

The background treatment differs depending on the particular region and sample. For  $Z+jets$ , a designated control region is used and so the yield in the signal region can be found by including a transfer factor which maps the yield in the  $Z+jets$  control region to its yield in the signal region. The ggF signal strengths are split and derived as shown in Chapter 5. Transfer factors to extrapolate yields from one region to another are derived by normalizing to the MC simulation. For backgrounds estimated within the signal region (Top/WW) the transfer factor is used only to extrapolate using simulations in the control regions where the contributions enter (other than the signal region) and so uncertainties associated with transfer factors are reduced. For the backgrounds included as fixed parameters in the fit, their predicted MC value is used (in the case of  $V\gamma$ ,  $VH$ , and  $tth$ ) and for fakes their data-driven estimate is used.

Events in the  $e\mu$  and  $\mu e$  channels are combined in the analysis and the likelihood is defined as the product of the Poissonians over BDT output bins in the signal region multiplied

## CHAPTER 7. RESULTS

by the product of Poissonians over discriminant distributions in the controls regions (ggF CRs,  $Z+jets$  CR and Top/ $WW$  CR- within the signal region). The likelihood can then be calculated:

$$\begin{aligned} \mathcal{L}(\mu) &= \prod_{j=0}^{N_{\text{sig BDT bins}}} \mathcal{P}(n_j | \lambda_r) \times \prod_{k=1}^{N_{\text{CR-bins}}} P(n_k | \lambda_b) \\ &= \prod_{j=0}^{N_{\text{sig BDT bins}}} \mathcal{P}(n_j | \mu_s \lambda_{sig,j} + \sum_n \mu_b^n \lambda_{bkg}^{nj}) \times \prod_{k=1}^{N_{\text{CR-bins}}} P(n_k | \mu_s \lambda_{sig,k} + \sum_n \mu_b^n \lambda_{bkg}^{nk}) \end{aligned} \quad (7.4)$$

Here the signal and background strengths are denoted  $\mu$  and yields  $\lambda$ . Sums over all backgrounds  $n$  include  $Z+jets$ , top/ $WW$ , and ggF; the minor backgrounds and data-driven  $W+jets$  estimate are added to Poisson expectations as fixed parameters. The last important addition to this likelihood is that of the numerous nuisance parameters affecting the analysis. These parameters  $\Theta$  come in two general types: systematics which do not affect the shape of discriminating variables and those that do. Flat systematics are constrained with a unit Gaussian and their effects are measured for  $\Theta = \pm 1$ . Shape systematics are split into purely flat and shape components and are again constrained with a unit Gaussian though their effects are measured through  $\nu_{\text{shape}}(\Theta) = 1 + \epsilon\Theta$  where  $\epsilon$  is determined by measuring  $\nu_{\text{shape}}$  at  $\Theta = \pm 1$ . Adding nuisance parameters  $\Theta$  our final likelihood can be written

$$\mathcal{L}(\mu, \vec{\Theta}) = \prod_{j=0}^{N_{\text{sig BDT bins}}} \mathcal{P}(n_j | \mu_s \lambda_{sig,j} + \sum_n \mu_b^n \lambda_{bkg}^{nj}) \times \prod_{k=1}^{N_{\text{CR-bins}}} P(n_k | \mu_s \lambda_{sig,k} + \sum_n \mu_b^n \lambda_{bkg}^{nk}) \times \prod_{i=1}^{N_{\Theta_i}} G(\tilde{\Theta}_i | \Theta_i, 1) \quad (7.5)$$

where  $G(\tilde{\Theta}_i | \Theta_i, 1) = \frac{1}{\sqrt{2\pi}} e^{\frac{(\tilde{\Theta}_i - \Theta_i)^2}{2}}$  is the unit Gaussian and  $\tilde{\Theta}$  is an auxiliary measurement of  $\Theta$ .

The fit is performed for a signal region, three ggF control regions, a  $Z+jets$  control region,

## CHAPTER 7. RESULTS

and a shared top/ $WW$  control defined within the signal region estimated in one half of the binned region defined by the signal region discriminant. Table 7.1 summarizes the regions used with in the fit and their cuts. These are discussed in more detail in Chapters 4 and 5.

Signal region	Top/ $WW$ control region	$Z+jets$ control region	ggF CR 1	ggF CR 2	ggF CR 3
			$n_{jets} \geq 2$		$n_{jets} < 2$
			b-veto		
CJV $\geq 20\text{GeV}$	CJV $\geq 20\text{GeV}$ and OLV = 1 or CJV $< 20\text{GeV}$ and OLV $\neq 1$		CJV $\geq 20\text{GeV}$	-	-
OLV = 1			-	OLV $\neq 1$	-
$m_{\tau\tau} < m_Z - 25\text{GeV}$	$ m_{\tau\tau}  - m_Z \leq 25\text{GeV}$		$m_{\tau\tau} < m_Z - 25\text{GeV}$		
$m_{jj} > 00\text{ GeV}$	$m_{\ell\ell} < 80\text{GeV}$		$m_{jj} > 00\text{ GeV}$		
$\Delta Y_{jj} > 2.1$	-		$\Delta Y_{jj} > 2.1$		
BDT <sub>VBF</sub> $\geq 0$	BDT <sub>VBF</sub> $< 0$	-	-	-	-

Table 7.1: Summary of all signal and control regions included in simultaneous fit

Each of the regions included in the fit uses a different discriminant to best characterize that sample’s contribution. These discriminants are almost all BDT outputs trained on various kinematic distributions to characterize certain background and signal samples. These are discussed in Chapters 4 and 5. The  $Z+jets$  control region is defined with  $m_T$  instead of a designated BDT after studies showed its success at mitigating uncertainty on  $\mu_{Z+jets}$ . These studies are included in Appendix A. Table 7.2 summarizes the regions used in the fit and shows their discriminant and number of bins. Table 7.3 shows MC event yields in all regions included in the fit.

Category	SR	Top/ $WW$ CR	$Z+jets$ CR	ggF-CR1	ggF-CR2	ggF-CR3
Discriminant	BDT <sub>VBF</sub>	BDT <sub>TopWWAll</sub>	$m_T$	BDT <sub>ggF1</sub>	BDT <sub>ggF2</sub>	BDT <sub>ggF1</sub>
Number of bins	25	8	10	5	5	5

Table 7.2: Fit categories, including SR and CRs, distributions and number of bins used in the fit.

## CHAPTER 7. RESULTS

Category	SR	Top/ $WW$ CR	$Z$ +jets CR	ggF-CR1	ggF-CR2	ggF-CR3
VBF	149.7	14.1	16.1	100.4	8.6	192.3
Top	385.9	2878.7	331.3	34110.4	10546.4	40206.3
$WW$	202.0	969.0	122.6	9653.7	2608.5	86387.7
ggF	72.4	30.3	11.7	543.1	114.1	4069.1
$Z$ +jets	313.1	476.4	1391.1	5288.5	1395.6	131014
Fakes	63.5	123.4	16.7	1752.9	570.0	15166.2
$V\gamma$	23.5	33.7	36.7	490.7	94.9	4652.9
$VH$	2.0	2.4	1.4	89.7	18.5	164.6
$t\bar{t}H$	3.8	5.9	19.8	35.5	8.5	437.6

Table 7.3: MC event yields in each signal and control region pre-fit

### 7.1.2 Asimov results

The results contained in this thesis are preliminary and data for this analysis has not yet been unblinded. Results shown will use the Asimov dataset approach to deduce expected performance of the fit. Using this approach, a representative dataset is formed (with Monte Carlo pseudo-experiments) which returns the true value for each estimated parameter. In this way we gain an understanding for the constraints on our nuisance parameters and so determine and correct any over-constraints from inputs into the fit. Using the Asimov datasets all floating parameters should be calculated to be 1 and all nuisance parameters found to be 0. Results of the fitted nuisance parameters to the Asimov dataset are made under the hypothesis that VBF Higgs boson cross-sections follow our Standard Model predictions. The distributions shown in 7.1 demonstrate the post-fit distributions for each of the signal and control regions included in the fit.

The results from here on are not the final ones I'll include but dummy plots/tables as more finalizations/developments are made.

The first fits performed use no nuisance parameters, only statistical uncertainties, to show both how the overall fit performs at its most basic level and to deduce the overall impact of

## CHAPTER 7. RESULTS




Figure 7.1: Binned distributions for each signal and control region shown after Asimov fit where ratio between Asimov data and MC predictions is one by design. Red points show VBF signal significance per bin.

adding systematics to the fit. The plot 7.2 shows correlations between floating parameters in the fit using a stat-only scheme where only the six floating signal strength parameters are fit.

Fit results for each of the signal strength parameters in our statistical fit are shown for both stat-only and statistical + systematic uncertainties. The current fit including all systematic uncertainties gives an error on the  $\mu_{VBF}$  or  $\pm 0.20$ . This corresponds to a null p-value of  $1.5 \times 10^{-9}$  or a significance of  $5.9\sigma$  which is more than twice the expected significance in the last published ATLAS VBF Higgs  $\rightarrow \ell\nu\ell\nu$  results [15]. The table below 7.4 shows the stat-only and combined systematic resulting signal strengths and their expected uncertainties

## CHAPTER 7. RESULTS



Figure 7.2: Correlations between fit after a stat-only fit

based on Asimov fits.

Floating parameter	Estimated uncertainty (stat-only)	Estimated uncertainty (stat+sys)
$\mu_{VBF}$	17.1 %	20.2%
$\mu_{top/WW}$	1.2 %	1.8%
$\mu_{ggF}$	48.4 %	64.5%
$\mu_{ggF_1}$	44.9 %	56.5%
$\mu_{ggF_2}$	15.1 %	40.7%
$\mu_{Z+jets}$	0.5 %	1.8%

Table 7.4: Asimov fit results for all floating parameters in stat-only and full systematic fits.

We can visualize and better understand these results through a 1D scan of our parameter of interest ( $\mu_{VBF}$ ). Using the inputs and distributions of of our fit we can view the minimization of the total likelihood and and its width the see the impact of systematics and constraints from the other floating parameters. Plots 7.3 shows fit results for  $\mu_{VBF}$  if we use all systematics, only statistical uncertainties, and finally if were to leave only  $\mu_{VBF}$  as a floating parameter and all other constant. These results demonstrate the substantial role systematics play in our analysis. We break down these impacts to assess which nuisance parameters have the greatest effects on each measured parameter so that we can find any that seem unduly high and find ways to mitigate their effects through our event selection and fit

## CHAPTER 7. RESULTS

techniques. The following plots 7.4 show the 40 largest impact parameters for each measured signal strength. These effects are derived post-fit which means that at the maximum of the likelihood each parameter is changed by  $\pm 1\sigma$  individually. The measured effects of these changes are shown in blue on the plots and ranked in order of total combined positive and negative impact.




Figure 7.3: Scan of  $\mu_{\text{VBF}}$  negative log-likelihood using Asimov data and using only one floating parameter, all  $\mu$  values floating, and all  $\mu$  and NPs floating.

As expected using the Asimov dataset, nuisance parameters are centered at zero and floating  $\mu$  values center at 1. There are a number of systematic uncertainties that play large roles in this measurement including jet flavor NPs, which play a large role in rejecting dominant top background events, and jet pile-up which causes potential errors in the jet kinematics which are key to this analysis.

Add expected post fit yields and comparisons with observed yields and distributions.

The fit results shown in this thesis demonstrate the most recent calculations and developments from the VBF  $H \rightarrow WW$  differential measurement but they are not yet complete. Additional theoretical uncertainties (as described in the previous chapter) need to be con-

CHAPTER 7. RESULTS




Figure 7.4: Impacts of top 40 most impactful nuisance parameters for each fit  $\mu$  using Asimov data.

sidered and evaluated and final fit parameters may continue to change before the analysis is unblinded.

## 7.2 Fiducial and inclusive cross sections

This analysis aims to measure the total VBF  $H \rightarrow WW^* \rightarrow \ell\nu\ell\nu$  cross-section in both a particular fiducial region and in a more inclusive region in addition to the fiducial differential

## CHAPTER 7. RESULTS

cross-section over a range of kinematic variables. In each case the measurement must be translated from one that only represents conditions in the ATLAS detector to values that can be understood with broad range of theoretical models. For fiducial measurements this can be done simply with a  $C$  factor. First a fiducial region must be defined that is applicable at both reconstruction and truth level and will define under what conditions the cross-section is measured. The fiducial cross-section is defined:

$$\sigma_{\text{VBF}H \rightarrow WW^* \rightarrow \ell\nu\ell\nu}^{\text{fid}} = \frac{N_{\text{obs}} - N_{\text{bkg}}}{C \times \mathcal{L}} \quad (7.6)$$

where  $\mathcal{L}$  is the integrated luminosity, and  $N$  (obs,bkg) is the observed and estimated background number of events respectively.  $C$  is a factor the accounts for detector effects that would lower the overall cross-section like detector inefficiencies and resolution effects. To minimize any necessary theoretical extrapolation, the fiducial phase space is defined as close the reconstruction-level signal region as possible. The table 7.5 lists all fiducial cuts that describe the phase space. These include requirements on lepton number, flavor, sign, and kinematics, missing energy, jet number and kinematics, and other VBF specific requirements (OLV and CJV cuts). The cuts represent the space within which both reconstruction-level and truth-level cross-sections are measured. Using yields extracted from the statistical fit and a  $C$ -factor we define with

$$C = \frac{N_{\text{fid}}}{N_{\text{reco}}} \quad (7.7)$$

where  $N_{\text{fid}}$  is extacted from theoretical simulation. **Include Jiayi C-factor calculations and result here. Then cross-section value and error (expected value, will be compared to observation).**

The inclusive cross-section requires one additional acceptance correction factor  $A$ , which is estimated from theoretical calculations and extrapolates the fiducial cross section to a

## CHAPTER 7. RESULTS

Fiducial Requirement
$ \eta(\ell)  < 2.5$
$p_T^{\text{lead}} > 22\text{GeV}$
$p_T^{\text{sublead}} > 15\text{GeV}$
$N_{\text{leptons}} \geq 2$
Leptons required to be opposite flavor and sign
$\Delta R(\ell, \ell) > 0.1$
$m_{\ell\ell} > 10\text{GeV}$
$E_T^{\text{miss}} > 20\text{GeV}$
$p_T(\text{jet}) > 30\text{ GeV}$
$ \eta(\text{jet})  < 4.5$
$N_{\text{jets}} \geq 2$
$N_{b\text{-jet}} < 1$
$m_{jj} > 200\text{GeV}$
$\Delta Y_{jj} > 2.1$
$\Delta R(\ell, \text{jet}) > 0.4$
OLV = 1
$\text{CJV} > 20\text{GeV}$

Table 7.5: Fiducial phase space definition

more inclusive phase space defined solely by a requirement for  $N_{\text{jets}} \geq 2$  where each jet has  $p_T > 30\text{GeV}$ . This will be calculated from

$$\sigma_{VBF}^{\text{incl}} = \frac{\sigma_{VBF}^{\text{fid}}}{A}. \quad (7.8)$$

The inclusive cross-section widens the applicable phase space of the measurement and provides another useful metric to compare data with expected Standard Model results. Factors  $A$  and  $C$  are found through theoretical calculations which use VBFNLO to estimate VBF processes at next-to-leading order in QCD. Add factor A from Jiayi as well as inclusive cross-section expected value, will be compared to observation.

## CHAPTER 7. RESULTS

### 7.3 Differential cross-section measurements

The differential cross-section measurement requires a more rigorous unfolding procedure than the total fiducial and inclusive cross-sections. Monte Carlo simulations processed in a pseudo-detector with all its expected inefficiencies and limits (reconstruction-level) are compared to Monte Carlo simulations processed solely at the particle level (truth-level). The detector effects that influence reconstruction-level events are unfolded so that they match their truth-level counterparts. This reco-truth matching is a test that the unfolding process works as expected and does not bias results. After the analysis is unblinded, data will be unfolded using the same mechanisms to create a differential measurement that can be utilized beyond ATLAS. This analysis employs the iterative bayesian unfolding method[59] whose goal is to determine the probability that each bin  $j$  in a reconstruction-level distribution corresponds to bin  $i$  in a truth-level distribution. Using Bayes' theorem, this probability can be attained with knowledge of the true spectrum  $T$  and the measured or reco-level spectrum  $R$ . We begin with Bayes' theorem:

$$P(T_i|R) = \frac{P(R, T_i)P(T_i)}{\sum_t P(R, T_t)P(T_t)} \quad (7.9)$$

where the denominator is a normalization factor,  $P(R, T_i)$  represents the likelihood and  $P(T_i)$  the prior. The prior here is vague and so we begin with the assumption that  $P(T_i)$  is constant. Hence the most probable spectrum for  $T$  maximizes the likelihood. If we observe  $n$  reco-level events, we can assign the probability of their true distributions through

$$\hat{n}(T_i) = n(R) \prod P(T_i|R). \quad (7.10)$$

## CHAPTER 7. RESULTS

We then use resultant possibilities in the Bayes formula to evaluate  $P(T_i|R_j)$ . These values constitute the smearing matrix  $M$ . This matrix can then be used in turn to estimate the truth-level events in each bin of a distribution with

$$\hat{n}(T_i) = \frac{1}{\epsilon_i} \sum_{j=1}^{n_R} n(R_j) \prod P(T_i|R_j). \quad (7.11)$$

where  $\epsilon$  is the inefficiency, or ratio of events that pass both truth and reco-level selection by those that pass only truth. Finally the truth can be determined by the reconstructed bin-by-bin yields through

$$n(T_i) = \sum_j M_{ij} \prod n(R_j). \quad (7.12)$$

The migration matrix thus directly maps reco-level distributions to truth-level. In order to calculate this matrix iteratively we first allow  $P(T_i)$  to be a constant distribution, say  $1/n_T$ , which gives an expected number of truth events  $n_0(C_i) = P_0(C_i) \prod n_R$ . Next, calculate  $\hat{n}(C)$  using the efficiency equation, and finally use  $\chi^2$  to compare  $\hat{n}(C)$  and  $n_0(C)$ . In the next iteration,  $n_0$  is replaced by  $\hat{n}$  and  $P_0$  by  $\hat{P}$  ( $P(\hat{C}_i) = \hat{n}(C_i)/N_{true}$ ) and the procedure continues until  $\chi^2$  falls below a set threshold.

While the final unfolding results fall beyond the scope of thesis, results for four kinematic variables are shown here. First, figure 7.5 shows a validation performed by using the unfolding mechanism on the same Monte Carlo events. The figures below demonstrate that truth distribution (blue) and unfolded truth distributions (black) perfectly match as expected. Reco-level distributions are similarly shown, though these clearly differ from truth distributions.

After validation, we use reconstruction-level Monte Carlo events and their truth counterparts to calculate migration matrices for each differential observable through the iterative Bayesian unfolding process. Four example migration matrices are shown in 7.6.

## CHAPTER 7. RESULTS




Figure 7.5: Validation for unfolding mechanism, truth (blue) and unfolded truth distributions (black) overlap while reconstruction level distributions (green) fall separately. Distributions shown for  $p_H^T$ ,  $m_T^{\ell\ell}$ ,  $\Delta Y_{\ell\ell}$  and  $m_{jj}^{**}$

Further developments in unfolding are underway currently and include estimating error added by the procedure (both statistical and systematic in the case of potential mismodelling), testing for potential biases toward the truth distribution, and understanding unfolding reconstruction level uncertainties as well as nominal distributions.

## CHAPTER 7. RESULTS




Figure 7.6: Unfolding matrices shown for  $p_H^T$ ,  $m_T^{\ell\ell}$ ,  $\Delta Y_{\ell\ell}$  and  $m_{jj}$  distributions. Each bin value corresponds to normalized Bayesian probabilities and the x-axis represents reconstruction-level distributions while the y-axis shows truth distributions. \*\*

## 7.4 Future results and measurements

This thesis detailed the HWW VBF differential analysis as it currently stands, but some work remains before final differential cross-section results can be measured. Chapter 5 dis-

## CHAPTER 7. RESULTS

cussed some theoretical uncertainties (VBF and ggF Higgs as well as top, WW, and  $Z \rightarrow \tau\tau$ ) but these have not been added to the overall statistical fit. These additional uncertainties are expected to play a significant role in the overall results, particularly the large shower uncertainties. With theory uncertainties and fit parameters finalized, we will estimate the signal and background yield. This is directly estimated from the fit signal strength described early in this chapter through multiplication by the theoretical cross-section of each process, total experimental efficiency ( $C$ -factor) and integrated luminosity ( $\mathcal{L}$ ). Finally, these yields are used to measure fiducial and inclusive cross-sections. The fiducial phase space definition was described previously in 7.5 as well as the fiducial and inclusive cross-section definitions. The expected MC cross-sections are reported and will be compared to observed results from data once the analysis is unblinded.

Signal and background yields will be used within the unfolding mechanism to extract final differential distributions. The variables used in the differential analysis will also be finalized in coming months. We aim to unfold 14 variables which are listed with their planned bin edges in 7.6. These variables probe the kinematics of the final state particles measured through this channel and add sensitivity to the total fiducial and inclusive cross-sections. The invariant mass of the two jets that accompany the  $W$  bosons ( $m_{jj}$ ) and the angular separation between them ( $\Delta y_{jj}$ ) are particularly sensitive to the electroweak symmetry breaking mechanism. Deviations between data and theoretical predictions in the distribution of these variables may be signs of new physics that alter the HWW coupling, like new resonances at high energy scales not accessible by direct searches.

This analysis uses the full Run-2 dataset for VBF HWW cross-sections and new statistical methods. These have led to expected results which show that large increases in sensitivity with respect to the most recent Run-1 measurements. The VBF differential cross-sections measured for  $H \rightarrow WW^*\ell\nu\ell\nu$  will be the first analyzed with the ATLAS experiment.

## CHAPTER 7. RESULTS

Observable	Bin Edges
$MT^{l,l,MET}$	15,40,65,80,95,110,115,120,180
$M_{ll}$	0,20,25,30,35,40,45, 50, 60, 70,200
$M_{jj}$	200,450,700,950,1200,1500,2200,3000,5000
$Cos(\theta^*)$	0,0.125,0.25,0.375,0.5,0.625,1.0
$DY_{ll}$	0,0.15,0.3,0.45,0.6,0.75,0.9,1.05,1.2,1.5,4
$DY_{jj}$	0,2.5,3.25,3.62,4,4.35,4.75,5,5.5,6.25,7,8.5
$P_t$ Total	150,200,250,350,500,600,900
$DPhi_{ll}$	0,0.1,0.2,0.3,0.4,0.6,0.8,1.0,1.3,1.7,2.1,2.6,3.2
$P_{tll}$	0,20,40,50,60,70,80,90,100,120,160,500
Higgs $P_t$	0,45,80,120,160,200,260,350,1000
Leading Lepton $P_t$	20,25,35,42,50,65,80,90,2000
Subleading Lepton $P_t$	20,25,35,42,50,65,80,90,2000
Leading Jet $P_t$	30,60,90,120,190,260,350,500
Subleading Jet $P_t$	30,60,90,120,190,260,350

Table 7.6: Planned observables and bin edges for differential cross-section measurements

This analysis has an additional motivation as well. It is the first step in a larger measurement of both the VBF HWW differential cross-section and the Standard Model VBF  $W + jj$  production cross-section. In the  $W +$  jets production in the  $\ell +$  jets final state, the invariant mass of the two jets that accompany the  $W$  bosons and the rapidity separation between them are particularly sensitive to the electroweak symmetry breaking mechanism just as in the VBF HWW cross-section. The final-state jets also have similar kinematics. Therefore the ratio of cross sections measurement allows for the cancellations of correlated systematic uncertainties. As both process are characterized by large (10% – 20%) jet-related uncertainties, a reduction via the ratio cancellation would greatly improve the sensitivity to the VBF Higgs process and to the search for new phenomena in the context of EFT. This ratio measurement falls beyond the scope of this thesis but represents a complementary motivation to the measurements shown here.

# **Chapter 8**

## **Conclusions**

# Bibliography

- [1] M. Tanabashi, K. Hagiwara, K. Hikasa, K. Nakamura, Y. Sumino, F. Takahashi, J. Tanaka, K. Agashe, G. Aielli, C. Amsler, M. Antonelli, D. M. Asner, H. Baer, Sw. Banerjee, R. M. Barnett, T. Basaglia, C. W. Bauer, J. J. Beatty, V. I. Belousov, J. Beringer, S. Bethke, A. Bettini, H. Bichsel, O. Biebel, K. M. Black, E. Blucher, O. Buchmuller, V. Burkert, M. A. Bychkov, R. N. Cahn, M. Carena, A. Cecucci, A. Cerri, D. Chakraborty, M.-C. Chen, R. S. Chivukula, G. Cowan, O. Dahl, G. D'Ambrosio, T. Damour, D. de Florian, A. de Gouv  a, T. DeGrand, P. de Jong, G. Dissertori, B. A. Dobrescu, M. D'Onofrio, M. Doser, M. Drees, H. K. Dreiner, D. A. Dwyer, P. Eerola, S. Eidelman, J. Ellis, J. Erler, V. V. Ezhela, W. Fettscher, B. D. Fields, R. Firestone, B. Foster, A. Freitas, H. Gallagher, L. Garren, H.-J. Gerber, G. Gerbier, T. Gershon, Y. Gershtein, T. Gherghetta, A. A. Godizov, M. Goodman, C. Grab, A. V. Gritsan, C. Grojean, D. E. Groom, M. Grunewald, A. Gurtu, T. Gutsche, H. E. Haber, C. Hanhart, S. Hashimoto, Y. Hayato, K. G. Hayes, A. Hebecker, S. Heinemeyer, B. Heltsley, J. J. Hern  ndez-Rey, J. Hisano, A. H  cker, J. Holder, A. Holtkamp, T. Hyodo, K. D. Irwin, K. F. Johnson, M. Kado, M. Karliner, U. F. Katz, S. R. Klein, E. Klemp, R. V. Kowalewski, F. Krauss, M. Kreps, B. Krusche, Yu. V. Kuyanov, Y. Kwon, O. Lahav, J. Laiho, J. Lesgourgues, A. Liddle, Z. Ligeti, C.-J. Lin, C. Lippmann, T. M. Liss, L. Littenberg, K. S. Lugovsky, S. B. Lugovsky, A. Lusiani, Y. Makida,

## BIBLIOGRAPHY

F. Maltoni, T. Mannel, A. V. Manohar, W. J. Marciano, A. D. Martin, A. Massoni, J. Matthews, U.-G. Meißner, D. Milstead, R. E. Mitchell, K. Mönig, P. Molaro, F. Moortgat, M. Moskovic, H. Murayama, M. Narain, P. Nason, S. Navas, M. Neubert, P. Nevski, Y. Nir, K. A. Olive, S. Pagan Griso, J. Parsons, C. Patrignani, J. A. Peacock, M. Pennington, S. T. Petcov, V. A. Petrov, E. Pianori, A. Piepke, A. Pomarol, A. Quadt, J. Rademacker, G. Raffelt, B. N. Ratcliff, P. Richardson, A. Ringwald, S. Roesler, S. Rolli, A. Romanouk, L. J. Rosenberg, J. L. Rosner, G. Rybka, R. A. Ryutin, C. T. Sachrajda, Y. Sakai, G. P. Salam, S. Sarkar, F. Sauli, O. Schneider, K. Scholberg, A. J. Schwartz, D. Scott, V. Sharma, S. R. Sharpe, T. Shutt, M. Silari, T. Sjöstrand, P. Skands, T. Skwarnicki, J. G. Smith, G. F. Smoot, S. Spanier, H. Spieler, C. Spiering, A. Stahl, S. L. Stone, T. Sumiyoshi, M. J. Syphers, K. Terashi, J. Terning, U. Thoma, R. S. Thorne, L. Tiator, M. Titov, N. P. Tkachenko, N. A. Tornqvist, D. R. Tovey, G. Valencia, R. Van de Water, N. Varelas, G. Venanzoni, L. Verde, M. G. Vincenter, P. Vogel, A. Vogt, S. P. Wakely, W. Walkowiak, C. W. Walter, D. Wands, D. R. Ward, M. O. Wascko, G. Weiglein, D. H. Weinberg, E. J. Weinberg, M. White, L. R. Wiencke, S. Willocq, C. G. Wohl, J. Womersley, C. L. Woody, R. L. Workman, W.-M. Yao, G. P. Zeller, O. V. Zenin, R.-Y. Zhu, S.-L. Zhu, F. Zimmermann, P. A. Zyla, J. Anderson, L. Fuller, V. S. Lugovsky, and P. Schaffner. Review of particle physics. *Phys. Rev. D*, 98:030001, Aug 2018.

- [2] F. Halzen and A. Martin. *Quarks and Leptons: An Introductory Course in Modern Particle Physics*. 1984.
- [3] John Ellis. Higgs Physics. (KCL-PH-TH-2013-49. KCL-PH-TH-2013-49. LCTS-2013-36. CERN-PH-TH-2013-315):117–168. 52 p, Dec 2013. 52 pages, 45 figures, Lectures

## BIBLIOGRAPHY

presented at the ESHEP 2013 School of High-Energy Physics, to appear as part of the proceedings in a CERN Yellow Report.

- [4] Jonathan M. Butterworth, Guenther Dissertori, and Gavin P. Salam. Hard Processes in Proton-Proton Collisions at the Large Hadron Collider. *Ann. Rev. Nucl. Part. Sci.*, 62:387–405, 2012.
- [5] John M. Campbell, J. W. Huston, and W. J. Stirling. Hard Interactions of Quarks and Gluons: A Primer for LHC Physics. *Rept. Prog. Phys.*, 70:89, 2007.
- [6] W.J. Stirling. Private communication. 2018.
- [7] Abdelhak Djouadi. The Anatomy of electro-weak symmetry breaking. I: The Higgs boson in the standard model. *Phys. Rept.*, 457:1–216, 2008.
- [8] D. de Florian, C. Grojean, F. Maltoni, C. Mariotti, A. Nikitenko, M. Pieri, P. Savard, M. Schumacher, R. Tanaka, R. Aggleton, M. Ahmad, B. Allanach, C. Anastasiou, W. Astill, S. Badger, M. Badziak, J. Baglio, E. Bagnaschi, A. Ballestrero, A. Banfi, D. Barducci, M. Beckingham, C. Becot, G. Bélanger, J. Bellm, N. Belyaev, F.U. Bernlochner, C. Beskidt, A. Biekötter, F. Bishara, W. Bizon, N.E. Bomark, M. Bonvini, S. Borowka, V. Bortolotto, S. Boselli, F.J. Botella, R. Boughezal, G.C. Branco, J. Brehmer, L. Brenner, S. Bressler, I. Brivio, A. Broggio, H. Brun, G. Buchalla, C.D. Burgard, A. Calandri, L. Caminada, R. Caminal Armadans, F. Campanario, J. Campbell, F. Caola, C.M. Carloni Calame, S. Carrazza, A. Carvalho, M. Casolino, O. Cata, A. Celis, F. Cerutti, N. Chanon, M. Chen, X. Chen, B. Chokoufé Nejad, N. Christensen, M. Ciuchini, R. Contino, T. Corbett, Raul Costa, D. Curtin, M. Dall’Osso, A. David, S. Dawson, J. de Blas, W. de Boer, P. de Castro Manzano, C. Degrande, R.L. Delgado, F. Demartin, A. Denner, B. Di Micco, R. Di Nardo, S. Dittmaier, A. Dobado, T. Dorigo,

## BIBLIOGRAPHY

F.A. Dreyer, M. Dührssen, C. Duhr, F. Dulat, K. Ecker, K. Ellis, U. Ellwanger, C. Englert, D. Espriu, A. Falkowski, L. Fayard, R. Feger, G. Ferrera, A. Ferroglio, N. Fidanza, T. Figy, M. Flechl, D. Fontes, S. Forte, P. Francavilla, E. Franco, R. Frederix, A. Freitas, F.F. Freitas, F. Frensch, S. Frixione, B. Fuks, E. Furlan, S. Gadatsch, J. Gao, Y. Gao, M.V. Garzelli, T. Gehrmann, R. Gerosa, M. Ghezzi, D. Ghosh, S. Gieseke, D. Gillberg, G.F. Giudice, E.W.N. Glover, F. Goertz, D. Gonçalves, J. Gonzalez-Fraile, M. Gorbahn, S. Gori, C.A. Gottardo, M. Gouzevitch, P. Govoni, D. Gray, M. Grazzini, N. Greiner, A. Greljo, J. Grigo, A.V. Gritsan, R. Gröber, S. Guindon, H.E. Haber, C. Han, T. Han, R. Harlander, M.A. Harrendorf, H.B. Hartanto, C. Hays, S. Heinemeyer, G. Heinrich, M. Herrero, F. Herzog, B. Hespel, V. Hirschi, S. Hoeche, S. Honeywell, S.J. Huber, C. Hugonie, J. Huston, A. Ilnicka, G. Isidori, B. Jäger, M. Jaquier, S.P. Jones, A. Juste, S. Kallweit, A. Kaluza, A. Kardos, A. Karlberg, Z. Kassabov, N. Kauer, D.I. Kazakov, M. Kerner, W. Kilian, F. Kling, K. Köneke, R. Kogler, R. Konoplich, S. Kortner, S. Kraml, C. Krause, F. Krauss, M. Krawczyk, A. Kulesza, S. Kuttimalai, R. Lane, A. Lazopoulos, G. Lee, P. Lenzi, I.M. Lewis, Y. Li, S. Liebler, J. Lindert, X. Liu, Z. Liu, F.J. Llanes-Estrada, H.E. Logan, D. Lopez-Val, I. Low, G. Luisoni, P. Maierhöfer, E. Maina, B. Mansoulié, H. Mantler, M. Mantoani, A.C. Marini, V.I. Martinez Outschoorn, S. Marzani, D. Marzocca, A. Massironi, K. Mawatari, J. Mazzitelli, A. McCarn, B. Mellado, K. Melnikov, S.B. Menari, L. Merlo, C. Meyer, P. Milenovic, K. Mimasu, S. Mishima, B. Mistlberger, S.-O. Moch, A. Mohammadi, P.F. Monni, G. Montagna, M. Moreno Llácer, N. Moretti, S. Moretti, L. Motyka, A. Mück, M. Mühlleitner, S. Munir, P. Musella, P. Nadolsky, D. Napoletano, M. Nebot, C. Neu, M. Neubert, R. Nevzorov, O. Nicrosini, J. Nielsen, K. Nikolopoulos, J.M. No, C. O'Brien, T. Ohl, C. Oleari, T. Orimoto, D. Pagani, C.E. Pandini, A. Papaefstathiou, A.S. Papanastasiou, G. Passarino, B.D. Pecjak, M. Pelliccioni, G. Perez, L. Per-

## BIBLIOGRAPHY

rozzi, F. Petriello, G. Petrucciani, E. Pianori, F. Piccinini, M. Pierini, A. Pilkington, S. Plätzer, T. Plehn, R. Podskubka, C.T. Potter, S. Pozzorini, K. Prokofiev, A. Pukhov, I. Puljak, M. Queitsch-Maitland, J. Quevillon, D. Rathlev, M. Rauch, E. Re, M.N. Rebelo, D. Rebuzzi, L. Reina, C. Reuschle, J. Reuter, M. Riembau, F. Riva, A. Rizzi, T. Robens, R. Röntsch, J. Rojo, J.C. Romão, N. Rompotis, J. Roskes, R. Roth, G.P. Salam, R. Salerno, M.O.P. Sampaio, R. Santos, V. Sanz, J.J. Sanz-Cillero, H. Sargsyan, U. Sarica, P. Schichtel, J. Schlenk, T. Schmidt, C. Schmitt, M. Schönher, U. Schubert, M. Schulze, S. Sekula, M. Sekulla, E. Shabalina, H.S. Shao, J. Shelton, C.H. Shepherd-Themistocleous, S.Y. Shim, F. Siegert, A. Signer, J.P. Silva, L. Silvestrini, M. Sjodahl, P. Slavich, M. Slawinska, L. Soffi, M. Spannowsky, C. Speckner, D.M. Sperka, M. Spira, O. Stål, F. Staub, T. Stebel, T. Stefaniak, M. Steinhauser, I.W. Stewart, M.J. Strassler, J. Streicher, D.M. Strom, S. Su, X. Sun, F.J. Tackmann, K. Tackmann, A.M. Teixeira, R. Teixeira de Lima, V. Theeuwes, R. Thorne, D. Tommasini, P. Torrielli, M. Tosi, F. Tramontano, Z. Trócsányi, M. Trott, I. Tsinikos, M. Ubiali, P. Vanlaer, W. Verkerke, A. Vicini, L. Viliani, E. Vryonidou, D. Wackerlo, C.E.M. Wagner, J. Wang, S. Wayand, G. Weiglein, C. Weiss, M. Wiesemann, C. Williams, J. Winter, D. Winterbottom, R. Wolf, M. Xiao, L.L. Yang, R. Yohay, S.P.Y. Yuen, G. Zanderighi, M. Zaro, D. Zeppenfeld, R. Ziegler, T. Zirke, and J. Zupan. *Handbook of LHC Higgs Cross Sections: 4. Deciphering the Nature of the Higgs Sector*. CERN Yellow Reports: Monographs. Oct 2016. 869 pages, 295 figures, 248 tables and 1645 citations. Working Group web page: <https://twiki.cern.ch/twiki/bin/view/LHCPhysics/LHCHXSWG>.

- [9] Jon Butterworth et al. PDF4LHC recommendations for LHC Run II. *J. Phys.*, G43:023001, 2016.
- [10] Georges Aad et al. Observation of a new particle in the search for the Standard Model

## BIBLIOGRAPHY

- Higgs boson with the ATLAS detector at the LHC. *Phys. Lett.*, B716:1–29, 2012.
- [11] Tom W. B. Kibble. The Standard Model of Particle Physics. 2014.
- [12] N.R. Council, D.E.P. Sciences, B.P. Astronomy, and C.E.P. Physics. *Elementary-Particle Physics: Revealing the Secrets of Energy and Matter*. Physics in a New Era: A Series. National Academies Press, 1998.
- [13] Georges Aad et al. Combined measurements of Higgs boson production and decay using up to  $80 \text{ fb}^{-1}$  of proton-proton collision data at  $\sqrt{s} = 13 \text{ TeV}$  collected with the ATLAS experiment. *Phys. Rev.*, D101(1):012002, 2020.
- [14] Georges Aad et al. Measurement of fiducial differential cross sections of gluon-fusion production of Higgs bosons decaying to  $WW^* \rightarrow e\nu\mu\nu$  with the ATLAS detector at  $\sqrt{s} = 8 \text{ TeV}$ . *JHEP*, 08:104, 2016.
- [15] M. Aaboud, G. Aad, B. Abbott, O. Abdinov, B. Abeloos, D.K. Abhayasinghe, S.H. Abidi, O.S. AbouZeid, N.L. Abraham, H. Abramowicz, and et al. Measurements of gluon-gluon fusion and vector-boson fusion higgs boson production cross-sections in the  $h \rightarrow ww^* \rightarrow e\nu\mu\nu$  decay channel in  $pp$  collisions at  $\sqrt(s)=13\text{tev}$  with the atlas detector. *Physics Letters B*, 789:508–529, Feb 2019.
- [16] Rende Steerenberg. Lhc report: Protons: mission accomplished, 2018.
- [17] M. Aaboud, G. Aad, B. Abbott, O. Abdinov, B. Abeloos, D.K. Abhayasinghe, S.H. Abidi, O.S. AbouZeid, N.L. Abraham, H. Abramowicz, and et al. Observation of  $h \rightarrow bb^-$  decays and  $vh$  production with the atlas detector. *Physics Letters B*, 786:59–86, Nov 2018.

## BIBLIOGRAPHY

- [18] M. Aaboud, G. Aad, B. Abbott, O. Abdinov, B. Abelos, S. H. Abidi, O. S. AbouZeid, N. L. Abraham, H. Abramowicz, and et al. Measurement of the higgs boson coupling properties in the  $h \rightarrow zz^* \rightarrow 4\ell$  decay channel at  $\sqrt{s} = 13$  tev with the atlas detector. *Journal of High Energy Physics*, 2018(3), Mar 2018.
- [19] M. Aaboud, G. Aad, B. Abbott, O. Abdinov, B. Abelos, S. H. Abidi, O. S. AbouZeid, N. L. Abraham, H. Abramowicz, H. Abreu, and et al. Measurements of higgs boson properties in the diphoton decay channel with  $36\text{ fb}^{-1}$  of  $pp$  collision data at  $\sqrt{s} = 13$  tev with the atlas detector. *Physical Review D*, 98(5), Sep 2018.
- [20] M. Aaboud, G. Aad, B. Abbott, O. Abdinov, B. Abelos, D. K. Abhayasinghe, S. H. Abidi, O. S. AbouZeid, N. L. Abraham, H. Abramowicz, and et al. Cross-section measurements of the higgs boson decaying into a pair of  $\tau$ -leptons in proton-proton collisions at  $\sqrt{s}=13$  tev with the atlas detector. *Physical Review D*, 99(7), Apr 2019.
- [21] Search for squarks and gluinos in final states with jets and missing transverse momentum using  $139\text{ fb}^{-1}$  of  $\sqrt{s} = 13$  TeV  $pp$  collision data with the ATLAS detector. Technical Report ATLAS-CONF-2019-040, CERN, Geneva, Aug 2019.
- [22] CERN. A new schedule for the lhc and its successor, 2019.
- [23] M. Benedikt, P. Collier, V. Mertens, J. Poole, and K. Schindl. LHC Design Report. 3. The LHC injector chain. 2004.
- [24] Radiofrequency cavities. Sep 2012.
- [25] Lyndon Evans. The large hadron collider. *New Journal of Physics*, 9(9):335–335, sep 2007.

## BIBLIOGRAPHY

- [26] G Aad et al. The ATLAS Experiment at the CERN Large Hadron Collider. *JINST*, 3:S08003. 437 p, 2008. Also published by CERN Geneva in 2010.
- [27] Apollinari G., Béjar Alonso I., Brüning O., Fessia P., Lamont M., Rossi L., and Tavian L. *High-Luminosity Large Hadron Collider (HL-LHC): Technical Design Report V. 0.1*. CERN Yellow Reports: Monographs. CERN, Geneva, 2017.
- [28] ATLAS Collaboration. Technical Design Report for the ATLAS Inner Tracker Strip Detector. Technical Report CERN-LHCC-2017-005. ATLAS-TDR-025, CERN, Geneva, Apr 2017.
- [29] Early Inner Detector Tracking Performance in the 2015 data at  $\sqrt{s} = 13$  TeV. Technical Report ATL-PHYS-PUB-2015-051, CERN, Geneva, Dec 2015.
- [30] Performance of the ATLAS Silicon Pattern Recognition Algorithm in Data and Simulation at  $\sqrt{s} = 7$  TeV. Technical Report ATLAS-CONF-2010-072, CERN, Geneva, Jul 2010.
- [31] ATLAS Collaboration. 14(12):P12006–P12006, dec 2019.
- [32] G. Aad, B. Abbott, J. Abdallah, O. Abdinov, B. Abelos, R. Aben, M. Abolins, O. S. AbouZeid, N. L. Abraham, and et al. Muon reconstruction performance of the atlas detector in proton–proton collision data at  $\sqrt{s}=13$  tev. *The European Physical Journal C*, 76(5), May 2016.
- [33] Georges Aad et al. Measurement of the muon reconstruction performance of the ATLAS detector using 2011 and 2012 LHC proton–proton collision data. Measurement of the muon reconstruction performance of the ATLAS detector using 2011 and 2012 LHC proton-proton collision data. *Eur. Phys. J. C*,

## BIBLIOGRAPHY

- 74(CERN-PH-EP-2014-151. CERN-PH-EP-2014-151):3130. 34 p, Jul 2014. Comments: 21 pages plus author list + cover pages (34 pages total), 21 figures, 2 tables, submitted to EPJC, All figures including auxiliary figures are available at <http://atlas.web.cern.ch/Atlas/GROUPS/PHYSICS/PAPERS/PERF-2014-05/>.
- [34] Johannes Josef Junggeburth. Muon identification and reconstruction efficiencies in full Run-2 dataset. Technical Report ATL-COM-PHYS-2019-1001, CERN, Geneva, Aug 2019.
- [35] G. Aad, B. Abbott, J. Abdallah, A. A. Abdelalim, A. Abdesselam, O. Abdinov, B. Abi, M. Abolins, H. Abramowicz, and et al. Jet energy measurement with the atlas detector in proton-proton collisions at  $\sqrt{s} = 7$  tev. *The European Physical Journal C*, 73(3), Mar 2013.
- [36] Miha Zgubič, Johannes Josef Junggeburth, and K. Technical report.
- [37] Jet global sequential corrections with the ATLAS detector in proton-proton collisions at  $\sqrt{s} = 8$  TeV. Technical Report ATLAS-CONF-2015-002, CERN, Geneva, Mar 2015.
- [38] Morad Aaboud et al. Jet reconstruction and performance using particle flow with the ATLAS Detector. *Eur. Phys. J.*, C77(7):466, 2017.
- [39] JET ETmiss group. Public plots, 2019.
- [40] Matteo Cacciari, Gavin P. Salam, and Gregory Soyez. The anti- $k_t$  jet clustering algorithm. *JHEP*, 04:063, 2008.
- [41] Morad Aaboud et al. Determination of jet calibration and energy resolution in proton-proton collisions at  $\sqrt{s} = 8$  TeV using the ATLAS detector. 2019.

## BIBLIOGRAPHY

- [42] Object-based missing transverse momentum significance in the ATLAS detector. Technical Report ATLAS-CONF-2018-038, CERN, Geneva, Jul 2018.
- [43] Morad Aaboud et al. Performance of missing transverse momentum reconstruction with the ATLAS detector using proton-proton collisions at  $\sqrt{s} = 13$  TeV. *Eur. Phys. J.*, C78(11):903, 2018.
- [44] Luminosity public results. Public plots, 2019.
- [45] S. Agostinelli et al. GEANT4: A Simulation toolkit. *Nucl. Instrum. Meth. A*, 506:250, 2003.
- [46] Simone Alioli, Paolo Nason, Carlo Oleari, and Emanuele Re. A general framework for implementing NLO calculations in shower Monte Carlo programs: the POWHEG BOX. *JHEP*, 06:043, 2010.
- [47] M. Ciccolini, Ansgar Denner, and S. Dittmaier. Strong and electroweak corrections to the production of Higgs+2jets via weak interactions at the LHC. *Phys. Rev. Lett.*, 99:161803, 2007.
- [48] K. Arnold et al. VBFNLO: A parton level Monte Carlo for processes with electroweak bosons. *Comput. Phys. Commun.*, 180:1661–1670, 2009.
- [49] Paolo Bolzoni, Fabio Maltoni, Sven-Olaf Moch, and Marco Zaro. Higgs production via vector-boson fusion at NNLO in QCD. *Phys. Rev. Lett.*, 105:011801, 2010.
- [50] Torbjorn Sjostrand, Stephen Mrenna, and Peter Z. Skands. PYTHIA 6.4 Physics and Manual. *JHEP*, 05:026, 2006.
- [51] Torbjorn Sjostrand, Stephen Mrenna, and Peter Z. Skands. A Brief Introduction to PYTHIA 8.1. *Comput. Phys. Commun.*, 178:852–867, 2008.

## BIBLIOGRAPHY

- [52] A. Hoecker, P. Speckmayer, J. Stelzer, J. Therhaag, E. von Toerne, H. Voss, M. Backes, T. Carli, O. Cohen, A. Christov, D. Dannheim, K. Danielowski, S. Henrot-Versille, M. Jachowski, K. Kraszewski, A. Krasznahorkay Jr., M. Kruk, Y. Mahalalel, R. Ospanov, X. Prudent, A. Robert, D. Schouten, F. Tegenfeldt, A. Voigt, K. Voss, M. Wolter, and A. Zemla. Tmva - toolkit for multivariate data analysis, 2007.
- [53] Travis Martin and W Martin. Analyzing new neutral gauge bosons at the lhc using third generation fermions. 05 2020.
- [54] Fabio Cardillo, Otilia Anamaria Ducu, Julien Maurer, Claudia Merlassino, Joseph Reichert, Erich Varnes, Pepijn Johannes Bakker, and Marcus Morgenstern. Tools for estimating fake/non-prompt lepton backgrounds in ATLAS. Technical Report ATL-COM-PHYS-2019-1071, CERN, Geneva, Sep 1999.
- [55] Luminosity determination in  $pp$  collisions at  $\sqrt{s} = 13$  TeV using the ATLAS detector at the LHC. Technical Report ATLAS-CONF-2019-021, CERN, Geneva, Jun 2019.
- [56] Wg1 scheme: meeting discussions.
- [57] Stefano Frixione, Eric Laenen, Patrick Motylinski, Chris White, and Bryan R Webber. Single-top hadroproduction in association with awboson. *Journal of High Energy Physics*, 2008(07):029–029, Jul 2008.
- [58] Kyle Cranmer. Practical statistics for the lhc, 2015.
- [59] G. D’Agostini. A Multidimensional unfolding method based on Bayes’ theorem. *Nucl. Instrum. Meth. A*, 362:487–498, 1995.