

2D Array - DS

Given a 6×6 2D Array *arr*:

```
1 1 1 0 0 0
0 1 0 0 0 0
1 1 1 0 0 0
0 0 0 0 0 0
0 0 0 0 0 0
0 0 0 0 0 0
```

We define an hourglass in *A* to be a subset of values with indices falling in this pattern in *arr*'s graphical representation:

```
a b c
  d
e f g
```

There are **16** hourglasses in *arr*, and an hourglass sum is the sum of an hourglass' values. Calculate the hourglass sum for every hourglass in *arr*, then print the maximum hourglass sum.

For example, given the 2D array:

```
-9 -9 -9 1 1 1
0 -9 0 4 3 2
-9 -9 -9 1 2 3
0 0 8 6 6 0
0 0 0 -2 0 0
0 0 1 2 4 0
```

We calculate the following **16** hourglass values:

```
-63, -34, -9, 12,
-10, 0, 28, 23,
-27, -11, -2, 10,
9, 17, 25, 18
```

Our highest hourglass value is **28** from the hourglass:

```
0 4 3
  1
8 6 6
```

Note: If you have already solved the Java domain's `Java 2D Array` challenge, you may wish to skip this challenge.

Function Description

Complete the function `hourglassSum` in the editor below. It should return an integer, the maximum hourglass sum in the array.

`hourglassSum` has the following parameter(s):

- `arr` an array of integers

Input Format

Each of the **6** lines of inputs *arr*[*i*] contains **6** space-separated integers *arr*[*i*][*j*].

Constraints

- $-9 \leq arr[i][j] \leq 9$
- $0 \leq i, j \leq 5$

Output Format

Print the largest (maximum) hourglass sum found in *arr*.

Sample Input

```
1 1 1 0 0 0
0 1 0 0 0 0
1 1 1 0 0 0
0 0 2 4 4 0
0 0 0 2 0 0
0 0 1 2 4 0
```

Sample Output

```
19
```

Explanation

arr contains the following hourglasses:

```
1 1 1 1 1 0 1 0 0 0 0 0
1 0 0 0
1 1 1 1 1 0 1 0 0 0 0 0

0 1 0 1 0 0 0 0 0 0 0 0
1 1 0 0
0 0 2 0 2 4 2 4 4 4 4 0

1 1 1 1 1 0 1 0 0 0 0 0
0 2 4 4
0 0 0 0 0 2 0 2 0 2 0 0

0 0 2 0 2 4 2 4 4 4 4 0
0 0 2 0
0 0 1 0 1 2 1 2 4 2 4 0
```

The hourglass with the maximum sum (**19**) is:

```
2 4 4
2
1 2 4
```