Zombie model with human behavior equations

Luke Botti Course number: MATH 502

December 8, 2020

$$\frac{dS}{dt} = bS(t) - mS(t) - \alpha S(t)Z(t) + r(Q(t) + E(t))$$

$$C(t) = \begin{cases} 0 & Q(t) < q_{\text{max}} \lor X_S(t) < 1/2 \\ C_0 & Q(t) \ge q_{\text{max}} \land X_S(t) \ge 1/2 \end{cases}$$

$$\frac{dE}{dt} = \begin{cases} \alpha S(t)Z(t) - zE(t) - rE(t) - (C_0 + X_E(t))E(t) - C(t)E(t) & Q(t) < q_{\text{max}} \\ \alpha S(t)Z(t) - zE(t) - rE(t) - C(t)E(t) & Q(t) < q_{\text{max}} \end{cases}$$

$$\frac{dQ}{dt} = \begin{cases} (C_0 + X_E)E - rQ(t) - zQ(t) & Q(t) < q_{\text{max}} \\ rQ(t) - zQ(t) & Q(t) \ge q_{\text{max}} \end{cases}$$

$$\frac{dZ}{dt} = zE(t) - kS(t)Z(t)$$

$$\frac{dD}{dt} = kS(t)Z(t) + mS(t) + zQ(t) + C(t)E$$

$$\frac{dX_S}{dt} = k_SX_S(t)(1 - X_S(t))[Z(t) + Q(t) - \epsilon_SL_S(t)]$$

$$\frac{dX_E}{dt} = \begin{cases} k_IX_E(t)(1 - X_E(t))\{(z - r)[E(t) + Q(t)] - \epsilon_I\} & Q(t) < q_{\text{max}} \\ Q(t) \ge q_{\text{max}} \end{cases}$$

$$\frac{dL_S}{dt} = \beta C(t) - \mu L_S(t)$$