

Informational Dissipative Structures

Relative-Entropy Slope Geometry and Resolution Ledgers

Laurent Caraffa

Univ. Gustave Eiffel, IGN-ENSG, LaSTIG

Preliminary version — comments welcome

Abstract

We propose Informational Dissipative Structures (IDS), a geometry-first architecture for modelling physical systems from finite observations. The primitive is relative information to a declared reference, read on three distinct spaces. On process paths it measures irreversibility. On observable laws, every channel obeys an exact visible/hidden ledger, whose second-order differential is the Fisher-information ledger of the quotient of experimentally identifiable laws. On probability measures over that quotient, Bayesian updating is an information projection, and the Wasserstein slope of relative entropy defines a calibrated geometry of belief revision, the relative-entropy slope geometry (RESG). The kernel is free of any graph or clique structure: finite resolution is an ordered family of channels, each level carrying a resolution ledger, visible plus hidden. Along an exhaustive tower the ledger recovers the fine values of relative information, Fisher information and Bayesian information monotonically, and so does the calibrated radius under the radial hypotheses; an almost-sufficient projection is almost isometric; a finite description closes when its predictive ledger says so. Graphs, cells and symbols enter only in realizations of finite resolution. The Brownian gyrator, a driven ring and a metastable double well separate path irreversibility, local relative-information balance, thermodynamic entropy production and predictive closure. The contribution is the assembly of classical ingredients into one reference-explicit reduction calculus with a common information ledger.

1 Introduction

The problem. Scientific models are finite descriptions of systems richer than any description. The description may be a sensor stream, a reduced state, a graph, an alphabet of symbols or a parametric family. The usual question is whether the description predicts well. We ask a second question: what information did the reduction hide, and can that loss be carried consistently from the physical process to the geometry of the model and to the observer’s belief?

The thesis. Informational Dissipative Structures (IDS) starts from a deliberately small datum: candidate processes, declared observation channels and declared references. A channel first determines which candidates are experimentally indistinguishable. On a smooth stratum of the identifiable quotient, relative entropy then provides the local Fisher ruler. Probability measures over that stratum describe residual uncertainty, and the Wasserstein slope of relative entropy supplies a calibrated length for revising those beliefs. A finite structure enters afterwards, as one more channel in a resolution tower, together with the exact ledger of what it hides. The chain of the paper is

process $\rightarrow$ channel $\rightarrow$ identifiable law $\rightarrow$ geometry $\rightarrow$ resolution $\rightarrow$ finite structure,

and the kernel is free of any graph or clique structure (Figure 1). A model is a channel-relative geometry first; a finite structure is accepted together with the ledger of what its resolution hides.

Uncertainty and information. Bayesian inference represents what is known about a law λ as a prior π and updates it with data $\mathcal{D}$ of likelihood $L_{\mathcal{D}}$ into a posterior $\mu_{\mathcal{D}} \propto L_{\mathcal{D}} \pi$ (Bayes, 1763), the law of least relative entropy to the prior under the evidence (Williams, 1980; Shore and Johnson, 1980; Zellner, 1988); Lindley (1956) measures the information an experiment provides as the relative entropy of the posterior from the prior, averaged over the outcome. Shannon’s source of law p carries the entropy $H(X) = -\sum_x p(x) \log p(x)$, in nats; the mutual information




Figure 1: The architecture. One primitive, relative information to a declared reference, read on three spaces and differentiated twice: the Hessian in the space of identifiable laws gives the Fisher metric h , the Wasserstein slope on beliefs gives the Fisher functional $\mathcal{I}_{h,m}$, and the two meet in the relative-entropy slope geometry; a tower of channels carries the visible and hidden ledgers down to cells, graphs and symbols.

$I(X; Y) = \sum_{x,y} p(x, y) \log \frac{p(x,y)}{p(x)p(y)}$ measures what an output tells about an input and stays equal or decreases under post-processing; the rate–distortion function $R(D) = \inf\{I(X; \hat{X}) : \mathbb{E}\|X - \hat{X}\|^2 \leq D\}$ is the least information a reconstruction of distortion at most D carries about the source (Shannon, 1948, 1959; Cover and Thomas, 2006).

Distinguishing and moving laws. The relative entropy of a law P from a law Q , $D(P \parallel Q) = \int \log \frac{dP}{dQ} dP \geq 0$, vanishes exactly when $P = Q$ and stays equal or decreases under every channel, with equality under a sufficient one (Kullback and Leibler, 1951; Csiszár, 1967; Blackwell (1953) orders experiments by garbling, and the projection of Csiszár (1975) selects, in a convex set, the law of least relative entropy to a reference. Between neighbouring laws of a smooth family, relative entropy is at second order a quadratic form, the Fisher information $D(p_\theta \parallel p_{\theta+d\theta}) = \frac{1}{2}g_{ij}(\theta)d\theta^i d\theta^j + o(\|d\theta\|^2)$, which Čencov (1982) characterized, up to a factor, as the metric invariant under sufficient Markov morphisms (Amari and Nagaoka, 2000; Ay et al., 2017). The quadratic Wasserstein distance between two probability measures is the least kinetic energy of a flow carrying one onto the other, $W_2^2(\mu_0, \mu_1) = \inf \int_0^1 \int \|v_t\|^2 d\mu_t dt$ under $\partial_t \mu_t + \nabla \cdot (\mu_t v_t) = 0$ (Benamou and Brenier, 2000; Villani, 2009); in that geometry the heat equation is the gradient flow of the entropy $\int \rho \log \rho$ (Jordan et al., 1998; Otto, 2001), the entropy decreases at the rate of the Fisher information of ρ , de Bruijn’s identity (Stam, 1959), and the relative entropy to a reference whose Bakry–Émery curvature is bounded below by K is K -convex along displacement interpolation (Bakry and Émery, 1985; McCann, 1997; von Renesse and Sturm, 2005), with a squared metric slope equal to the relative Fisher information (Otto and Villani, 2000; Ambrosio et al., 2008).

Organization away from equilibrium: dissipative structures. A dissipative structure is an organization maintained away from equilibrium by fluxes exchanged with reservoirs (Nicolis and Prigogine, 1977). At equilibrium every transition is balanced by its reverse and every current vanishes; a structure embedded in reservoirs at distinct parameters sustains stationary




Figure 2: The Brownian gyrator. (a) Two coordinates, one coupling spring, two walls, two baths. (b) The stationary probability current at $T_1 = 2$, $T_2 = 0.5$ (orange) and the level sets of the stationary density (grey). (c) The entropy production and the heat delivered to the hot bath against the temperature gap, for $k > 0$: the structure exists exactly when $k(T_1 - T_2) \neq 0$.

currents and produces entropy. Stochastic thermodynamics makes this quantitative: on a Markov process with rates k_{ij} and stationary law p the entropy production is $\sigma = \frac{1}{2} \sum_{ij} J_{ij} A_{ij} \geq 0$ with $J_{ij} = p_i k_{ij} - p_j k_{ji}$ and $A_{ij} = \log \frac{p_i k_{ij}}{p_j k_{ji}}$, current times affinity (Schnakenberg, 1976); on a path ω of a process of law P and reversed law $P^\dagger$ the entropy production is $\Sigma(\omega) = \log(dP/dP^\dagger)(\omega)$, whose exponential averages to one and whose mean is a relative entropy (Seifert, 2005, 2012; Esposito and Van den Broeck, 2010); local detailed balance ties the asymmetry of each transition to the entropy received by the reservoir that mediates it.

The Brownian gyrator, a running benchmark. A minimal, exactly solvable nonequilibrium heat-engine model (Filliger and Reimann, 2007) has two overdamped coordinates x_1, x_2 coupled by a spring of stiffness k , each tied to a wall by a spring k_1, k_2 and each immersed in its own bath at temperature T_1, T_2 :

$$dx_1 = [-k_1 x_1 - k(x_1 - x_2)] dt + \sqrt{2T_1} dW_1, \quad dx_2 = [-k_2 x_2 - k(x_2 - x_1)] dt + \sqrt{2T_2} dW_2.$$

Everything is Gaussian and in closed form: the stationary covariance solves a Lyapunov equation and, for $k > 0$ and $T_1 \neq T_2$, the stationary state carries a circulating probability current, heat flows from the hot bath to the cold one through the spring and entropy is produced at a rate proportional to $k^2(T_1 - T_2)^2$; at $k = 0$ or at equal temperatures the current and the production vanish (Figure 2). The gyrator is the object on which the kernel is stress-tested; a driven ring and a metastable double well complete it where the gyrator is degenerate.

Contributions. The ingredients are established; the contribution is their assembly into one reference-explicit calculus of reduction, organized around four pillars and three consequences. The twelve theorems of the body are named here, with their identifier, their status and their section.

- **A, the channel ledger.** Every channel splits relative information into a visible and a hidden part, exactly, and the ledger characterizes relative entropy.
 - T1 · Declared-Reference Principle (K0) · principle · Section 3
 - T2 · Channel Ledger Theorem (K2, with C1) · imported · Section 3
 - T3 · Retention Theorem (K3) · imported, proof recalled · Section 3
 - T4 · Recovery-Kernel Theorem (K4) · imported · Section 3
- **B, the differential ledger.** On the quotient of identifiable laws the second-order differential of the ledger is the Fisher metric of the observation, and the missing Fisher information is the expected conditional variance of the score.
 - T5 · Operational Fisher Theorem (O1) · imported · Section 4
 - T6 · Missing-Fisher Ledger Theorem (O3) · proved · Section 4
- **C, the slope geometry.** On a length space the metric slope of a lower semicontinuous functional is a ruler with a floor; for relative entropy under Wasserstein transport the ruler

is the Fisher functional, certainty lies on a horizon, the weight is pointwise rigid and the prior-centred distance is relative entropy itself.

- T7 · Slope-Geometry Proposition · proved · Section 4
- T8 · RESG Theorem: Entropy Floor, Constant Slope, Radial Identity (G3, G4, G5) · proved · Section 4
- **D, resolution recovery.** Along an exhaustive tower of channels, relative information, Fisher information, Bayesian information and calibrated radius recover their fine values monotonically.
 - T9 · Resolution Ledger Theorem (R1, R2) · proved · Section 5
 - T10 · Resolution Recovery Theorem (R5) · proved · Section 5
- **Three consequences.** A δ -almost sufficient projection is a δ -bi-Lipschitz map of the slope geometry; a finite description obeys an exact predictive ledger, $I_{\text{pred}} = I_{\text{ret}} + I_{\text{hid}}$, which defines its closure; an informational dissipative structure is a realization that closes, is irreversible and is sustained by declared ports.
 - T11 · Almost-Sufficient Stability Theorem (G10, with its corollary G11) · proved · Section 5
 - T12 · Predictive Closure Ledger Theorem (E6) · proved · Section 6
 - Informational Dissipative Structure (H1) · definition; current and ports measured on the gyrator and the driven ring, closure on the double well · Section 7

Organization. A statement lives at one of two levels, printed after its identifier: the *kernel* is what follows from the declared datum in the language of relative information alone; a *realization* instantiates the kernel on a named space, where statements gain their closed forms; a *projection*, a resolution of a realization, keeps the map to the realization and the ledger of what it hides, and is stated at the level of the realization it resolves. A numerical check on a benchmark is cited as a gate, such as X1.1; every gate was declared in writing before its first computation, and Appendix S.D lists them with their measured values. Each statement prints its level and its status: IMPORTED names a standard theorem with its reference, PROVED a proof in this paper or its supplement, MEASURED a fact established on a realization by gates, PRINCIPLE and DEFINITION what they say. Section 2 fixes the primitive, the three spaces and what is primitive; Section 3 states the formal core; Section 4 the two differentials of the primitive; Section 5 resolution as an ordered family of channels; Section 6 the emergent finite structure; Section 7 the physical realization and the numerical experiments; Section 8 the relation to prior work, the scope and the limitations; Section 9 concludes. The twelve theorems T1 to T12 carry the body; every other statement is stated in the supplement (Appendix S.C).

2 One primitive, three spaces

Problem. Describe, from relative information to declared references, the laws an observer can tell apart, the beliefs over those laws and the cost of revising them, the information that every finite description hides, and the structure that a driven physical process maintains, in one architecture whose every statement is tested on a physical object.

The relative information of a law P from a declared reference R ,

$$D(P \parallel R) = \int \log \frac{dP}{dR} dP \geq 0, \quad (2.1)$$

is the primitive. A problem is a datum $\mathfrak{J} = (\mathcal{M}_{\text{adm}}, \mathcal{S}, \mathcal{E}, R, \pi; m_{\Lambda})$: the admissible candidates, their semantics, the operational access $\mathcal{E}$ (the declared experiments), the path reference R , the prior π and, when the geometry is invoked, the calibration measure m_{Λ} on a stratum. A problem is well posed when its references are declared. The primitive is read on three spaces (Table 1).

Two objects share a symbol in much of the literature and are typed apart here: the ensemble law $p_t \in \mathcal{P}(\mathbf{X})$ of a physical state, read against a reference $m_{\mathbf{X}}$ on the state space, and a belief

space	object	declared reference	channel	reading of D
physical paths	path law P	reversal $P^\dagger$, path reference R	observation of paths	irreversibility (K5)
identifiable laws	$K_\lambda, \lambda \in \bar{\Lambda}_\mathcal{E}$	the law of a neighbouring class	experimental channel K	distinguishability and its ledger (K2, O1)
beliefs on laws	$\mu \in \mathcal{P}(\bar{\Lambda}_\mathcal{E})$	prior π , calibration m_Λ	Bayesian experiment	update, information geometry (K3, G3)

Table 1: One primitive read on three spaces, each with its declared reference and its channel.

$\mu \in \mathcal{P}(\bar{\Lambda}_\mathcal{E})$ over identifiable laws, read against π or m_Λ . The slope construction of Section 4 is abstract before it is epistemic: in the informational reading it acts on beliefs over identifiable laws; in the physical benchmarks the same construction is instantiated on ensemble laws over a state space. The two uses share the calculus and differ in semantics. On beliefs the reading is a geometry: the transport distance built on the law ruler h , weighted by the slope of relative entropy,

$$d\ell_1 = \ell_\star \sqrt{\mathcal{I}_{h,m}} d\ell_{W_{2,h}}$$

the *relative-entropy slope geometry* (RESG, T8, Section 4.2), where $\ell_\star$ is the length of one nat, $\mathcal{I}_{h,m}$ the relative Fisher information of a belief and $W_{2,h}$ the quadratic transport distance built on h (Section 4.2); a finite description is a channel in a tower, each level carrying its *resolution ledger* (T9, Section 5).

The path reading. On the paths of a process the primitive reads irreversibility: when a reversal $P^\dagger$ equivalent to P exists,

$$\Sigma = \log \frac{dP}{dP^\dagger}, \quad \mathbb{E}_P[e^{-\Sigma}] = 1, \quad \mathbb{E}_P[\Sigma] = D(P \parallel P^\dagger) \geq 0, \quad (2.2)$$

and every quotient of the paths that commutes with time reversal reads at most the fine irreversibility, $\langle \Sigma_{\text{visible}} \rangle = D(KP \parallel (KP)^\dagger) \leq \langle \Sigma_{\text{fine}} \rangle$ (K5, a consequence of the ledger of Section 3; Kawai et al., 2007; Gomez-Marin et al., 2008). On the gyrator every scalar linear observable is a stationary Gaussian process, hence reversible: an observer of one coordinate reads zero arrow, the observer of both reads σ (Section 7).

What is primitive. Table 2 says in one glance what is declared, what is derived and what belongs to a realization.

3 The formal core

3.1 Declared data

Four definitions fix the objects that the principles use; the notation follows, and every statement carries a status.

Definition D1 (Processes and paths · kernel · DEFINITION). A *process* is a probability law on the paths of a state space $\mathbf{X}$, with the family of its finite-dimensional laws. Its *path law* is written P ; its *reversal* $P^\dagger$ is the law of the time-reversed paths, a probability law equivalent to P when it exists. Its ensemble law at time t is $p_t \in \mathcal{P}(\mathbf{X})$. Information is counted in nats.

Definition D2 (Observable laws and strata · kernel · DEFINITION). Under an experiment $e = (a, K)$ of the operational access $\mathcal{E}$, an action and a channel, the *observable law* of a candidate M is $p_M(\cdot \mid e)$. For a class λ of the identifiable quotient $\bar{\Lambda}_\mathcal{E}$ (K1), K_λ denotes the observable law of the class under the channel K , the same for every M in the class. A *stratum* is a smooth part of $\bar{\Lambda}_\mathcal{E}$, a *regular stratum* a stratum on which the signature is differentiable; h is the Fisher metric of the observable laws on a regular stratum (O1), and $\preceq$ the order of quadratic forms.

object	primitive	derived	realization-dependent
relative information to a declared reference	yes		
observation channel, operational access $\mathcal{E}$	declared		
identifiable quotient $\bar{\Lambda}_{\mathcal{E}}$		yes (K1)	
Fisher metric h		yes (O1)	
Bayesian posterior		yes, under retention (K3)	
relative-entropy slope geometry		yes, under HR0 (G3)	
resolution tower	declared, ordered		
harmonic modes			optional (R3)
cells, graphs, symbols			finite resolution (E2, E3, E5)
cliques, factorizations			generator realization (E3)
ports, temperature, energy			thermodynamic typing (H2, K6)

Table 2: What is primitive, what is derived, what belongs to a realization.

Definition D3 (References and the datum $\cdot$ kernel $\cdot$ DEFINITION). Every informational quantity of the kernel is referenced explicitly, and the relative information $D(P \| R)$ of a law P from a declared reference is the primitive from which the differential and geometric quantities are derived. Three reference measures are declared, the *path reference* R of a process, the *epistemic reference* π , the prior on the identifiable quotient, and, when the geometry is invoked, the *calibration measure* m_{Λ} on a stratum; the operational access $\mathcal{E}$ is a family of experiments, declared beside them. The datum is $\mathfrak{J} = (\mathcal{M}_{\text{adm}}, \mathbf{S}, \mathcal{E}, R, \pi; m_{\Lambda})$, and zero internal structure is imposed on a candidate M .

Definition D4 (Beliefs and information $\cdot$ kernel $\cdot$ DEFINITION). A *belief* is a probability measure $\mu \in \mathcal{P}(\bar{\Lambda}_{\mathcal{E}})$ on the identifiable quotient. The *random law* Λ is a random element of $\Lambda_{\mathcal{E}}$ distributed as the prior π ; for the outcome Y of an experiment, π_Y is the posterior after Y , and $I(\Lambda; Y)$ the mutual information between the law and the outcome, equal to $\mathbb{E}_Y D(\pi_Y \| \pi)$ (B1).

Notation. $\log$ is the natural logarithm; $\mathbb{E}_P$ is the expectation under P , written $\mathbb{E}$ when the law is clear; $\mathcal{P}(\cdot)$ is the set of probability measures on a space; $K : \mathbf{X} \rightsquigarrow \mathbf{Y}$ is a Markov kernel, a channel, and KP the image of a law P by K ; ∇_h , $\|\cdot\|_h$, Δ_h , div_h are the gradient, norm, Laplace–Beltrami operator and divergence of h ; $\text{div}_m Y = e^V \text{div}_h(e^{-V} Y)$ for $m = e^{-V} dV_h$; $\mathbf{D}$ is the diffusion matrix of a diffusion, $\mathcal{D}$ a data set; visible and hidden parts of a ledger at resolution N carry the superscripts *vis* and *hid*. An f -divergence is $\int f(dP/dQ) dQ$ for a convex f with $f(1) = 0$ (Csiszár, 1967).

3.2 Channel and quotient

Postulate K0 (T1, Declared-Reference Principle $\cdot$ kernel $\cdot$ PRINCIPLE). Every information is relative to a reference declared on the space where it is computed. The path reference R , the prior π and the calibration m_{Λ} stay distinct, and an ensemble law on $\mathbf{X}$ is read against its own reference $m_{\mathbf{X}}$.

Definition K1 (Channels, signature and identifiable quotient $\cdot$ kernel $\cdot$ DEFINITION). Every observation, reduction or resolution is a channel. The signature of M under $\mathcal{E}$ is $F_{\mathcal{E}}(M) = \{p_M(\cdot | e)\}_{e \in \mathcal{E}}$; two candidates are equivalent when their signatures coincide,

$$M \sim_{\mathcal{E}} M' \iff F_{\mathcal{E}}(M) = F_{\mathcal{E}}(M'), \quad \bar{\Lambda}_{\mathcal{E}} = \mathcal{M}_{\text{adm}} / \sim_{\mathcal{E}}, \quad (3.1)$$

and $\bar{\Lambda}_{\mathcal{E}}$ is the space of the laws that the experiments tell apart. On the gyrator with $\theta = (k_1, k_2, k, T_1, T_2)$, a window of one observed coordinate leaves four identifiable combinations of the five parameters, two coordinates leave five; at $k = 0$ one coordinate sees two, the stiffness $k_1 + k$ and T_1 , and its null space holds $(1, 0, -1, 0, 0)$, e_{k_2} and e_{T_2} (gates X1.1, X1.4b).

Theorem K2 (T2, Channel Ledger Theorem: the exact ledger and data processing · kernel · IMPORTED). *For every channel K ,*

$$D(P \parallel R) = D(KP \parallel KR) + \mathbb{E}_{KP} D(P_{X|Y} \parallel R_{X|Y}), \quad (3.2)$$

on standard Borel spaces, the visible part plus the hidden part carried by the conditionals; hence $D(KP \parallel KR) \leq D(P \parallel R)$, and when $D(P \parallel R) < \infty$ equality holds exactly when the channel is sufficient for the pair (Kullback and Leibler, 1951, Thm 4.1), (Csiszár, 1975; Cover and Thomas, 2006).

Corollary C1 (The ledger characterizes relative entropy, finite alphabets · kernel · IMPORTED). *Let I assign a real number to each pair (p, r) of laws on finitely many points with $p \ll r$. Suppose I is measurable in r at fixed p , invariant under a common relabelling of both laws, vanishes on (p, p) , and obeys the exact ledger of every partition channel (Theorem K2), $I(w \circ (p^1, \dots, p^n) \parallel \tilde{w} \circ (\tilde{p}^1, \dots, \tilde{p}^n)) = I(w \parallel \tilde{w}) + \sum_i w_i I(p^i \parallel \tilde{p}^i)$, the visible part plus the hidden part averaged under the visible law. Then $I = c D(\cdot \parallel \cdot)$ for a constant c (Leinster, 2019, Thm 2.1); the unit of one nat fixes $c = 1$, and the functorial form of the statement is Baez and Fritz (2014), extended to standard Borel spaces by Gagné and Panangaden (2018); the characterization is a finite-alphabet statement, and the kernel's use of the ledger on general spaces rests on Equation (3.2) itself. Negative controls: on fifty random joints on 3×3 states the residual of the ledger is $2 \cdot 10^{-16}$ for relative entropy and reaches 0.05 for the squared Hellinger distance, 0.37 for the total variation and 10.8 for χ^2 ; the telescoping over two channels behaves the same way (gates C1.1, C1.2).*

3.3 Retention

Theorem K3 (T3, Retention Theorem: information projection and Bayes · kernel · IMPORTED). *Under hard constraints the retained law is the information projection of the declared reference onto the constraint set (Csiszár, 1975). Under likelihood evidence L with $Z = \int L d\pi \in (0, \infty)$, the least-deformation rule from the prior is Bayes,*

$$\mu_{\mathcal{D}} = \arg \min_{\mu} \left\{ \mathbb{E}_{\mu}[-\log L] + D(\mu \parallel \pi) \right\} = \pi_{\mathcal{D}} := \frac{L \pi}{Z}, \quad (3.3)$$

Bayes' rule in its Gibbs variational form (Zellner, 1988).

Proof. For every belief μ , $\mathbb{E}_{\mu}[-\log L] + D(\mu \parallel \pi) = \int \log \frac{d\mu}{L d\pi} d\mu = \int \log \frac{d\mu}{Z d\pi_{\mathcal{D}}} d\mu = D(\mu \parallel \pi_{\mathcal{D}}) - \log Z \geq -\log Z$, and relative entropy vanishes exactly on equal laws (gate C2.3). Among the f -divergences with f differentiable, relative entropy is the one whose least-deformation update coincides with $\pi_{\mathcal{D}}$ for every prior and every likelihood (C2, proved in the supplement, Appendix S.B); Bissiri and Walker (2012) and Bissiri et al. (2016) reach the same selection through a coherence requirement. $\square$

Corollary B1 (Belief and the information of an experiment, Lindley · kernel · IMPORTED). *The expected information of an experiment is the mean realized relative information of its posterior, $I(\Lambda; Y) = \mathbb{E}_Y D(\pi_Y \parallel \pi)$ (Lindley, 1956); post-processing contracts it, by Equation (3.2) applied to the joint law against the product (Theorem K2).*

3.4 Sufficiency

Theorem K4 (T4, Recovery-Kernel Theorem: sufficiency and recovery · kernel · IMPORTED). *When $D(P \parallel R) < \infty$, a zero hidden term in the ledger Equation (3.2) of a channel K is sufficiency (Kullback and Leibler, 1951, Thm 4.1). For the declared family $\{P_\lambda\}$, sufficiency is equivalent to the existence of a recovery Markov kernel L with $L K P_\lambda = P_\lambda$ for every law of the family and for the reference (Blackwell, 1953; Petz, 1988), and a sufficient channel preserves the Fisher metric of a regular stratum.*

4 Information calculus and relative-entropy slope geometry

The primitive is differentiated twice, and Fisher information appears twice, as two objects: the Hessian of D in the space of identifiable laws is a tensor h on the stratum; the Wasserstein slope of D on measures over the stratum is a scalar functional $\mathcal{I}_{h,m}$ on beliefs. The two meet in the relative-entropy slope geometry (Figure 3).




Figure 3: Two differentials of one primitive. The Hessian gives the tensor h on identifiable laws; the slope gives the functional $\mathcal{I}_{h,m}$ on beliefs; the transport distance built on h , weighted by the slope, is the calibrated length.

4.1 The differential in law space

Theorem O1 (T5, Operational Fisher Theorem · kernel · IMPORTED). *On a regular stratum of the quotient,*

$$D(K_\lambda \parallel K_{\lambda+d\lambda}) = \frac{1}{2} h_K(d\lambda, d\lambda) + o(\|d\lambda\|^2), \quad h_{T \circ K} \preceq h_K. \quad (4.1)$$

The metric is induced by the observational access; it contracts under every channel and stays under a sufficient one; up to a factor it is the metric invariant under sufficient Markov morphisms, on finite sample spaces by Čencov (1982) and on parametrized measure models by Ay et al. (2017).

Theorem O3 (T6, Missing-Fisher Ledger Theorem: score projection and hidden Fisher · kernel · PROVED). *For an observation $Y = K(X)$ of a family with scores $S_v \in L^2$, the observable score is the conditional expectation of the fine score, $S_v^Y = \mathbb{E}[S_v \mid Y]$ (O2), and*

$$h = h_Y + h_Y^{\text{hid}}, \quad h_Y^{\text{hid}}(v, v) = \mathbb{E} \text{Var}(S_v \mid Y) \geq 0, \quad (4.2)$$

the hidden Fisher being the Hessian of the hidden relative information of Equation (3.2). Along the null direction of the observable Fisher all of the fine Fisher is hidden (gates X1.2, X1.3b, X2.2; Figure 4).

The identities are classical: Fisher’s identity for the score of an observation, the missing-information principle of Orchard and Woodbury (1972), Louis’s formula for the observed information (Louis, 1982), and the fraction of missing information of Meng and Rubin (1991). IDS reads




Figure 4: The channel builds the ground. Left: the singular values of the Fisher metric of $\theta = (k_1, k_2, k, T_1, T_2)$ from a window of the gyrator’s trajectory, 64 samples at step 0.2 without observation noise, two coordinates observed (blue) and one coordinate observed (orange): rank five against rank four. Right: the components of the direction that one coordinate is blind to; along it the whole Fisher information is hidden. [K1, O3]

them as the second-order differential of the exact channel ledger and places them in the same resolution tower as relative information, mutual information and calibrated radius; the hidden share of the Fisher information along a direction is the fraction of missing information, and a δ -almost sufficient projection is one whose largest fraction of missing information is at most $1 - \delta^2$ (G10).

4.2 The relative-entropy slope geometry

RESG extends the belief-cost geometry of Caraffa (2026), whose reference is the volume, to a declared reference.

Proposition S (T7, Slope-Geometry Proposition · kernel · PROVED). *Let (X, d) be a length space, $H : X \rightarrow (-\infty, +\infty]$ lower semicontinuous, $g = |\partial H|_d$ a strong upper gradient of H (Ambrosio et al., 2008, Def. 1.2.1) and $\ell_\star > 0$ a constant. For $L_H(\gamma) = \ell_\star \int g(\gamma_t) |\dot{\gamma}_t| dt$, every absolutely continuous curve with $L_H(\gamma) < \infty$ satisfies $L_H(\gamma) \geq \ell_\star |H(\gamma_1) - H(\gamma_0)|$; and when a curve of maximal slope from x , with $-\frac{d}{dt}H = g^2 = |\dot{x}|^2$, converges to a minimizer $x_\star$ of H with $H(x_t) \rightarrow H(x_\star)$, in finite time or as $t \rightarrow \infty$, the least length from x to $x_\star$ is $\ell_\star [H(x) - H(x_\star)]$.*

Proof. The strong upper gradient gives $|H(\gamma_1) - H(\gamma_0)| \leq \int g(\gamma_t) |\dot{\gamma}_t| dt$ (Ambrosio et al., 2008, Def. 1.2.1); multiply by $\ell_\star$. Along the curve of maximal slope (Ambrosio et al., 2008, Def. 1.3.2), $\ell_\star \int_0^T g(x_t) |\dot{x}_t| dt = \ell_\star \int_0^T (-\frac{d}{dt}H(x_t)) dt = \ell_\star [H(x) - H(x_T)]$, which tends to $\ell_\star [H(x) - H(x_\star)]$; the first part makes this length the least one. $\square$

The construction is abstract before it is epistemic. Its epistemic realization is $X = \mathcal{P}_2(\bar{\Lambda}_\mathcal{E})$, $d = W_{2,h}$, $H = D(\cdot \| m_\Lambda)$: the points are beliefs over identifiable laws. Its physical realization is $X = \mathcal{P}_2(\mathbf{X})$, $d = W_{2,g_\mathbf{X}}$, $H = D(\cdot \| m_\mathbf{X})$: the points are ensemble laws over a state space, and the gyrator benchmarks it. Both are invoked under one contract.

Contract HR0 (Geometric contract · kernel · DEFINITION). The stratum (Λ, h) is a complete Riemannian manifold; the reference is $m = w dV_h$ with $\text{Ric}_h + \text{Hess}_h(-\log w) \geq Kh$ for some real K (Bakry–Émery); beliefs are regular. Under HR0 the metric slope of $\mathcal{H}_m(\mu) = \int f \log f dm$, $f = d\mu/dm$, in $W_{2,h}$ is $|\partial \mathcal{H}_m|^2 = \mathcal{I}_{h,m} = \int \|\nabla_h \log f\|_h^2 d\mu$, its gradient flow is the Fokker–Planck equation and $\mathcal{H}_m$ is K -convex along displacement interpolation (G1, imported in Appendix S.B); for $K > 0$ a logarithmic Sobolev inequality holds and gives exponential decay and Talagrand’s inequality (Bakry and Émery, 1985; Otto and Villani, 2000). The same contract, written for the state space $(\mathbf{X}, g_\mathbf{X}, m_\mathbf{X})$ with $g_\mathbf{X}$ its Euclidean metric, covers the physical realization.

Definition G2 (The calibrated length $\cdot$ kernel $\cdot$ DEFINITION).

$$d\ell_I = \ell_\star \sqrt{\mathcal{I}_{h,m}} d\ell_{W_{2,h}} \quad (4.3)$$

the slope of relative entropy is the ruler, $\ell_\star$ is a gauge, the length of one nat; L_I is the length of a curve and d_I the induced length pseudometric, symmetric with the triangle inequality under HR0 alone and a metric wherever points are separated, which the logarithmic Sobolev inequality guarantees on the extended space (Appendix S.B.2); the name calibrated distance is kept for it.

Theorem G3 (T8, RESG Theorem: Entropy Floor, Constant Slope, Radial Identity $\cdot$ kernel $\cdot$ PROVED).
Under HR0:

- (i) floor and horizon: every path pays at least $\ell_\star$ times the entropy difference of its endpoints, $d_I(\mu_0, \mu_1) \geq \ell_\star |\mathcal{H}_m(\mu_1) - \mathcal{H}_m(\mu_0)|$; the states of infinite relative entropy, exact certainty among them, lie at infinite calibrated distance and at finite transport distance: $W_{2,h}(\mu_n, \delta_\lambda) \rightarrow 0$ with $m(\{\lambda\}) = 0$ forces $d_I(\mu_0, \mu_n) \rightarrow \infty$;
- (ii) pointwise rigidity: for a scalar-conformal deformation of $W_{2,h}$ by a relief a , the slope of relative entropy at a regular belief where a is continuous is $\sqrt{\mathcal{I}_{h,m}}/a$, so it equals $1/\ell_\star$ there exactly when $a = \ell_\star \sqrt{\mathcal{I}_{h,m}}$ there; the weight $\ell_\star \sqrt{\mathcal{I}_{h,m}}$ gives the constant slope $1/\ell_\star$ at every regular belief (G4);
- (iii) radial identity: with the prior as reference and an ergodic gradient flow, $\rho_1(\mu) = \ell_\star D(\mu \parallel \pi)$, relative entropy is the radial coordinate and the gradient flow is the radial geodesic (G5).

On the gyrator, in the physical realization with $(k_1, k_2, k) = (1, 0.7, 0.6)$: cooled to $T \rightarrow 0$ against the reference at $T = 1$, D grows like $\log(1/T)$ with slope 0.998 while W_2 stays below 1.30 (gate X3.3); the relaxation runs at equal temperatures $T_1 = T_2 = 1$ from the displaced stationary law $\mathcal{N}(m_0, C)$, with C the stationary covariance at $T_1 = 2$, $T_2 = 0.5$ and the offset $m_0 = (1.5, -1)$, towards the Gibbs law at $T = 1$, with the gauge $\ell_\star = \varepsilon/2 = 1$ of G8 ($\varepsilon = 2T$); on it the bare transport length is 1.857 and the length with weight $\mathcal{I}_{h,m}$ in place of $\sqrt{\mathcal{I}_{h,m}}$ is 9.15 against a drop of relative information of 3.554 (gate X3.4); with isotropic diffusion and a symmetric drift the relaxation is the gradient flow of $D(\cdot \parallel \pi)$, hence radial: its calibrated length is 3.55407, the drop of relative information 3.55402 and the entropy production measured on $2 \cdot 10^4$ paths 3.535 ± 0.023 (gates X3.1, X3.2, LIB.8). The proofs are in Appendix S.B; Figure 5 shows the Gaussian leaf under the two calibration references.

Corollary G7 (Scale invariance $\cdot$ kernel $\cdot$ PROVED). Rescaling the metric of Definition G2, $h' = \alpha h$ with the same reference measure, or with the volume of h' when the reference is the volume of h , leaves the calibrated length invariant (cf. Caraffa, 2026, Thm 4.23); n repetitions of an experiment turn h into nh : distinguishability is extensive, nats are intensive.

Corollary G8 (Diffusive-Unit Corollary $\cdot$ kernel $\cdot$ PROVED). For a diffusive reference of noise ε and the gauge $\ell_\star = \varepsilon/2$, the Girsanov relative entropy of a controlled path measure P against the reference R , completed by the entropy boundary term, bounds the calibrated length of its marginals (Definition G2), $\varepsilon D(P \parallel R) - \frac{\varepsilon}{2} [\mathcal{H}_m(\mu_1) - \mathcal{H}_m(\mu_0)] \geq L_I(\mu)$, with equality along the reference flow; $\varepsilon/2$ is the largest gauge for which this bound holds, and the energy form of the entropic cost behind it is that of Gentil et al. (2017, Cor. 5.3) and Chen et al. (2016): a path-space information price for diffusive revision. A reading of that price as heat requires a thermal realization under local detailed balance (H2).

Corollary G9 (Shannon as posterior geometry $\cdot$ kernel $\cdot$ PROVED). Under the radial identity of Theorem G3, $\ell_\star$ times the mutual information is the mean calibrated radius of the posteriors, and $\ell_\star$ times the rate-distortion function is the least mean calibrated radius of the posterior cells of a test channel under a mean squared transport radius to the code points; for a source without atoms zero distortion is the horizon, $R(0) = \infty$, while a finite alphabet gives $R(0) = H(X)$ (Shannon,




Figure 5: One leaf, two calibration references. (a) The volume as reference, drawn in the Poincaré disk of the Gaussian leaf: the geometry is hyperbolic and homogeneous, the prior $\pi = \mathcal{N}(0, 1)$ at the centre carries zero privilege and the certainties are the boundary; two beliefs, the geodesic between them, solid with one mark per nat, and the chord at fixed width, dashed; below, the belief along each route, one snapshot per nat of cost. (b) The prior as reference, drawn in the half-plane (a, s) of the same leaf: the geometry acquires a centre. Purple: the level sets of $D(\cdot \| \pi)$, the calibrated circles about π , half a nat apart. Green: the traces of the gradient flow, the radial geodesics. The horizon $s = 0$ is at infinite calibrated distance and at finite transport distance, the Dirac mass $\delta_{0.6}$ at $W_2 = 1.17$. The dashed line $s = \sqrt{D}$ carries the cells $\mathcal{N}(a, D)$ of Shannon’s channel at distortion $D = \frac{1}{4}$: their transport radius to the code point is s , their calibrated radius is the level they touch, and the mean level along the line under $a \sim \mathcal{N}(0, 1 - D)$ is the rate $\frac{1}{2} \log(1/D) = 0.69$ nat (gate S1.1). [G3, G5, G6, G9]

1959). For the Gaussian source the mean radius of the cells equals $\frac{1}{2} \log(\sigma^2/D)$ to 10^{-16} (gate S1.1); on Figure 5(b) the cells at distortion D are the line $s = \sqrt{D}$ and the rate is the mean level along it.

The hyperbolic geometry of the location–scale leaves (G6) is stated in the supplement (Appendix S.C); Figure 5(b) draws the radial geodesics and the calibrated circles under the prior. The Fisher-weighted transport geometry $\ell_{\star}^2 \mathcal{I}_{h,m} g_{W_{2,h}}$ is a distinct object from the Wasserstein–Fisher–Rao metric of unbalanced transport, where the Fisher term enters the transport itself.

5 Resolution as an ordered family of channels

A finite description is a channel, and resolutions are ordered by post-processing. Figure 6 shows the four ledgers on nested sensors of the gyrator’s state; the statements follow.

Theorem R1 (T9, Resolution Ledger Theorem: the tower and its ledgers $\cdot$ kernel $\cdot$ PROVED). *A resolution K_N is coarser than K_M when*

$$K_N = T_{N \leftarrow M} \circ K_M \quad (5.1)$$

for a channel T , Blackwell’s garbling order (Blackwell, 1953). Along a nested tower, $D_N \leq D_M$, $h_N \leq h_M$ and $I_N \leq I_M$, by Equation (3.2), Equation (4.2) and B1, and at every level the primitive, its second order, its Bayesian reading and its geometric reading split into a visible and a hidden part (Table 3, R2). Two resolutions unrelated by post-processing are unordered: a sensor on one coordinate at noise 0.3 and a sensor on both at noise 0.6 dominate each other in different directions (gate X2.3); in time, the entropy production seen by an observer of the gyrator’s two coordinates grows monotonically along the dyadic steps $\Delta = 2^{-n}$ (gate LIB.7).




Figure 6: The resolution ledger. Nested sensors $Y_N = x + \mathcal{N}(0, 2^{-2N}I)$ on the gyrator's state with an unknown force offset: the relative information between two offsets, the Fisher information of the offset, the mutual information and the mean calibrated radius of the posteriors fill monotonically with N (dark: visible, light: hidden) towards the fine value (line). [R1, R2, R5]

Proof. With $K_N = T \circ K_M$, the data-processing inequality of Equation (3.2) gives $D_N \leq D_M$, the score projection of Equation (4.2) gives $h_N \preceq h_M$, and B1 gives $I_N \leq I_M$; the four splits are Equation (3.2), Equation (4.2), the chain rule of mutual information and, under the radial identity, $\mathbb{E}\rho_{I,N} = \ell_* I_N$ (Theorem G3); they are collected in R2. $\square$

level	visible	hidden	needs
relative information	D_N	$D_N^{\text{hid}} = D - D_N$	any channel (K2)
Fisher	h_N	$h_N^{\text{hid}} = \mathbb{E} \text{Var}(S Y_N)$	a regular stratum, scores in L^2 (O3)
Bayesian	$I_N = I(\Lambda; Y_N)$	$I_N^{\text{hid}} = I - I_N$	a latent law under the prior (B1)
calibrated radius	$\mathbb{E}\rho_{I,N} = \ell_* I_N$	$\ell_* I_N^{\text{hid}}$	the prior as reference and the radial identity (G3)

Table 3: The Resolution Ledger: the four lines of a tower and what each needs (R2; on the nested sensors of Figure 6 the four quantities increase strictly and $\mathbb{E}\rho_I = \ell_* I_N$ to 10^{-16} , gate X2.1).

Step by step, the increments of the ledger are the conditional ledgers: $D_{n+1} - D_n = \mathbb{E} D(P_{Y_{n+1}|Y_n} \| R_{Y_{n+1}|Y_n})$, $h_{n+1} - h_n = \mathbb{E}[(S_{n+1} - S_n)^{\otimes 2}]$, $I_{n+1} - I_n = I(\Lambda; Y_{n+1} | Y_n)$ and $\mathbb{E}\rho_{I,n+1} - \mathbb{E}\rho_{I,n} = \ell_* I(\Lambda; Y_{n+1} | Y_n)$, by the chain rules of Equation (3.2), Equation (4.2) and B1: a tower carries an additive spectrum of information revealed from coarse to fine (R2; on the sensors the increments equal the conditional quantities to $3 \cdot 10^{-12}$ in absolute value, gate X2.8b).

Theorem R5 (T10, Resolution Recovery Theorem · kernel · PROVED). *Let $\mathcal{F}_1 \subset \mathcal{F}_2 \subset \dots$ be a filtration generating $\mathcal{F}$ up to $(P + R)$ -null sets, R a probability. Then $D(P|_{\mathcal{F}_N} \| R|_{\mathcal{F}_N}) \uparrow D(P \| R)$ in $[0, \infty]$ and the hidden term decreases to zero. For a family differentiable in quadratic mean with scores in L^2 , $h_N \uparrow h$ in the Loewner order with $h_M - h_N = \text{Cov}(S_M - S_N)$. For observations Y_N generating Y , $I(\Lambda; Y_N) \uparrow I(\Lambda; Y)$, and under the radial identity $\mathbb{E}\rho_{I,N} \uparrow \mathbb{E}\rho_I$. The proof, in Appendix S.B, rests on Lévy's upward theorem and on relative entropy as a supremum over finite partitions. On the gyrator the dyadic tower in time is a filtration on path space: the observer of both coordinates sees 99.4% of σ at $\Delta = 2^{-8}$ (gate X5.2); the tower of sensors is a garbling with $K_N P \rightarrow P$, whose limit follows from the lower semicontinuity of relative entropy: the four ledgers are within 0.5% of their fine values from $N = 5$, with a gap linear in the noise variance (gate X2.4).*

Theorem G10 (T11, Almost-Sufficient Stability Theorem · kernel · PROVED). *Let $\delta^2 h \preceq h_K \preceq h$ on the same stratum, $0 < \delta \leq 1$, with the same reference measure m and the same gauge ℓ_* , the coarse metric h_K being that of a δ -almost sufficient statistic (Yamaguchi and Nozawa, 2024).*

Then $\delta W_{2,h} \leq W_{2,h_K} \leq W_{2,h}$, $\mathcal{I}_{h,m} \leq \mathcal{I}_{h_K,m} \leq \delta^{-2} \mathcal{I}_{h,m}$, and

$$\delta L_1^h(\gamma) \leq L_1^{h_K}(\gamma) \leq \delta^{-1} L_1^h(\gamma), \quad \delta d_1^h \leq d_1^{h_K} \leq \delta^{-1} d_1^h : \quad (5.2)$$

almost sufficient in Fisher means almost isometric in the slope geometry, and $\delta^2 = 1 - \varphi$ with φ the largest fraction of missing information of the projection. Under HR0 for (Λ, h_K, m) as well, $L_1^{h_K}$ is the calibrated length of the coarse experiment and the radial identity holds with the same right-hand side. On the gyrotor the 32-step sub-window of the 64-step window is δ -almost sufficient with $\delta^2 = 0.49$, while one coordinate alone gives $\delta = 0$ (gate X1.5); on the tower of sensors $\delta_N^2 = \min_i \lambda_i(C)/(\lambda_i(C) + 2^{-2N})$ in closed form and the calibrated length of a translation of a Gaussian belief stays within $[\delta_N, \delta_N^{-1}]$ of its fine value and tends to it (gate X2.5). Proof in Appendix S.B.

Corollary G11 (Uniform almost-sufficiency convergence and composition $\cdot$ kernel $\cdot$ PROVED). Under the hypotheses of Theorem G10: (i) if a tower (h_N) on the same stratum satisfies $\delta_N^2 h \preceq h_N \preceq h$ with $\delta_N \rightarrow 1$, then $L_1^{h_N}(\gamma) \rightarrow L_1^h(\gamma)$ for every common admissible curve of finite length and $d_1^{h_N}(\mu, \nu) \rightarrow d_1^h(\mu, \nu)$ for every pair at finite distance: non-radial convergence is automatic along uniformly almost-sufficient towers, and what remains open is the convergence of minimizing geodesics and the discrete realizations that leave the stratum (Q2); (ii) if K_1 is δ_1 -almost sufficient and K_2 is δ_2 -almost sufficient for the model induced by K_1 , then $K_2 \circ K_1$ is $\delta_1 \delta_2$ -almost sufficient, $(\delta_1 \delta_2)^2 h \preceq h_{K_2 K_1} \preceq h$, so that $-\log \delta$ is an additive upper budget of geometric distortion along a reduction pipeline and the fractions of missing information compose as $\varphi_{\text{tot}} \leq 1 - \prod_i (1 - \varphi_i)$. Both follow from Equation (5.2), by passing to the limit and by chaining the Loewner sandwiches. On the tower of sensors $|L_N/L_\infty - 1| \leq \delta_N^{-1} - 1 \rightarrow 0$, and δ^2 composes multiplicatively as a lower bound (gates X2.6, X2.7).

Proof. (i) Equation (5.2) with $\delta = \delta_N$ gives $\delta_N L_1^h \leq L_1^{h_N} \leq \delta_N^{-1} L_1^h$ and the same sandwich for d_1 ; let $\delta_N \rightarrow 1$. (ii) $h_{K_2 K_1} \succeq \delta_2^2 h_{K_1} \succeq \delta_2^2 \delta_1^2 h$ by the two Loewner sandwiches, and $1 - \varphi_{\text{tot}} \geq \prod_i (1 - \varphi_i)$ is the same inequality read with $\delta_i^2 = 1 - \varphi_i$. $\square$

The harmonic Markov resolution. A resolution tower is any ordered family of channels. Under HR0 the heat semigroup of the reference, $K_t = e^{tL_m}$ with $L_m = \Delta_h + \nabla_h \log w \cdot \nabla_h$, is one: each K_t is a Markov channel, $K_{t_2} = K_{t_2-t_1} \circ K_{t_1}$ for $t_2 \geq t_1$, so that increasing t is a Blackwell coarsening, and along $\mu_t = K_t \mu$, $-\frac{d}{dt} D(\mu_t \| m) = \mathcal{I}_{h,m}(\mu_t)$ (G1): relative Fisher information is the instantaneous relative-information loss rate of the canonical harmonic coarse-graining. When the generator has a compact resolvent the semigroup is diagonal in its eigenbasis, $K_t \phi_k = e^{-\lambda_k t} \phi_k$, coarse to fine as $\lambda_1 \leq \lambda_2 \leq \dots$; the finite spectral cutoff is the analysis, with the χ^2 ledger of Parseval and a tail bounded by $e^{-2\lambda_{N+1} t}$, and the exponential family of the first N modes is the positive reconstruction (R3, R4, in Appendix S.C; diffusion maps build multiscale coordinates from the same operators, Coifman and Lafon, 2006). The kernel is free of this choice.

6 Emergent finite structure

Here, and only here, appear meshes, cells, facets, graphs, hypergraphs, symbols and Markov factorizations: they are what a resolution reads on a geometry, and a graph used as a finite representation of a continuum geometry is determined by the measure, the declared resolution and the discretization rule as much as by the geometry.

Cells and facets. Quantised on cells of step ε , the current of a process crosses facets and the balance of each cell is a discrete Kirchhoff law; the chain with facet rates k_{ij} and stationary law p has the entropy production

$$\sigma_\varepsilon = \frac{1}{2} \sum_{ij} J_{ij} A_{ij}, \quad J_{ij} = p_i k_{ij} - p_j k_{ji}, \quad A_{ij} = \log \frac{p_i k_{ij}}{p_j k_{ji}}, \quad (6.1)$$




Figure 7: Emergence. (a) The stationary gyration quantised at $\varepsilon = 0.4 \sigma_{\min}$, the current through the facets. (b) The production of the four-neighbour chain converges to σ ; the eight-neighbour chain keeps an excess. (c) Three graphs on one object: the proximity graph of the cells, the interaction hyperedge of the spring, the conditional-independence graph of the stationary law. [E2, E3]

which converges to the production of the process as $\varepsilon \rightarrow 0$ when the rates are those of the facets; under the tested eight-neighbour mobility convention, the diagonal links add a persistent excess production, so the ledger is a property of the facet conductances of the stencil (E2, Figure 7).

Three graphs, and the ruler that bends. On one object live three graphs: the proximity graph of the cells, the interaction hyperedges of the generator and the conditional-independence graph of the law; on the gyration the first ignores the coupling k while the two others depend on it (E3). The law ruler joins them: on the stratum $(\log k_1, \log k_2)$ at fixed k , the stationary law factorizes at $k = 0$ with one parameter per factor, so the Fisher metric is the direct sum of two one-dimensional metrics, $\frac{1}{2}I$ in log-stiffness coordinates and flat; it curves for $k > 0$, with Gaussian curvature -0.274 at the nominal point for $k = 0.6$; the generator edge A_{12} , the conditional-independence edge $(C^{-1})_{12}$ and the curvature of the ruler leave zero together with the coupling, three concomitant signatures of the same coupling in this Gaussian realization, the same loss of factorization that the coupling term of Equation (6.2) measures (gates SYM.7, X6.6; Figure 8). A graphical factorization is a realization property (Lauritzen, 1996).

Two axes. Relative to a product reference, what a belief has learned splits into what concentrates each coordinate and the total correlation that couples them,

$$D(\mu \parallel m) = \sum_i D(\mu_i \parallel m_i) + D(\mu \parallel \prod_i \mu_i), \quad (6.2)$$

and a sensor on one coordinate moves the first part only, a joint sensor the second (E4).

Theorem E6 (T12, Predictive Closure Ledger Theorem · kernel · PROVED). *Let (x_t) be stationary, q measurable with values in a finite alphabet, $X_t = q(x_t)$, and at the time resolution Δ let $I_{\text{pred}} = I(X_{t+\Delta}; x_t)$, $I_{\text{ret}} = I(X_{t+\Delta}; X_t)$ and $I_{\text{hid}} = I(X_{t+\Delta}; x_t \mid X_t)$. Then*

- (i) $I_{\text{pred}} = I_{\text{ret}} + I_{\text{hid}}$, with $I_{\text{hid}} = \mathbb{E}_{x_t} D(P_{X_{t+\Delta}|x_t} \parallel P_{X_{t+\Delta}|X_t})$;
- (ii) $I_{\text{hid}} = 0$ exactly when $X_{t+\Delta}$ and x_t are conditionally independent given X_t : X_t is a predictively sufficient statistic of x_t for $X_{t+\Delta}$;
- (iii) $0 \leq I_{\text{hid}} \leq I_{\text{pred}} \leq H(X_t)$, so that the loss $\varepsilon_{\text{cl}} = I_{\text{hid}}/I_{\text{pred}}$, the fidelity $\eta_{\text{ret}} = I_{\text{ret}}/I_{\text{pred}} = 1 - \varepsilon_{\text{cl}}$ and the memory $\eta_{\text{mem}} = I_{\text{pred}}/H(X_t)$ lie in $[0, 1]$;
- (iv) $\mathbb{E} \|P_{X_{t+\Delta}|x_t} - P_{X_{t+\Delta}|X_t}\|_{\text{TV}} \leq \sqrt{I_{\text{hid}}/2}$;




Figure 8: One coupling, three objects. (a) Seven covariance-shape sheets of the stationary gyrotator, $k = 0$ (blue) to $k = 0.6$ (red), each the image of the chart $(\log k_1, \log k_2)$ in the coordinates $(\frac{1}{2} \log \det C, (S_{11} - S_{22})/2, S_{12})$, $S = C/\sqrt{\det C}$: visual covariance-shape coordinates, free of any isometry claim; the curvature is computed intrinsically from the Fisher metric. (b) Tissot indicatrices of the law ruler h on the chart, $T_1 = 2$, $T_2 = 0.5$: at $k = 0$ the metric is $\frac{1}{2}I$, circles everywhere, curvature zero; at $k = 0.6$ the indicatrices stretch and turn, curvature -0.274 at the cross. (c) Against the coupling k : the curvature of h , the partial correlation and the mutual information of the two coordinates, each relative to its value at $k = 0.8$, leave zero together. The curvature is a local quantity, -0.35 and -0.20 at two other points of the chart. [E3, O1]

(v) when the paths are continuous and x_t avoids the boundaries of q almost surely, $I_{\text{ret}}, I_{\text{pred}} \rightarrow H(X_t)$ and $I_{\text{hid}} \rightarrow 0$ as $\Delta \rightarrow 0$.

A level closes at tolerance (η, α) when $\varepsilon_{\text{cl}} \leq \eta$ and $\eta_{\text{mem}} \geq \alpha$. On the double well $U(x) = E_b(x^2 - 1)^2$ with the quotient $\{\text{left}, \text{right}\}$: at $T = 0.15$ the loss stays at or below 0.005 and the memory above 0.79 on $\Delta \in [0.5, 20]$; at $T = 1$ the fidelity stays near 0.88 while the memory falls to 0.04 at $\Delta = 2$ and to the floor of the estimator beyond; the mean total-variation error of predicting from the symbol is 0.001 at $T = 0.15$ and 0.075 at $T = 1$, $\Delta = 0.5$; the short-lag limit is the occupation entropy (gates X8.5, X8.6b, X8.7, X8.8, X8.2b; Figure 9). Proof in Appendix S.B; the retained information is the relevant information of the information bottleneck (Tishby et al., 1999), $I_{\text{hid}} = 0$ is the informational closure of Bertschinger et al. (2006) and Pfante et al. (2014), and predictive sufficiency is the causal-state construction of Shalizi and Crutchfield (2001); see Geiger (2026) for the reduction of Markov chains.

Realization E5 (Symbol $\cdot$ realization $\cdot$ MEASURED). The number of live cells of a belief at resolution ε is its perplexity e^{H_ε} , equal to $2\pi e \sqrt{\det C}/\varepsilon^2$ for a Gaussian to 0.1 % (gate X6.4b), and informative experiments reduce its logarithm, the entropy H_ε , on average. A symbol at resolution $(\varepsilon, \Delta, \eta, \alpha)$ is a coarse state that is predictively closed to tolerance (η, α) over the horizon Δ and whose exit time exceeds Δ by the declared factor. With atoms in the reference, exact certainty has a finite relative-information cost, $-\log m(\{x\})$, where the continuous horizon has an infinite one; on one well of the double well the count of cells above 10^{-3} of the largest cell mass shrinks from 16 to 7 as T goes from 0.4 to 0.05, the perplexity of the cell masses from 11.1 to 3.5 (gates X8.3, PUIITS.1), and the exit rate follows Arrhenius (E7).




Figure 9: The double well. (a) The predictive ledger of the two-symbol quotient against the time resolution, retained and hidden information at $T = 0.15$ (closes) and $T = 1$ (drowned barrier). (b) The closure ratios, memory η_{mem} and loss ε_{cl} : at $T = 1$ the loss stays small while the memory dies. (c) The count of cells of one well above 10^{-3} of the largest cell mass at fixed spatial resolution, against the temperature. [E5, E6]

7 Physical realization and numerical experiments

7.1 Informational dissipative structures

Definition H1 (Informational dissipative structure · realization · DEFINITION). A realization observed at a declared resolution (ε, Δ) is an informational dissipative structure when (i) its quotient closes at tolerance (η, α) , $\varepsilon_{\text{cl}} \leq \eta$ and $\eta_{\text{mem}} \geq \alpha$ (E6); (ii) the path law of the level is irreversible, $D(P \| P^\dagger) > 0$, which, for the stationary overdamped diffusions used here under the standard time reversal, is equivalent to a nonvanishing stationary current (K5); (iii) the irreversibility is sustained by declared environmental ports whose conservation ledger and entropy-flow ledger close: in the stationary state $\dot{W} = \sum_i \dot{Q}_i$ and $\sigma = \sum_i \dot{Q}_i / T_i$, with $\dot{W}$ the work received from the work ports, $\dot{Q}_i$ the heat delivered to bath i , and $\dot{W} = 0$ for a purely thermal realization (H2). Its revision is priced by $\sqrt{\mathcal{I}_{h,m}}$ (G2). The gyrator satisfies clauses (ii) and (iii) exactly when $k(T_1 - T_2) \neq 0$, and everything is a closed form: with $\Gamma = \mathbf{D}C^{-1} - A$ the current is $j = \Gamma x p$, $\Gamma C = sJ$ with

$$s = \frac{k(T_1 - T_2)}{2k + k_1 + k_2}, \quad \sigma = \frac{k^2(T_1 - T_2)^2}{T_1 T_2 (2k + k_1 + k_2)}, \quad Q_1 = -Q_2 = -\frac{k^2(T_1 - T_2)}{2k + k_1 + k_2}, \quad (7.1)$$

the current is the rotation of the stream function $\psi = sp$, a sufficient and exactly solvable realization of clause (ii), and the three quantities vanish exactly when $k(T_1 - T_2) = 0$ (gates X4.2, X4.3, X4.4, LIB.1, LIB.2, LIB.4, LIB.5, SYM.1–SYM.4, SYM.6). The driven ring carries a harmonic current, $\sigma = F^2/T$ at uniform potential, so clause (ii) covers coexact and harmonic currents alike; its drive is a work port, $\dot{W} = FjL$, delivered as heat to its single bath, $\sigma = \dot{W}/T$ (gate RING.3).

Realization H2 (Ports · realization · MEASURED). A port is a declared environmental channel through which a realization exchanges a conserved quantity or a control signal: a thermal port, a work or control port, a measurement port. In the canonical thermal realization a bath at temperature T imposes the reference $e^{-E/k_B T}$, with $k_B = 1$ in the units of the gyrator, and for an isothermal port under local detailed balance the heat delivered is $k_B T$ times the nats given to the port; a work port carries $\dot{W}$ with zero entropy flow, and the stationary energy ledger reads $\dot{W} = \sum_i \dot{Q}_i$, with $\dot{W}$ received from the work ports and $\dot{Q}_i$ delivered to bath i . On the gyrator the heats delivered to the baths satisfy $Q_1 + Q_2 = 0$ and $\sigma = Q_1/T_1 + Q_2/T_2$, Clausius, exactly (gate LIB.3); a geometric boundary is one way to realize a port. The reference potential $-\log(dm/d\nu)$ of the kernel becomes $\beta E + \text{const}$ in this realization alone (K6).




Figure 10: The gyrotor: current, ports and the two ledgers. (a) At $T_1 = 2, T_2 = 0.5$ the stationary current (orange) runs along the level sets of the density (grey); on the disk (dashed) the interface flux and the bulk term of the relative-information balance vanish while the entropy production is positive, $\Phi_A^{\text{rel}} = 0, B_A^{\text{rel}} = 0, \sigma_A = 0.172$; the two baths are the ports. (b) At equal temperatures $T_1 = T_2 = 1$, from the displaced stationary law $\mathcal{N}(m_0, C)$ of Theorem G3, the ensemble relaxes along the radial geodesic: the calibrated length of its path, the drop of relative information and the measured entropy production coincide. (c) At equal temperatures the current and the production vanish. [H1, G3, E1]

Realization E1 (Two-Ledger Identity: the balance of a zone · realization · PROVED). On a weighted manifold with a density f relative to m and a current $j = fv$, every zone A obeys

$$\frac{d}{dt} \int_A f \log f \, dm = -\Phi_A^{\text{rel}} + B_A^{\text{rel}}, \quad \Phi_A^{\text{rel}} = \oint_{\partial A} (1 + \log f) \langle j, n \rangle \, d\sigma_m, \quad B_A^{\text{rel}} = \int_A \langle v, \nabla \log f \rangle \, d\mu, \quad (7.2)$$

content, interface flux and bulk relative-information term; for the gradient flow $v = -\nabla \log f$ the bulk term is $-\mathcal{I}_A$. For a diffusion of drift b and constant diffusion matrix $\mathbf{D}$, the thermodynamic entropy production of the zone is

$$\sigma_A = \int_A \langle v, \mathbf{D}^{-1} v \rangle \, d\mu = M_A - B_{m,A}^{\text{rel}} - \int_A \langle v, \nabla \log m \rangle \, d\mu, \quad M_A = \int_A \langle v, \mathbf{D}^{-1} b \rangle \, d\mu, \quad (7.3)$$

with M_A the entropy flow to the medium and a reference term that vanishes for the volume reference, since $v = b - \mathbf{D} \nabla \log p$ and $\nabla \log p = \nabla \log f + \nabla \log m$ (Appendix S.B.1): the relative-information balance and the thermodynamic balance are two ledgers, and the bulk term depends on the declared reference. On the stationary gyrotor Φ_A^{rel} and B_A^{rel} vanish on every zone, the current running along the level sets of the density, while $\sigma_A = 0.172$ on the disk of Figure 10 equals the medium term (gates X4.1b, X4.5); on the driven ring $dx = (F - U'(x))dt + \sqrt{2T}dW$ with $U = \cos x, F = 1, T = 0.5$, the zone $[0, \pi]$ has $\Phi_A^{\text{rel}} = B_A^{\text{rel}} = 0.109$ and $\sigma_A = 0.916$ equal to the medium term minus the bulk term (gates RING.1, RING.2; with the Gibbs reference the bulk term changes and the identity holds with its reference term, gate RING.4; closed forms in Appendix S.A, Figure 12).

7.2 Numerical experiments

Protocol. Three libraries fix the closed forms of the gyrotor, of the double well and of the driven ring (LIB.1–LIB.8, PUIITS.1–PUIITS.2, SYM.1–SYM.6); C1, C2 the characterization controls; X1 the quotient, the hidden Fisher and the almost-sufficient sub-window; X2 the tower, its limit and its stability; X3 the calibrated geometry; X4 the structure, the two ledgers and the control; X5 the hidden arrow; X6 the emergence; X7 the two axes; X8 the predictive closure and the symbols; X9 the ring; S1 Shannon; S2 the harmonics; S3 the Gaussian leaf. Every gate was declared in writing




Figure 11: The quotient hides the arrow. Left: the entropy production seen by the observer of both coordinates grows with its time resolution and tends to σ , an instance of exhaustive recovery on path space; the observer of one coordinate reads zero at every Δ . Right: the entropy production of the fine process on $2 \cdot 10^4$ paths. [K5, R5]

before its first computation; twelve declarations were falsified at their first run and are kept in the archive with their revised statements. Everything runs on ten CPU cores. The gates are closed forms or exact propagations, with two sampled exceptions: the entropy productions of K5 and G3, averaged over $2 \cdot 10^4$ Euler–Maruyama paths, and the double-well statistics of E5, E6 and E7, time averages along single Euler–Maruyama trajectories with fixed seeds, quoted without error bars. Appendix S.D lists every gate with its measured value.

What the benchmarks test. The hidden arrow: every scalar linear observable is reversible, the window covariance being symmetric Toeplitz, while the two coordinates read $\sigma = 0.2793$, measured as 0.2839 ± 0.0037 on $2 \cdot 10^4$ paths (K5, Figure 11). Fisher identifiability: rank four from one coordinate, five from two (Figure 4). The stationary current and the two ledgers of a zone (Realization E1, Figure 10). The finite-volume projection (E2). The relaxation at equal temperatures as a radial geodesic (G3). The double well tests the predictive closure and the symbols (E6, E5, E7). The gyrator is deliberately minimal and therefore degenerate in useful ways: its stationary bulk term vanishes pointwise and every scalar linear marginal is reversible; these are stress tests of observability, and the driven ring supplies the missing nondegenerate balance.

8 Discussion

8.1 Relation to prior work

Table 4 says, for each named theorem, what is classical, what is read anew and what is proved here.

8.2 Scope and limitations

Table 5 says what each statement needs.

The benchmarks leave untested, by construction, generic non-Gaussian beliefs, non-radial geodesics of the slope geometry and nonlinear observation channels. The clauses of H1 are measured on two objects: the current and the ports on the gyrator and the driven ring, the closure on the double well; an object that closes and carries a sustained current is the next benchmark. The open problems are stated in Appendix S.E.

result	status	provenance
T1 Declared-Reference Principle	principle	this work (the datum and its references)
T2 Channel Ledger Theorem, C1	imported	Kullback, Csiszár; Leinster, Baez–Fritz for the characterization
T3 Retention Theorem	imported, proof recalled	Csiszár, Zellner; the one-line Gibbs identity
T4 Recovery-Kernel Theorem	imported	Blackwell
T5 Operational Fisher Theorem	imported	Čencov, Ay–Jost–Lê–Schwachhöfer
T6 Missing-Fisher Ledger Theorem	classical identity, IDS reading	Fisher, Orchard–Woodbury, Louis, Meng–Rubin
T7 Slope-Geometry Proposition	proved here, two lines	strong upper gradients and curves of maximal slope of Ambrosio–Gigli–Savaré
T8 RESG Theorem	proved here	extends Caraffa (2026) (Def. 3.6, Thms 4.10, 4.15) from the entropy of a density to relative entropy under a declared reference, with the radial identity; the Wasserstein calculus of entropy imported
T9 Resolution Ledger Theorem	assembly proposed here	the four chain rules are classical, their common ledger is this work
T10 Resolution Recovery Theorem	classical for D and I , proved here for Fisher and radius	Pinsker’s supremum, Barron; Lévy’s upward theorem
T11 Almost-Sufficient Stability Theorem	proved here	a lift of the bi-Lipschitz equivalence of Yamaguchi–Nozawa to $W_{2,h}$, $\mathcal{I}_{h,m}$ and d_I
T12 Predictive Closure Ledger Theorem	classical chain rule	informational closure of Bertschinger et al. and Pfante et al.; Shalizi–Crutchfield; new here: the tolerance pair (η, α) and the Pinsker control
Hyperbolic-Leaf Theorem (G6), G11	G6 restated, proof recalled; G11 proved here	Caraffa (2026, Thms 4.18–4.19); flat W_2 leaf, J_0/s^2 scaling, Stam; Loewner sandwiches
realizations, graphs, cells, symbols	measured	this work, gyrator, ring and double well

Table 4: Relation to prior work: status and provenance of each named theorem.

9 Conclusion

The proposed contribution is an architecture in which one declared-reference ledger is carried consistently across the three spaces and through finite-resolution projections. Learning, diffusion models, scaling laws, Turing machines, the compiler, gauge theories, gravity and quantum channels are outside this paper.

IDS begins from a reduction, and a representation comes second. A declared channel determines which candidate laws are distinguishable; relative entropy records exactly what the channel preserves and hides. On regular identifiable families, differentiating that ledger yields the Fisher metric and its missing-information decomposition. Lifting the same operational metric to probability measures over identifiable laws turns the Wasserstein slope of relative entropy into a calibrated geometry of revision. Finite descriptions then form a tower of channels in place of a primitive graph: their validity is quantified by the information that survives the projection, and their closure by the predictive ledger. This separation changes the role of discrete structure. Cells, graphs, conditional-independence relations and physical ports are realization-level objects, outside the axioms of the informational kernel. The Brownian gyrator, the driven ring and the double well show how the same bookkeeping reads hidden irreversibility, finite-volume projections, metastable symbols and predictive closure while keeping thermodynamic entropy production distinct from local relative-information balance. The remaining mathematical frontier is precise: the convergence of non-radial calibrated lengths under discretization, the Hodge decomposition

statement	needs
exact channel ledger (K2)	a measurable channel
Fisher geometry (O1, O3)	a regular identifiable stratum, scores in L^2
slope geometry (G3)	the contract HR0
radial identity (G5)	the prior as reference and an entropy-ergodic flow
exhaustive recovery (R5)	a generating filtration; scores in L^2 for Fisher
stability (G10)	a δ -almost sufficient projection, the same reference measure
predictive closure (E6)	a stationary process and a finite alphabet
harmonic resolution (R3)	a compact resolvent
thermodynamic heat reading (H2)	a physical realization under local detailed balance

Table 5: What each statement needs.

of hidden irreversibility, and the extension of the operational geometry to stratified identifiable quotients (Appendix S.E). These are questions about the fidelity of finite representations of a continuum geometry; the kernel assumes zero privileged graph.

References

- S. Amari and H. Nagaoka. *Methods of Information Geometry*, volume 191 of *Translations of Mathematical Monographs*. AMS and Oxford University Press, 2000. doi: 10.1090/mmono/191.
- L. Ambrosio, N. Gigli, and G. Savaré. *Gradient Flows in Metric Spaces and in the Space of Probability Measures*. Birkhäuser, 2008.
- L. Ambrosio, N. Gigli, and G. Savaré. Calculus and heat flow in metric measure spaces and applications to spaces with Ricci bounds from below. *Invent. Math.*, 195:289–391, 2014. doi: 10.1007/s00222-013-0456-1.
- V. I. Arnold. *Mathematical Methods of Classical Mechanics*. Springer, 2 edition, 1989.
- Nihat Ay, Jürgen Jost, Hông Vân Lê, and Lorenz Schwachhöfer. *Information Geometry*. Springer, 2017.
- John C. Baez and Tobias Fritz. A Bayesian characterization of relative entropy. *Theory Appl. Categ.*, 29:421–456, 2014.
- D. Bakry and M. Émery. Diffusions hypercontractives. In *Séminaire de probabilités XIX*, pages 177–206. Springer, 1985.
- Andrew R. Barron. The strong ergodic theorem for densities: generalized Shannon–McMillan–Breiman theorem. *Ann. Probab.*, 13:1292–1303, 1985.
- T. Bayes. An essay towards solving a problem in the doctrine of chances. *Phil. Trans. R. Soc.*, 53: 370–418, 1763.
- J.-D. Benamou and Y. Brenier. A computational fluid mechanics solution to the Monge–Kantorovich mass transfer problem. *Numer. Math.*, 84:375–393, 2000.
- Nils Bertschinger, Eckehard Olbrich, Nihat Ay, and Jürgen Jost. Information and closure in systems theory. In Stefan Artmann and Peter Dittrich, editors, *Explorations in the Complexity of Possible Life. Proceedings of the 7th German Workshop on Artificial Life*, pages 9–21, Amsterdam, 2006. IOS Press. ISBN 978-1-58603-644-7.

- Pier Giovanni Bissiri and Stephen G. Walker. Converting information into probability measures with the Kullback–Leibler divergence. *Ann. Inst. Statist. Math.*, 64(6):1139–1160, 2012. doi: 10.1007/s10463-012-0350-4.
- Pier Giovanni Bissiri, Chris C. Holmes, and Stephen G. Walker. A general framework for updating belief distributions. *J. R. Stat. Soc. Ser. B*, 78(5):1103–1130, 2016. doi: 10.1111/rssb.12158.
- D. Blackwell. Equivalent comparisons of experiments. *Ann. Math. Statist.*, 24:265–272, 1953.
- Leon Bungert and Dejan Slepčev. Convergence of graph Dirichlet energies and graph Laplacians on intersecting manifolds of varying dimensions. *Calc. Var. Partial Differential Equations*, 2026. arXiv:2509.24458.
- Laurent Caraffa. A transport-based geometry of belief-cost, 2026. URL <https://arxiv.org/abs/2606.21585v5>. arXiv:2606.21585v5, 6 July 2026; theorem numbers of v5.
- Yongxin Chen, Tryphon T. Georgiou, and Michele Pavon. On the relation between optimal transport and Schrödinger bridges: a stochastic control viewpoint. *J. Optim. Theory Appl.*, 169(2):671–691, 2016. doi: 10.1007/s10957-015-0803-z. arXiv:1412.4430.
- R. R. Coifman and S. Lafon. Diffusion maps. *Appl. Comput. Harmon. Anal.*, 21:5–30, 2006.
- T. M. Cover and J. A. Thomas. *Elements of Information Theory*. Wiley, 2 edition, 2006.
- I. Csiszár. Information-type measures of difference of probability distributions and indirect observations. *Studia Sci. Math. Hungar.*, 2:299–318, 1967.
- I. Csiszár. I -divergence geometry of probability distributions and minimization problems. *Ann. Probab.*, 3:146–158, 1975.
- Julius Degünther, Jann van der Meer, and Udo Seifert. Fluctuating entropy production on the coarse-grained level: inference and localization of irreversibility. *Phys. Rev. Research*, 6:023175, 2024.
- K. D. Elworthy. *Stochastic Differential Equations on Manifolds*. Cambridge University Press, 1982.
- M. Erbar. The heat equation on manifolds as a gradient flow in the Wasserstein space. *Ann. Inst. H. Poincaré Probab. Statist.*, 46:1–23, 2010.
- M. Esposito and C. Van den Broeck. Three faces of the second law. I. Master equation formulation. *Phys. Rev. E*, 82:011143, 2010.
- R. Filliger and P. Reimann. Brownian gyrator: a minimal heat engine on the nanoscale. *Phys. Rev. Lett.*, 99:230602, 2007.
- D. Forkert, J. Maas, and L. Portinale. Evolutionary Γ -convergence of entropic gradient flow structures for Fokker–Planck equations in multiple dimensions. *SIAM J. Math. Anal.*, 54:4297–4333, 2022.
- A. Fukushi, Y. Nakanishi-Ohno, and T. Matsuda. Flatness of location-scale-shape models under the Wasserstein metric. *Information Geometry*, 2026. arXiv:2511.09959.
- Nicolas Gagné and Prakash Panangaden. A categorical characterization of relative entropy on standard Borel spaces. *Electron. Notes Theor. Comput. Sci.*, 336:135–153, 2018.
- Bernhard C. Geiger. Information-theoretic reduction of Markov chains. *Computer Science Review*, 59, 2026. arXiv:2204.13896.

- Ivan Gentil, Christian Léonard, and Luigia Ripani. About the analogy between optimal transport and minimal entropy. *Ann. Fac. Sci. Toulouse Math. (6)*, 26(3):569–600, 2017. doi: 10.5802/afst.1546. arXiv:1510.08230.
- A. Gomez-Marin, J. M. R. Parrondo, and C. Van den Broeck. Lower bounds on dissipation upon coarse graining. *Phys. Rev. E*, 78(1):011107, 2008. doi: 10.1103/PhysRevE.78.011107.
- E. P. Hsu. *Stochastic Analysis on Manifolds*. AMS, 2002.
- R. Jordan, D. Kinderlehrer, and F. Otto. The variational formulation of the Fokker–Planck equation. *SIAM J. Math. Anal.*, 29:1–17, 1998.
- R. Kawai, J. M. R. Parrondo, and C. Van den Broeck. Dissipation: the phase-space perspective. *Phys. Rev. Lett.*, 98(8):080602, 2007. doi: 10.1103/PhysRevLett.98.080602.
- H. A. Kramers. Brownian motion in a field of force and the diffusion model of chemical reactions. *Physica*, 7:284–304, 1940.
- S. Kullback and R. A. Leibler. On information and sufficiency. *Ann. Math. Statist.*, 22:79–86, 1951.
- S. L. Lauritzen. *Graphical Models*. Oxford University Press, 1996.
- Tom Leinster. A short characterization of relative entropy. *J. Math. Phys.*, 60:023302, 2019.
- C. Léonard. Girsanov theory under a finite entropy condition. In *Séminaire de Probabilités XLIV*, pages 429–465. Springer, 2012.
- C. Léonard. A survey of the Schrödinger problem and some of its connections with optimal transport. *Discrete Contin. Dyn. Syst.*, 34:1533–1574, 2014.
- D. V. Lindley. On a measure of the information provided by an experiment. *Ann. Math. Statist.*, 27:986–1005, 1956.
- Thomas A. Louis. Finding the observed information matrix when using the EM algorithm. *J. R. Stat. Soc. Ser. B*, 44:226–233, 1982.
- J. Maas. Gradient flows of the entropy for finite Markov chains. *J. Funct. Anal.*, 261:2250–2292, 2011.
- Robert J. McCann. A convexity principle for interacting gases. *Adv. Math.*, 128(1):153–179, 1997. doi: 10.1006/aima.1997.1634.
- Xiao-Li Meng and Donald B. Rubin. Using EM to obtain asymptotic variance-covariance matrices: the SEM algorithm. *J. Amer. Statist. Assoc.*, 86:899–909, 1991.
- G. Nicolis and I. Prigogine. *Self-Organization in Nonequilibrium Systems*. Wiley, 1977.
- Terence Orchard and Max A. Woodbury. A missing information principle: theory and applications. In *Proc. Sixth Berkeley Symp. Math. Statist. Prob.*, volume 1, pages 697–715, 1972.
- F. Otto. The geometry of dissipative evolution equations: the porous medium equation. *Comm. PDE*, 26:101–174, 2001.
- F. Otto and C. Villani. Generalization of an inequality by Talagrand and links with the logarithmic Sobolev inequality. *J. Funct. Anal.*, 173:361–400, 2000.
- Liam Paninski. Estimation of entropy and mutual information. *Neural Computation*, 15:1191–1253, 2003.

- Dénes Petz. Sufficiency of channels over von Neumann algebras. *Q. J. Math.*, 39(1):97–108, 1988. doi: 10.1093/qmath/39.1.97.
- Oliver Pfante, Eckehard Olbrich, Nils Bertschinger, Nihat Ay, and Jürgen Jost. Closure measures for coarse-graining of the tent map. *Chaos*, 24(1):013136, 2014. doi: 10.1063/1.4869075.
- Mark S. Pinsker. *Information and Information Stability of Random Variables and Processes*. Holden-Day, 1964.
- Hannes Risken. *The Fokker–Planck Equation*. Springer, 2 edition, 1989.
- J. Schnakenberg. Network theory of microscopic and macroscopic behavior of master equation systems. *Rev. Mod. Phys.*, 48:571–585, 1976.
- U. Seifert. Entropy production along a stochastic trajectory and an integral fluctuation theorem. *Phys. Rev. Lett.*, 95:040602, 2005.
- U. Seifert. Stochastic thermodynamics, fluctuation theorems and molecular machines. *Rep. Prog. Phys.*, 75:126001, 2012.
- Cosma Rohilla Shalizi and James P. Crutchfield. Computational mechanics: pattern and prediction, structure and simplicity. *J. Stat. Phys.*, 104:817–879, 2001.
- C. E. Shannon. A mathematical theory of communication. *Bell Syst. Tech. J.*, 27:379–423, 623–656, 1948.
- C. E. Shannon. Coding theorems for a discrete source with a fidelity criterion. In *IRE Nat. Conv. Rec.*, volume 7, pages 142–163, 1959.
- J. E. Shore and R. W. Johnson. Axiomatic derivation of the principle of maximum entropy and the principle of minimum cross-entropy. *IEEE Trans. Inform. Theory*, 26(1):26–37, 1980. doi: 10.1109/TIT.1980.1056144.
- A. J. Stam. Some inequalities satisfied by the quantities of information of Fisher and Shannon. *Inform. Control*, 2:101–112, 1959.
- K.-T. Sturm. On the geometry of metric measure spaces I. *Acta Math.*, 196:65–131, 2006.
- Naftali Tishby, Fernando C. Pereira, and William Bialek. The information bottleneck method. In *Proc. 37th Allerton Conf. Communication, Control and Computing*, pages 368–377, 1999.
- Aad W. van der Vaart. *Asymptotic Statistics*. Cambridge University Press, 1998.
- N. N. Čencov. *Statistical Decision Rules and Optimal Inference*. AMS, 1982.
- C. Villani. *Optimal Transport: Old and New*. Springer, 2009.
- M.-K. von Renesse and K.-T. Sturm. Transport inequalities, gradient estimates, entropy and Ricci curvature. *Comm. Pure Appl. Math.*, 58:923–940, 2005.
- G. Weiss. Time-reversibility of linear stochastic processes. *J. Appl. Probab.*, 12:831–836, 1975.
- David Williams. *Probability with Martingales*. Cambridge University Press, 1991.
- P. M. Williams. Bayesian conditionalisation and the principle of minimum information. *Brit. J. Philos. Sci.*, 31(2):131–144, 1980. doi: 10.1093/bjps/31.2.131.
- Kaori Yamaguchi and Hiraku Nozawa. On statistics which are almost sufficient from the viewpoint of the Fisher metrics. *Information Geometry*, 7:543–553, 2024.
- A. Zellner. Optimal information processing and Bayes’s theorem. *Amer. Statist.*, 42:278–280, 1988.

Supplement

S.A Closed forms

The gyrator. With $A = \begin{pmatrix} k_1+k & -k \\ -k & k_2+k \end{pmatrix}$ and $\mathbf{D} = \text{diag}(T_1, T_2)$, the stationary covariance solves $AC + CA = 2\mathbf{D}$; writing $\Delta = (2k + k_1 + k_2)(kk_1 + kk_2 + k_1k_2)$,

$$C_{11} = \frac{T_1k^2 + T_1kk_1 + 3T_1kk_2 + T_1k_1k_2 + T_1k_2^2 + T_2k^2}{\Delta}, \quad C_{12} = \frac{k(T_1k + T_1k_2 + T_2k + T_2k_1)}{\Delta},$$

and C_{22} by exchange of the two sides. $\Gamma = \mathbf{D}C^{-1} - A$ satisfies $\Gamma C = sJ$ with $J = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}$; the stream function of the current is $\psi = sp$; the entropy production, the heats and Clausius are Equation (7.1) (gates SYM.1–SYM.4, symbolic, exact). The three quantities factor as $k(T_1 - T_2)$, $k^2(T_1 - T_2)^2$ and $k(T_1 - T_2)$ times a cofactor of constant sign on the positive orthant, so they vanish exactly when $k(T_1 - T_2) = 0$ (gate SYM.6). The window covariance of a scalar stationary process is symmetric Toeplitz, hence invariant under time reversal (gate SYM.5).

The driven ring. On the circle of length L , $dx = (F - U'(x))dt + \sqrt{2T}dW$ with $U(x) = a \cos(2\pi x/L)$ has the stationary density $p = g/Z$, $g(x) = e^{-\varphi(x)} \int_x^{x+L} e^{\varphi(y)} dy$, $\varphi = (U - Fx)/T$, and the constant current $j = T(1 - e^{-FL/T})/Z$ (Risken, 1989); the periodic extension gives $\int_x^{x+L} e^\varphi = (E(L) - E(x)) + e^{-FL/T}E(x)$ with $E(x) = \int_0^x e^\varphi$. For a zone $A = [a, b]$: $\Phi_A^{\text{rel}} = j \log(p(b)/p(a))$, $B_A^{\text{rel}} = \int_A j(\log p)' dx = \Phi_A^{\text{rel}}$, the content being stationary; $\sigma_A = \int_A j^2/(Tp) dx$ and the medium term $M_A = (j/T)[F(b-a) - U(b) + U(a)]$ satisfy $\sigma_A = M_A - B_A^{\text{rel}}$ (Equation (7.3)). At $a = 0$ the density is uniform, the current F/L is harmonic and $\sigma = F^2/T$ (gate SYM.6). With $a = 1$, $F = 1$, $T = 0.5$, $L = 2\pi$: the closed form is stationary to $2 \cdot 10^{-7}$ on 65 536 points with a spectral derivative, $j = 0.0997$, and on the zone $[0, \pi]$, $\Phi_A^{\text{rel}} = B_A^{\text{rel}} = 0.109$, $\sigma_A = 0.916$, $M_A = 1.025$ (gates RING.1, RING.2; Figure 12).

S.B Proofs

This section proves the theorems of the body from the imported results named in Proposition G1 and in the text; everything else is proved here.

S.B.1 The kernel: retention, sufficiency, the score

Proof of the second claim of Theorem K3, another divergence, another rule (C2). Let f be convex, differentiable on $(0, \infty)$, $f(1) = 0$, $D_f(\mu||\pi) = \sum_i \pi_i f(\mu_i/\pi_i)$ on a simplex with π of full support and likelihood L with positive entries. Suppose the least-deformation update equals $\pi_{\mathcal{D}}$ for every prior of full support and every positive likelihood. The minimizer $\pi_{\mathcal{D}}$ is interior, and the stationarity condition of $\min_{\mu} \{\mathbb{E}_{\mu}[-\log L] + D_f(\mu||\pi)\}$ on the simplex reads $f'(\mu_i/\pi_i) = \log L_i + c$ for every i , with a multiplier c that depends on the instance. At $\mu = \pi_{\mathcal{D}}$, with $u_i = L_i/Z$, it reads $g(u_i) = c + \log Z$ for every i , where $g(u) := f'(u) - \log u$: g takes one value on the ratios of each instance. Take two states, $0 < u < 1 < w$, the prior $(p, 1-p)$ with $p = (w-1)/(w-u)$ and the likelihood (u, w) : then $Z = pu + (1-p)w = 1$, the ratios are u and w , and $g(u) = g(w)$. Fixing $w > 1$ gives $g(u) = g(w)$ for every $u \in (0, 1)$, and fixing $u < 1$ gives $g(w) = g(u)$ for every $w > 1$; f' is continuous, f being convex and differentiable, so g is constant on $(0, \infty)$, $f'(u) = \log u + c'$, $f(t) = t \log t + a(t-1)$ and $D_f = D$ on probabilities. For relative entropy the residual $f'(u) - \log u$ is constant; for the squared Hellinger distance, $f'(u) = 1 - u^{-1/2}$, and for χ^2 , $f'(u) = 2(u-1)$, it varies by 1.85 and $1.8 \cdot 10^3$ in variance on $[10^{-2}, 10^2]$, and the least-deformation update differs from Bayes by 0.03 to 0.13 in total variation on random evidences (gates C2.4b, C2.2; the optimizer control C2.1 finds Bayes to 10^{-9} with relative entropy). $\square$

Proof of Theorem K4, sufficiency and the recovery kernel. When $D(P \parallel R) < \infty$, a zero hidden term in Equation (3.2) means $\mathbb{E}_{KP} D(P_{X|Y} \parallel R_{X|Y}) = 0$, that is $P_{X|Y} = R_{X|Y}$ KP -almost surely: the conditional law of the fine variable given the observation is the same under P and under R , which is sufficiency of K for the pair. For the declared family, let L be the Markov kernel $y \mapsto P_{X|Y=y}$, the same for every λ by sufficiency; then $LKP_\lambda = \int P_{X|Y=y}(KP_\lambda)(dy) = P_\lambda$, so L recovers each law of the family from its observed law. Conversely, if such an L exists, the observed experiment K and the original one are each a garbling of the other, so they are equivalent in Blackwell's order and K is sufficient (Blackwell, 1953). On a regular stratum the ledger of Equation (4.2) then has zero hidden term, and the Fisher metric of the observed laws equals h . $\square$

Proof of Theorem O3 and of the score projection O2. Let p_θ be dominated and differentiable in θ with a locally dominated derivative, $Y = q(X)$ and p_θ^Y the density of Y . Differentiating $p_\theta^Y(y) = \int_{q^{-1}(y)} p_\theta d\nu_y$ under the integral gives $\partial_v p_\theta^Y(y) = p_\theta^Y(y) \mathbb{E}_\theta[S_v \mid Y = y]$, that is $S_v^Y = \mathbb{E}[S_v \mid Y]$, Fisher's identity for the score of an observation (O2). With $S_v \in L^2(P_\theta)$ and $\mathbb{E}S_v = 0$, the law of total variance gives $h(v, v) = \mathbb{E}(\mathbb{E}[S_v \mid Y])^2 + \mathbb{E} \text{Var}(S_v \mid Y) = h_Y(v, v) + h_Y^{\text{hid}}(v, v)$, the expectation over Y of Louis's identity between observed, complete and missing information (Louis, 1982); polarization gives the bilinear form. The hidden term is the second derivative at λ of the hidden relative information $\lambda' \mapsto \mathbb{E}_{KP_{\lambda'}} D(P_{X|Y} \parallel P_{\lambda|Y})$, by Equation (3.2) and Equation (4.1). On the gyrator the variance of the conditional score equals the Fisher metric of the marginal in the five directions to 10^{-14} (gate X1.2). $\square$

S.B.2 The slope geometry

Setting. (Λ, h) is a smooth, connected, complete Riemannian manifold of dimension n , with geodesic distance d_h , volume dV_h , gradient ∇_h and Laplace–Beltrami operator Δ_h . The reference is $m = e^{-V} dV_h$ with $V \in C^2(\Lambda)$, finite or σ -finite, under the Bakry–Émery condition

$$\text{CD}(K, \infty) : \quad \text{Ric}_h + \text{Hess}_h V \geq K h \quad \text{for some } K \in \mathbb{R} \quad (\text{S.B.1})$$

(Bakry and Émery, 1985; von Renesse and Sturm, 2005); this is the contract HR0 of Section 4.2. For $\mu = f m$,

$$\mathcal{H}_m(\mu) := \int_\Lambda f \log f \, dm, \quad \mathcal{I}_{h,m}(\mu) := \int_\Lambda \|\nabla_h \log f\|_h^2 \, d\mu = 4 \int_\Lambda \|\nabla_h \sqrt{f}\|_h^2 \, dm,$$

the second when $\sqrt{f} \in W_{\text{loc}}^{1,2}(m)$ with a finite integral and $+\infty$ otherwise, both $+\infty$ when μ charges a set of m -measure zero. Under Equation (S.B.1) the reference has at most Gaussian volume growth (Sturm, 2006), so that $\mathcal{H}_m$ is well defined with values in $(-\infty, +\infty]$ on the space $\mathcal{P}_2(\Lambda)$ of probability measures with finite second moment (Ambrosio et al., 2008, §9.4). The regular domain is $\mathcal{P}_{\text{reg}} := \{\mu \in \mathcal{P}_2(\Lambda) : \mu = f m, \mathcal{H}_m(\mu) < \infty, \mathcal{I}_{h,m}(\mu) < \infty\}$. $W_{2,h}$ is the quadratic Wasserstein distance of (Λ, d_h) ; a curve $\gamma : [0, 1] \rightarrow \mathcal{P}_2(\Lambda)$ is absolutely continuous (AC) when $W_{2,h}(\gamma_s, \gamma_t) \leq \int_s^t g(r) \, dr$ for some $g \in L^1(0, 1)$, and its metric derivative $|\dot{\gamma}_t| := \lim_{s \rightarrow t} W_{2,h}(\gamma_s, \gamma_t)/|s - t|$ exists for a.e. t and is the least such g (Ambrosio et al., 2008, §1.1). For a functional F on a metric space (X, d) , the descending metric slope at x with $F(x) < \infty$ is $|\partial F|_d(x) := \limsup_{y \rightarrow x} (F(x) - F(y))^+ / d(x, y)$.

Proposition G1 (The imported Wasserstein calculus of entropy · kernel · IMPORTED). *Under Equation (S.B.1): (i) slope: for $\mu \in \mathcal{P}_{\text{reg}}$, $|\partial \mathcal{H}_m|_{W_{2,h}}^2(\mu) = \mathcal{I}_{h,m}(\mu)$; (ii) strong upper gradient: for every AC curve γ in $\mathcal{P}_{\text{reg}}$ with $\int_0^1 \sqrt{\mathcal{I}_{h,m}(\gamma_t)} |\dot{\gamma}_t| \, dt < \infty$, the map $t \mapsto \mathcal{H}_m(\gamma_t)$ is absolutely continuous and*

$$\left| \frac{d}{dt} \mathcal{H}_m(\gamma_t) \right| \leq \sqrt{\mathcal{I}_{h,m}(\gamma_t)} |\dot{\gamma}_t| \quad \text{for a.e. } t; \quad (\text{S.B.2})$$

(iii) gradient flow: for $\mu \in \mathcal{P}_{\text{reg}}$ the law $(\mu_t)_{t \geq 0}$ of the m -reversible diffusion with generator $\Delta_h - \nabla_h V \cdot \nabla_h$ started from μ is an AC curve in $\mathcal{P}_{\text{reg}}$, it is the gradient flow of $\mathcal{H}_m$ in $(\mathcal{P}_2(\Lambda), W_{2,h})$,

$t \mapsto \mathcal{H}_m(\mu_t)$ is nonincreasing, and for a.e. t

$$\frac{d}{dt} \mathcal{H}_m(\mu_t) = -\mathcal{I}_{h,m}(\mu_t) = -\sqrt{\mathcal{I}_{h,m}(\mu_t)} |\dot{\mu}_t|, \quad |\dot{\mu}_t| = \sqrt{\mathcal{I}_{h,m}(\mu_t)}; \quad (\text{S.B.3})$$

(iv) lower semicontinuity: $\mathcal{H}_m$ is lower semicontinuous on $(\mathcal{P}_2(\Lambda), W_{2,h})$ (Jordan et al., 1998; Otto, 2001; Ambrosio et al., 2008; Erbar, 2010).

Sources. (i) is Otto's identity, proved in the metric setting for λ -convex functionals (Otto, 2001; Ambrosio et al., 2008); under Equation (S.B.1) the entropy is K -convex along Wasserstein geodesics (von Renesse and Sturm, 2005), and the manifold case is treated by Erbar (2010). (ii) is the fact that the slope of a λ -convex lower semicontinuous functional is a strong upper gradient (Ambrosio et al., 2008, §§1.2, 2.4). (iii) is the identification of the heat flow with the entropy gradient flow (Jordan et al., 1998; Otto, 2001; Erbar, 2010), with the energy identity that characterizes gradient flows in metric spaces (Ambrosio et al., 2008, §1.3). (iv) is Ambrosio et al. (2008, §9.4) under the Gaussian volume growth. For a weighted reference, $m = e^{-V} dV_h$, the heat flow of (iii) is the gradient flow of the entropy on $\text{RCD}(K, \infty)$ spaces (Ambrosio et al., 2014). These four facts are the imports of the slope geometry; the logarithmic Sobolev and Talagrand inequalities used below are those of Bakry and Émery (1985) and Otto and Villani (2000).

Length and distance. For an AC curve γ in $\mathcal{P}_{\text{reg}}$, $L_I(\gamma) := \ell_\star \int_0^1 \sqrt{\mathcal{I}_{h,m}(\gamma_t)} |\dot{\gamma}_t| dt \in [0, \infty]$, and $d_I(\mu, \nu) := \inf\{L_I(\gamma) : \gamma \text{ AC in } \mathcal{P}_{\text{reg}}, \gamma_0 = \mu, \gamma_1 = \nu\}$, with $\inf \emptyset = +\infty$ (Definition G2). The integrand is measurable because $\mathcal{I}_{h,m}$ is lower semicontinuous; L_I is invariant under reparametrization, so d_I is symmetric and satisfies the triangle inequality by concatenation. By Proposition G1 (i) the conformal factor is the metric slope of the entropy itself; it vanishes exactly at $\mu = m$ when m is a probability measure.

Proof of Theorem G3 (i), the floor and the horizon. If $L_I(\gamma) = \infty$ the inequality holds. Otherwise Proposition G1 (ii) applies: $t \mapsto \mathcal{H}_m(\gamma_t)$ is absolutely continuous and, integrating Equation (S.B.2),

$$L_I(\gamma) = \ell_\star \int_0^1 \sqrt{\mathcal{I}_{h,m}(\gamma_t)} |\dot{\gamma}_t| dt \geq \ell_\star \int_0^1 \left| \frac{d}{dt} \mathcal{H}_m(\gamma_t) \right| dt \geq \ell_\star |\mathcal{H}_m(\gamma_1) - \mathcal{H}_m(\gamma_0)|.$$

Taking the infimum over curves gives $d_I(\mu, \nu) \geq \ell_\star |\mathcal{H}_m(\nu) - \mathcal{H}_m(\mu)|$: $\mathcal{H}_m$ is $(1/\ell_\star)$ -Lipschitz for d_I . For the horizon, let (μ_k) be a sequence in $\mathcal{P}_{\text{reg}}$ with $\mathcal{H}_m(\mu_k) \rightarrow +\infty$; then $d_I(\mu_0, \mu_k) \rightarrow +\infty$ by the floor. When $\mu_k \rightarrow \delta_\lambda$ in $W_{2,h}$ and $m(\{\lambda\}) = 0$, the Dirac mass charges a set of m -measure zero, so $\mathcal{H}_m(\delta_\lambda) = +\infty$, and Proposition G1 (iv) gives $\liminf_k \mathcal{H}_m(\mu_k) \geq \mathcal{H}_m(\delta_\lambda) = +\infty$, while $W_{2,h}(\mu_0, \mu_k) \rightarrow W_{2,h}(\mu_0, \delta_\lambda) = (\int d_h(x, \lambda)^2 d\mu_0)^{1/2} < \infty$. With an atomic reference, $D(\delta_\lambda \| m) = -\log m(\{\lambda\})$ is finite and certainty is at finite distance: the horizon is a statement of the continuous sector. $\square$

Conformal reliefs. For a lower semicontinuous relief $a : \mathcal{P}_2(\Lambda) \rightarrow [0, \infty]$, write $\mathcal{O}_a := \{0 < a < \infty\}$ and $L_a(\gamma) := \int_0^1 a(\gamma_t) |\dot{\gamma}_t| dt$, $d_a(\mu, \nu) := \inf\{L_a(\gamma)\}$ over AC curves from μ to ν , so that $L_I = L_a$ and $d_I = d_a$ for $a = \ell_\star \sqrt{\mathcal{I}_{h,m}}$. $\mathcal{H}_m$ has uniform slope $1/\ell_\star$ for a when $|\partial \mathcal{H}_m|_{d_a}(\mu) = 1/\ell_\star$ at every $\mu \in \mathcal{O}_a \cap \{0 < \mathcal{I}_{h,m} < \infty\}$.

Proposition L2 (Conformal contraction $\cdot$ kernel $\cdot$ PROVED). *Let a be continuous at $\mu \in \mathcal{O}_a$ in the $W_{2,h}$ topology, with $\mathcal{O}_a$ open. Then $d_a(\mu, \nu) = a(\mu) W_{2,h}(\mu, \nu) (1 + o(1))$ as $W_{2,h}(\mu, \nu) \rightarrow 0$.*

Proof. Fix $\delta \in (0, a(\mu))$ and $r > 0$ such that $B := \{\nu : W_{2,h}(\mu, \nu) < r\} \subset \mathcal{O}_a$ and $|a - a(\mu)| \leq \delta$ on B , and let $W_{2,h}(\mu, \nu) < r(a(\mu) - \delta)/(a(\mu) + \delta)$. Upper bound: $(\mathcal{P}_2(\Lambda), W_{2,h})$ is a geodesic space since (Λ, d_h) is complete (Ambrosio et al., 2008, ch. 7); a constant-speed geodesic from μ to ν stays in B , so $d_a(\mu, \nu) \leq (a(\mu) + \delta) W_{2,h}(\mu, \nu)$. Lower bound: an AC curve from μ to ν that stays

in B has $L_a \geq (a(\mu) - \delta) \int |\dot{\gamma}| \geq (a(\mu) - \delta) W_{2,h}(\mu, \nu)$; one that leaves B has $W_{2,h}$ -length at least r before leaving, hence $L_a \geq (a(\mu) - \delta)r > (a(\mu) + \delta)W_{2,h}(\mu, \nu)$, which is longer than the geodesic. Thus $(a(\mu) - \delta) \leq d_a(\mu, \nu)/W_{2,h}(\mu, \nu) \leq (a(\mu) + \delta)$ for ν close to μ , and δ is arbitrary. $\square$

Proof of Theorem G3 (ii), the constant slope is unique (G4). (a) *Only the entropy slope has uniform slope.* Let a be continuous and positive on the open set $\mathcal{O}_a$. By Proposition L2, d_a and $W_{2,h}$ are locally equivalent at $\mu \in \mathcal{O}_a \cap \mathcal{P}_{\text{reg}}$, so the lim sup defining $|\partial \mathcal{H}_m|_{d_a}(\mu)$ may be taken along $\nu \rightarrow \mu$ in $W_{2,h}$, and

$$|\partial \mathcal{H}_m|_{d_a}(\mu) = \limsup_{\nu \rightarrow \mu} \frac{(\mathcal{H}_m(\mu) - \mathcal{H}_m(\nu))^+}{a(\mu)W_{2,h}(\mu, \nu)(1 + o(1))} = \frac{|\partial \mathcal{H}_m|_{W_{2,h}}(\mu)}{a(\mu)} = \frac{\sqrt{\mathcal{I}_{h,m}(\mu)}}{a(\mu)} \quad (\text{S.B.4})$$

by Proposition G1 (i). Constancy at $1/\ell_\star$ on $\mathcal{O}_a \cap \{0 < \mathcal{I}_{h,m} < \infty\}$ is then exactly $a = \ell_\star \sqrt{\mathcal{I}_{h,m}}$ there. (b) *The entropy slope has uniform slope*, without any continuity assumption. Upper bound: for every $\nu \in \mathcal{P}_{\text{reg}}$ the floor gives $(\mathcal{H}_m(\mu) - \mathcal{H}_m(\nu))^+ \leq d_I(\mu, \nu)/\ell_\star$, hence $|\partial \mathcal{H}_m|_{d_I}(\mu) \leq 1/\ell_\star$. Lower bound: let (μ_t) be the gradient flow of Proposition G1 (iii) from μ with $\mathcal{I}_{h,m}(\mu) > 0$. By Equation (S.B.3), $d_I(\mu, \mu_t) \leq L_I(\mu|_{[0,t]}) = \ell_\star \int_0^t \mathcal{I}_{h,m}(\mu_r) dr = \ell_\star(\mathcal{H}_m(\mu) - \mathcal{H}_m(\mu_t))$, which tends to 0 as $t \rightarrow 0$ since $\mathcal{H}_m \circ \mu$ is continuous. Moreover $\mathcal{H}_m(\mu_t) < \mathcal{H}_m(\mu)$ for every $t > 0$: otherwise $\mathcal{I}_{h,m}(\mu_r) = 0$ for a.e. r in some $(0, t_0)$, i.e. μ_r is the normalized reference $\bar{m}$ for a.e. $r \in (0, t_0)$, hence for all $r \in (0, t_0]$ by continuity of the curve, and $\mu = \lim_{r \rightarrow 0} \mu_r = \bar{m}$, in contradiction with $\mathcal{I}_{h,m}(\mu) > 0 = \mathcal{I}_{h,m}(\bar{m})$. So $\mu_t \neq \mu$, $\mu_t \rightarrow \mu$ in d_I , and $(\mathcal{H}_m(\mu) - \mathcal{H}_m(\mu_t))/d_I(\mu, \mu_t) \geq 1/\ell_\star$, whence $|\partial \mathcal{H}_m|_{d_I}(\mu) \geq 1/\ell_\star$. Therefore the relief $\ell_\star \sqrt{\mathcal{I}_{h,m}}$ gives relative entropy the constant metric slope $1/\ell_\star$ at every regular belief, by (b), and a relief continuous at a regular belief gives it the slope $1/\ell_\star$ there exactly when it equals $\ell_\star \sqrt{\mathcal{I}_{h,m}}$ there, by (a): the weight is pointwise rigid. The statement is pointwise because $\sqrt{\mathcal{I}_{h,m}}$ is only lower semicontinuous on $(\mathcal{P}_2(\Lambda), W_{2,h})$, and small oscillations of a density make it arbitrarily large; on a smooth finite-dimensional family of regular beliefs, such as the leaves below, where $\mathcal{I}_{h,m}$ is continuous for the topology of the family, (a) applied to the length structure of the family makes the weight unique. $\square$

The radial distance. Let the reference be a probability measure π , so that $\mathcal{H}_\pi = D(\cdot \| \pi)$. A curve *reaches π in relative entropy* when it is defined on $[0, 1)$, AC on every $[0, t]$, and $D(\gamma_t \| \pi) \rightarrow 0$ as $t \rightarrow 1$; its length is $L_I(\gamma) := \lim_{t \rightarrow 1} L_I(\gamma|_{[0,t]})$. For $\mu \in \mathcal{P}_{\text{reg}}$, $\rho_I(\mu) := \inf\{L_I(\gamma) : \gamma \text{ reaches } \pi \text{ in relative entropy}, \gamma_0 = \mu\}$.

Proof of Theorem G3 (iii), the radial identity (G5). Assume the gradient flow (μ_t) of Proposition G1 (iii) started from μ is ergodic in relative entropy, $D(\mu_t \| \pi) \rightarrow 0$. *Lower bound.* Let γ reach π in relative entropy from μ . For $t < 1$ the floor on $[0, t]$ gives $L_I(\gamma|_{[0,t]}) \geq \ell_\star(D(\mu \| \pi) - D(\gamma_t \| \pi))$; letting $t \rightarrow 1$, $L_I(\gamma) \geq \ell_\star D(\mu \| \pi)$. *Upper bound.* Along the flow, Equation (S.B.3) gives for every T

$$\begin{aligned} L_I(\mu|_{[0,T]}) &= \ell_\star \int_0^T \sqrt{\mathcal{I}_{h,\pi}(\mu_t)} |\dot{\mu}_t| dt = -\ell_\star \int_0^T \frac{d}{dt} D(\mu_t \| \pi) dt \\ &= \ell_\star (D(\mu \| \pi) - D(\mu_T \| \pi)) \xrightarrow{T \rightarrow \infty} \ell_\star D(\mu \| \pi). \end{aligned}$$

The reparametrization $t = T/(1 + T)$ turns the flow into a curve on $[0, 1)$ reaching π in relative entropy, with the same length. Hence $\rho_I(\mu) = \ell_\star D(\mu \| \pi)$ and the gradient-flow trace realizes the infimum: it is the radial geodesic. Along every gradient-flow trace, Equation (S.B.3) reads $d\ell_I = |dr|$ for $r(\mu) = \ell_\star D(\mu \| \pi)$: relative entropy is the arc-length coordinate from the reference along the radial geodesics, and its level sets are the calibrated spheres about π . *A sufficient condition.* If π satisfies the logarithmic Sobolev inequality $D(\mu \| \pi) \leq \mathcal{I}_{h,\pi}(\mu)/(2K)$ for all $\mu \ll \pi$, which holds under $\text{CD}(K, \infty)$ with $K > 0$ (Bakry and Émery, 1985; Otto and Villani, 2000), then $\frac{d}{dt} D(\mu_t \| \pi) = -\mathcal{I}_{h,\pi}(\mu_t) \leq -2KD(\mu_t \| \pi)$ and Grönwall's lemma gives $D(\mu_t \| \pi) \leq e^{-2Kt} D(\mu \| \pi) \rightarrow 0$; by Talagrand's inequality $W_{2,h}(\mu_t, \pi)^2 \leq \frac{2}{K} D(\mu_t \| \pi) \rightarrow 0$ as well. *A metric.*

Under the logarithmic Sobolev inequality, extend the length structure to $\bar{\mathcal{P}} := \mathcal{P}_{\text{reg}} \cup \{\pi\}$ by admitting the curves that reach π in relative entropy, their reversals and their concatenations; the resulting infimum of lengths $\bar{d}_I$ is symmetric and satisfies the triangle inequality, $\bar{d}_I(\pi, \mu) = \rho_I(\mu)$, $\bar{d}_I(\mu, \nu) \leq \ell_\star(D(\mu\|\pi) + D(\nu\|\pi)) < \infty$ through the centre, and for $\mu \neq \nu$ with $D_\mu := D(\mu\|\pi) > 0$,

$$\bar{d}_I(\mu, \nu) \geq \min \left\{ \frac{1}{2} \ell_\star D_\mu, \ell_\star \sqrt{K D_\mu} W_{2,h}(\mu, \nu) \right\} > 0,$$

because a curve through π has length at least $\ell_\star D_\mu$, while an AC curve in $\mathcal{P}_{\text{reg}}$ from μ to ν of length below $\frac{1}{2} \ell_\star D_\mu$ keeps $D(\gamma_t\|\pi) > \frac{1}{2} D_\mu$ by the floor, hence $\mathcal{I}_{h,\pi}(\gamma_t) > K D_\mu$ along the whole curve by the logarithmic Sobolev inequality, and $L_I(\gamma) \geq \ell_\star \sqrt{K D_\mu} \int_0^1 |\dot{\gamma}_t| dt \geq \ell_\star \sqrt{K D_\mu} W_{2,h}(\mu, \nu)$; so $\bar{d}_I$ separates points and is a metric on $\bar{\mathcal{P}}$. $\square$

Location–scale leaves. Here $\Lambda = \mathbb{R}$ with its Euclidean metric and m is Lebesgue measure ($V = 0$, Equation (S.B.1) with $K = 0$); $\mathcal{H}_m = \text{Ent}(\rho) = \int \rho \log \rho$ and $\mathcal{I}_{h,m} = J(\rho) = \int (\rho'/\rho)^2 \rho$ is the classical Fisher information of a density. Let f be a positive, absolutely continuous probability density with $\int f |\log f| < \infty$, $zf(z) \rightarrow 0$ as $|z| \rightarrow \infty$, centred and standardized, $\int zf = 0$, $\int z^2 f = 1$, $J_0 := \int (f'/f)^2 f < \infty$. Its leaf is $\mathcal{L}_f := \{\rho_{a,s}\}$, $\rho_{a,s}(x) = s^{-1} f((x-a)/s)$, $a \in \mathbb{R}$, $s > 0$, a subset of $\mathcal{P}_{\text{reg}}$.

Proposition L3 (The leaf is a Euclidean half-plane in $W_2 \cdot \text{kernel} \cdot \text{PROVED}$). *For two members of $\mathcal{L}_f$, $W_2(\rho_{a_0,s_0}, \rho_{a_1,s_1})^2 = (a_1 - a_0)^2 + (s_1 - s_0)^2$, the displacement interpolation between them stays in $\mathcal{L}_f$ with a_t, s_t affine in t , and an AC curve $t \mapsto \rho_{a(t),s(t)}$ in $\mathcal{L}_f$ has metric derivative $|\dot{\gamma}_t| = \sqrt{\dot{a}^2 + \dot{s}^2}$.*

Proof. The map $T(x) = a_1 + \frac{s_1}{s_0}(x - a_0)$ is increasing and pushes ρ_{a_0,s_0} forward to ρ_{a_1,s_1} ; in one dimension the increasing push-forward is the quantile coupling, which is optimal for the quadratic cost (Villani, 2009). With $Z \sim f$, $W_2^2 = \mathbb{E}[(a_1 - a_0 + (s_1 - s_0)Z)^2] = (a_1 - a_0)^2 + (s_1 - s_0)^2$ by the standardization. The displacement interpolation $((1-t)\text{id} + tT)_{\#} \rho_{a_0,s_0}$ is ρ_{a_t,s_t} with $a_t = (1-t)a_0 + ta_1$, $s_t = (1-t)s_0 + ts_1$. Hence $(\mathcal{L}_f, W_2)$ is isometric to the Euclidean half-plane and the metric derivative of a curve in the leaf is its Euclidean speed. $\square$

Proof of Theorem G6, hyperbolic leaves. (i) $\partial_x \log \rho_{a,s}(x) = s^{-1}(f'/f)((x-a)/s)$, so $J(\rho_{a,s}) = J_0/s^2$ after the substitution $x = a + sz$. (ii) By Proposition L3 and (i), for an AC curve in the leaf $L_I(\gamma) = \ell_\star \int_0^1 \frac{\sqrt{J_0}}{s(t)} \sqrt{\dot{a}^2 + \dot{s}^2} dt$, the length of the Riemannian metric $G_I|_{\mathcal{L}_f} = \ell_\star^2 J_0 (da^2 + ds^2)/s^2$. Writing it as $e^{2\varphi}(da^2 + ds^2)$ with $\varphi = \frac{1}{2} \log(\ell_\star^2 J_0) - \log s$, its Gaussian curvature is $K = -e^{-2\varphi} \Delta \varphi = -1/(\ell_\star^2 J_0)$. (iii) The metric is the constant multiple $\ell_\star^2 J_0$ of the Poincaré metric $(da^2 + ds^2)/s^2$, whose distance is $\text{arcosh}(1 + |z - w|^2/(2 \text{Im} z \text{Im} w))$ and whose geodesics are the vertical lines and the half-circles orthogonal to the boundary; a constant factor λ on the metric multiplies distances by $\sqrt{\lambda}$ and leaves the geodesics unchanged, so

$$d_{\mathcal{L}_f}((a_0, s_0), (a_1, s_1)) = \ell_\star \sqrt{J_0} \text{arcosh} \left(1 + \frac{(a_1 - a_0)^2 + (s_1 - s_0)^2}{2s_0 s_1} \right).$$

(iv) Since $zf(z) \rightarrow 0$ at infinity, integration by parts gives $\int z f' = -1$; by Cauchy–Schwarz, $1 = (\int z \frac{f'}{f})^2 \leq \int z^2 f \cdot \int (\frac{f'}{f})^2 f = J_0$, with equality iff $f'/f = -cz$, i.e. iff f is Gaussian, and $c = 1$ by the normalization (Stam, 1959). Hence $-1/\ell_\star^2 \leq K < 0$ and the Gaussian leaf is the most curved. Along a vertical geodesic from s_0 to s_1 the length is $\ell_\star \sqrt{J_0} |\log(s_1/s_0)|$ while $\text{Ent}(\rho_{a,s}) = \text{Ent}(f) - \log s$ makes the floor equal to $\ell_\star |\log(s_1/s_0)|$: the floor is attained along the vertical geodesics iff $J_0 = 1$, iff f is Gaussian, and then they are global minimizers of d_I , being the traces of the heat flow of Proposition G1 (iii). On the Gaussian leaf, with the prior $\pi = \mathcal{N}(0, \sigma^2)$ as reference instead, the factor becomes $\mathcal{I}_{h,\pi}(\mathcal{N}(a, s^2)) = a^2/\sigma^4 + (s/\sigma^2 - 1/s)^2$, which vanishes at π : the geometry acquires a centre (Figure 5b). $\square$

Proof of Corollary G7, scale invariance. Let $h' = \alpha h$ with $\alpha > 0$ constant, and either $m' = m$, or $m' = dV_{h'}$ when $m = dV_h$. Then $\nabla_{h'} = \alpha^{-1} \nabla_h$ and $\|\cdot\|_{h'}^2 = \alpha \|\cdot\|_h^2$ give $\|\nabla_{h'} \log f\|_{h'}^2 = \alpha^{-1} \|\nabla_h \log f\|_h^2$, so $\mathcal{I}_{h',m'} = \alpha^{-1} \mathcal{I}_{h,m}$; in the second case $dV_{h'} = \alpha^{n/2} dV_h$, so $f' = \alpha^{-n/2} f$, $\nabla \log f' = \nabla \log f$ and $\mathcal{H}_{m'} = \mathcal{H}_m - \frac{n}{2} \log \alpha$. Distances, hence Wasserstein distances and metric derivatives, scale by $\sqrt{\alpha}$. Then $\sqrt{\mathcal{I}_{h',m'}} |\dot{\gamma}| = \sqrt{\mathcal{I}_{h,m}} |\dot{\gamma}|$: L_I, d_I, G_I and the differences of $\mathcal{H}$ are invariant, and Equation (S.B.1) holds for (h', m') with $K' = K/\alpha$. When (Λ, h) is the Fisher manifold of an experiment, n independent repetitions turn h into nh . $\square$

The diffusive reference. Let $\varepsilon > 0$ and let the reference diffusion be the m -reversible process $dX_t = -\frac{\varepsilon}{2} \nabla_h V(X_t) dt + \sqrt{\varepsilon} dB_t^h$, with generator $\frac{\varepsilon}{2}(\Delta_h - \nabla_h V \cdot \nabla_h)$, which is non-explosive under Equation (S.B.1) (Bakry and Émery, 1985; Hsu, 2002); its law on $C([0, 1]; \Lambda)$ started from $\mu_0 \in \mathcal{P}_{\text{reg}}$ is R . A controlled diffusion is $dX_t = (-\frac{\varepsilon}{2} \nabla_h V + u_t)(X_t) dt + \sqrt{\varepsilon} dB_t^h$ with the same initial law, u a smooth bounded time-dependent vector field; P is its law and $\mu_t = f_t m$ its marginals, with smooth positive densities. Nelson's current velocity is $v_t := u_t - \frac{\varepsilon}{2} \nabla_h \log f_t$.

Proof of Corollary G8, the diffusive unit. (i) Girsanov's theorem on a manifold (Elworthy, 1982; Hsu, 2002; Léonard, 2012) gives $D(P\|R) = \frac{1}{2\varepsilon} \int_0^1 \int \|u_t\|_h^2 d\mu_t dt$. The Fokker-Planck equation of the controlled process, for densities with respect to dV_h , is $\partial_t \mu = \frac{\varepsilon}{2} [\Delta_h \mu + \nabla_h \cdot (\mu \nabla_h V)] - \nabla_h \cdot (\mu u)$, and $\Delta_h \mu + \nabla_h \cdot (\mu \nabla_h V) = \nabla_h \cdot (\mu \nabla_h \log f)$ since $f = \mu e^V$: this is the continuity equation $\partial_t \mu_t + \nabla_h \cdot (\mu_t v_t) = 0$ with the current velocity. Expanding $\|u\|_h^2 = \|v\|_h^2 + \varepsilon \langle v, \nabla_h \log f \rangle_h + \frac{\varepsilon^2}{4} \|\nabla_h \log f\|_h^2$ and integrating against μ_t , with $\frac{d}{dt} \int f \log f dm = \int \langle v, \nabla_h \log f \rangle_h d\mu$ for smooth positive densities,

$$\varepsilon D(P\|R) = \int_0^1 \left[\frac{1}{2} \int_{\Lambda} \|v_t\|_h^2 d\mu_t + \frac{\varepsilon^2}{8} \mathcal{I}_{h,m}(\mu_t) \right] dt + \frac{\varepsilon}{2} [\mathcal{H}_m(\mu_1) - \mathcal{H}_m(\mu_0)]. \quad (\text{S.B.5})$$

(ii) For an AC curve solving the continuity equation with a field v_t , $|\dot{\mu}_t| \leq \|v_t\|_{L^2(\mu_t)}$ for a.e. t , with equality iff v_t is the tangent field, the one of minimal norm (Benamou and Brenier, 2000; Ambrosio et al., 2008); and the elementary $\frac{1}{2}a^2 + b \geq \sqrt{2b}a$ for $a, b \geq 0$, with equality iff $a = \sqrt{2b}$, applied to $a = |\dot{\mu}_t|$ and $b = \frac{\varepsilon^2}{8} \mathcal{I}_{h,m}(\mu_t)$, gives with $\ell_\star = \varepsilon/2$

$$\varepsilon D(P\|R) - \frac{\varepsilon}{2} [\mathcal{H}_m(\mu_1) - \mathcal{H}_m(\mu_0)] \geq \int_0^1 \left[\frac{1}{2} |\dot{\mu}_t|^2 + \frac{\varepsilon^2}{8} \mathcal{I}_{h,m}(\mu_t) \right] dt \geq L_I(\mu), \quad (\text{S.B.6})$$

with equality throughout iff v_t is the tangent field and $\frac{1}{2} |\dot{\mu}_t|^2 = \frac{\varepsilon^2}{8} \mathcal{I}_{h,m}(\mu_t)$ for a.e. t , zero Jacobi energy. (iii) For $u = 0$, $P = R$ and $v = -\frac{\varepsilon}{2} \nabla_h \log f$ is the tangent field of the reference flow, whose speed is $\frac{\varepsilon}{2} \sqrt{\mathcal{I}_{h,m}(\mu_t)}$ by Proposition G1 (iii) after the time change $t \mapsto \varepsilon t/2$: both equalities hold. When m is a probability measure and $\mu_0 = m$, Equation (S.B.6) and the floor give $\varepsilon D(P\|R) \geq \varepsilon D(\mu_1\|m)$, the data-processing inequality for the terminal marginal. The functional $\int [\frac{1}{2} |\dot{\mu}|^2 + c \mathcal{I}_{h,m}]$ with $c = \varepsilon^2/8$ is the action of the Schrödinger problem (Léonard, 2014); its Maupertuis length at energy e is $\int \sqrt{2(e + c \mathcal{I}_{h,m})} |\dot{\mu}|$ (Arnold, 1989), and the slope geometry is its member at zero energy, $\ell_\star^2 = 2c$: a diffusive reference prices every revision above its calibrated length and fixes $\ell_\star = \varepsilon/2$. $\square$

Proof of Corollary G9, Shannon as posterior geometry. Let $\Lambda \sim \pi$ with π satisfying the logarithmic Sobolev inequality, Y an observation with a likelihood and π_Y its posterior, regular for a.e. y . (i) *Lindley.* $I(\Lambda; Y) = \iint \log \frac{dP_{\Lambda Y}}{d(\pi \otimes P_Y)} dP_{\Lambda Y} = \int D(\pi_y \| \pi) P_Y(dy)$ by disintegration, and Theorem G5 for each π_y gives $\mathbb{E}_Y \rho_I(\pi_Y) = \ell_\star I(\Lambda; Y)$. (ii) *Post-processing.* For a channel $Y \rightarrow Z$, $\pi_Z = \mathbb{E}[\pi_Y | Z]$ and the convexity of $D(\cdot \| \pi)$ give $D(\pi_Z \| \pi) \leq \mathbb{E}[D(\pi_Y \| \pi) | Z]$; taking expectations, $\mathbb{E}_Z \rho_I(\pi_Z) \leq \mathbb{E}_Y \rho_I(\pi_Y)$. (iii) *Chain rule.* $I(\Lambda; Y_1, Y_2) = I(\Lambda; Y_1) + I(\Lambda; Y_2 | Y_1)$ with $I(\Lambda; Y_2 | Y_1) = \mathbb{E}_{Y_1} \mathbb{E}_{Y_2|Y_1} D(\pi_{Y_1 Y_2} \| \pi_{Y_1})$: each conditioning re-centres the geometry at the intermediate posterior. (iv) *Rate-distortion.* Let $X \sim \pi$ be a source on Λ , a test channel a Markov kernel producing a reconstruction $\hat{X}$ of law ν , μ_a the law of X given $\hat{X} = a$ (the cells), and

$c(x, \hat{x}) = d_h(x, \hat{x})^2$. The tower property gives $\pi = \int \mu_a \nu(da)$; the disintegration of the mutual information gives $I(X; \hat{X}) = \int D(\mu_a \| \pi) \nu(da)$; and $W_{2,h}^2(\mu_a, \delta_a) = \int d_h(x, a)^2 \mu_a(dx)$, the only coupling with a Dirac mass being the product, gives $\mathbb{E} c(X, \hat{X}) = \int W_{2,h}^2(\mu_a, \delta_a) \nu(da)$. With regular cells, Theorem G5 applied to each cell turns Shannon's $R(D) = \inf\{I(X; \hat{X}) : \mathbb{E} c \leq D\}$ (Shannon, 1959) into

$$\ell_\star R(D) = \inf \left\{ \int \rho_I(\mu_a) \nu(da) : \int \mu_a \nu(da) = \pi, \int W_{2,h}^2(\mu_a, \delta_a) \nu(da) \leq D \right\} :$$

rate is the mean calibrated radius of the cells from the source, distortion the mean transport radius of the cells to their code points. (v) *Zero distortion*. If $\mathbb{E} c = 0$ then $\mu_a = \delta_a$ for ν -a.e. a , so $D(\mu_a \| \pi) = +\infty$ and $I(X; \hat{X}) = +\infty$ for every such channel: $R(0) = +\infty$, and the cells sit on the horizon of Theorem G3. *The Gaussian source*. For $X \sim \mathcal{N}(0, \sigma^2)$ and $0 < D \leq \sigma^2$, the channel with $\hat{X} \sim \mathcal{N}(0, \sigma^2 - D)$ and $X | \hat{X} = a \sim \mathcal{N}(a, D)$ has barycentre $\mathcal{N}(0, \sigma^2)$, cells with $W_2^2(\mathcal{N}(a, D), \delta_a) = D$, and $\mathbb{E}_a \rho_I(\mathcal{N}(a, D)) = \ell_\star \mathbb{E}_a \frac{1}{2} [(a^2 + D)/\sigma^2 - 1 + \log(\sigma^2/D)] = \ell_\star \frac{1}{2} \log(\sigma^2/D) = \ell_\star R(D)$, the Gaussian rate-distortion function (gate S1.1); Shannon's theorem provides the optimality of this channel, the geometry provides the reading. $\square$

S.B.3 Recovery, stability, closure

Derivation of Realization E1, the balance of a zone and its two ledgers. With $\mu = f m$ and $j = f v$, $\partial_t(f \log f) = (1 + \log f) \partial_t f = -(1 + \log f) \operatorname{div}_m(f v)$; the product rule $\operatorname{div}_m((1 + \log f) f v) = (1 + \log f) \operatorname{div}_m(f v) + f \langle v, \nabla \log f \rangle$ and the divergence theorem for div_m give Equation (7.2) for every declared m . For a diffusion of drift b and diffusion matrix $\mathbf{D}$ with density p relative to the volume, the Fokker-Planck velocity is $v = b - \mathbf{D} \nabla \log p$, so $\langle v, \mathbf{D}^{-1} v \rangle = \langle v, \mathbf{D}^{-1} b \rangle - \langle v, \nabla \log p \rangle$ and, with $\nabla \log p = \nabla \log f + \nabla \log m$, integration over the zone gives Equation (7.3); the reference term vanishes for the volume reference. On the gyrator $\langle v, \nabla \log p \rangle = -x^\top \Gamma^\top C^{-1} x = 0$ pointwise because $\Gamma^\top C^{-1}$ is antisymmetric (gates X4.1b, X4.5); on the driven ring with the Gibbs reference the bulk term changes and the identity holds with its reference term (gate RING.4). $\square$

Proof of Theorem R5, exhaustive resolution recovery. Relative information. Restriction to $\mathcal{F}_N \subset \mathcal{F}_{N+1}$ is a channel, so $D_N \leq D_{N+1} \leq D(P \| R)$ by Equation (3.2). If $P \ll R$, put $Z = dP/dR$ and $Z_N = \mathbb{E}_R[Z | \mathcal{F}_N] = dP|_{\mathcal{F}_N}/dR|_{\mathcal{F}_N}$; Lévy's upward theorem gives $Z_N \rightarrow Z$ R -almost surely and in $L^1(R)$ (Williams, 1991), and Fatou's lemma for $\varphi(Z_N) \geq 0$, $\varphi(z) = z \log z - z + 1$, gives $\liminf_N D_N \geq D(P \| R)$; hence $D_N \uparrow D(P \| R)$, and the hidden term $D(P \| R) - D_N$ decreases to zero. If P charges an R -null set A , choose $A_N \in \mathcal{F}_N$ with $(P + R)(A \triangle A_N) \rightarrow 0$; the binary data-processing bound $D_N \geq P(A_N) \log \frac{P(A_N)}{R(A_N)} + (1 - P(A_N)) \log \frac{1 - P(A_N)}{1 - R(A_N)}$ tends to $+\infty = D(P \| R)$. This is relative entropy as a supremum over finite partitions (Pinsker, 1964; Barron, 1985). *Fisher*. For a family differentiable in quadratic mean (van der Vaart, 1998, §7), $S_{v,N} = \mathbb{E}[S_v | \mathcal{F}_N]$ is a martingale bounded in L^2 , so $S_{v,N} \rightarrow \mathbb{E}[S_v | \mathcal{F}_\infty] = S_v$ in L^2 ; orthogonality of the increments gives $h_M(v, v) - h_N(v, v) = \mathbb{E}(S_{v,M} - S_{v,N})^2 \geq 0$ and $h_N(v, v) = \|S_{v,N}\|_2^2 \uparrow \|S_v\|_2^2 = h(v, v)$; polarize. *Bayesian information*. The σ -algebras $\mathcal{A} \otimes \mathcal{B}_N$ increase to $\mathcal{A} \otimes \mathcal{B}$ up to null sets; apply the first part to $P = P_{\Lambda Y}$ and $R = \pi \otimes P_Y$. *Radius*. Theorem G3 (iii) and B1 give $\mathbb{E} \rho_{I,N} = \ell_\star I(\Lambda; Y_N)$, and the previous step concludes. For a garbling tower which is a filtration in law only, $K_N P \rightarrow P$, the monotonicity is Blackwell's and the limit follows from the lower semicontinuity of relative entropy under weak convergence (Ambrosio et al., 2008, Lemma 9.4.3); on the Gaussian sensors of the gyrator the limit is the closed form (gate X2.4). $\square$

Proof of Theorem G10, stability under almost-sufficient projection. (a) For a piecewise C^1 curve c on the stratum, $\sqrt{h_K(\dot{c}, \dot{c})} \in [\delta, 1] \sqrt{h(\dot{c}, \dot{c})}$ pointwise; integrating and taking the infimum over curves joining two points gives $\delta d_h \leq d_{h_K} \leq d_h$, and completeness and finite second moments transfer. The couplings of two measures form one set, so the Wasserstein distances are sandwiched in the same way, and so are the difference quotients that define the metric derivative: the absolutely

continuous curves coincide and $\delta|\dot{\gamma}|_h \leq |\dot{\gamma}|_{h_K} \leq |\dot{\gamma}|_h$ almost everywhere. (b) $\mathcal{H}_m$ depends on m and μ alone; with $f = d\mu/dm$ one function, $\|\nabla_h \log f\|_h^2 = h^{-1}(d \log f, d \log f)$, and inversion reverses the Loewner order, $h^{-1} \preceq h_K^{-1} \preceq \delta^{-2}h^{-1}$; integrating against μ gives $\mathcal{I}_{h,m} \leq \mathcal{I}_{h_K,m} \leq \delta^{-2}\mathcal{I}_{h,m}$, and the regular domains coincide. (c) Multiply the two sandwiches inside $\ell_\star \int \sqrt{\mathcal{I}} |\dot{\gamma}| dt$ and take the infimum over the common class of curves. (d) Under HR0 for (Λ, h, m) , (c) and the floor of Theorem G3 give $L_I^{h_K}(\gamma) \geq \delta \ell_\star |\mathcal{H}_m(\gamma_1) - \mathcal{H}_m(\gamma_0)|$. (e) Under HR0 for (Λ, h_K, m) the proofs above apply verbatim to (Λ, h_K, m) , and the radial identity has a right-hand side free of the metric. The identification $\delta^2 = 1 - \varphi$ is Theorem 1.4 of Yamaguchi and Nozawa (2024): a statistic is δ -almost sufficient exactly when $h^{\text{hid}}(v, v) \leq (1 - \delta^2)h(v, v)$ for every direction, and the largest such ratio is the fraction of missing information of Meng and Rubin (1991). A calibration that changes with the metric, such as the Riemannian volume, is outside the statement. $\square$

Proof of Theorem E6, the predictive closure ledger. Write $X' = X_{t+\Delta}$, $x = x_t$, $X = q(x)$; everything is finite, bounded by $\log |\mathcal{A}|$, and regular conditional laws exist. Since X is $\sigma(x)$ -measurable, $I(X'; x, X) = I(X'; x)$, and the chain rule $I(X'; x, X) = I(X'; X) + I(X'; x | X)$ gives (i) (Cover and Thomas, 2006, Thm 2.5.2); the conditional mutual information equals $\mathbb{E}_x D(P_{X'|x} \| P_{X'|X})$ because $P_{X'|x,X} = P_{X'|x}$ and $P_{X'|X} = \mathbb{E}[P_{X'|x} | X]$, and it vanishes exactly when $P_{X'|x} = P_{X'|X}$ almost surely, which is (ii). Nonnegativity of I_{ret} and I_{hid} and $I(X'; x) \leq H(X') = H(X)$, by stationarity, give (iii). Pinsker's inequality for each x and Jensen's inequality for the concave square root give (iv). For (v), $p_\Delta = \mathbb{P}(X' \neq X) \rightarrow 0$ by path continuity, Fano's inequality gives $H(X' | X) \leq h_2(p_\Delta) + p_\Delta \log(|\mathcal{A}| - 1) \rightarrow 0$, hence $I_{\text{ret}} \rightarrow H(X)$, and $I_{\text{ret}} \leq I_{\text{pred}} \leq H(X)$ squeezes I_{pred} and I_{hid} . On the empirical measure of a trajectory binned on a grid whose cells refine the symbols, (i) holds exactly, so the numerical content of the gates is in the ratios, the floors by circular shifts and the tightness of (iv) (gates X8.5–X8.8; estimator bias after Paninski (2003)). $\square$

S.C Statements outside the body

Each statement below cites its gates; the theorems of the body are complete without them.

Theorem K5 (Path irreversibility · kernel · PROVED). *When a reversal $P^\dagger$ equivalent to P exists, $\Sigma = \log dP/dP^\dagger$ satisfies $\mathbb{E} e^{-\Sigma} = 1$ and $\mathbb{E} \Sigma = D(P \| P^\dagger) \geq 0$; under every quotient K of the paths that commutes with time reversal, $K(P^\dagger) = (KP)^\dagger$, the visible irreversibility obeys $\langle \Sigma_{\text{visible}} \rangle = D(KP \| (KP)^\dagger) \leq \langle \Sigma_{\text{fine}} \rangle$ (Kawai et al., 2007; Gomez-Marín et al., 2008). On the gyrtator every scalar linear observable is a stationary Gaussian process, hence reversible (Weiss, 1975), the window covariance being symmetric Toeplitz: an observer of one coordinate reads zero, and the two coordinates read $\sigma = 0.2793$, measured as 0.2839 ± 0.0037 on $2 \cdot 10^4$ paths (gates X5.1, X5.3, X5.4, LIB.6, SYM.5). Hidden irreversibility under coarse graining is a subject of its own (Degünther et al., 2024).*

Proof. $\mathbb{E}_P e^{-\Sigma} = \int \frac{dP^\dagger}{dP} dP = 1$ and $\mathbb{E}_P \Sigma = D(P \| P^\dagger) \geq 0$. When K commutes with time reversal, $D(KP \| (KP)^\dagger) = D(KP \| KP^\dagger) \leq D(P \| P^\dagger)$ by Equation (3.2) applied to the pair $(P, P^\dagger)$. $\square$

Proposition K6 (The tilt · kernel · IMPORTED). *Every law equivalent to the reference reads $dP = e^V dR$ (Williams, 1991); the potential V is the logarithm of the Radon–Nikodym derivative, and $-\log(dm/d\nu)$ is the reference potential associated with m ; it becomes $\beta E + \text{const}$ in a thermal canonical realization alone (H2).*

Proposition C2 (Least deformation selects Bayes, the controls · kernel · PROVED). *The Gibbs identity of Theorem K3 holds to 10^{-16} on random beliefs, priors and likelihoods on four states (gate C2.3); the stationarity residual $f'(u) - \log u$ is constant for relative entropy and varies for the squared Hellinger distance and for χ^2 (gate C2.4b); the least-deformation update with those two divergences differs from Bayes by 0.03 to 0.13 in total variation, while an optimizer with relative entropy finds Bayes to 10^{-9} (gates C2.2, C2.1).*

Proposition O2 (The score projection $\cdot$ kernel $\cdot$ PROVED). *For an observation $Y = K(X)$ the observable score is the conditional expectation of the fine score, $S_v^Y = \mathbb{E}[S_v | Y]$, and $h_Y(v, w) = \mathbb{E}[S_v^Y S_w^Y]$ (proof in Appendix S.B.1). On the gyrator the variance of the conditional score equals the Fisher metric of the marginal in the five directions to 10^{-14} (gate X1.2).*

Theorem G4 (The constant slope is unique $\cdot$ kernel $\cdot$ PROVED). *The weight $\ell_\star \sqrt{\mathcal{I}_{h,m}}$ gives relative entropy the constant slope $1/\ell_\star$ at every regular belief, and a relief continuous at a regular belief gives it the slope $1/\ell_\star$ there exactly when it equals $\ell_\star \sqrt{\mathcal{I}_{h,m}}$ there (Theorem G3, proof in Appendix S.B.2). On the relaxation path of the gyrator, the bare transport length is 1.857 and the length with weight $\mathcal{I}_{h,m}$ is 9.15, against a drop of relative information of 3.554 (gate X3.4).*

Theorem G5 (The radial identity $\cdot$ kernel $\cdot$ PROVED). *With the prior as reference and an ergodic gradient flow, $\rho_1(\mu) = \ell_\star D(\mu \| \pi)$: relative entropy is the radial coordinate and the gradient flow is the radial geodesic (proof in Appendix S.B.2). Physical realization on the gyrator: at equal temperatures $T_1 = T_2 = 1$, from the displaced stationary law $\mathcal{N}(m_0, C)$ with $m_0 = (1.5, -1)$ and C the stationary covariance at $T_1 = 2$, $T_2 = 0.5$, the relaxation of the ensemble law is a Fokker–Planck flow with isotropic diffusion and a symmetric drift, the gradient flow of $D(\cdot \| \pi)$, hence radial, with $\ell_\star = 1$; the calibrated length of the path is 3.55407, the drop of relative information 3.55402, and the entropy production measured on $2 \cdot 10^4$ paths 3.535 ± 0.023 (gates X3.1, X3.2, LIB.8). Figure 5(b) draws the radial geodesics and the calibrated circles on the Gaussian leaf.*

Theorem G6 (Hyperbolic-Leaf Theorem $\cdot$ kernel $\cdot$ PROVED). *With Lebesgue reference, every location–scale family carries the metric $\ell_\star^2 J_0 (da^2 + ds^2)/s^2$, a Poincaré half-plane of curvature $-1/(\ell_\star^2 J_0)$ set by the Fisher information of the standardized profile, the Gaussian being extremal by Stam’s inequality (proof in Appendix S.B.2, from four classical ingredients: the leaf is flat and totally geodesic in W_2 , $J(\rho_{a,s}) = J_0/s^2$, the conformal factor of Equation (4.3), and Stam, 1959). The leaves first appear, with the volume as reference, in Caraffa (2026, Thms 4.18–4.19). This hyperbolicity is distinct from the Fisher–Rao hyperbolicity of Gaussian families: the underlying leaf is flat in W_2 (Fukushi et al., 2026), and the negative curvature is created by the factor J_0/s^2 . On the Gaussian leaf the geodesic and the chord of Figure 5 cost 4.5 and 9.6 nats, the curvature is $-1/(\ell_\star^2 J_0)$ symbolically and $J_0 = 1, 2$ for the Gaussian and Laplace profiles (gate S3.1).*

Theorem R2 (The quadruple ledger and its increments $\cdot$ kernel $\cdot$ PROVED). *Along a tower, $D = D_N + D_N^{\text{hid}}$, $h = h_N + h_N^{\text{hid}}$, $I = I_N + I_N^{\text{hid}}$ and $\mathbb{E}\rho_I = \mathbb{E}\rho_{I,N} + \ell_\star I_N^{\text{hid}}$: the global, the infinitesimal, the Bayesian and the geometric ledgers agree, each under its own scope (Table 3); from one level to the next the increments are the conditional ledgers, $D_{n+1} - D_n = \mathbb{E} D(P_{Y_{n+1}|Y_n} \| R_{Y_{n+1}|Y_n})$, $h_{n+1} - h_n = \mathbb{E}[(S_{n+1} - S_n)^{\otimes 2}]$, $I_{n+1} - I_n = I(\Lambda; Y_{n+1} | Y_n)$, $\mathbb{E}\rho_{I,n+1} - \mathbb{E}\rho_{I,n} = \ell_\star I(\Lambda; Y_{n+1} | Y_n)$. On nested Gaussian sensors of the gyrator’s state the four quantities increase strictly, $\mathbb{E}\rho_I = \ell_\star I_N$ to 10^{-16} , and the increments equal the conditional quantities to $3 \cdot 10^{-12}$ in absolute value (gates X2.1, X2.8b).*

Proof. The four splits are Equation (3.2), Equation (4.2), the chain rule of mutual information and, with the prior as reference, $\mathbb{E}\rho_{I,N} = \ell_\star I_N$ by B1 and Theorem G3 (iii). The increments are the same identities applied to the pair (Y_n, Y_{n+1}) of a nested tower: the chain rule of relative entropy for the joint law of (Y_n, Y_{n+1}) , the orthogonality of the martingale increments of the score, and $I(\Lambda; Y_{n+1}) = I(\Lambda; Y_n) + I(\Lambda; Y_{n+1} | Y_n)$. $\square$

Realization R3 (Harmonic Markov resolution $\cdot$ realization $\cdot$ MEASURED). *Under HR0 with m a probability, let $L_m = \Delta_h + \nabla_h \log w \cdot \nabla_h$ be the reversible generator and $K_t = e^{tL_m}$. Then $(K_t)_{t \geq 0}$ is a Markov semigroup, $K_{t+s} = K_t K_s$, so $t_2 \geq t_1$ makes K_{t_2} a garbling of K_{t_1} ; with a compact resolvent and $-L_m \phi_k = \lambda_k \phi_k$, $K_t \phi_k = e^{-\lambda_k t} \phi_k$; along $\mu_t = K_t \mu$, $-\frac{d}{dt} D(\mu_t \| m) = \mathcal{I}_{h,m}(\mu_t)$ (G1): the harmonic spectrum supplies a canonical channel-valued resolution scale, and relative Fisher information is its instantaneous relative-information loss rate. For $d\mu/dm - 1 \in L^2(m)$ the spectral cutoff is the analysis, $\chi^2(\mu \| m) = \sum_k a_k^2$ (Parseval) with a tail after diffusion bounded by*

$e^{-2\lambda_{N+1}t}\|g\|^2$; it is free of any positivity claim, the reconstruction being Realization R4. On the circle the tail bound holds with ratios below 0.06, and the semigroup is positive, mass-preserving, a semigroup to 10^{-16} , diagonal with damping e^{-k^2t} , with $-dD/dt = \mathcal{I}$ to 10^{-7} (gates S2.1, S2.3).

Realization R4 (The exponential reconstruction $\cdot$ realization $\cdot$ MEASURED). When the partition function $Z(\theta)$ of the first N modes is finite, the information projection onto their exponential family reconstructs a probability and obeys $D(\mu \| m) = D(\mu \| \tilde{\mu}_N) + D(\tilde{\mu}_N \| m)$ (gate S2.2). The spectral projector analyses; the exponential projection reconstructs.

Realization E2 (The interface and the facet ledger $\cdot$ realization $\cdot$ MEASURED). Quantised on a square grid of step ε , the continuum current obeys Kirchhoff on the facets to 10^{-15} ; the finite-volume chain with four neighbours and Scharfetter–Gummel rates reproduces σ with errors 2.1, 0.6, 0.2, 0.1 % at $\varepsilon/\sigma_{\min} = 0.4, 0.2, 0.1, 0.05$; under the tested eight-neighbour mobility convention, the diagonal links create a persistent excess production, above a factor three at every ε (Figure 7; gates X6.1, X6.2, X6.3b). The ledger is a property of the facet conductances of the stencil; entropic gradient-flow structures of discretizations converge to the continuum while their transport distances may converge slowly (Maas, 2011; Forkert et al., 2022).

Realization E3 (Three graphs, and the ruler that bends $\cdot$ realization $\cdot$ MEASURED). The proximity graph (geometric simplices), the generator graph (interaction hyperedges) and the conditional-independence graph (Markov cliques) are three objects. On the gyrator the first is the same lattice at $k = 0$ and $k = 0.6$, while the precision entry $(C^{-1})_{12}$ and the drift entry A_{12} vanish exactly at $k = 0$ (gate X6.5). The law ruler of the stratum $(\log k_1, \log k_2)$ at fixed k is $\frac{1}{2}I$ at $k = 0$, symbolically, and $(C^{-1})_{12} = -k$ times a positive rational function (gate SYM.7); its Gaussian curvature at $(k_1, k_2) = (1, 0.7)$ decreases from 0 at $k = 0$ to -0.274 at $k = 0.6$ and -0.320 at $k = 0.8$, while the partial correlation and the mutual information of the two coordinates increase from zero (gate X6.6; Figure 8).

Realization E4 (The two axes of information $\cdot$ realization $\cdot$ MEASURED). Relative to a product reference, Equation (6.2) holds exactly, the second term being the total correlation. A sensor on one coordinate moves the first part only, a joint sensor the second, and conditioning on one coordinate of a coupled prior drives the coupling part to zero (gates X7.1, X7.2, X7.3).

Realization E7 (The lifetime $\cdot$ realization $\cdot$ MEASURED). The escape rate of a well follows Arrhenius, $e^{-E_b/T}$ (Kramers, 1940): with buffered basin passages the measured rates at $T = 0.25, 0.2, 0.16$ give a log-slope of -0.968 against $1/T$ for a barrier $E_b = 1$ (gate X8.4); a crude zero-crossing count, inflated by recrossings, gives -0.854 (gate PUTS.2). The lifetime of a symbol grows as its exchange capacity shrinks.

S.D Figures and the table of gates

The table of gates follows, one line per gate with its measurement.

gate	measurement (green)
C1.1	chain-rule residual on 50 random joints : KL 2.2e-16 ; Hellinger2 0.048, TV 0.365, chi2 10.774
C1.2	telescoping over $X \rightarrow Y \rightarrow Z$: KL 2.3e-16 ; Hellinger2 0.062, TV 0.379, chi2 11.341
C2.1	least-deformation update with KL against Bayes, three evidences : total-variation gaps ['3.9e-09', '1.5e-09', '5.8e-09']
C2.2	the same variation with Hellinger2 or chi2 : total-variation gaps to Bayes ['0.029', '0.118', '0.134']
C2.3	Gibbs identity $E_{\mu}[-\log L] + D(\mu \pi) = D(\mu \pi_D) - \log Z$ on 20 random (μ, π, L) : max residual 2.2e-16
C2.4b	redeclared : $f'(u) - \log u$ on fifty points of $[1e-2, 1e2]$, variance : KL 1.6e-32 ; Hellinger2 1.85, chi2 1.8e+03 : the stationarity equation $f'(u) = \log u + c$ selects $f = t \log t + a(t - 1)$

gate	measurement (green)
LIB.1	Lyapunov residual $A C + C A = 2D : 4.4e-16$
LIB.2	Gamma C antisymmetric : residual $7.5e-16$, $s = 0.310345$
LIB.3	first law and Clausius : $Q1 + Q2 = -7.8e-16$, $\sigma = 0.279310 = Q1/T1 + Q2/T2 = 0.279310$
LIB.4	$\sigma > 0$ at $T1 \neq T2$ (0.279310) ; at $T1 = T2$: $\sigma = 3.4e-31$, current = $6.7e-16$
LIB.5	σ against $(T1 - T2)$: log-slope 2.0022 (expected 2)
LIB.6	scalar window KL for $n = 2, 8, 32$: $[0.0e+00', 0.0e+00', 0.0e+00']$ (reversible)
LIB.7	2D chain entropy production per unit time, dyadic steps $dt = 2^{-n}$, $n = 0..10$: $[0.04691', 0.12265', 0.18945', 0.23150', 0.25470', 0.26683', 0.27303', 0.27616', 0.27773', 0.27852', 0.27892'] \rightarrow \sigma = 0.27931$ (monotone, 0.14%)
LIB.8	relaxation at $T = 1$: $D(\mu_t \pi)$ decreasing from 3.5540 to $3.5e-07$
PUITS.1	Gibbs density normalised and symmetric
PUITS.2	Arrhenius : measured log-rate against $1/T$ has slope -0.854 (barrier $h = 1$; crude zero-crossing count, recrossings inflate it)
S1.1	mean calibrated radius of the cells against $R(D) = (1/2) \log(\sigma^2/D)$, $D = 0.1, 0.3, 0.6$: gaps $[2.2e-16', 1.1e-16', 0.0e+00']$
S2.1	spectral tail / bound $e^{-(2(N+1)^2 t)} g ^2$ for $N = 2, 4, 8$ and $t = 0, 0.1$: $[0.0593', 0.0522', 0.0007', 0.0006', 0.0000', 0.0000']$
S2.2	exponential reconstruction $N = 4$: $D(\mu m) = 0.198954 = D(\mu \mu_N) + D(\mu_N m) = 0.000075 + 0.198879$, gap $0.0e+00$; moments matched to $1.2e-15$
S2.3	heat semigroup K_t on the circle : $\min K_t p = 3.601e-02 > 0$, mass 1 to $2.2e-16$, $K_{0.1} K_{0.05} = K_{0.15}$ to $5.6e-17$, modes damped by $e^{-(k^2 t)}$ to $2.5e-16$, $-dD/dt = \text{Fisher}$ to $1.3e-07$ at $t = 0.02, 0.05, 0.1, 0.5$
S3.1	Gaussian leaf : RESG length along the hyperbolic geodesic between $(-2.4, 0.5)$ and $(2.4, 0.5) = 4.544883 = \text{arcosh}(1 + q) = 4.544883$ (relative gap $1.6e-10$), chord 9.60 ; Gaussian curvature of $l^2 J_0$ ($da^2 + ds^2$)/ $s^2 = -1/(J_0 \ell_{\text{star}}^2)$; Stam : $J_0 = 1.00000$ (Gaussian), 1.99973 (Laplace), 1.8077 (symmetric two-Gaussian mixture) ≥ 1
SYM.1	Lyapunov solved : $C11 = (T1^*k^2 + T1^*k^*k1 + 3^*T1^*k^*k2 + T1^*k1^*k2 + T1^*k2^2 + T2^*k^2)/((2^*k + k1 + k2)^*(k^*k1 + k^*k2 + k1^*k2))$, $C12 = k^*(T1^*k + T1^*k2 + T2^*k + T2^*k1)/((2^*k + k1 + k2)^*(k^*k1 + k^*k2 + k1^*k2))$, $C22 = (T1^*k^2 + T2^*k^2 + 3^*T2^*k^*k1 + T2^*k^*k2 + T2^*k1^2 + T2^*k1^*k2)/((2^*k + k1 + k2)^*(k^*k1 + k^*k2 + k1^*k2))$
SYM.2	Gamma C antisymmetric ; $s = (\text{Gamma C})_{21} = k^*(T1 - T2)/(2^*k + k1 + k2)$
SYM.3	$\sigma = k^2(T1 - T2)^2/(T1^*T2^*(2^*k + k1 + k2))$; vanishes at $T1 = T2$ and at $k = 0$; symmetric under $(k1, T1) \leftrightarrow (k2, T2)$
SYM.4	$Q1 = -k^2(T1 - T2)/(2^*k + k1 + k2)$; $Q1 + Q2 = 0$; $\sigma = Q1/T1 + Q2/T2$ exactly
SYM.5	the window covariance of a scalar stationary process is symmetric Toeplitz, $J S J = S$: the reversed window has the same law
SYM.6	$s = k(T1 - T2) * (1/(2^*k + k1 + k2))$, $\sigma = k^2(T1 - T2)^2 * (1/(2^*T1^*T2^*k + T1^*T2^*k1 + T1^*T2^*k2))$, $Q1 = k(T1 - T2) * (-k/(2^*k + k1 + k2))$ with sign-definite cofactors : current, production and heat vanish exactly when $k(T1 - T2) = 0$; uniform driven ring : $p = 1/L$, $j = F/L$, $\sigma = F^2/T$
SYM.7	Fisher metric of $(\log k1, \log k2)$ at $k = 0$: $[[1/2, 0], [0, 1/2]] = I/2$, flat, curvature zero ; $A_{12} = -k$; $(C^{-1})_{12} = -C_{12}/\det C$ with $C_{12}/k = (T1^*k + T1^*k2 + T2^*k + T2^*k1)/(2^*k^2*k1 + 2^*k^2*k2 + k^*k1^2 + 4^*k^*k1^*k2 + k^*k2^2 + k1^2*k2 + k1^*k2^2)$ and $\det C = (T1^2*k^2 + 2^*T1^*T2^*k^2 + 4^*T1^*T2^*k^*k1 + 4^*T1^*T2^*k^*k2 + T1^*T2^*k1^2 + 2^*T1^*T2^*k1^*k2 + T1^*T2^*k2^2 + T2^2*k^2)/((2^*k + k1 + k2)^2(k^*k1 + k^*k2 + k1^*k2))$, both positive : the generator edge, the conditional-independence edge and the curvature of the law ruler vanish together at $k = 0$ and appear together for $k > 0$
X1.1	Fisher from $x1$ alone : singular values relative $[1.0, 0.28329832, 0.03046298, 0.00114225, 0.0]$ (rank 4) ; from $(x1, x2)$: $[1.0, 0.138216, 0.061115, 0.048273, 0.021142]$ (rank 5)
X1.2	$\text{Var}(E[S_v Y])$ against the marginal Fisher, five directions : relative gaps $[2.0e-16', 8.8e-16', 6.8e-16', 2.9e-15', 2.5e-14']$

gate	measurement (green)
X1.3b	redeclared : $h - h_Y \geq 0$ and the fine Fisher along the null direction of h_Y is entirely hidden (share 1.00000000)
X1.4b	$k = 0$: from x_1 alone the Fisher has rank 2 (relative singular values [1.0, 0.34095508, 0.0, 0.0, 0.0]) ; its null space holds (1,0,-1,0,0), e_{k2} , e_{T2} (residuals ['1.1e-10', '0.0e+00', '0.0e+00']) ; from both coordinates rank 5, cross block $(k_1, T_1) \times (k_2, T_2) = 0.0e+00$
X1.5	sub-window of 32 steps against 64 (both coordinates) : $h_{32} \leq h_{64}$ (min eigenvalue of the gap 1.3e+00), $\Delta^2 = \lambda_{\min}(h_{64}^{-1} h_{32}) = 0.4921$; control x_1 alone : $\Delta^2 = -1.3e-17$ (blind direction, outside the statement)
X2.1	$D_N = [0.1123, 0.2405, 0.3367, 0.3741, 0.3848, 0.3876, 0.3883, 0.3884, 0.3885]$; $I_N = [0.3258, 0.5012, 0.594, 0.6252, 0.6337, 0.6358, 0.6364, 0.6365, 0.6366]$; $E \rho_I = I^* I_N$ to 2.2e-16 ; h_N strictly increasing in the Loewner order
X2.2	$h_M - h_N = \text{Cov}(S_M - S_N)$ for $(N, M) = (0, 4), (2, 8)$: gaps ['2.2e-16', '4.2e-17']
X2.3	sensor P (x_1 only, $s = 0.3$) against Q (both, $s = 0.6$) : eigenvalues of $h_P - h_Q = [-0.8725, 0.0474]$: neither dominates
X2.4	exhaustive limit $Q_N \rightarrow Q_{\inf} = \text{ledgers}(0)$ for $(D, \text{tr } h, I, E \rho)$: relative gaps at $N = 5$ ['2.4e-03', '1.8e-03', '1.1e-03', '1.1e-03'], $N = 8$ ['3.8e-05', '2.8e-05', '1.8e-05', '1.8e-05'] ; $(Q_{\inf} - Q_N)/s_N^2$ at $N = 5..8$ over its value at $N = 8$: ['1.00', '1.00', '1.00', '1.00'] (D), linear in s^2
X2.5	$\Delta_N^2 = \lambda_{\min}(h_{\inf}^{-1} h_N) = \min_i \lambda_i(C)/(\lambda_i(C) + s_N^2)$ to 2.2e-16 : [0.2886, 0.6187, 0.8665, 0.9629, 0.9905, 0.9976, 0.9994, 0.9998, 1.0] ; translation of a Gaussian belief on the leaf : $L_N/L_{\inf} = [0.9663, 0.9683, 0.9871, 0.9963, 0.999, 0.9998, 0.9999, 1.0, 1.0]$ within $[\Delta_N, 1/\Delta_N]$, gap to 1 decreasing, 3.8e-06 at $N = 8$
X2.6	$ L_N/L_{\inf} - 1 = ['3.4e-02', '3.2e-02', '1.3e-02', '3.7e-03', '9.6e-04', '2.4e-04', '6.1e-05', '1.5e-05', '3.8e-06'] \leq 1/\Delta_N - 1 = ['8.6e-01', '2.7e-01', '7.4e-02', '1.9e-02', '4.8e-03', '1.2e-03', '3.0e-04', '7.5e-05', '1.9e-05']$; the bound tends to zero
X2.7	composition : $\Delta^2(\text{fine} \rightarrow \text{coarse}) \geq \Delta^2(\text{fine} \rightarrow \text{mid}) \Delta^2(\text{mid} \rightarrow \text{coarse})$ for $(\text{fine}, \text{mid}, \text{coarse}) = [(8, 4, 2), (8, 6, 1), (6, 3, 0)]$: ['0.8665 $\geq$ 0.8665', '0.6188 $\geq$ 0.6188', '0.2888 $\geq$ 0.2888']
X2.8b	incremental ledger for $(\text{coarse}, \text{fine}) = (2, 3), (4, 5), (7, 8)$: $D_f - D_c = E D(P_Y f Y_c R_Y f Y_c)$: ['0.037415 = 0.037415', '0.002769 = 0.002769', '0.000044 = 0.000044'] ; $I_f - I_c = I(\Lambda ; Y_f Y_c)$: ['0.031139 = 0.031139', '0.002173 = 0.002173', '0.000034 = 0.000034'] ; $E \rho$ increments = I^* times the latter
X3.1	calibrated length of the ensemble path $L_I = 3.554071$; $D_0 - D_T = 3.554023$ (1.3e-05) ; $ v ^2 = I$ along the path : max rel gap 2.0e-15
X3.2	$\langle \Sigma_{\text{tot}} \rangle$ on 20000 paths = 3.5348 $\pm$ 0.0228 vs $D_0 - D_T = 3.5540$ (0.54 %)
X3.3	horizon : D against $\log(1/T)$ has slope 0.9982 (expected 1) ; W_2 stays in [1.1686, 1.2855] $< \sqrt{\text{tr } C_{\inf}}$ = 1.2985
X3.4	other weights on the same path : W_2 length 1.8571, weight I (not sqrt) 9.1525, against $\Delta D = 3.5540$
X4.3	$j = \text{grad}^{\perp} \psi$ with $\psi = s p$, $s = 0.310345$: relative residual 7.5e-16
X4.1b	information ledger of a zone : disk $r = 1.880$: flux term -2.3e-16, bulk term -3.1e-16 ; level ellipse : flux -2.9e-16, bulk term -1.6e-16 ; pointwise $\langle j, \text{grad } \log p \rangle = 0$: 1.3e-16 (the current runs along the level sets of p)
X4.2	$\sigma_A(\text{disk } r) = 0.171849 > 0$; $\sigma_A(r = 6 \sqrt{\lambda_{\max}}) = 0.279310$ vs $\sigma = 0.279310$ (3.6e-08)
X4.4	$T_1 = T_2$: $\sigma_A = 2.1e-31$, $ j _{\max} = 7.3e-17$: the structure is gone
X4.5	two ledgers : disk $\sigma_A = 0.171849$, $B_A = -3.1e-16$, $M_A = 0.171849$; ellipse $\sigma_A = 0.086618$, $B_A = -1.6e-16$, $M_A = 0.086618$; $\sigma_A + B_A - M_A$ relative residual 1.6e-16
X5.1	window KL ($n = 16$) for $x_1, x_2, x_1 + x_2$: ['0.0e+00', '8.9e-16', '8.9e-16']
X5.2	$\Sigma_{\text{visible}}(\Delta)/\Delta$ for $\Delta = 2 \dots 2^8$: ['0.0062', '0.0469', '0.1227', '0.1895', '0.2315', '0.2547', '0.2668', '0.2730', '0.2762', '0.2777'] $\rightarrow \sigma = 0.2793$ (last : 0.57 %)
X5.3	$\langle \Sigma \rangle / t$ on 20000 paths ($dt = 0.001$, $t = 4.0$) : 0.2839 $\pm$ 0.0037 vs $\sigma = 0.2793$ (1.66 %)
X5.4	two coordinates at $\Delta = 2^8$ see 99.43 % of σ ; one coordinate sees 0

gate	measurement (green)
X6.1	discrete Kirchhoff of the continuum current at $\text{eps} = 0.1$ $\sigma_{\min} : \max \text{sum of facet fluxes} / \max \text{flux} = 6.7\text{e-}16$
X6.2	entropy production of the four-neighbour chain, $\text{eps}/\sigma_{\min} = 0.4, 0.2, 0.1, 0.05$: [0.27331, 0.27762, 0.27871, 0.27898] vs $\sigma = 0.27931$; relative errors ['0.021', '0.006', '0.002', '0.001'], log-slope 1.40
X6.3b	the eight-neighbour chain (corner cliques) : production [0.8683, 0.89005, 0.89599, 0.89735] against $\sigma = 0.27931$, excess ['2.13', '2.19', '2.21', '2.21'] : above 100 % at every eps and flat in eps ; the corner links are mobility, they break the facet ledger
X6.4b	perplexity e^H against $2 \pi e \sqrt{\det C} / \text{eps}^2$: ratios ['1.0005', '0.9993', '0.9990', '0.9989'] (the canonical count of live cells)
X6.5	cell graph : 28363 facets on the same lattice at $k = 0$ and $k = 0.6$; conditional-independence graph : precision $C^{-1}_{12} = 0.0\text{e}+00$ at $k = 0$, -0.6416 at $k = 0.6$; generator hyperedge $A_{12} = -0.0$ at $k = 0$, -0.6 at $k = 0.6$
X6.6	law ruler of $(\log k_1, \log k_2)$ at $(k_1, k_2) = (1, 0.7)$: Gaussian curvature $K(k) = [0.0, -0.031, -0.0909, -0.1515, -0.2018, -0.2422, -0.2751, -0.2996, -0.3196]$ for $k = 0.0.8$, flat at $k = 0$ and decreasing ; partial correlation [0.0, 0.12, 0.217, 0.295, 0.359, 0.411, 0.456, 0.494, 0.526], $I(x_1;x_2) = [-0.0, 0.0073, 0.024, 0.0454, 0.0688, 0.0927, 0.1165, 0.1397, 0.1622]$ nats, both increasing from zero ; control points at $k = 0.6$: $K = [-0.350', -0.201']$
X7.1	$D(\mu \mu) = \sum_i D(\mu_i \mu_i) + D(\mu \mu_1 \times \mu_2)$ on ten posteriors : max gap 3.9e-16
X7.2	sensor on x_1 : coupling part ['0.0e+00', '0.0e+00', '0.0e+00', '0.0e+00'], cell part [0.2704, 1.2063, 2.2754, 3.476] increasing ; joint sensor $y = x_1 + x_2 + \text{noise}$: coupling part [0.1627, 0.9481, 1.9893, 3.1863]
X7.3	coupled prior (multi-information 0.1165), sensor on x_1 of increasing precision : coupling part ['1.02e-01', '5.18e-02', '7.85e-03', '9.27e-04', '8.41e-05', '9.35e-06'] decreasing to 0
X8.5	ledger $I_{\text{pred}} = I_{\text{ret}} + I_{\text{hid}}$, $\Delta = [0.5, 1, 2, 5, 10, 20]$: $T = 0.15$: $\text{eps}_{\text{cl}} = ['0.005', '0.004', '0.003', '0.002', '0.002', '0.002']$, $\text{eta}_{\text{mem}} = ['0.994', '0.985', '0.971', '0.934', '0.882', '0.794']$ ($H(X) = 0.308$) ; $T = 1$: $\text{eta}_{\text{ret}} = ['0.884', '0.899', '0.878', '0.703', '0.594', '0.635']$, $\text{eta}_{\text{mem}} = ['0.351', '0.158', '0.040', '0.001', '0.000', '0.000']$ ($H(X) = 0.692$)
X8.6b	redeclared : $R \geq 10$ floors where memory is claimed, $T = 0.15$ ['39.4', '39.1', '38.4', '36.8', '34.3', '30.1'], $T = 1$ $\Delta \leq 2$ ['2806.3', '811.6', '197.9'] ; within a decade of the floor at $T = 1$, $\Delta \geq 5$: ['5.8', '0.3', '3.4'] ; D/floor_D at $T = 0.15$ ['11.1', '8.7', '7.0', '5.0', '4.1', '2.9'] with $D \leq 0.01$ P
X8.7	estimator tower on nested grids $30 < 60 < 120$: P monotone at every (T, Δ) ; increments $P_{120} - P_{60}$ relative to P_{60} : $T = 0.15$ ['0.000', '0.000', '0.000', '0.000', '0.000', '0.000'], $T = 1$ ['0.000', '0.000', '0.001', '0.013', '0.119', '0.034']
X8.8	Pinsker : $e_{\text{TV}} \leq \sqrt{D/2}$ everywhere ; e_{TV} at $T = 0.15$ ['0.001', '0.001', '0.001', '0.001', '0.001', '0.001'], at $T = 1$ ['0.075', '0.053', '0.032', '0.008', '0.003', '0.005'] ; tightness $e_{\text{TV}}/\sqrt{D/2}$: $T = 0.15$ ['0.03', '0.03', '0.03', '0.04', '0.06', '0.08'], $T = 1$ ['0.63', '0.72', '0.78', '0.81', '0.71', '0.72']
X8.2b	$R(\text{dt}) = H(\text{occupation})$: 0.3077 vs 0.3080 at $T = 0.15$ (occupation 0.908), 0.6883 vs 0.6923 at $T = 0.25$ (occupation 0.520, close to $\log 2$) ; $R(500)$ at $T = 0.25 = 0.0028 \rightarrow 0$ (Kramers time 61)
X8.3	live cells (width 0.1) of the right-well belief for $T = 0.4, 0.2, 0.1, 0.05$: [16, 15, 10, 7], decreasing ; the calibrated distance from the two-atom reference to the certainty 'right' is $D(\delta_R \text{uniform}) = \log 2 = 0.6931$ nats, finite
X8.4	Kramers : escape rates ['8.81e-03', '3.44e-03', '1.00e-03'] at $T = [0.25, 0.2, 0.16]$, log-slope against $1/T = -0.968$ (barrier $h = 1$)
RING.1	closed-form density : $(F - U) p - T p' = j$ to $2.4\text{e-}07$ relative (spectral derivative, 65536 points ; $1.6\text{e-}08$ on 4000 points), mass 1.0000000000, $j = 0.099714 > 0$; control $F = 0$: $p = \text{Gibbs}$ to $2.2\text{e-}16$, $j = 0.0\text{e}+00$
RING.2	zone $[0, \pi]$: $\Phi_A = 0.109239$, $B_A = 0.109239$ (both nonzero, equal to $2.6\text{e-}10$), $\sigma_A = 0.916141 = M_A - B_A$ with $M_A = 1.025380$ (residual $3.2\text{e-}11$) ; total $\sigma = 1.253021$ against $F^2/T = 2.000$ at uniform potential
RING.3	port ledger : mean velocity $0.626523 = j L = 0.626523$; work of the drive $W = F j L = 0.626523 = \text{heat delivered to the bath}$; $\sigma_{\text{total}} = 1.25304638 = W/T = 1.25304638$ (residual $5.7\text{e-}11$)

gate	measurement (green)
RING.4	weighted reference $m = \text{Gibbs on } [0, \pi] : \sigma_A = 0.916141 = M_A - B_{m,A} - \int \langle v, \text{grad log } m \rangle d\mu = 1.025380 - (-0.289618) - 0.398857$ (residual $3.2e-11$) ; with the Lebesgue reference the bulk term is $B_A = 0.109239$: the bulk term depends on the reference, the identity carries the reference term

S.E Open problems

Open problem Q1 (The Hodge decomposition of hidden irreversibility). Can the hidden path-space relative entropy of a coarse-graining be decomposed into contributions of the exact, coexact and harmonic components of the current? The gyrator carries a coexact current, the driven ring a harmonic one.

Open problem Q2 (Beyond uniformly almost-sufficient towers). The radial convergence is acquired by exhaustive recovery (R5) and the non-radial convergence of lengths and distances along uniformly almost-sufficient towers by Corollary G11. What remains: the compactness and convergence of minimizing geodesics, the towers without uniform almost sufficiency, and the finite-state realizations whose coarse metrics leave the smooth stratum, where the evolutionary Γ -convergence of entropic gradient-flow structures (Forkert et al., 2022) is the available tool.

Open problem Q3 (Port references and a common thermodynamic statistic). Under what conditions can a family of port references be represented by a common finite-dimensional thermodynamic statistic, energy being the canonical candidate? The Darmois–Koopman–Pitman theorem is a lead, and the formalization of the port family is open.

Open problem Q4 (Stratified quotients). The operational geometry on quotients whose strata have different dimensions, and its spectral limit. Recent convergence results for graph Dirichlet energies and graph Laplacians on unions of intersecting manifolds of varying dimensions (Bungert and Slepčev, 2026) suggest an analytic backend once the metric, the measure and the transmission conditions across strata are fixed; the operational problem is to derive those conditions from the channel quotient.




Figure 12: The driven ring. Left: the stationary density (blue) and the potential (grey) for $U = \cos x$, $F = 1$, $T = 0.5$; the current is constant. Right: the ledger of the zone $[0, \pi]$: the interface flux and the bulk term are equal and nonzero, the entropy production equals the medium term minus the bulk term. [E1, H1]

S.F Notation

$D(P \parallel R)$	relative information of P from the declared reference R , in nats
$\mathfrak{J} = (\mathcal{M}_{\text{adm}}, \mathcal{S}, \mathcal{E}, R, \pi; m_\Lambda)$	the datum: candidates, semantics, operational access, path reference, prior, calibration
$p_t \in \mathcal{P}(\mathbf{X})$, m_X	ensemble law of a physical state and its reference (K0)
$\bar{\Lambda}_{\mathcal{E}}$, h	identifiable quotient and its Fisher metric (K1, O1)
$\mu \in \mathcal{P}(\bar{\Lambda}_{\mathcal{E}})$, π_Y , $I(\Lambda; Y)$	belief, posterior, mutual information (B1)
$\mathcal{H}_m$, $\mathcal{I}_{h,m}$, $W_{2,h}$, $d\ell_1$, $\ell_\star$	relative entropy, relative Fisher information, transport distance, calibrated length, gauge (G1, G2)
Σ , σ , Q_i	path irreversibility, entropy production rate, heat delivered to bath i (K5, H2)
A , $\mathbf{D}$, C , Γ , s	drift, diffusion matrix, stationary covariance, current matrix and vorticity of the gyrator (H1)
K_N , D_N , D_N^{hid} , h_N , I_N , ρ_I	resolution, visible and hidden ledgers, calibrated radius (R1, R2)
I_{pred} , I_{ret} , I_{hid} , ε_{cl} , η_{mem}	predictive, retained, hidden information of a quotient at time resolution Δ , loss and memory (E6)
Φ_A^{rel} , B_A^{rel} , σ_A , M_A	interface flux, bulk term, entropy production and medium term of a zone (E1)
k_B , $\dot{W}$, $\dot{Q}_i$	Boltzmann's constant, set to one in the units of the gyrator; work rate received from the work ports, heat rate delivered to bath i (H2)