

Errata (Updated May 2026)

Page 38 (Section 2.7), Equation 2.2 and 2.3, where μ_3 and μ_4 denote the third and fourth moments, respectively.

Page 67 (**R Note**)

Diff <- Diving2017\$Final - Diving2017\$Semifinal

Page 203

Figure 7.10 Tick marks on x-axis should be q_1, q_2 and not $-q, q$.

Paragraph before Example 7.12: If $P(q_1 < \hat{\theta}/\theta < q_2)$, then $P(\hat{\theta}/q_2 < \theta < \hat{\theta}/q_1) = 1 - \alpha$ and $(\hat{\theta}/q_2, \hat{\theta}/q_1)$ is a $1 - \alpha$ confidence interval.

Example 7.12, second paragraph: ... then the confidence interval is $(\bar{X}/q_2, \bar{X}/q_1)$.

Page 225 (Section 7.7), in the second bullet,

$$1\mu_X$$

should be

$$1/\mu_X$$

Page 232, Exercise 7.10 R code Second line of code `counter <- 0`

Page 254 (Section 3.3.1.1) Third paragraph of this section:

In Example 3.4, ... one sample had 1664 observations... Using a permutation test,... P -value 0.0178...

Page 453 (Section 13.3.1) currently has

$$\hat{\mu} = \sum_{j=1}^J \pi_j \bar{x}_j$$

That should be

$$\hat{\mu} = \sum_{j=1}^J \pi_j \bar{y}_j$$

for consistency with earlier sections and to avoid confusion in Exercise 13.6.

Page 454 (Section 13.3.2) This equation

$$n_k^* = n \frac{\pi_k n_k}{\sum_{j=1}^n \pi_j n_j}.$$

should be

$$n_k^* = n \frac{\pi_k \sigma_k}{\sum_{j=1}^n \pi_j \sigma_j}.$$

Problem Solutions

2.19 a) 3, b) 3, c) 3.

4.17 (a) $N(15 - 2 \cdot 4, 3^2 + (-2)^2 2^2)$.

Thank you to readers, including Wesley Burr, Hessel Jonker, Derin Gezgin, for notifying us about these errors.