

Errata (Updated July 2026)

Page 28 Two bullets indicate to “place a cap” at the end of the whiskers. Caps are optional, and the figure (produced by ggplot2) does not include caps.

Page 38 (Section 2.7), Equation 2.2 and 2.3, where μ_3 and μ_4 denote the third and fourth moments, respectively.

Page 67 (**R Note**)

```
Diff <- Diving2017$Final - Diving2017$Semifinal
```

Page 203

Figure 7.10 Tick marks on x-axis should be q_1, q_2 and not $-q, q$.

Paragraph before Example 7.12: If $P(q_1 < \hat{\theta}/\theta < q_2)$, then $P(\hat{\theta}/q_2 < \theta < \hat{\theta}/q_1) = 1 - \alpha$ and $(\hat{\theta}/q_2, \hat{\theta}/q_1)$ is a $1 - \alpha$ confidence interval.

Example 7.12, second paragraph: ... then the confidence interval is $(\bar{X}/q_2, \bar{X}/q_1)$.

Page 225 (Section 7.7), in the second bullet,

$$1\mu_X$$

should be


$$1/\mu_X$$

Page 232, Exercise 7.10 R code Second line of code `counter <- 0`

Page 254 (Section 3.3.1.1) Third paragraph of this section:

In Example 3.4, ... one sample had 1664 observations. ... Using a permutation test, ... P -value 0.0178. ...

Page 339, Figure 9.16 is the wrong figure. Here is the correct figure



Page 445 (Figure 13.1 caption)

Page 448 (Figure 13.3 caption) (same correction for both
 ... (b) The **median** and (c) the **mean**...

Page 453 (Section 13.3.1) currently has

$$\hat{\mu} = \sum_{j=1}^J \pi_j \bar{x}_j$$

That should be

$$\hat{\mu} = \sum_{j=1}^J \pi_j \bar{y}_j$$

for consistency with earlier sections and to avoid confusion in Exercise 13.6.

Page 454 (Section 13.3.2) This equation

$$n_k^* = n \frac{\pi_k n_k}{\sum_{j=1}^n \pi_j n_j}.$$

should be

$$n_k^* = n \frac{\pi_k \sigma_k}{\sum_{j=1}^n \pi_j \sigma_j}.$$

Problem Solutions

2.19 a) 3, b) **3**, c) **3**.

4.17 (a) $N(15 - 2 \cdot 4, 3^2 + (-2)^2 2^2)$.

Thank you to readers, including Wesley Burr, Hessel Jonker, Derin Gezgin, Roy St Laurent, for notifying us about errors.