

## Stochastic Games<sup>1</sup>)

By *J.-F. Mertens*, Louvain-la-Neuve<sup>2</sup>), and *A. Neyman*, Berkeley<sup>3</sup>)

**Abstract:** Stochastic Games have a value.

### 1. Introduction

A stochastic game is played in stages. At each stage, the game is in one of finitely many states and every player observes the current state  $z$  and chooses one of finitely many actions. The pair of actions at stage  $i$ , together with  $z$  determines the payoff  $x_i$  to be made by player II to player I at stage  $i$ , and the probability used by the referee to select the next state. All the referee's choices are made independently of the past. A player's (behavioural) strategy is a specification of a probability distribution over his actions at each stage conditional on the current state and the sequence of moves up to that stage. Any pair of strategies,  $\sigma$  of player I and  $\tau$  of player II, induces together with the initial state  $z_1$ , a probability distribution on the stream  $(x_1, x_2, \dots)$  of payoffs. The definition of a value depends on how the players evaluate a distribution of streams of payoffs. *Shapley* [1953] proved that the  $\lambda$ -discounted game, i.e.,

the game with "evaluation"  $E(\sum_{i=1}^{\infty} \lambda(1-\lambda)^{i-1} x_i)$  for  $0 < \lambda \leq 1$ , has a value and that

both players have optimal stationary strategies. *Bewley/Kohlberg* [1976], proved that the value  $v_{\lambda}(z)$  (respectively the optimal stationary strategy  $\sigma_{\lambda}$ ) of the  $\lambda$ -discounted game with initial state  $z$ , has a convergent expansion in fractional powers of  $\lambda$ , and

---

<sup>1</sup>) This work was initiated during an informal workshop at CORE on repeated games in the Winter of 78–79. Helpful comments and stimulus from the participants, especially S. Sorin and S. Zamir, is gratefully acknowledged. The work was continued in the Summer of 1979, first in Stanford, during the IMSSS Seminar, next at the Institute of Advanced Studies, at the Hebrew University of Jerusalem. Finally, a first draft could be put together during a visit of A. Neyman to the Mathematics Department at the Université Catholique de Louvain, and appeared as CORE DP 8001. The considerable extension and simplification we present here was carried out at the Institute for Advanced Studies in Jerusalem.

The second author's research was also supported throughout by National Science Foundation Grant MCS-79-06634 at U.C. Berkeley. Support from all those institutions is gratefully acknowledged.

<sup>2</sup>) Prof. *J.-F. Mertens*, C.O.R.E. and Department of Mathematics, Université Catholique de Louvain, Louvain-la-Neuve, Belgium.

<sup>3</sup>) Prof. *A. Neyman*, Department of Mathematics, University of California, Berkeley 94720; USA.

that the limit  $v_\infty(z)$  of  $v_\lambda(z)$  as  $\lambda \rightarrow 0$  exists. The question as to whether or not the undiscounted stochastic games, i.e., the games with "evaluation"  $E(\lim_{N \rightarrow \infty} \inf \bar{x}_N)$  where  $\bar{x}_N = (1/N) \sum_{i=1}^N x_i$ , always have a value, was open for many years. *Gillette* [1957]

proved the existence of the value in two cases: first when all games have perfect information and also in the so called cyclic case. *Blackwell/Ferguson* [1968] found in a particular example ("The Big Match") two strategies that would prove to be basic for further generalizations. Extending their second strategy *Kohlberg* [1974] proved that all "games with absorbing states" have a value. Recently we answered in the affirmative the question as to whether or not all stochastic games have a value [*Mertens/Neyman*]. The proof however was long and involved and the present paper presents a simplified proof of the main theorem.

**Theorem.** For every stochastic game and for every  $\epsilon > 0$  there exists a strategy  $\sigma$  of player I and  $N > 0$  such that for every  $n, n = N, N + 1, \dots, \infty$ , and for every strategy  $\tau$  of Player II,

$$E_{\sigma, \tau}(\bar{x}_n) \geq v_\infty - \epsilon$$

where by  $\bar{x}_\infty$  we mean  $\lim_{n \rightarrow \infty} \inf \bar{x}_n$ .

This form of statement means that the strategy  $\sigma$  is  $\epsilon$ -optimal both in the infinite game and in all sufficiently long finite games. The second implies in particular that the strategy  $\sigma$  is  $2\epsilon$ -optimal in all  $\lambda$ -discounted games with  $\lambda$  sufficiently small ( $\lambda \leq \sqrt{\epsilon/N}$ ).

Independently of ourselves, *Monash* [1980] announced a weaker version of the present result: it is not claimed that the strategy  $\sigma$  is good, neither in the infinite game nor in sufficiently long finite games, but only that for every strategy  $\tau$  of player II, there exists  $N$  such that for all finite  $n \geq N$  ( $n \neq \infty$ ),  $E_{\sigma, \tau}(\bar{x}_n) \geq v_\infty - \epsilon$ .

In section 2 we present the new short proof of the main theorem, and in section 3 a simplified proof of the  $\epsilon$ -optimality of our original strategy is given. In section 4 we present sufficient conditions on stochastic games with infinite sets of states and actions to have a value. For instance if payoffs are uniformly bounded, the value  $v_\lambda$  of the  $\lambda$ -discounted games exists, and for every  $\epsilon > 0$  there exists a sequence  $0 < x_i \leq 1$  converging to zero and such that  $x_{i+1} \geq (1 - \epsilon)x_i$  and

$$\sum \|v_{x_{i+1}} - v_{x_i}\| < \infty$$

where  $\| \cdot \|$  denotes the supremum norm (over the state space) then the undiscounted stochastic game has a value.

## 2. The Proof

In what follows we assume a fixed stochastic game. We will define a sequence  $(\lambda_i)_{i=1}$  so that  $\lambda_i$  is a function of the past history, i.e., measurable with respect to the

$\sigma$ -algebra  $F_i$  of all events preceding time  $i$  (including the choice of a new state  $z_i$  after time  $i-1$ ). The  $(\lambda_i)_{i=1}^\infty$  strategy of player I is to play on time  $i$  an optimal strategy in the  $\lambda_i$ -discounted game. For such a strategy,  $v_{\lambda_i}(z_i) \leq E(\lambda_i x_i + (1 - \lambda_i) v_{\lambda_i}(z_{i+1}) | F_i)$ , i.e.,

$$E(v_{\lambda_i}(z_{i+1}) - v_{\lambda_i}(z_i) + \lambda_i(x_i - v_{\lambda_i}(z_{i+1}))) | F_i \geq 0. \quad (2.1)$$

The basic result of *Bewley/Kohlberg* [1976, Theorem 3.4] implies the existence of  $0 < \lambda_0 < 1$ ,  $0 < r < 1$ ,  $B > 0$ , so that for any state  $z$ ,  $v_\lambda(z)$  is differentiable on  $(0, \lambda_0]$  and  $|(dv_\lambda(z))/(d\lambda)| \leq B\lambda^{-r}$  for  $0 < \lambda \leq \lambda_0$ . On  $(\lambda_0, 1]$ ,  $v_\lambda$  is Lipschitz ( $\|v_\lambda - v_\eta\| \leq (|\lambda - \eta|/\lambda) 2A$ , e.g. by lemma 4.2), where  $\|\cdot\|$  denotes the maximum norm on  $\mathbf{R}^S$ ,  $S$  the state space, and  $A$  denotes the largest absolute value of pay-offs appearing in the game matrices). Therefore, there exists a positive integrable function  $\psi: (0, 1] \rightarrow \mathbf{R}_+$  such that for  $0 < \lambda < \bar{\lambda} \leq 1$ ,

$$\|v_\lambda - v_{\bar{\lambda}}\| \leq \int_{\bar{\lambda}}^{\lambda} \psi(x) dx. \quad (2.2)$$

Let  $\epsilon > 0$  be given. It suffices to consider  $0 < \epsilon < A$ . Define

$$s(y) = \frac{12A}{\epsilon} \int_y^1 \frac{\psi(x)}{x} dx + y^{-1/2}, \quad 0 < y \leq 1.$$

Observe that  $s(y)$  is a strictly decreasing continuous function from  $(0, 1]$  onto  $[1, \infty)$  so that an inverse continuous function  $\lambda: [1, \infty) \rightarrow (0, 1]$  exists. Our strategy will depend on the function  $\lambda$  and on two additional constants  $M, s_1$  ( $s_1 \geq M \geq 1$ ) sufficiently large to satisfy further requirements that will be specified later.

We now define inductively:

$$\lambda_i = \lambda(s_i)$$

$$s_{i+1} = \text{Max} [M, s_i + x_i - v_{\lambda_i}(z_{i+1}) + 4\epsilon].$$

We first observe the following:

$$|s_{i+1} - s_i| \leq 6A \quad (2.3)$$

$$x_i - v_{\lambda_i}(z_{i+1}) + 4\epsilon \leq s_{i+1} - s_i \leq x_i - v_{\lambda_i}(z_{i+1}) + 4\epsilon + 2A I(s_{i+1} = M) \quad (2.4)$$

where  $I$  denotes the indicator function.

Since  $s((1 - \epsilon)\lambda) - s(\lambda) \geq [(1 - \epsilon)^{-1/2} - 1] \lambda^{-1/2} \rightarrow \infty$  as  $\lambda \rightarrow 0$ , (2.3) implies that there is  $M_0 \geq 1$  such that for  $M \geq M_0$

$$|\lambda_{i+1} - \lambda_i| \leq \epsilon \lambda_i / (6A). \quad (2.5)$$

Thus for  $M \geq M_0$ , by using (2.3), the definition of  $s$ , (2.5) and (2.2),

$$\begin{aligned} 6A &\geq |s_{i+1} - s_i| \geq \frac{12A}{\epsilon} \left| \int_{\lambda_i}^{\lambda_{i+1}} \frac{\psi(x)}{x} dx \right| \geq \frac{12A}{\epsilon} \frac{1}{2\lambda_i} \left| \int_{\lambda_i}^{\lambda_{i+1}} \psi(x) dx \right| \geq \\ &\geq \frac{6A}{\epsilon\lambda_i} \|v_{\lambda_{i+1}} - v_{\lambda_i}\|. \end{aligned}$$

Hence

$$\|v_{\lambda_i} - v_{\lambda_{i+1}}\| \leq \epsilon \lambda_i. \quad (2.6)$$

We now verify the integrability of  $\lambda$  on  $(1, \infty)$ .

$$\begin{aligned} 1 + \int_1^\infty \lambda(s) ds &= \int_0^1 s(y) dy = \int_0^1 y^{-1/2} dy + \frac{12A}{\epsilon} \int_0^1 \int_0^1 I(y \leq x \leq 1) \frac{\psi(x)}{x} dy dx = \\ &= \int_0^1 y^{-1/2} dy + \frac{12A}{\epsilon} \int_0^1 \psi(x) dx < \infty. \end{aligned}$$

Define  $t: [1, \infty) \rightarrow \mathbf{R}_+$  by  $t(s) = \int_s^\infty \lambda(y) dy$ ,  $t_i = t(s_i)$ . Observe that  $t$  is differentiable,  $t'(s) = -\lambda(s)$  and that  $t(s)$  decreases to 0 as  $s$  goes to  $\infty$ . Using the mean value theorem together with (2.3) and (2.5),

$$t_i - t_{i+1} \geq \lambda_i (s_{i+1} - s_i) - \epsilon \lambda_i. \quad (2.7)$$

Replacing in (2.1),  $v_{\lambda_i}(z_{i+1})$  by  $v_{\lambda_{i+1}}(z_{i+1})$ ,  $\lambda_i(x_i - v_{\lambda_i}(z_{i+1}))$  by  $\lambda_i(s_{i+1} - s_i) - 4\epsilon\lambda_i$ , while using the left inequality of (2.4), and (2.6), we have for sufficiently large  $M$ ,

$$E(v_{\lambda_{i+1}}(z_{i+1}) - v_{\lambda_i}(z_i) + \lambda_i(s_{i+1} - s_i) | F_i) \geq 3\epsilon\lambda_i. \quad (2.8)$$

Applying (2.7),  $E(v_{\lambda_{i+1}}(z_{i+1}) - v_{\lambda_i}(z_i) - t_{i+1} + t_i | F_i) \geq 2\epsilon\lambda_i$ , or in other words, letting  $Y_i = v_{\lambda_i}(z_i) - t_i$ ,

$$E(Y_{i+1} - Y_i | F_i) \geq 2\epsilon\lambda_i. \quad (2.9)$$

Since  $\lambda_i \geq 0$ , (2.9) means that  $Y_i$  is a submartingale. Obviously  $Y_i$  is bounded and thus it converges a.s., say to  $Y_\infty$ , with  $E(Y_\infty | F_1) > Y_1$ . It follows also that for sufficiently large  $M$ ,

$$4A \geq 2t(M) + 2A \geq E(Y_k - Y_1) \geq 2\epsilon E\left(\sum_{1 \leq i < k} \lambda_i\right), \quad (2.10)$$

so that by the monotone convergence theorem

$$E\left(\sum_{i < \infty} \lambda_i\right) \leq 2A/\epsilon, \quad \text{and} \quad (2.11)$$

$$E(\#\{i \mid \lambda_i \geq \eta\}) \leq \frac{2A}{\epsilon\eta}. \quad (2.12)$$

So that a.s.,  $\lambda_i \rightarrow 0$ ,  $s_i \rightarrow \infty$ ,  $t_i \rightarrow 0$ , and therefore

$$\begin{aligned} v_{\lambda_i}(z_i) &= Y_i + t_i \rightarrow Y_\infty \text{ a.s. and thus by (2.6) if } \int_0^{\lambda(M)} \psi(x) dx \leq \epsilon, \\ v_{\lambda_i}(z_{i+1}) &\rightarrow Y_\infty \text{ with } E(Y_\infty \mid F_1) > Y_1 \geq v_\infty(z_1) - \epsilon - t_1. \end{aligned} \quad (2.13)$$

Also by (2.10) and (2.6) for every  $i$ ,

$$E(v_{\lambda_i}(z_{i+1})) \geq v_\infty(z_1) - 2\epsilon - t_1. \quad (2.14)$$

Now, summing the right hand inequalities of (2.4) for  $1 \leq i < n$ , we have

$$\sum_{i < n} x_i \geq \sum_{i < n} v_{\lambda_i}(z_{i+1}) + s_n - s_1 - 2A \sum_{i < n} I(s_{i+1} = M) - 4n\epsilon. \quad (2.15)$$

The result follows now by bounding the terms in (2.15) using (2.14), (2.13) and (2.12) and observing that for  $M$  sufficiently large  $t$  is sufficiently small.

### 3. Other $\epsilon$ -Optimal Strategies

Denote by  $A$  four times the largest absolute value of payoffs, and let  $\epsilon > 0$ . Assume without loss of generality  $\epsilon \leq A$ , and let  $\delta = \epsilon/(12A)$ . In what follows we denote  $v(z, \lambda) = v_\lambda(z)$ ,  $v(z, 0) = v_\infty(z)$ .

Take two functions  $L(s)$  and  $\lambda(s)$  of a real variable  $s$ , with values in the positive integers for  $L$  and in  $(0, 1]$  for  $\lambda$ . Choose them such that  $\lambda(s)$  is decreasing and such that, for all  $s$  sufficiently large and all  $\theta$  with  $|\theta| \leq A$ , and for every state  $z$ :

$$\frac{AL(s)}{s} \leq \delta \quad (i)$$

$$\left| \frac{\lambda(s + \theta L(s))}{\lambda(s)} - 1 \right| \leq \delta \quad (ii)$$

$$|v(z, \lambda(s + \theta L(s))) - v(z, \lambda(s))| \leq \delta AL(s)\lambda(s) \quad (\text{iii})$$

$$\int_0^\infty \lambda(s) ds < +\infty. \quad (\text{iv})$$

For instance, for a finite stochastic game where — using the *Bewley/Kohlberg* result [1976, Th. 3.4] —  $|v(z, \lambda) - v(z, 0)| \leq B\lambda^{1-r}$ , with  $0 \leq r < 1$ , one could take  $\lambda(s) = s^{-\beta}$ ,  $L(s)$  the minimal integer exceeding  $2(A\delta)^{-1}B(\lambda(s))^{-r}$  — where  $\beta > 1$  such that  $\beta r < 1$  — (this yields the strategy used in *Mertens/Neyman* [1980]).

Or alternatively, as in the previous proof,  $L(s) = 1$ ,  $\lambda(s) = \eta s^{-1/r}$  for  $\eta$  sufficiently small — or still:  $L(s) = 1$ ,  $\lambda(s) = 1/(s \ln^2 s)$ .

Our strategy will also depend on two constants  $M$  and  $s_0$  ( $s_0 \geq M$  — one can always choose  $s_0 = M$ ) sufficiently large such as to satisfy further requirements. To begin with,  $M$  will be assumed such that

$$\forall s \geq M, (i) - (iv) \text{ hold and } |v(z, \lambda(s)) - v_\infty(z)| \leq A\delta. \quad (3.0)$$

Define now inductively, using  $x_i$  (resp.  $z_i$ ) for the payoff (resp. state) at stage  $i$  ( $i = 1, 2, \dots$ ), starting with  $s_0 \geq M$ :

$$\lambda_k = \lambda(s_k), L_k = L(s_k);$$

$$B_0 = 1, B_{k+1} = B_k + L_k;$$

$$s_{k+1} = \text{Max} [M, s_k + \sum_{B_k \leq i < B_{k+1}} (x_i - v_\infty(z_{B_{k+1}})) + \epsilon/2].$$

The strategy is to start playing from time  $B_k$  up to time  $B_{k+1}$  a  $(\delta AL_k \lambda_k)$  optimal strategy in the  $\lambda_k$ -discounted game.

We denote by  $F_i$  ( $i = 1, \dots, \infty$ ) the  $\sigma$ -field of all events preceding stage  $i$  (including the choice of the state  $z_i$ ) and let  $G_k = F_{B_k}$ .

Observe that

$$|s_{k+1} - s_k| \leq AL_k \quad (3.1)$$

and that,  $\lambda$  being decreasing and integrable,

$$\lim_{s \rightarrow \infty} s\lambda(s) = 0,$$

and therefore by (i)

$$\lim_{s \rightarrow \infty} \lambda(s)L(s) = 0.$$

In particular, assuming  $M$  sufficiently large,

$$\lambda(s)L(s) \leq \delta \text{ for } s \geq M. \quad (3.2)$$

Let  $l_k = v(z_{B_k}, \lambda_k)$ , and note that by (iii)

$$|v(z_{B_{k+1}}, \lambda_k) - l_{k+1}| \leq \delta AL_k \lambda_k. \quad (3.3)$$

*Lemma 3.4.*

$$E(l_{k+1} - l_k + \lambda_k(s_{k+1} - s_k) | G_k) \geq 2\delta AL_k \lambda_k.$$

*Proof.* Assume without loss of generality  $k = 0$ , and write  $\lambda, L, v^1(\lambda)$  for  $\lambda_0, L_0, v(Z_{B_1}, \lambda_0)$  respectively. Then, by the  $\delta AL\lambda$ -optimality,

$$l_0 \leq E(\lambda \sum_{i < L} (1 - \lambda)^i x_{i+1} + (1 - \lambda)^L v^1(\lambda)) + \delta AL\lambda$$

or

$$(as \ 1 - \lambda \sum_{i < L} (1 - \lambda)^i = (1 - \lambda)^L)$$

$$E(v^1(\lambda) - l_0 + \lambda \sum_{i < L} (1 - \lambda)^i (x_{i+1} - v^1(\lambda))) \geq -\delta AL\lambda.$$

Using (3.3),  $1 - \lambda L \leq (1 - \lambda)^i \leq 1$  for  $i < L$ , (3.2) and (3.0) we get

$$E(l_1 - l_0 + \lambda \sum_{i < L} (x_{i+1} - v^1(0))) \geq -4\delta AL\lambda,$$

and therefore by the inequality  $s_1 - s_0 \geq \sum_{i < L} (x_{i+1} - v^1(0) + 6\delta A)$  we have:

$$E(l_1 - l_0 + \lambda(s_1 - s_0)) \geq 2\delta AL\lambda.$$

Let now  $t(s) = \int_s^\infty \lambda(x)dx$ ,  $t_k = t(s_k)$ . Observe that  $t(s)$  decreases to zero (by iv), in particular we can assume  $t(M) \leq \delta A$ .

*Lemma 3.5.*

$$E[(l_{k+1} - t_{k+1}) - (l_k - t_k) | G_k] \geq \delta AL_k \lambda_k.$$

*Proof.* Using (ii) and (3.1):

$$t_{k+1} - t_k = \int_{s_{k+1}}^{s_k} \lambda(s) ds \leq \lambda_k(s_k - s_{k+1}) + \delta AL_k \lambda_k$$

and the result follows from lemma 3.4.

Let  $k(i)$  be the  $(G_k)_{k=0}^\infty$ -stopping time  $\inf\{k \mid B_k > i\}$ , and let  $\bar{\lambda}_i = \lambda_{k(i)-1}$ ,  $\bar{l}_i = v_\infty(z_{B_{k(i)}})$ . ( $\bar{\lambda}_i$  is the discount rate "used at stage  $i$ ".)

*Proposition 3.6.*

- a)  $l_k$  converges a.s., say to  $l_\infty$ ; and  $s_k$  converges a.s. to  $+\infty$ .  
 b) For any  $(G_k)_{k=0}^\infty$ -stopping time  $T$ ,

$$E(l_T \mid G_0) \geq l_0 - t_0 (\geq l_0 - \delta A)$$

$$c) E\left(\sum_{i=1}^{\infty} \bar{\lambda}_i\right) \leq \delta^{-1}.$$

Note in particular the following consequences (using (3.0) and the uniform convergence of  $v_\lambda$  to  $v_\infty$  (cfr. section 4)):

$$\bar{l}_i \text{ converges a.s. to } \bar{l}_\infty = l_\infty$$

$$\forall i = 0, 1, 2, \dots, \infty, E(\bar{l}_i \mid G_0) \geq \bar{l}_0 - 3\delta A (= v_\infty(z_1) - 3\delta A)$$

$$E \sum_k I(s_k = M) \leq 1/(\delta \lambda(M)).$$

*Proof.* By lemma 3.5,  $Y_k = l_k - t_k$  is a bounded (by  $A$ ) submartingale, and therefore converges a.s. to  $Y_\infty$ .

Since  $Y_k - Y_0 \leq A$ , the same lemma implies further

$$A \geq E(Y_k - Y_0) \geq \delta A E\left(\sum_{i < B_k} \bar{\lambda}_i\right).$$

c) Follows now by the monotone convergence theorem, and implies in particular that a.s.  $\lambda_k \rightarrow 0$ , thus  $s_k \rightarrow \infty$ , thus  $t_k \rightarrow 0$  and therefore  $l_k = Y_k + t_k$  converges a.s. to  $l_\infty = Y_\infty$ .

The stopping theorem for bounded submartingales implies now  $E(l_T \mid G_0) \geq E(Y_T \mid G_0) \geq Y_0 = l_0 - t_0$ , which completes the proof.

*Lemma 3.7.*

$$\sum_{i=1}^n x_i \geq \sum_{i=1}^n \bar{l}_i - 2s_0 - 8\delta A n - \delta M \sum_{k=1}^{\infty} I(s_k = M).$$

*Proof.* The definition of  $s_k$  implies that

$$s_{k+1} - s_k \leq \sum_{B_k \leq i < B_{k+1}} (x_i - \bar{l}_i) + 6\delta A L_k + I(s_{k+1} = M) A L_k / 2.$$



Since  $AL(s)/s \leq \delta$ , (3.1) implies that, when  $s_{k+1} = M$ ,  $AL_k \leq \delta M/(1 - \delta)$ ;

$$s_{k+1} - s_k \leq \sum_{B_k \leq i < B_{k+1}} (x_i - \bar{l}_i) + 6\delta AL_k + \delta MI(s_{k+1} = M).$$

By summing

$$s_k - s_0 \leq \sum_{i < B_k} (x_i - \bar{l}_i) + 6\delta AB_k + \delta M \sum_{l=1}^{\infty} I(s_l = M),$$

and thus

$$\begin{aligned} \sum_1^n x_i &\geq s_{k(n)} - s_0 - A(B_{k(n)} - n) + \sum_1^n \bar{l}_i - 6\delta AB_{k(n)} - \delta M \sum_{k=1}^{\infty} I(s_k = M) \\ &\geq \sum_1^n \bar{l}_i - s_0 - 6\delta An - 2A(B_{k(n)} - n) - \delta M \sum_k I(s_k = M). \end{aligned}$$

But  $B_{k(n)} - n \leq L(s_{k(n)-1}) \leq A^{-1}\delta s_{k(n)-1} \leq \delta(A^{-1}s_0 + n)$ , so the result follows.

The  $\epsilon$ -optimality of the strategy follows now immediately by bounding the terms in lemma 3.7 using the consequences of proposition 3.6.

#### 4. Infinite Stochastic Games

The finiteness hypothesis on the state space and the action sets we made in the definition of stochastic games are by no means necessary: the only thing required by our proof is

- (1) that payoffs are uniformly bounded
- (2) that the value  $v_\lambda$  of the  $\lambda$ -discounted games exists, and
- (3) that for any  $\delta > 0$  one can find functions  $\lambda(s)$  and  $L(s)$

satisfying the conditions of section 3, i.e. for  $s$  sufficiently large the following hold (all  $|\theta| \leq A$ );

a)  $\lambda(s)$  has values in  $(0, 1]$  and  $L(s)$  in  $1, 2, 3, \dots$

b)  $\lambda(s)$  is monotone and integrable

c)  $\frac{AL(s)}{s} \leq \delta$

d)  $|\frac{\lambda(s + \theta L(s))}{\lambda(s)} - 1| \leq \delta$

e)  $|v_z[\lambda(s + \theta L(s))] - v_z[\lambda(s)]| \leq \delta AL(s)\lambda(s)$ .

In this section we formulate (3) explicitly as a condition on  $v$  only.

Note that the monotone character of  $\lambda(s)$  was introduced in section 3 as a matter of convenience, and that we used it only to guarantee such easy properties as

$\lim_{s \rightarrow \infty} s\lambda(s) = 0$ . For this reason we will show at the same time that this convenience

was no restriction.

*Theorem 4.1.* If

- (1) payoffs are uniformly bounded
- (2) the values  $v_\lambda(z)$  of the  $\lambda$ -discounted games exists
- (3\*)  $\forall \alpha < 1$  there exists a sequence  $\lambda_i$  ( $0 < \lambda_i \leq 1$ ) such that  $\lambda_{i+1} \geq \alpha \lambda_i$ ,  
 $\lim_{i \rightarrow \infty} \lambda_i = 0$  and  $\sum_i \sup_z |v_{\lambda_i}(z) - v_{\lambda_{i+1}}(z)| < \infty$

then  $v_\infty(z) = \lim_{\lambda \rightarrow 0} v_\lambda(z)$  exists (uniformly in  $z$ ) and the undiscounted game has  $v_\infty$  as value, in the sense that

$$\forall \epsilon > 0 \exists \sigma \text{ (a strategy of player 1)} \exists N: \forall \tau \text{ (strategy of player 2)} \forall z \text{ (initial state)}, \forall n = N, N+1, N+2, \dots, \infty, E_{\sigma, \tau}^z(\bar{x}_n) \geq v_\infty(z) - \epsilon$$

where  $\bar{x}_n$  denotes the average payoff up to stage  $n$ ,  $\bar{x}_\infty = \liminf_{n \rightarrow \infty} \bar{x}_n$ .

A dual statement holds interchanging the two players.

*Remark:* The third condition is for example obviously satisfied in each of the following cases:

- $v_\lambda$  is of bounded variation — the variation being computed using the supremum norm  $\|\cdot\|$  over the state space.
- For some sequence  $\lambda_i$  with  $\lambda_i > 0$ ,  $\inf \lambda_{i+1} / \lambda_i > 0 = \lim \lambda_i$  one has

$$\sum \Delta v[\lambda_{i+1}, \lambda_i] < \infty$$

where  $\Delta v[x, y] = \sup \{ \|v_{\lambda_1}(\cdot) - v_{\lambda_2}(\cdot)\| \mid \lambda_i \in (0, 1], x \leq \lambda_i \leq y \}$  (and  $\sup \phi = 0$ ).

- there exists a function  $v_\infty(z)$  such that  $\|v_\lambda - v_\infty\| / \lambda$  is integrable (indeed, this is equivalent to the integrability of  $\|v_\lambda - v_\infty\|$  as a function of  $\ln(\lambda)$ , so that taking for  $\lambda_i$  the minimizer of  $\|v_\lambda - v_\infty\|$  in the interval  $i(1-\alpha) \leq -\ln \lambda \leq (i+1)(1-\alpha)$  yields the condition).

A property of  $v$  that is always valid is the following:

*Lemma 4.2.* If (1) and (2) hold then

$$|v_\lambda / \lambda - v_\eta / \eta| \leq A |\lambda^{-1} - \eta^{-1}|$$

where  $A$  is the upper bound of the absolute value of payoffs. In particular  $v_\lambda$  is Lipschitz in  $\ln \lambda$  (and also  $\|v_\lambda - v_\eta\| \leq 2A\lambda^{-1} |\lambda - \eta|$ ).

*Proof.* The difference of the payoff functions  $|\sum (1-\lambda)^i x_i - \sum (1-\eta)^i x_i|$  is at most  $A |\lambda^{-1} - \eta^{-1}|$  and thus the difference in the values  $|v_\lambda / \lambda - v_\eta / \eta|$  is bounded by the same constant.

The theorem follows from the results of section 3 and the following (note that (4.7), together with lemma (4.2), implies the norm convergence of  $v_\lambda$ ):

*Proposition 4.3.* The following conditions on  $v_\lambda$  are equivalent:

- (4.4) For some  $A > 0$ , there exist for every  $\delta > 0$  functions  $L(s)$  and  $\lambda(s)$  (not necessarily monotonic) satisfying conditions a) – e).  
 (4.5) There exist functions  $L(s)$  and  $\lambda(s)$  (strictly decreasing) that satisfy a) – e) whatever be  $A > 0$  and  $\delta > 0$ .  
 (4.6)  $\forall \alpha < 1$ , there exists a sequence  $0 < \lambda_i \leq 1$  converging to 0 and such that  $\lambda_{i+1} \geq \alpha \lambda_i$  and

$$\sum_i \|v_{\lambda_{i+1}} - v_{\lambda_i}\| < \infty.$$

- (4.7) There exists a strictly decreasing sequence  $\lambda_i$  such that  $\lambda_0 = 1$ ,  $\lim \lambda_i = 0$ ,  $\lim \lambda_{i+1} / \lambda_i = 1$  and

$$\sum_i \|v_{\lambda_{i+1}} - v_{\lambda_i}\| < \infty.$$

*Proof.* We will prove (4.4)  $\rightarrow$  (4.6)  $\rightarrow$  (4.7)  $\rightarrow$  (4.5)  $\rightarrow$  (4.4). The last implication is obvious.

(4.4)  $\rightarrow$  (4.6): Take  $s_0 = 0$ ,  $\lambda_i = \lambda(s_i)$ ,  $s_{i+1} = s_i + L_i$  where  $L_i = AL(s_i)$ . Then

$$\|v_{\lambda_{i+1}} - v_{\lambda_i}\| \leq \delta \lambda_i L_i \text{ (by e)) and } \lambda_i L_i = \lambda_i(s_{i+1} - s_i) \leq (1 - \delta)^{-1} \int_{s_i}^{s_{i+1}} \lambda(s) ds$$

(by (d)). Thus  $\sum \|v_{\lambda_{i+1}} - v_{\lambda_i}\| \leq \delta \sum \lambda_i L_i \leq \delta (1 - \delta)^{-1} \sum \int_{s_i}^{s_{i+1}} \lambda(s) ds < \infty$ . Since

$L_i$  is bounded away from 0,  $\lambda_i \rightarrow 0$ . Finally by d)  $\lambda_{i+1} \geq \lambda_i (1 - \delta)$ .

(4.6)  $\rightarrow$  (4.7): Remark first that there is no loss in assuming the sequence  $\lambda_i$  of (4.6) to be strictly decreasing: let  $i_0 = 0$ ,  $i_{k+1} = \inf \{i : \lambda_i < \lambda_{i_k}\}$  and  $\tilde{\lambda}_k = \lambda_{i_k}$ . Then

$\tilde{\lambda}_k \rightarrow 0$ ,  $\tilde{\lambda}_{k+1} \geq \epsilon \tilde{\lambda}_k$  and by the triangle inequality the variation on the subsequence

$(\tilde{\lambda}_k)_{k=0}^\infty$  is at most the variation on the sequence. Observe also that there is no loss in assuming  $\lambda_0 = 1$ .

Thus, for every  $n \geq 1$ , there are strictly decreasing sequences  $(x_i^n)_{i=0}^\infty$  with  $x_0^n = 1$ ,

$\ln(x_i^n / x_{i+1}^n) \leq 2^{-n}$ ,  $x_i^n \rightarrow 0$  as  $i \rightarrow \infty$ , so that

$$\sum_i \|v_{x_{i+1}^n} - v_{x_i^n}\| < \infty.$$

Therefore, there is a sequence  $(a_n)_{n=1}^\infty$  with  $a_1 = 1$ ,  $\ln(a_n / a_{n+1}) > 1$  so that

$$\sum_{n=1}^{\infty} \sum_{i=0}^{\infty} \| \nu_{x_{i+1}^n} - \nu_{x_i^n} \| I(x_i^n \leq a_n) < \infty.$$

The decreasing sequence  $\lambda_i$  generated by the union of the nonempty sets

$\Lambda_n = \{x_i^n \mid a_{n+1} < x_i^n \leq a_n\}$  does the job. Obviously  $\lambda_0 = 1$ , and  $\lambda_i \rightarrow 0$ . To verify the other conditions on  $(\lambda_i)$ , let  $\bar{a}_n = \min \{x_i^{n-1} \mid x_i^{n-1} > a_n\}$ ,  $\underline{a}_n = \max \{x_i^n \mid x_i^n \leq a_n\}$ .

As  $\ln \bar{a}_n / \underline{a}_n \leq \ln \bar{a}_n / a_n + \ln a_n / \underline{a}_n \leq 2^{-n+1} + 2^{-n}$ ,  $\lambda_{i+1} / \lambda_i \rightarrow 1$ , and also

$\| \nu_{\bar{a}_n} - \nu_{\underline{a}_n} \| \leq K (2^{-n+1} + 2^{-n})$  (by lemma 4.2), which is summable, and thus

$$\sum \| \nu_{\lambda_{i+1}} - \nu_{\lambda_i} \| \leq \sum_{n=1}^{\infty} \sum_{x_i^n \leq a_n} \| \nu_{x_{i+1}^n} - \nu_{x_i^n} \| + \sum_{n=1}^{\infty} \| \nu_{\bar{a}_n} - \nu_{\underline{a}_n} \| < \infty.$$

(4.7)  $\rightarrow$  (4.5): Let  $(\lambda_i)_{i=1}^{\infty}$  be the sequence of (4.7), and define  $\bar{\nu}(\lambda)$  by linear interpolation from the values  $\bar{\nu}(\lambda_i) = \nu_{\lambda_i}$ . Let  $n \geq 1$ ,  $y = n/(n+1)$ ,  $l_i = \Delta \bar{\nu}[y^{i+1}, y^i]$  (thus

$l_i = 0$  for  $i < 0$ ). Observe that  $\sum l_i \leq \sum \| \nu_{\lambda_{k+1}} - \nu_{\lambda_k} \| < \infty$ . Let also  $\bar{l}_i = \sum_{|j| \leq 2} l_{i+j}$ ,

$g(x) = 2n[I(x \leq 1) + \bar{l}_i y^{-i}]$  for  $y^{i+1} < x \leq y^i$ , and let  $h$  be defined by linear interpolation from the values  $h(y^i) = n \sum_{j < i} [g(y^{j+1}) + g(y^j) + g(y^{j-1})]$ . Then  $g \geq n$  on

$(0, 1]$  and  $h$  is continuous, decreasing,  $\geq n g$  and integrable (e.g.

$$\begin{aligned} \int h dx &\leq \sum h(y^i) (y^{i-1} - y^i) = n (y^{-1} - 1) \sum y^i [g(y^{i+1}) + g(y^i) + g(y^{i-1})] \\ &= n \sum_j y^j [g(y^{j+1}) + g(y^j) + g(y^{j-1})] \leq \frac{3n}{y} \sum_{j < i} y^j g(y^j) = \frac{6n^2}{y} [1/(1-y) + \sum \bar{l}_i] < \infty. \end{aligned}$$

Further, for  $y^{i+1} < x \leq y^i$ ,  $\Delta \bar{\nu}[xy^2, xy^{-2}] \leq \Delta \bar{\nu}[y^{i+3}, y^{i-2}] \leq \bar{l}_i \leq (1/2n) y^i g(y^i) \leq (1/n) x g(x)$  (recall that  $y \geq 1/2$ ).

Since on the interval  $(y^{i+1}, y^i]$ ,  $h(xy^k)$  is linear (in  $x$ ), and  $g(x) = g(y^i)$ , the equality  $h(y^{i+1}) = h(y^i) + n[g(y^{i+1}) + g(y^i) + g(y^{i-1})]$  implies that

$$h(x) + n g(x) \leq h(xy) \quad \text{and} \quad h(x) - n g(x) \geq h(xy^{-1})$$

hold in every interval  $(y^{i+1}, y^i]$ . Altogether we found for each  $n \geq 1$  a pair of functions on  $(0, \infty)$ ,  $h_n$  and  $g_n$  (the integer part of  $g$ ), such that  $g_n$  is integer valued, and  $\geq n$  on  $(0, 1]$ , while  $h_n$  is continuous, decreasing,  $\geq n g_n$ , integrable, and such that letting  $y_n = n/(n+1)$

$$\Delta \bar{\nu}[x y_n^2, x y_n^{-2}] \leq \frac{1}{n} x g_n(x)$$

and

$$h_n(x) + n g_n(x) \leq h_n(x y_n), \quad h_n(x) - n g_n(x) \geq h_n(x y_n^{-1}).$$

Take a sequence  $(a_n)_{n=1}^{\infty}$  satisfying  $1/2 \geq a_n > a_{n+1} > 0$ ,

$$\inf \{ \lambda_i / \lambda_{i-1} \mid \lambda_i \leq 2 a_n \} \geq y_n,$$

and

$$\sum_{1 \leq n < \infty} \int_0^{a_n} h_n(x) dx < \infty; \text{ and set } \bar{h}_n(x) = h_n(x) (2 - x/a_n)^+.$$

Then  $\bar{h}(x) = \sum_1^{\infty} \bar{h}_n(x)$  is continuous, decreasing (strictly on  $(0, 2a_1]$ ) and integrable.

Thus, if we let  $g_0 = 1$ ,  $y_0 = 1/2$ , and  $n(x) = \# \{n \mid a_n \geq x\}$ ,  $\bar{g}(x) = g_{n(x)}(x)$ , we have  $\bar{g}(x) \geq n(x) \rightarrow \infty$  as  $x \rightarrow 0$ .

$$\Delta \bar{v}[xy_{n(x)}^2, xy_{n(x)}^{-2}] \leq \frac{1}{n(x)} x \bar{g}(x),$$

and

$$\begin{aligned} \bar{h}(x) - \bar{h}(xy_{n(x)}^{-1}) &\geq \bar{h}_{n(x)}(x) - \bar{h}_{n(x)}(xy_{n(x)}^{-1}) \geq h_{n(x)}(x) - h_{n(x)}(xy_{n(x)}^{-1}) \geq \\ &\geq n(x) g_{n(x)}(x) = n(x) \bar{g}(x) \text{ (thus } \bar{h}(x)/\bar{g}(x) \geq h(x) \rightarrow \infty \text{ as } x \rightarrow 0) \end{aligned}$$

and similarly  $\bar{h}(xy_{n(x)}) - \bar{h}(x) \geq n(x) \bar{g}(x)$ .

Therefore for  $x \leq a_N$ ,  $\bar{h}(x) - N\bar{g}(x) \leq \bar{h}(z) \leq \bar{h}(x) + N\bar{g}(x)$  implies  $z \in [xy_{n(x)}, xy_{n(x)}^{-1}]$ ,

$$\text{so if further } \lambda_{i+1} \leq z \leq \lambda_i, \text{ then } \lambda_i, \lambda_{i+1} \in [xy_{n(x)}^2, xy_{n(x)}^{-2}] \quad (*)$$

$$\lambda_{i+1} \leq xy_{n(x)}^{-1} \leq a_{n(x)} y_{n(x)}^{-1} \leq 2a_{n(x)}, \text{ and thus } \frac{\lambda_{i+1}}{\lambda_i} \geq y_{n(x)}.$$

By our above results,  $\bar{h}^{-1}(s)$  is well defined on  $R_+$ , and is continuous, strictly decreasing, and integrable.

Let  $\lambda(s)$  be the closest  $\lambda_i$  to  $\bar{h}^{-1}(s)$  (selected in a monotonic way), and  $L(s) = \bar{g}(\bar{h}^{-1}(s))$ . By (\*), if we let  $\bar{h}(x) = s$ , then for all  $|\theta| \leq n(x)$ ,  $\lambda(s + \theta L(s)) \in [xy_{n(x)}^2, xy_{n(x)}^{-2}]$ .

It follows now immediately that, whatever be  $A > 0$  and  $\delta > 0$ , conditions a) to e) are satisfied for  $s$  sufficiently large.

Finally, make  $\lambda(s)$  strictly decreasing — without changing  $L(s)$  — using the continuity of  $v_\lambda$  at each  $\lambda_i$  (lemma 4.2). This finishes the proof.

One might wonder whether the discontinuities in the function  $\lambda(s)$  are really necessary. Note however that our proof proved at the same time the following:

**Proposition 4.8.** The following conditions on  $v_\lambda$  are equivalent:

(4.9) For some  $A > 0$  and  $1 > \delta > 0$  there exist a function  $L(s)$  and a continuous function  $\lambda(s)$  (not necessarily monotonic) satisfying conditions a) to e);

- (4.10) There exist functions  $L(s)$  and  $\lambda(s)$  (continuous and strictly decreasing) that satisfy conditions a) – e) whatever be  $A > 0$ ,  $\delta > 0$ ;
- (4.11) There exists a sequence such that  $\lambda_i$ ,  $\liminf \lambda_{i+1} / \lambda_i > 0 = \lim \lambda_i$  and that  $\sum_i \Delta v[\lambda_{i+1}, \lambda_i] < \infty$ ;
- (4.12) For any sequence satisfying  $\limsup \lambda_{i+1} / \lambda_i < 1$ ,  $\lambda_i > 0$  one has  $\sum_i \Delta v[\lambda_{i+1}, \lambda_i] < \infty$ .

The proof is  $4.9 \Rightarrow 4.11 \Rightarrow 4.12 \Rightarrow 4.10 \Rightarrow 4.9$ . Each of those is essentially the same as the corresponding step in the proof of proposition 4.3 – except for the implication  $4.11 \Rightarrow 4.12$ , which is trivial: just note that there is a bounded number of terms in the sequence of 4.12 that can fall between any two successive terms of the sequence of 4.11 (first made decreasing). Also in the implication  $4.12 \Rightarrow 4.10$ , the sequence  $\lambda_i$  is not used; one works directly with the  $\Delta v$ 's, and uses 4.12 to guarantee that  $\sum \Delta v[y^{i+1}, y^i] < \infty$ .  $\lambda(s)$  is defined simply as  $\bar{h}^{-1}(s)$ .

## References

- Bewley, T., and E. Kohlberg: The Asymptotic Theory of Stochastic Games. *Math. Oper. Res.* 1, 1976, 197–208.
- Blackwell, D., and T.S. Ferguson: The Big Match. *Ann. Math. Statist.* 39, 1968, 159–163.
- Gillette, D.: Stochastic Games with Zero Stop Probabilities. *Contributions to the Theory of Games*, Vol. III (Annals of Mathematics Studies, No. 39). Princeton, N.J. 1957, 179–187.
- Kohlberg, E.: Repeated Games with Absorbing States. *The Annals of Statistics* 2, 1974, 724–738.
- Mertens, J.-F., and A. Neyman: Stochastic Games. Core Discussion Paper 8001, Université Catholique de Louvain, 1980.
- Monash, C.A.: Stochastic Games: The Minmax Theorem. Preprint, 1980.
- Shapley, L.: Stochastic Games. *Proc. Nat. Acad. Sci. USA* 39, 1953, 1095–1100.