

ARTICLE

Modal dependent type theory and dependent right adjoints

Lars Birkedal¹, Ranald Clouston², Bassel Manna^{3,*} , Rasmus Ejlers Møgelberg⁴, Andrew M. Pitts⁵ , and Bas Spitters¹

¹Department of Computer Science, Aarhus University, Denmark, ²Research School of Computer Science, Australian National University, Australia, ³eToroX Labs, Denmark, ⁴IT University of Copenhagen, Denmark and ⁵Department of Computer Science and Technology, University of Cambridge, UK

*Corresponding author. Email: bassel.manna@gmail.com

(Received 12 October 2018; revised 13 August 2019; accepted 18 October 2019)

Abstract

In recent years, we have seen several new models of dependent type theory extended with some form of modal necessity operator, including nominal type theory, guarded and clocked type theory and spatial and cohesive type theory. In this paper, we study *modal dependent type theory*: dependent type theory with an operator satisfying (a dependent version of) the K axiom of modal logic. We investigate both semantics and syntax. For the semantics, we introduce categories with families with a *dependent right adjoint* (CwDRA) and show that the examples above can be presented as such. Indeed, we show that any category with finite limits and an adjunction of endofunctors give rise to a CwDRA via the local universe construction. For the syntax, we introduce a dependently typed extension of Fitch-style modal λ -calculus, show that it can be interpreted in any CwDRA, and build a term model. We extend the syntax and semantics with universes.

Keywords: Dependent type theory; modal logic; category theory

1. Introduction

Dependent types are a powerful technology for both programming and formal proof. In recent years, we have seen several new models of dependent type theory extended with a type former resembling modal necessity,¹ such as nominal type theory (Pitts et al. 2015), guarded (Birkedal et al. 2012, 2018 Bizjak et al. 2016; Bizjak and Møgelberg 2018) and clocked type theory (cloTT) (Manna and Møgelberg 2018), and spatial and cohesive type theory (Shulman 2018). These examples all satisfy the K axiom of modal logic

$$\Box(A \rightarrow B) \rightarrow \Box A \rightarrow \Box B \quad (1)$$

but are not all (co)monads, the more extensively studied construction in the context of dependent type theory (Krishnaswami et al. 2015; de Paiva and Ritter 2016; Shulman 2018; Vákár 2017). Motivated in part by these examples, in this paper, we study *modal dependent type theory*: dependent type theory with an operator satisfying (a dependent generalisation of) the K axiom² of modal logic. We investigate both semantics and syntax.

For the semantics, we introduce categories with families with a *dependent right adjoint* (CwDRA) and show that this dependent right adjoint models the modality in the examples mentioned above. Indeed, we show that any finite limit category with an adjunction of endofunctors³

gives rise to a CwDRA via the local universe construction (Lumsdaine and Warren 2015). In particular, by applying the local universe construction to a locally cartesian closed category with an adjunction of endofunctors, we get a model of modal dependent type theory with Π - and Σ -types.

For the syntax, we adapt the simply typed Fitch-style modal λ -calculus introduced by Borghuis (1994) and Martini and Masini (1996), inspired by Fitch's proof theory for modal logic (Fitch 1952). In such, a calculus \Box is introduced by 'shutting' a strict subordinate proof and eliminating by 'opening' one. For example, the K axiom (1) is inhabited by the term

$$\lambda f. \lambda x. \text{shut}((\text{open } f)(\text{open } x)) \quad (2)$$

The nesting of subordinate proofs can be tracked in sequent style by a special symbol in the context which we call a *lock*, and write \blacksquare ; the open lock symbol is intended to suggest we have access to the contents of a box. Following Clouston (2018), the lock can be understood as an operation on contexts *left adjoint* to \Box ; hence, Fitch-style modal λ -calculus has a model in any cartesian closed category equipped with an adjunction of endofunctors. Here, we show, in work inspired by cloTT (Bahr *et al.* 2017), that Fitch-style λ -calculus lifts with a minimum of difficulty to dependent types. In particular, the term (2), where f is a dependent function, has type

$$\Box(\Pi y : A. B) \rightarrow \Pi x : \Box A. \Box B[\text{open } x/y]$$

This dependent version of the K axiom, not obviously expressible without the *open* construct of a Fitch-style calculus, allows modalised functions to be applied to modalised data even in the dependent case. This capability is known to be essential in at least one example, namely, proofs about guarded recursion (Bizjak *et al.* 2016)⁴. We show that our calculus can be soundly interpreted in any CwDRA and construct a term model.

We also extend the syntax and semantics of modal dependent type theory with universes. Here, we restrict attention to models based on (pre)sheaves, for which Coquand has proposed a particularly simple formulation of universes (Coquand 2012). We show how to extend Coquand's notion of a category with universes with dependent right adjoint (CwUDRA) and observe that a construction encoding the modality on the universe, introduced for guarded type theory by Bizjak *et al.* (2016), in fact arises for more general reasons.

Another motivation for the present work is that it can be understood as providing a notion of a *dependent* adjunction between endofunctors. An ordinary adjunction $L \dashv R$ on a category \mathbf{C} is a natural bijective correspondence $\mathbf{C}(LA, B) \cong \mathbf{C}(A, RB)$. With dependent types, one might consider *dependent* functions from LA to B , where B may depend on LA , and similarly from A to RB . Our notion of CwDRA then defines what it means to have an adjoint correspondence in this dependent case. Our Fitch-style modal dependent type theory can therefore also be understood as a term language for dependent adjoints.

Outline. We introduce CwDRAs in Section 2 and present the syntax of modal dependent type theory in Section 3. In Section 4, we show how to construct a CwDRA from an adjunction on a category with finite limits. In Section 5, we show how various models in the literature can be presented as CwDRAs. The extension with universes is defined in Section 6. We end with a discussion of related and future work in Section 7.

2. Categorical Semantics of Modal Dependent Type Theory

The notion of *category with families* (CwF) (Dybjer 1995; Hofmann 1997) provides a semantics for the development of dependent type theory that elides some difficult aspects of syntax, such as variable binding, as well as the coherence problems of simpler notions of model. It can be connected to syntax by a soundness argument and term model construction, and to more mathematical models via 'strictification' constructions (Hofmann 1994; Lumsdaine and Warren 2015). In this section, we extend this notion to introduce CwDRA. We first recall the standard definition:

Definition 1. (CwF). A CwF is specified by:

1. A category \mathbf{C} with a terminal object \top . Given objects $\Gamma, \Delta \in \mathbf{C}$, write $\mathbf{C}(\Delta, \Gamma)$ for the set of morphisms from Δ to Γ in \mathbf{C} . The identity morphism on Γ is just written id with Γ implicit. The composition of $\gamma \in \mathbf{C}(\Delta, \Gamma)$ with $\delta \in \mathbf{C}(\Phi, \Delta)$ is written as $\gamma \circ \delta$.
2. For each object $\Gamma \in \mathbf{C}$, a set $\mathbf{C}(\Gamma)$ of families over Γ .
3. For each object $\Gamma \in \mathbf{C}$ and family $A \in \mathbf{C}(\Gamma)$, a set $\mathbf{C}(\Gamma \vdash A)$ of elements of the family A over Γ .
4. For each morphism $\gamma \in \mathbf{C}(\Delta, \Gamma)$, re-indexing functions $A \in \mathbf{C}(\Gamma) \mapsto A[\gamma] \in \mathbf{C}(\Delta)$ and $a \in \mathbf{C}(\Gamma \vdash A) \mapsto a[\gamma] \in \mathbf{C}(\Delta \vdash A[\gamma])$, satisfying $A[\text{id}] = A$, $A[\gamma \circ \delta] = A[\gamma][\delta]$, $a[\text{id}] = a$ and $a[\gamma \circ \delta] = a[\gamma][\delta]$.
5. For each object $\Gamma \in \mathbf{C}$ and family $A \in \mathbf{C}(\Gamma)$, a comprehension object $\Gamma.A \in \mathbf{C}$ equipped with a projection morphism $p_A \in \mathbf{C}(\Gamma.A, \Gamma)$, a generic element $q_A \in \mathbf{C}(\Gamma.A \vdash A[p_A])$ and a pairing operation mapping $\gamma \in \mathbf{C}(\Delta, \Gamma)$ and $a \in \mathbf{C}(\Delta \vdash A[\gamma])$ to $(\gamma, a) \in \mathbf{C}(\Delta, \Gamma.A)$ satisfying $p_A \circ (\gamma, a) = \gamma$, $q_A[(\gamma, a)] = a$, $(\gamma, a) \circ \delta = (\gamma \circ \delta, a[\delta])$ and $(p_A, q_A) = \text{id}$.

A dependent right adjoint then extends the definition of CwF with a functor on contexts L and an operation on families R , intuitively understood to be left and right adjoints:

Definition 2. (Category with a dependent right adjoint). A CwDRA is a CwF \mathbf{C} equipped with the following extra structure:

1. An endofunctor $L : \mathbf{C} \rightarrow \mathbf{C}$ on the underlying category of the CwF.
2. For each object $\Gamma \in \mathbf{C}$ and family $A \in \mathbf{C}(L\Gamma)$, a family $R_\Gamma A \in \mathbf{C}(\Gamma)$, stable under re-indexing in the sense that for all $\gamma \in \mathbf{C}(\Delta, \Gamma)$, we have

$$(R_\Gamma A)[\gamma] = R_\Delta (A[L\gamma]) \in \mathbf{C}(\Delta) \quad (3)$$

3. For each object $\Gamma \in \mathbf{C}$ and family $A \in \mathbf{C}(L\Gamma)$, a bijection

$$\mathbf{C}(L\Gamma \vdash A) \cong \mathbf{C}(\Gamma \vdash R_\Gamma A) \quad (4)$$

We write the effect of this bijection on $a \in \mathbf{C}(L\Gamma \vdash A)$ as $\bar{a} \in \mathbf{C}(\Gamma \vdash R_\Gamma A)$ and write the effect of its inverse on $b \in \mathbf{C}(\Gamma \vdash R_\Gamma A)$ also as $\bar{\bar{b}} \in \mathbf{C}(L\Gamma \vdash A)$. Thus

$$\bar{\bar{a}} = a \quad (a \in \mathbf{C}(L\Gamma \vdash A)) \quad (5)$$

$$\bar{\bar{b}} = b \quad (b \in \mathbf{C}(\Gamma \vdash R_\Gamma A)) \quad (6)$$

The bijection is required to be stable under re-indexing in the sense that for all $\gamma \in \mathbf{C}(\Delta, \Gamma)$, we have

$$\bar{a}[\gamma] = \overline{\bar{a}[L\gamma]} \quad (7)$$

Note that equation (7) is well typed by (3). Equation (7) also implies that the opposite direction of the isomorphism (4) is natural, that is, that the equation

$$\bar{\bar{b}}[L\gamma] = \overline{\bar{b}[\gamma]} \quad (8)$$

also holds, since $\overline{\bar{b}[\gamma]} = \overline{\bar{\bar{b}}[L\gamma]} = \overline{\bar{\bar{b}}[L\gamma]} = \bar{\bar{b}}[L\gamma]$.

3. Syntax of Modal Dependent Type Theory

In this section, we extend Fitch-style modal λ -calculus (Borghuis 1994) to dependent types and connect this to the notion of CwDRA via a soundness proof and term model construction.

We define our dependent types broadly in the style of ECC (Luo 1989), as this is close to the implementation of some proof assistants (Norell 2007).

We define the raw syntax of contexts, types, and terms as follows:

$$\begin{aligned}\Gamma &\triangleq \diamond \mid \Gamma, x : A \mid \Gamma, \text{lock} \\ A &\triangleq \Pi x : A. B \mid \Box A \\ t &\triangleq x \mid \lambda x : A. t \mid t t \mid \text{shut } t \mid \text{open } t\end{aligned}$$

We omit the leftmost ‘ \diamond ’, where the context is non-empty. We will usually omit the type annotation on the λ for brevity. Π -types are included in the grammar as an example to show that standard type formers can be defined as usual, without reference to the locks in the context. One could similarly add an empty type, unit type, booleans, Σ -types, W -types, universes (of which more in Section 6), and so forth.

Judgements have forms

$$\begin{array}{ll}\Gamma \vdash & \text{‘}\Gamma \text{ is a well-formed context’} \\ \Gamma \vdash A & \text{‘}A \text{ is a well-formed type in context } \Gamma\text{’} \\ \Gamma \vdash A = B & \text{‘}A \text{ and } B \text{ are equal types in context } \Gamma\text{’} \\ \Gamma \vdash t : A & \text{‘}t \text{ is a term with type } A \text{ in context } \Gamma\text{’} \\ \Gamma \vdash t = u : A & \text{‘}t \text{ and } u \text{ are equal terms with type } A \text{ in context } \Gamma\text{’}\end{array}$$

Figure 1 presents the typing rules of the calculus. The syntactic results below follow easily by induction on these rules. We remark only that exchange of variables with locks, and weakening of locks, are not admissible, and that the (lock-free) weakening Γ' in the open rule is essential to prove variable weakening.

Lemma 3. *Let \mathcal{J} range over the possible strings to the right of a turnstile in a judgement.*

1. If $\Gamma, x : A, y : B, \Gamma' \vdash \mathcal{J}$ and x is not free in B , then $\Gamma, y : B, x : A, \Gamma' \vdash \mathcal{J}$;
2. If $\Gamma, \Gamma' \vdash \mathcal{J}$, and $\Gamma \vdash A$, and x is a fresh variable, then $\Gamma, x : A, \Gamma' \vdash \mathcal{J}$;
3. If $\Gamma, x : A, \Gamma' \vdash \mathcal{J}$ and $\Gamma \vdash u : A$, then $\Gamma, \Gamma'[u/x] \vdash \mathcal{J}[u/x]$;
4. If $\Gamma \vdash t : A$ then $\Gamma \vdash A$;
5. If $\Gamma \vdash t = u : A$ then $\Gamma \vdash t : A$ and $\Gamma \vdash u : A$.

3.1 Sound interpretation in CwDRAs

In this section, we show that the calculus of Figure 1 can be soundly interpreted in any CwDRA. We wish to give meaning to contexts, types and terms, but (via the type conversion rule) these can have multiple derivations, so it is not possible to work by induction on the formation rules. Instead, following, for example, Hofmann (1997), we define a partial map from *raw syntax* to semantics by induction on the grammar and then prove that this map is defined for well-formed syntax. By ‘raw syntax’, we mean contexts, types accompanied by a context, and terms accompanied by context and type, defined via the grammar. The *size* of a type or term is the number of connectives and variables used to define it, and the size of a context is the sum of the sizes of its types.

Well-defined contexts Γ will be interpreted as objects $\llbracket \Gamma \rrbracket$ in \mathbf{C} , types in context $\Gamma \vdash A$ as families in $\mathbf{C}(\llbracket \Gamma \rrbracket)$, and typed terms in context $\Gamma \vdash t : A$ as elements in $\mathbf{C}(\llbracket \Gamma \rrbracket \vdash \llbracket \Gamma \vdash A \rrbracket)$. Where there is no confusion, we write $\llbracket \Gamma \vdash A \rrbracket$ as $\llbracket A \rrbracket$ and $\llbracket \Gamma \vdash t : A \rrbracket$ as $\llbracket \Gamma \vdash t \rrbracket$ or $\llbracket t \rrbracket$.

The partial interpretation of raw syntax is as follows, following the convention that ill-formed expressions (e.g., where a subexpression is undefined) are undefined. We omit the details for Π -types and other standard constructions, which are as usual.

$$\begin{aligned}- \llbracket \diamond \rrbracket &= \top; \\ - \llbracket \Gamma, x : A \rrbracket &= \llbracket \Gamma \rrbracket. \llbracket A \rrbracket;\end{aligned}$$

Context formation rules:

$$\frac{}{\diamond \vdash} \quad \frac{\Gamma \vdash \quad \Gamma \vdash A}{\Gamma, x : A \vdash} x \notin \Gamma \quad \frac{\Gamma \vdash}{\Gamma, \blacksquare \vdash} \quad \frac{\Gamma, x : A, y : B, \Gamma' \vdash}{\Gamma, y : B, x : A, \Gamma' \vdash} x \text{ NOT FREE IN } B$$

Type formation rules:

$$\frac{\Gamma \vdash A \quad \Gamma, x : A \vdash B}{\Gamma \vdash \Pi x : A. B} \quad \frac{\Gamma, \blacksquare \vdash A}{\Gamma \vdash \Box A}$$

Type equality rules are as standard, asserting equivalence, and congruence with respect to all type formers.

Term formation rules:

$$\frac{\Gamma \vdash t : A \quad \Gamma \vdash A = B}{\Gamma \vdash t : B} \quad \frac{\Gamma, x : A, \Gamma' \vdash}{\Gamma, x : A, \Gamma' \vdash x : A} \blacksquare \notin \Gamma' \quad \frac{\Gamma, x : A \vdash t : B}{\Gamma \vdash \lambda x. t : \Pi x : A. B}$$

$$\frac{\Gamma \vdash t : \Pi x : A. B \quad \Gamma \vdash u : A}{\Gamma \vdash t u : B[u/x]} \quad \frac{\Gamma, \blacksquare \vdash t : A}{\Gamma \vdash \text{shut } t : \Box A} \quad \frac{\Gamma \vdash t : \Box A \quad \Gamma, \blacksquare, \Gamma' \vdash}{\Gamma, \blacksquare, \Gamma' \vdash \text{open } t : A} \blacksquare \notin \Gamma'$$

Term equality rules, omitting equivalence and congruence:

$$\frac{\Gamma \vdash (\lambda x. t) u : A}{\Gamma \vdash (\lambda x. t) u = t[u/x] : A} \quad \frac{\Gamma \vdash \text{open shut } t : A}{\Gamma \vdash \text{open shut } t = t : A} \quad \frac{\Gamma \vdash t : \Pi x : A. B}{\Gamma \vdash t = \lambda x. t x : \Pi x : A. B} x \notin \Gamma$$

$$\frac{\Gamma \vdash t : \Box A}{\Gamma \vdash t = \text{shut open } t : \Box A}$$

Figure 1. Typing rules for a dependent Fitch-style modal λ -calculus.

- $\llbracket \Gamma, \blacksquare \rrbracket = \mathbb{L}[\llbracket \Gamma \rrbracket]$;
- $\llbracket \Gamma \vdash \Box A \rrbracket = R_{\llbracket \Gamma \rrbracket}(\llbracket A \rrbracket)$;
- $\llbracket \Gamma, x : A, x_1 : A_1, \dots, x_n : A_n \vdash x : A \rrbracket = q_{\llbracket A \rrbracket} [p_{\llbracket A_1 \rrbracket} \circ \dots \circ p_{\llbracket A_n \rrbracket}]$;
- $\llbracket \Gamma \vdash \text{shut } t : \Box A \rrbracket = \llbracket t \rrbracket$;
- $\llbracket \Gamma, \blacksquare, x_1 : A_1, \dots, x_n : A_n \vdash \text{open } t : A \rrbracket = \llbracket t \rrbracket [p_{\llbracket A_1 \rrbracket} \circ \dots \circ p_{\llbracket A_n \rrbracket}]$.

In Figure 2, we define expressions $P(\Gamma; A; \Gamma')$, $E(\Gamma; A; B; \Gamma')$, and $S(\Gamma; A; \Gamma'; t)$ that, where defined, define morphisms in \mathbf{C} corresponding, respectively, to weakening, exchange, and substitution in contexts.

Lemma 4. Suppose $\llbracket \Gamma, \Gamma' \rrbracket$ and $\llbracket \Gamma, x : A, \Gamma' \rrbracket$ are defined. Then, the following properties hold:

1. $\llbracket \Gamma, x : A, \Gamma' \vdash X \rrbracket \simeq \llbracket \Gamma, \Gamma' \vdash X \rrbracket [P(\Gamma; A; \Gamma')]$, where \simeq is the Kleene equality, and X is a type or typed term;
2. $P(\Gamma; A; \Gamma')$ is a well-defined morphism from $\llbracket \Gamma, x : A, \Gamma' \rrbracket$ to $\llbracket \Gamma, \Gamma' \rrbracket$.

Proof. The proof proceeds by mutual induction on the size of Γ' (for statement 2) and the size of Γ' plus the size of X (for statement 1). We present only the cases particular to \Box .

$$\begin{aligned}
P(\Gamma; A; \diamond) &= p_{[A]} \\
P(\Gamma; A; \Gamma', y : B) &= (P(\Gamma; A; \Gamma') \circ p_{[\Gamma, x:A, \Gamma' \vdash B]}, q_{[\Gamma, x:A, \Gamma' \vdash B]}) \\
P(\Gamma; A; \Gamma', \blacksquare) &= L P(\Gamma; A; \Gamma') \\
E(\Gamma; A; B; \diamond) &= ((p_{[\Gamma \vdash B]} \circ p_{[\Gamma, y:B \vdash A]}, q_{[\Gamma, y:B \vdash A]}), q_{[\Gamma \vdash B]} [p_{[\Gamma, y:B \vdash A]}]) \\
E(\Gamma; A; B; \Gamma', z : C) &= (E(\Gamma; A; B; \Gamma') \circ p_{[\Gamma, y:B, x:A, \Gamma' \vdash C]}, q_{[\Gamma, y:B, x:A, \Gamma' \vdash C]}) \\
E(\Gamma; A; B; \Gamma', \blacksquare) &= L E(\Gamma; A; B; \Gamma') \\
S(\Gamma; A; \diamond; t) &= (id, [t]) \\
S(\Gamma; A; \Gamma', y : B; t) &= (S(\Gamma; A; \Gamma'; t) \circ p_{[\Gamma, \Gamma' [t/x] \vdash B[t/x]]}, q_{[\Gamma, \Gamma' [t/x] \vdash B[t/x]]}) \\
S(\Gamma; A; \Gamma', \blacksquare; t) &= L S(\Gamma; A; \Gamma'; t)
\end{aligned}$$

Figure 2. Morphisms in \mathbf{C} corresponding to weakening, exchange and substitution.

We start with statement 1. We use the mutual induction with statement 2 at the smaller size of Γ' alone to ensure that $P(\Gamma; A; \Gamma')$ is well formed with the correct domain and codomain and then proceed by induction on the construction of X .

The \square case follows because

$$\begin{aligned}
[\Gamma, x : A, \Gamma' \vdash \square B] &\simeq R_{[\Gamma, x:A, \Gamma']} [\Gamma, x : A, \Gamma', \blacksquare \vdash B] \\
&\simeq R_{[\Gamma, x:A, \Gamma']} ([\Gamma, \Gamma', \blacksquare \vdash B] [P(\Gamma; A; \Gamma', \blacksquare)]) \quad (\text{induction}) \\
&\simeq R_{[\Gamma, x:A, \Gamma']} ([\Gamma, \Gamma', \blacksquare \vdash B] [LP(\Gamma; A; \Gamma')]) \\
&\simeq (R_{[\Gamma, \Gamma']} [\Gamma, \Gamma', \blacksquare \vdash B]) [P(\Gamma; A; \Gamma')] \quad (3) \\
&= [\Gamma, \Gamma' \vdash \square B] [P(\Gamma; A; \Gamma')]
\end{aligned}$$

The shut case follows immediately from (7) and induction. For open, the case where the deleted variable x is to the right of the lock follows by Definition 1 part 5. Suppose, instead it is to the left. Then,

$$\begin{aligned}
&[\Gamma, \Gamma', \blacksquare, y_1 : B_1, \dots, y_n : B_n \vdash \text{open } t] [P(\Gamma; A; \Gamma', \blacksquare, y_1 : B_1, \dots, y_n : B_n)] \\
&\simeq \overline{[t]} [p_{[B_1]} \circ \dots \circ p_{[B_n]} \circ P(\Gamma; A; \Gamma', \blacksquare, y_1 : B_1, \dots, y_n : B_n)] \\
&\simeq \overline{[t]} [P(\Gamma; A; \Gamma', \blacksquare) \circ p_{[B_1]} \circ \dots \circ p_{[B_n]}] \quad (\text{Definition 1 part 5}) \\
&\simeq \overline{[t]} [LP(\Gamma; A; \Gamma')] [p_{[B_1]} \circ \dots \circ p_{[B_n]}] \\
&\simeq \overline{[t]} [P(\Gamma; A; \Gamma')] [p_{[B_1]} \circ \dots \circ p_{[B_n]}] \quad (8) \\
&\simeq \overline{[\Gamma, x : A, \Gamma' \vdash t]} [p_{[B_1]} \circ \dots \circ p_{[B_n]}] \quad (\text{induction}) \\
&\simeq [\Gamma, x : A, \Gamma', \blacksquare, y_1 : B_1, \dots, y_n : B_n \vdash \text{open } t]
\end{aligned}$$

For statement 2, the lock case holds immediately by application of the functor L . Other cases follow as standard; for example, the base case holds because $p_{[A]}$ is indeed a morphism. \square

Lemma 5. Suppose $[\Gamma, x : A, y : B, \Gamma']$ and $[\Gamma \vdash B]$ are defined. Then, the following properties hold:

1. $[\Gamma, y : B, x : A, \Gamma' \vdash X] \simeq [\Gamma, x : A, y : B, \Gamma' \vdash X] [E(\Gamma; A; B; \Gamma')]$, where X is a type or typed term;
2. $E(\Gamma; A; B; \Gamma')$ is a well-defined morphism from $[\Gamma, y : B, x : A, \Gamma']$ to $[\Gamma, x : A, y : B, \Gamma']$;

Proof. The base case of statement 1 uses Lemma 4; the proof otherwise follows just as with Lemma 4. \square

Lemma 6. Suppose $\llbracket \Gamma \vdash t : A \rrbracket$ and $\llbracket \Gamma, x : A, \Gamma' \rrbracket$ are defined. Then, the following properties hold:

1. $\llbracket \Gamma, \Gamma'[t/x] \vdash X[t/x] \rrbracket \simeq \llbracket \Gamma, x : A, \Gamma' \vdash X \rrbracket [S(\Gamma; A; \Gamma'; t)]$, where X is a type or typed term;
2. $S(\Gamma; A; \Gamma'; t)$ is a well-defined morphism from $\llbracket \Gamma, \Gamma'[t/x] \rrbracket$ to $\llbracket \Gamma, x : A, \Gamma' \rrbracket$.

Proof. As with Lemma 4. \square

Theorem 7. (Soundness). Where a context, type or term is well formed, its denotation is well defined, and all types and terms identified by equations have the same denotation.

Proof. Most cases follow as usual, using Lemmas 4, 5 and 6 as needed. The well definedness of the formation rules for \square is straightforward, so we present only the equations for \square :

Starting with $\Gamma, \blacksquare \vdash t : A$, we have $\Gamma, \blacksquare, x_1 : A_1, \dots, x_n : A_n \vdash \text{open shut } t : A$ and wish to prove that its denotation is equal to that of t (with the weakening x_1, \dots, x_n). Then $\llbracket \text{open shut } t \rrbracket = \overline{\llbracket t \rrbracket} [p_{[A_1]} \circ \dots \circ p_{[A_n]}] = \llbracket t \rrbracket [p_{[A_1]} \circ \dots \circ p_{[A_n]}]$, which is the weakening of t by Lemma 4.

The equality of $\llbracket \text{shut open } t \rrbracket$ and $\llbracket t \rrbracket$ is straightforward. \square

3.2 Term model

We now develop as our first example of a CwDRA, a term model built from the syntax of our calculus. The objects of this category are contexts modulo equality, which is defined pointwise via type equality. We define an arrow $\Delta \rightarrow \Gamma$ as a sequence of substitutions of an equivalence class of terms for each variable in Γ :

- the empty sequence is an arrow $\Delta \rightarrow \cdot$;
- Given $f : \Delta \rightarrow \Gamma$, type $\Gamma \vdash A$ and term $\Delta \vdash t : Af$, where Af is the result of applying the substitutions f to A , then $[t/x] \circ f$ modulo equality on t is an arrow $\Delta \rightarrow \Gamma, x : A$;
- Given $f : \Delta \rightarrow \Gamma$ and a well-formed context $\Delta, \blacksquare, \Delta'$ with no locks in Δ' , then f is also an arrow $\Delta, \blacksquare, \Delta' \rightarrow \Gamma, \blacksquare$.

We usually refer to the equivalence classes in arrows via representatives. Note that substitution respects these equivalence classes because of the congruence rules.

We next prove that this defines a category. Identity arrows are easily constructed:

Lemma 8. If $f : \Delta \rightarrow \Gamma$, then $f : \Delta, x : A \rightarrow \Gamma$.

Proof. By induction on the construction on f . The base case is trivial.

Given $f : \Delta \rightarrow \Gamma$ and $\Delta \vdash t : Bf$, by induction, we have $f : \Delta, x : A \rightarrow \Gamma$ and by variable weakening, we have $\Delta, x : A \vdash t : Bf$ as required.

Supposing we have $f : \Delta \rightarrow \Gamma$ yielding $f : \Delta, \blacksquare, \Delta' \rightarrow \Gamma$, we could similarly get $f : \Delta, \blacksquare, \Delta', x : A \rightarrow \Gamma$. \square

The identity on Γ simply replaces all variables by themselves.

Lemma 9. The identity on each Γ is well defined as an arrow.

Proof. By induction on Γ . The identity on \cdot is the empty sequence of substitutions. Given $id : \Gamma \rightarrow \Gamma$, we have $id : \Gamma, x : A \rightarrow \Gamma$ by Lemma 8, and $\Gamma, x : A \vdash x : A$ as required. $id : \Gamma \rightarrow \Gamma$ immediately yields $id : \Gamma, \blacksquare \rightarrow \Gamma, \blacksquare$. \square

The composition case is slightly more interesting:

Lemma 10. *Given $\Gamma, \Gamma' \vdash \mathcal{J}$ and $f : \Delta \rightarrow \Gamma$, we have $\Delta, \Gamma' f \vdash \mathcal{J}f$.*

Proof. By induction on the construction on f . The base case requires that $\Gamma' \vdash \mathcal{J}$ implies $\Delta, \Gamma' \vdash \mathcal{J}$; this *left weakening* property is easily proved by induction on the typing rules.

Given $f : \Delta \rightarrow \Gamma$, $\Delta \vdash t : Af$ and $\Gamma, x : A, \Gamma' \vdash \mathcal{J}$, by induction $\Delta, x : Af, \Gamma' f \vdash \mathcal{J}f$. Then by Lemma 3 part 3, we have $\Delta, (\Gamma' f)[t/x] \vdash (\mathcal{J}f)[t/x]$ as required. The lock case is trivial. \square

The composition of $f : \Delta \rightarrow \Delta'$ and $g : \Delta' \rightarrow \Gamma$ involves replacing each $[t/x]$ in g with $[tf/x]$.

Lemma 11. *The composition of two arrows $f : \Delta \rightarrow \Delta'$ and $g : \Delta' \rightarrow \Gamma$ is a well-defined arrow.*

Proof. By induction on the definition of g . The base case is trivial, and extension by a new substitution follows via Lemma 10.

Now suppose we have $g : \Delta' \rightarrow \Gamma$ yielding $g : \Delta', \blacksquare, \Delta'' \rightarrow \Gamma, \blacksquare$. Now if we have $f : \Delta \rightarrow \Delta', \blacksquare, \Delta''$, this must have arisen via some $f' : \Delta_0 \rightarrow \Delta'$ generating $f' : \Delta_0, \blacksquare, \Delta_1 \rightarrow \Delta', \blacksquare$, where $\Delta = \Delta_0, \blacksquare, \Delta_1$. By induction, we have well-defined $g \circ f' : \Delta_0 \rightarrow \Gamma$. Hence, $g \circ f' : \Delta \rightarrow \Gamma, \blacksquare$. But $g \circ f' = g \circ f$ because the variables of Δ'' do not appear in g . \square

Checking the category axioms is straightforward. The category definitions then extend to a CwF in the usual way: the terminal object is \diamond , the families over Γ are the types modulo equivalence well defined in context Γ , the elements of any such type are the terms modulo equivalence, re-indexing is substitution, comprehension corresponds to extending a context with a new variable, the projection morphism is the replacement of variables by themselves, and the generic element is given by the variable rule.

Moving to the definition of a CwDRA, the endofunctor L acts by mapping $\Gamma \mapsto \Gamma, \blacksquare$, and does not change arrows. The family $R_\Gamma A$ is the type $\Gamma \vdash \square A$, which is stable under re-indexing by Lemma 3 part 3. The bijections between families are supplied by the shut and open rules, with all equations following from the definitional equalities.

We do not attempt to prove that the term model is the *initial* CwDRA; such a result for dependent type theories appears to require syntax be written in a more verbose style than is appropriate for a paper introducing a new type theory (Castellan 2014). Nonetheless, our type theory and notion of model are close enough that we conjecture that such a development is possible.

4. A General Construction of CwDRAs

In this section, we show how to construct a CwDRA from an adjunction of endofunctors on a category with finite limits. We will refer to categories with finite limits more briefly as *cartesian* categories. We will use this construction in Section 5 to prove that the examples mentioned in the introduction can indeed be presented as CwDRAs. Our construction is an extension of the local universe construction (Lumsdaine and Warren 2015), which maps cartesian categories to categories with families, and locally cartesian closed categories to categories with families with Π - and Σ -types. The local universe construction is one of the known solutions to the problem of constructing a strict model of type theory out of a locally cartesian closed category (see (Hofmann 1994, 1997; Kapulkin and Lumsdaine 2018; Lumsdaine and Warren 2015) for discussions of alternative approaches to ‘strictification’).

We first recall the local universe construction. Since it can be traced back to Giraud's work on fibred categories (Giraud 1965), we refer to it as the Giraud CwF associated to a cartesian category.

Definition 12. Let \mathbf{C} be a cartesian category. The **Giraud CwF of \mathbf{C}** ($\mathcal{G}\mathbf{C}$) is the CwF whose underlying category is \mathbf{C} , and where a family $A \in \mathcal{G}\mathbf{C}(\Gamma)$ is a pair of morphisms

$$\begin{array}{ccc} & E & \\ & \downarrow v & \\ \Gamma & \xrightarrow{u} & U \end{array} \quad (9)$$

and an element of $\mathcal{G}\mathbf{C}(\Gamma \vdash A)$, for $A = (u, v) \in \mathcal{G}\mathbf{C}(\Gamma)$, is a map $a : \Gamma \rightarrow E$ such that $v \circ a = u$. Re-indexing of $A = (u, v) \in \mathcal{G}\mathbf{C}(\Gamma)$ and $a \in \mathcal{G}\mathbf{C}(\Gamma \vdash A)$ along $\gamma \in \mathbf{C}(\Delta, \Gamma)$ are given by

$$A[\gamma] \triangleq (u \circ \gamma, v) \in \mathcal{G}\mathbf{C}(\Delta) \quad (10)$$

$$a[\gamma] \triangleq a \circ \gamma \in \mathcal{G}\mathbf{C}(\Delta \vdash A[\gamma]) \quad (11)$$

The comprehension $\Gamma.A \in \mathbf{C}$, for $A = (u, v) \in \mathcal{G}\mathbf{C}(\Gamma)$, is given by the pullback of diagram (9),

$$\begin{array}{ccc} \Gamma.A & \xrightarrow{q_A} & E \\ p_A \downarrow & & \downarrow v \\ \Gamma & \xrightarrow{u} & U \end{array}$$

with projection morphism p_A and generic element q_A as indicated in the diagram. Note that q_A is an element of $A[p_A] = (u \circ p_A, v)$ as required by commutativity of the pullback square. The pairing operation is obtained from the universal property of pullbacks.

Note that the local universe construction does indeed yield a CwF; in particular, re-indexing in $\mathcal{G}\mathbf{C}$ is strict as required, simply because re-indexing is given by composition.

Remark 13. The name 'local universe' derives from the similarity to Voevodsky's use of a (global) universe U to construct strict models of type theory (Voevodsky 2014; Kapulkin and Lumsdaine 2018) in which types in a context Γ are modelled as morphisms $\Gamma \rightarrow U$. In the local universe construction, the universe varies from type to type.

In fact, the local universe construction is functorial; a precise statement requires a novel notion of CwF morphism:

Definition 14. A weak CwF morphism R between CwFs consists of a functor $R : \mathbf{C} \rightarrow \mathbf{D}$ between the underlying categories preserving the terminal object, an operation on families mapping $A \in \mathbf{C}(\Gamma)$ to a family $RA \in \mathbf{D}(R\Gamma)$ and an operation on elements mapping $a \in \mathbf{C}(\Gamma \vdash A)$ to an element $Ra \in \mathbf{D}(R\Gamma \vdash RA)$, such that

1. The functor $R : \mathbf{C} \rightarrow \mathbf{D}$ preserves terminal objects (up to isomorphism)
2. The operations on families and elements commute with re-indexing in the sense that $RA[R\gamma] = R(A[\gamma])$ and $Rt[R\gamma] = R(t[\gamma])$.
3. The maps $(Rp_A, Rq_A) : R(\Gamma.A) \rightarrow R\Gamma.RA$ are isomorphisms for all Γ and A . We write $v_{\Gamma,A}$ for the inverse.

We note the following equalities as consequences of the axioms above.

$$R(p_A) \circ v_{\Gamma, A} = p_{RA} \quad (12)$$

$$R(q_A)[v_{\Gamma, A}] = q_{RA} \quad (13)$$

$$v_{\Gamma, A} \circ (R\gamma, Ra) = R(\gamma, a) \quad (14)$$

For example, the last of these is proved by post-composing with the inverse of $v_{\Gamma, A}$ and noting

$$\begin{aligned} (R p_A, R q_A) \circ R(\gamma, a) &= (R p_A \circ R(\gamma, a), R q_A[R(\gamma, a)]) \\ &= (R(p_A \circ (\gamma, a)), R(q_A[(\gamma, a)])) \\ &= (R\gamma, Ra) \end{aligned}$$

Note that a weak CwF morphism preserves comprehension and the terminal object only up to isomorphism instead of on the nose, as required by the stricter notion of morphism of Dybjer (1995, Definition 2). Weak CwF morphisms sit between strict CwF morphisms and pseudo-CwF morphisms (Castellan et al. 2017). The latter allow substitution to be preserved only up to isomorphism satisfying a number of coherence conditions. Since weak CwF morphisms preserve substitution on the nose, these are not needed here.

Theorem 15. \mathcal{G} extends to a functor from the category of cartesian categories and finite limit preserving functors, to the category of CwFs with weak morphisms.

Proof. Let $R : \mathbf{C} \rightarrow \mathbf{D}$ be a finite limit preserving functor. For each $\Gamma \in \mathbf{C}$ and $A = (u, v) \in \mathcal{G}\mathbf{C}(\Gamma)$, we simply let $RA \triangleq (Ru, Rv)$. Likewise, for an element $a \in \mathcal{G}\mathbf{C}(\Gamma \vdash A)$, we let Ra be the action of R on the morphism a . Finally, since comprehension is defined by pullback and R preserves pullbacks up to isomorphism, we obtain the required $v_{\Gamma, A}$. \square

We now embark on showing that if we apply the local universe construction to a cartesian category \mathbf{C} with a pair of adjoint endofunctors, then the resulting CwF $\mathcal{G}\mathbf{C}$ is in fact a CwDRA (Theorem 19). To this end, we introduce the auxiliary notion of a CwF with an adjunction:

Definition 16. A CwF+A consists of a CwF with an adjunction $L \dashv R$ on the category of contexts, such that R extends to a weak CwF endomorphism.

Lemma 17. If \mathbf{C} with the adjunction $L \dashv R$ is a CwF+A, then there is a CwDRA structure on \mathbf{C} with L as the required functor on \mathbf{C} .

Proof. We write η for the unit of the adjunction. For a family $A \in \mathbf{C}(L\Gamma)$, we define $R_\Gamma A \in \mathbf{C}(\Gamma)$ to be $(RA)[\eta]$. For an element $a \in \mathbf{C}(L\Gamma \vdash A)$, we define its transpose $\bar{a} \in \mathbf{C}(\Gamma \vdash R_\Gamma A)$ to be $(Ra)[\eta]$. For the opposite direction, suppose $b \in \mathbf{C}(\Gamma \vdash R_\Gamma A)$. Since $(\eta, b) : \Gamma \rightarrow R L\Gamma \cdot RA$, we have that $L(v_{L\Gamma, A} \circ (\eta, b)) : L\Gamma \rightarrow L R(L\Gamma \cdot A)$ and thus we can define $\bar{b} \in \mathbf{C}(L\Gamma \vdash A)$ to be the element $q_A[\varepsilon \circ L(v_{L\Gamma, A} \circ (\eta, b))]$. Note that this is well typed because q_A is an element of the family $A[p_A]$ and so \bar{b} is an element of

$$\begin{aligned} A[p_A \circ \varepsilon \circ L(v \circ (\eta, b))] &= A[\varepsilon \circ L(R p_A \circ v \circ (\eta, b))] \\ &= A[\varepsilon \circ L(p_{RA} \circ (\eta, b))] \\ &= A[\varepsilon \circ L(\eta)] \\ &= A \end{aligned}$$

using equation (12) in the second equality. These operations can be proved inverses of each other using the equations (13) and (14). \square

Note that the conditions for a $\text{CwF}+\text{A}$ are stronger than those for a CwDRA ; for instance, a CwDRA does not require R to be defined on the context category. We return to the relation between these constructions in Section 4.1

Lemma 18. *If \mathbf{C} is a cartesian category and $L \dashv R$ are the adjoint endofunctors on \mathbf{C} , then $\mathcal{G}\mathbf{C}$ with the adjunction $L \dashv R$ is a $\text{CwF}+\text{A}$.*

Proof. We are already given an adjunction on the underlying category of $\mathcal{G}\mathbf{C}$. Theorem 15 constructs the weak CwF morphism. \square

Theorem 19. *If \mathbf{C} is a cartesian category and $L \dashv R$ are the adjoint endofunctors on \mathbf{C} , then $\mathcal{G}\mathbf{C}$ has the structure of a CwDRA .*

Proof. By Lemmas 17 and 18. \square

The above Theorem 19 thus provides a general construction of CwDRAs . In Section 5, we use it to present examples from the literature. As mentioned earlier, the local universe construction interacts well with other type formers: If we start with a locally cartesian closed category \mathbf{C} (with W -types, Id -types and a universe), then $\mathcal{G}\mathbf{C}$ also models dependent products Π and sums Σ (and W -types, Id -types and a universe); see Lumsdaine and Warren (2015). In Section 6, we consider universes.

4.1 $\text{CwF}+\text{A}$ from a CwDRA

In this subsection, we show how to produce a $\text{CwF}+\text{A}$ from a CwDRA under the assumption that the CwF is *democratic*. Intuitively, a democratic CwF is one where every context comes from a type, and hence it is not surprising that for a democratic CwDRA one can use the action of the dependent right adjoint on families to define a right adjoint on contexts.

Definition 20. *A CwF is democratic (Clairambault and Dybjer 2014), if for every context Γ , there is a family $\widehat{\Gamma} \in \mathbf{C}(\top)$ and an isomorphism $\zeta_\Gamma : \Gamma \rightarrow \top.\widehat{\Gamma}$.*

Theorem 21. *Let \mathbf{C} be a democratic CwDRA . The endofunctor $L : \mathbf{C} \rightarrow \mathbf{C}$, part of the CwDRA structure, has a right adjoint R .*

Proof. For $\Gamma \in \mathbf{C}$, we define $R\Gamma \in \mathbf{C}$ by

$$R\Gamma \triangleq \top.R_\top(\widehat{\Gamma}[\cdot]_{L\top}) \quad (15)$$

We have a bijection, natural in Δ

$$\begin{aligned} \mathbf{C}(\Delta, R\Gamma) &\cong \mathbf{C}(\Delta \vdash R_\top(\widehat{\Gamma}[\cdot]_{L\top}))[\cdot]_{L\Delta}) \\ &\cong \mathbf{C}(\Delta \vdash R_\Delta(\widehat{\Gamma}[\cdot]_{L\Delta})) \\ &\cong \mathbf{C}(L\Delta \vdash \widehat{\Gamma}[\cdot]_{L\Delta}) \\ &\cong \mathbf{C}(L\Delta, \top.\widehat{\Gamma}) \\ &\cong \mathbf{C}(L\Delta, \Gamma) \end{aligned}$$

The last of the above bijections follows by composition with ζ_Γ^{-1} .

Let $\gamma : \Gamma' \rightarrow \Gamma$, we have then an action $\gamma^* : \mathbf{C}(-, R\Gamma') \rightarrow \mathbf{C}(-, R\Gamma)$ given by

$$\mathbf{C}(-, R\Gamma') \cong \mathbf{C}(L-, \Gamma') \xrightarrow{-\circ\gamma} \mathbf{C}(L-, \Gamma) \cong \mathbf{C}(-, R\Gamma)$$

Define $R\gamma = \gamma_{R\Gamma'}^*(\text{id}_{R\Gamma'})$. Then, the correspondence $\mathbf{C}(\Delta, R\Gamma) \cong \mathbf{C}(L\Delta, \Gamma)$ is natural in Γ , proving that R is a right adjoint to L . \square

Consider a democratic CwDRA, with \mathbf{C} as the underlying category, and $L \dashv R$ the adjunction obtained from the above theorem. We then extend R to a weak CwF morphism by defining, for a family $A \in \mathbf{C}(\Gamma)$ and an element $a \in \mathbf{C}(\Gamma \vdash A)$,

$$RA \triangleq R_{R\Gamma} (A[\varepsilon]) \qquad Ra \triangleq \overline{a[\varepsilon]}$$

where $\varepsilon : L R \Gamma \rightarrow \Gamma$ is the counit of the adjunction.

Lemma 22. *R as defined above is a weak CwF morphism. In particular, for $A \in \mathbf{C}(\Gamma)$, we have an isomorphism $v_{\Gamma,A} : R\Gamma.RA \rightarrow R(\Gamma.A)$, inverse to (Rp_A, Rq_A) .*

Proof. We will show a bijection $\mathbf{C}(\Delta, R\Gamma.RA) \cong \mathbf{C}(\Delta, R(\Gamma.A))$ natural in Δ . We have

$$\mathbf{C}(\Delta, R\Gamma.RA) \cong \prod_{\gamma : \mathbf{C}(\Delta, R\Gamma)} \mathbf{C}(\Delta \vdash (RA)[\gamma])$$

We have a bijection $-^\top : \mathbf{C}(\Delta, R\Gamma) \cong \mathbf{C}(L\Delta, \Gamma)$. But

$$(RA)[\gamma] = (R_{R\Gamma} A[\varepsilon])[\gamma] = R_\Delta (A[\varepsilon \circ L\gamma]) = R_\Delta (A[\gamma^\top])$$

Hence, we have a bijection $\mathbf{C}(\Delta \vdash (RA)[\gamma]) \cong \mathbf{C}(L\Delta \vdash A[\gamma^\top])$. So

$$\begin{aligned} \mathbf{C}(\Delta, R\Gamma.RA) &\cong \prod_{\gamma : \mathbf{C}(\Delta, R\Gamma)} \mathbf{C}(\Delta \vdash (RA)[\gamma]) \\ &\cong \prod_{\gamma' : \mathbf{C}(L\Delta, \Gamma)} \mathbf{C}(L\Delta \vdash A[\gamma']) \\ &\cong \mathbf{C}(L\Delta, \Gamma.A) \\ &\cong \mathbf{C}(\Delta, R(\Gamma.A)) \end{aligned}$$

By the Yoneda lemma, this implies $R\Gamma.RA \cong \mathbf{C}(\Delta, R(\Gamma.A))$, and it is easy to check that the direction $\mathbf{C}(\Delta, R(\Gamma.A)) \rightarrow R\Gamma.RA$ is given by (Rp_A, Rq_A) . \square

Corollary 23. *A democratic CwDRA has the structure of CwF+A.*

Remark 24. For a category \mathbf{C} with a terminal object, the CwF $\mathcal{G}\mathbf{C}$ is democratic with $\widehat{\Gamma}$ given by the diagram:

$$\begin{array}{ccc} & & \Gamma \\ & & \downarrow !_\Gamma \\ 1 & \xrightarrow{\quad} & 1 \\ & & \uparrow !_1 \end{array}$$

Remark 25. For ordinary dependent type theory, the term model is a democratic CwF (Castellan *et al.* 2017, Section 4). However, the term model for our modal dependent type theory is *not* democratic, since there is, for example, no type corresponding to the context \blacksquare consisting of just one lock.

5. Examples

We now present concrete examples of CwDRAs generated from cartesian categories with an adjunction of endofunctors, including those mentioned in the introduction.

Π type with closed domain. Consider a CwF where the underlying category of contexts \mathbf{C} is cartesian closed, and let A be a closed type. We have then an adjunction of endofunctors $- \times \top.A \dashv -^{\top.A}$ on \mathbf{C} and suppose that the right adjoint extends to a weak CwF endomorphism, giving the structure of a CwF+A. As we saw above, this happens, for example, when the CwF is of the form $\mathcal{G}\mathbf{C}$. In this case, $R_\Gamma B$ behaves as a type of the form $\Pi(x:A)B$ since $\mathbf{C}(\Gamma \vdash R_\Gamma B) \cong \mathbf{C}(\Gamma \times \top.A \vdash B) \cong \mathbf{C}(\Gamma.(A[!_\Gamma]) \vdash B)$.

Thus, the notion of dependent right adjoint generalises Π types with closed domain. This generalises to the setting where \mathbf{C} carries the structure of a monoidal closed category, in which case the adjunction $- \otimes \top.A \dashv \top.A \multimap (-)$ extends to give a dependent notion of linear function space with closed domain. The next example is an instance of this.

Dependent name abstraction. The notion of *dependent name abstraction* for families of nominal sets was introduced by Pitts et al. (2015, Section 3.6) to give a semantics for an extension of Martin-Löf Type Theory with names and constructs for freshness and name abstraction. It provides an example of a CwDRA that can be presented via Theorem 19. In this case, \mathbf{C} is the category \mathbf{Nom} of nominal sets and equivariant functions (Pitts 2013). Its objects are sets Γ equipped with an action of finite permutations of a fixed infinite set of atomic names \mathbb{A} , with respect to which the elements of Γ are finitely supported, and its morphisms are functions that preserve the action of name permutations. \mathbf{Nom} is a topos (it is equivalent to the Schanuel topos (Pitts 2013, Section 6.3)) and hence in particular is cartesian. We take the functor $L : \mathbf{Nom} \rightarrow \mathbf{Nom}$ to be separated product (Pitts 2013, Section 3.4) with the nominal set of atomic names. This has a right adjoint R that sends each $\Gamma \in \mathbf{Nom}$ to the nominal set of name abstractions $[\mathbb{A}]\Gamma$ (Pitts 2013, Section 4.2) whose elements are a generic form of α -equivalence class in the case that Γ is a nominal set of syntax trees for some language.

Applying Theorem 19, we get a CwDRA structure on $\mathcal{G}\mathbf{Nom}$. In fact, the CwF $\mathcal{G}\mathbf{Nom}$ has an equivalent, more concrete description in this case, in terms of *families of nominal sets* (Pitts et al. 2015, Section 3.1). Under this equivalence, the value $R_\Gamma A \in \mathcal{G}\mathbf{Nom}(\Gamma)$ of the dependent right adjoint at $A \in \mathcal{G}(\mathbf{L}\Gamma)$ corresponds to the family of *dependent name abstractions* defined by Pitts et al. (2015, Section 3.6). The bijection (4) is given in one direction by the name abstraction operation (Pitts et al. 2015, (40)) and in the other by concretion at a fresh name (Pitts et al. 2015, (42)).

Guarded and cloTT. Guarded recursion (Nakano 2000) is an extension of type theory with a modal *later* operator, denoted \triangleright , on types, an operation $\text{next} : A \rightarrow \triangleright A$ and a guarded fixed point operator $\text{fix} : (\triangleright A \rightarrow A) \rightarrow A$ mapping f to a fixed point for $f \circ \text{next}$. The standard model of guarded recursion is the topos of trees (Birkedal et al. 2012), that is, the category of presheaves on ω , with $\triangleright X(n+1) = X(n)$, $\triangleright X(0) = 1$. The later operator has a left adjoint \triangleleft , called *earlier*, given by $\triangleleft X(n) = X(n+1)$, so \triangleright yields a dependent right adjoint on the induced CwDRA.

Birkedal et al. (2012, Section 6.1) show that \triangleleft in a dependently typed setting does not commute with re-indexing. However, it does have a left adjoint, namely, the ‘stutter’ functor $!$ with $!X(0) = X(0)$ and $!X(n+1) = X(n)$, so \triangleleft does give rise to a well-behaved modality in the setting of this paper. This apparent contradiction is resolved by the use of locks in the context: $\Gamma \vdash A$ does not give rise to a well-behaved $\Gamma \vdash \triangleleft A$, but $\Gamma, \blacksquare \vdash A$ does. This is an intriguing example of the Fitch-style approach increasing expressivity.

Guarded recursion can be used to encode co-induction given a *constant* modality (Clouston et al. 2015), denoted \square , on the topos of trees, defined as $\square X(n) = \lim_k X(k)$. The \square functor is the right adjoint of the essential geometric morphism on $\hat{\omega}$ induced by $0 : \omega \rightarrow \omega$, the constant map to 0, and hence it also yields a dependent right adjoint. In Clouston et al. (2015), \square was used in

a simple type theory, employing ‘explicit substitutions’ following Bierman and de Paiva (2000). As we will discuss in Section 7, this approach proved difficult to extend to dependent types, and we wish to use the modal dependent type theory of the present paper to study \Box in dependent type theory.

An alternative to the constant modality are the *clock quantifiers* of Atkey and McBride (2013), which unlike the constant modality have already been combined successfully with dependent types (Bizjak *et al.* 2016; Møgelberg 2014). They are also slightly more general than the constant modality, as multiple clocks allow co-inductive data structures that unroll in multiple dimensions, such as infinitely wide infinitely deep trees. The denotational semantics, however, are more complicated, consisting of presheaves over a category of ‘time objects’, restricted to those fulfilling an ‘orthogonality’ condition (Bizjak and Møgelberg 2018). Nevertheless, the $\triangleleft \dashv \triangleright$ adjunction of the topos of trees lifts to this category, and so once again we may construct a CwDRA.

cloTT (CloTT) (Bahr *et al.* 2017) is a recent type theory for guarded recursion that has strongly normalising reduction semantics and has been shown to have semantics in the category discussed above (Mannaa and Møgelberg 2018). The operator \triangleright is refined to a form of dependent function type $\triangleright (\alpha : \kappa).A$ over ticks α on clock κ . Ticks can appear in contexts as $\Gamma, \alpha : \kappa$; these are similar to the locks of Fitch-style contexts, except that ticks have names, and can be weakened. The names of ticks play a crucial role in controlling fixed point unfoldings.

Finally, the modal operator \triangleright on the topos of trees can be generalised to the presheaf topos $\widehat{\mathbf{C} \times \omega}$ for any category \mathbf{C} , simply by using the identity on \mathbf{C} to extend the underlying functor (which generates the essential geometric morphism) on ω to $\mathbf{C} \times \omega$. In Birkedal *et al.* (2018), this topos, with \mathbf{C} the cube category, is used to model guarded *cubical* type theory, an extension of cubical type theory (Cohen *et al.* 2018). In more detail, one uses a CwF where families are certain *fibrations*, and since \triangleright preserves fibrations, it does indeed extend to a CwDRA.

Cohesive toposes Cohesive toposes have also recently been considered as models of a form of modal type theory (Rijke *et al.* 2018; Shulman 2018). Cohesive toposes carry a triple adjunction $\int \dashv \flat \dashv \sharp$ and hence induce two dependent right adjoints. Examples of cohesive toposes include simplicial sets $\hat{\Delta}$ and cubical sets $\hat{\square}$; since these are presheaf toposes, they also model universes. For example, for simplicial sets, the triple of adjoints are given by the essential geometric morphism induced by the constant functor $0 : \Delta \rightarrow \Delta$. In the category of cubical sets, \sharp has a further right adjoint, used by Nuyts *et al.* (2017) to reason about parametricity.

Tiny objects Licata *et al.* (2018) use a ‘tiny’ object \mathbb{I} to construct the fibrant universe in the cubical model of homotopy type theory. By definition, an object \mathbb{I} of a category \mathbf{C} is *tiny* if the exponentiation functor $(-)^{\mathbb{I}} : \mathbf{C} \rightarrow \mathbf{C}$ has a right adjoint, which they denote by \checkmark . As for \triangleleft above, the right adjoint functor \checkmark exists globally, but not locally; in other words, there is no right adjoint to $(-)^{\mathbb{I}}$ on each CwF over an object $\Gamma \in \mathbf{C}$, stable under re-indexing Γ (except in the trivial case that \mathbb{I} is terminal). Nevertheless, our present framework is still applicable: the corresponding *dependent* right adjoint for $(-)^{\mathbb{I}}$, constructed as in Section 4, plays an important part in the construction of the fibrant universe given in (Licata *et al.* 2018).

6. Universes

In this section, we extend our modal dependent type theory with universes. For the semantics, we start from Coquand’s notion of a category with universes (Coquand 2012), which covers all presheaf models of dependent type theory with universes. The notion of *category with universes* rests on the observation that in presheaf models one can interpret an inverse $\ulcorner \dashv \urcorner$ to the usual function El from codes to types and hence obtain a simpler notion of universe than usual (such as in Hofmann 1997, Section 2.1.6).

Definition 26. (Category with universes). A *CwU* is specified by:

1. A category \mathbf{C} with a terminal object \top .
2. For each object $\Gamma \in \mathbf{C}$ and natural number $n \in \mathbb{N}$, a set $\mathbf{C}(\Gamma, n)$ of families at universe level n over Γ .
3. For each object $\Gamma \in \mathbf{C}$, natural number n , and family $A \in \mathbf{C}(\Gamma, n)$, a set $\mathbf{C}(\Gamma \vdash A)$ of elements (at some level) of the family A over Γ .
4. For each morphism $\gamma \in \mathbf{C}(\Delta, \Gamma)$, re-indexing functions $A \in \mathbf{C}(\Gamma, n) \mapsto A[\gamma] \in \mathbf{C}(\Delta, n)$ and $a \in \mathbf{C}(\Gamma \vdash A) \mapsto a[\gamma] \in \mathbf{C}(\Delta \vdash A[\gamma])$, satisfying equations for associativity and identity as in a *CwF*.
5. For each object $\Gamma \in \mathbf{C}$, number n and family $A \in \mathbf{C}(\Gamma, n)$, a comprehension object $\Gamma.A \in \mathbf{C}$ equipped with projections and generic elements satisfying equations as in a *CwF*.
6. For each number n , a family $\mathbf{U}_n \in \mathbf{C}(\top, n+1)$, the universe at level n .
7. For each object $\Gamma \in \mathbf{C}$ and number n , a code function $A \in \mathbf{C}(\Gamma, n) \mapsto \ulcorner A \urcorner \in \mathbf{C}(\Gamma \vdash \mathbf{U}_n[!_\Gamma])$, and an element function $u \in \mathbf{C}(\Gamma \vdash \mathbf{U}_n[!_\Gamma]) \mapsto Eu \in \mathbf{C}(\Gamma, n)$, satisfying $\ulcorner A \urcorner[\gamma] = \ulcorner A[\gamma] \urcorner$, $E\ulcorner A \urcorner = A$, and $E\ulcorner Eu \urcorner = u$.

We will of course want the universes to be closed under various type-forming operations, but in this formalisation of universes these definitions are just as for *CwFs*, without having to explicitly reflect them into the universes.

Lemma 27. The element function is stable under re-indexing: $(Eu)[\gamma] = E(u[\gamma])$.

Proof. $(Eu)[\gamma] = E^\ulcorner(Eu)[\gamma]\urcorner = E(\ulcorner Eu \urcorner[\gamma]) = E(u[\gamma])$. □

Corollary 28. In a *CwU*, there is a generic family $E! \in \mathbf{C}(\top, \mathbf{U}_n, n)$ of types of level n (for each $n \in \mathbb{N}$), with the property that $E! [(\ulcorner \Gamma \urcorner, \ulcorner A \urcorner)] = A$, for all $A \in \mathbf{C}(\Gamma, n)$.

Proof. Since $p_{\mathbf{U}_n} = ! : \top, \mathbf{U}_n \rightarrow \top$, we have $q \in \mathbf{C}(\top, \mathbf{U}_n \vdash \mathbf{U}_n[!_{\top, \mathbf{U}_n}])$, and thus we can define $E!$ to be $E q$, and then the required property follows by Lemma 27. □

For a *CwU*, there is an underlying *CwF* with families over Γ given as $\mathbf{C}(\Gamma) = \bigcup_n \mathbf{C}(\Gamma, n)$. Using this, we can extend the definition of *CwDRA* to categories with universes in the obvious way, as follows.

Definition 29. (CwUDRA). A *CwUDRA* is a *CwU* with the structure of a *CwDRA* such that operation on types preserves universe levels in the sense that $A \in \mathbf{C}(\mathbf{L}\Gamma, n)$ implies $R_\Gamma A \in \mathbf{C}(\Gamma, n)$.

Similarly, one can extend the notion of *CwF+A* from Definition 16 to the setting of universes.

Definition 30. (CwU+A). A weak *CwU* morphism R is a weak *CwF* morphism on the underlying *CwFs* preserving size in the sense that $A \in \mathbf{C}(\Gamma, n)$ implies $RA \in \mathbf{C}(R\Gamma, n)$. A *CwU+A* consists of a *CwU* with an adjunction $L \dashv R$ on the category of contexts, such that R extends to a weak *CwU* morphism.

The construction of Lemma 17 extends to a construction of a *CwUDRA* from a *CwU+A*. We now show (Lemma 32) that the action of the right adjoint on families and elements can be defined by just defining it on the universe as in the following definition.

Definition 31. A universe endomorphism on a CwU is a finite limit preserving functor R on the category of contexts together with, for each n , a family $RI \in C(R(\top.U_n), n)$ and an element $r \in C(R(\top.U_n.El) \vdash RI[Rp])$ such that the morphism

$$\begin{array}{ccc} R(\top.U_n.El) & \xrightarrow{(Rp, r)} & R(\top.U_n).RI \\ & \searrow Rp \quad \swarrow p & \\ & R(\top.U_n) & \end{array} \quad (16)$$

over $R(\top.U_n)$ is an isomorphism; in other words, there is a morphism $\ell : R(\top.U_n).RI \rightarrow R(\top.U_n.El)$ satisfying $\ell \circ (Rp, r) = id$ and $(Rp, r) \circ \ell = id$.

This means that we have a universe category endomorphism in the sense of Voevodsky (2014, Section 4.1): a family $RI \in C(R(\top.U_n), n)$ gives a pullback square with the morphism $\top.U_n.El \rightarrow \top.U_n$ and the code function. The isomorphism above implies that the universe $R(\top.U_n.El) \rightarrow R(\top.U_n)$ is also pullback of $\top.U_n.El \rightarrow \top.U_n$ along the code function.

Given a CwU with a weak CwU morphism R , then clearly R is a universe endomorphism, with $RI \triangleq R(El)$, $r \triangleq Rq$ and $\ell \triangleq v$. Conversely:

Lemma 32. Any CwU with a universe endomorphism $R : C \rightarrow C$ extends to a weak CwU morphism.

Proof. Given $A \in C(\Gamma, n)$, since we have $(!_\Gamma, \ulcorner A \urcorner) : \Gamma \rightarrow \top.U_n$, we can define

$$RA \triangleq RI[R(!_\Gamma, \ulcorner A \urcorner)] \in C(R\Gamma, n) \quad (17)$$

This is stable under re-indexing, since for $\gamma : \Delta \rightarrow \Gamma$

$$\begin{aligned} R(A[\gamma]) &\triangleq RI[R(!_\Delta, \ulcorner A[\gamma] \urcorner)] \\ &= RI[R(!_\Delta, \ulcorner A \urcorner[\gamma])] \\ &= RI[R((!_\Gamma, \ulcorner A \urcorner) \circ \gamma)] \\ &= RI[R(!_\Gamma, \ulcorner A \urcorner) \circ R\gamma] \\ &= (RI[R(!_\Gamma, \ulcorner A \urcorner)])[R\gamma] \\ &\triangleq (RA)[R\gamma] \end{aligned}$$

Given $a \in C(\Gamma \vdash A)$, by Corollary 28, we have $a \in C(\Gamma \vdash El[(!_\Gamma, \ulcorner A \urcorner)])$ and hence

$$((!_\Gamma, \ulcorner A \urcorner), a) : \Gamma \rightarrow \top.U_n.El$$

Therefore,

$$r[R((!_\Gamma, \ulcorner A \urcorner), a)] \in C(R\Gamma \vdash (RI[Rp])[R((!_\Gamma, \ulcorner A \urcorner), a)])$$

But $(RI[Rp])[R((!_\Gamma, \ulcorner A \urcorner), a)] = RI[R(p \circ ((!_\Gamma, \ulcorner A \urcorner), a))] = RI[R(!_\Gamma, \ulcorner A \urcorner)] \triangleq RA$. We can therefore define

$$Ra \triangleq r[R((!_\Gamma, \ulcorner A \urcorner), a)] \in C(R\Gamma \vdash RA) \quad (18)$$

and this is stable under re-indexing, since for $\gamma : \Delta \rightarrow \Gamma$

$$\begin{aligned} (Ra)[R\gamma] &\triangleq r[R((!_\Gamma, \ulcorner A \urcorner), a)][R\gamma] \\ &= r[R((!_\Delta, \ulcorner A \urcorner[\gamma]), a[\gamma])] \\ &= r[R((!_\Delta, \ulcorner A[\gamma] \urcorner), a[\gamma])] \\ &\triangleq R(a[\gamma]) \end{aligned}$$

Finally, we must show that R commutes with comprehension. For this, note that there are pull-back squares

$$\begin{array}{ccc} R(\Gamma.A) & \xrightarrow{R(!_{\Gamma.A}, \ulcorner A \urcorner[p_A]), q)} & R(\top.U_n.EI) \\ R p \downarrow \lrcorner & & \downarrow R p \\ R \Gamma & \xrightarrow{R(!_{\Gamma}, \ulcorner A \urcorner)} & R(\top.U_n) \end{array} \quad \begin{array}{ccc} R \Gamma. R A & \xrightarrow{(R(!_{\Gamma}, \ulcorner A \urcorner) \circ p, q)} & R(\top.U_n). RI \\ p \downarrow \lrcorner & & \downarrow p \\ R \Gamma & \xrightarrow{R(!_{\Gamma}, \ulcorner A \urcorner)} & R(\top.U_n) \end{array}$$

the former because the functor R preserves finite limits and the latter by definition of RA . Applying (18) with $a = q$, we get that the pullback along $R(!_{\Gamma}, \ulcorner A \urcorner)$ of the morphism $(R p, r)$ in (16) is

$$\begin{array}{ccc} R(\Gamma.A) & \xrightarrow{(R p, R q)} & R \Gamma. R A \\ & \searrow R p \quad \swarrow p & \\ & R \Gamma & \end{array}$$

Then, since $(R p, r)$ is an isomorphism, so is its pullback $(R p, R q)$, as required. \square

Remark 33. We observe that for R as constructed above, the image under R of maps with U_n -small fibers is classified by $RI \in C(R(\top.U_n), n)$. That is to say that $(!_{R\Gamma}, \ulcorner RA \urcorner) = (!_{R\Gamma}, \ulcorner RI \urcorner) \circ R(!_{\Gamma}, \ulcorner A \urcorner)$ which is true by our choice of $RA = RI[R(!_{\Gamma}, \ulcorner A \urcorner)]$. Hence, the type of codes for such fibers is R applied to the codes for types. The same situation occurred for \triangleright in (Birkedal and Møgelberg 2013, V.5), but was not observed at the time.

Theorem 34. Any CwU equipped with an adjunction on the category of contexts whose right adjoint is a universe endomorphism can be given the structure of a $CwUDRA$.

Proof. Combine Lemmas 17 and 32. \square

For most of the presheaf examples considered in Section 5, the dependent right adjoint is obtained as the direct image of an essential geometric morphism arising from a functor on the category on which the presheaves are defined. We show that in this case, the right adjoint preserves universe levels and hence gives a $CwUDRA$. For simplicity, we will restrict to one universe and show that the right adjoint preserves smallness with respect to this.

Let U be a universe in an ambient set theory. We call elements of U , U -sets. A U -small category is the one where both the sets of objects and the set of morphisms are U -small. Let us assume that U is U -complete – it is closed under limits of U -small diagrams. A Grothendieck universe in Zermelo Fraenkel set theory with the axiom of Choice (ZFC) would satisfy these conditions.

Proposition 35. Let C, D be U -small categories and $f : C \rightarrow D$ a functor between them. The direct image f_* of the induced geometric morphism preserves size. In particular, for each endofunctor f , the direct image is a weak CwU morphism.

Proof. Since f_* is a right adjoint, we know that it induces a weak CwF morphism, and we just need to show that it maps U -small families to U -small families. Recall first that the direct image f_* is the (pointwise) right Kan extension (Johnstone 2002, A4.1.4) defined on objects by the limit of the diagram

$$(\text{Ran}_f F)d \triangleq \lim (f \downarrow d) \xrightarrow{\pi_1} C^{op} \xrightarrow{F} U\text{-set},$$

for $F \in \widehat{C}$ and $d \in D$. Here, $(f \downarrow d)$ denotes the comma category consisting of pairs $(c; g : f(c) \rightarrow d)$.

A family $\alpha : F \rightarrow G$, for $F, G \in \hat{\mathcal{C}}$, is U -small if for each c and each $x \in G(c)$, the set $\alpha_c^{-1}(x)$ is in U . Given $(x_g)_{g \in f \downarrow d} \in f_* G(d) = (\text{Ran}_f G)d$, the pre-image $(f_* \alpha)^{-1}((x_g)_{g \in f \downarrow d})$ is the set

$$\{(y_g)_{g \in f \downarrow d} \in (\text{Ran}_f G)d \mid \forall g. \alpha_c(y_g) = x_g\}$$

which is the limit of the diagram associating to each g the set $\alpha_c^{-1}(x_g)$. Since each of these sets are in U by assumption and since also $f \downarrow d$ is in U , by the assumption of U being closed under limits, also $(f_* \alpha)^{-1}((x_g)_{g \in f \downarrow d})$ is in U as desired. \square

Syntax. At this stage, it should hopefully be clear that one can refine and extend the syntax of modal dependent type theory from Section 3, so that the resulting syntactic type theory can be modelled in a CwU+A. The idea is, of course, to refine the judgement for well-formed types and to include a level n , so that it has the form $\Gamma \vdash_n A$, and likewise for type equality judgements. For example,

$$\frac{\Gamma, \blacksquare \vdash_n A}{\Gamma \vdash_n \Box A}$$

In addition to the existing rules for types (indexed with a level) and terms, we then also include:

$$\frac{}{\diamond \vdash_{n+1} \mathbf{U}_n} \quad \frac{\Gamma \vdash_n A}{\Gamma \vdash \ulcorner A \urcorner : \mathbf{U}_n} \quad \frac{\Gamma \vdash u : \mathbf{U}_n}{\Gamma \vdash_n E u}$$

Finally, we add the following type and term equality rules:

$$\frac{\Gamma \vdash_n A}{\Gamma \vdash_n E \ulcorner A \urcorner = A} \quad \frac{\Gamma \vdash u : \mathbf{U}_n}{\Gamma \vdash \ulcorner E u \urcorner = u : \mathbf{U}_n}$$

As an example, there is a term

$$\hat{\Box} \triangleq \lambda x. \ulcorner \Box E (\text{open } x) \urcorner : \Box \mathbf{U}_n \rightarrow \mathbf{U}_n$$

which encodes the \Box type constructor on the universe in the sense that

$$E(\hat{\Box}(\text{shut } u)) = \Box(E u)$$

This is similar to the $\hat{\triangleright}$ operator of Guarded Dependent Type Theory (Bizjak et al. 2016), which is essential to defining guarded recursive types. Thus, $\hat{\triangleright}$ arises for general reasons quite unconnected to the specifics of guarded recursion.

7. Discussion

7.1 Related Work

Modal dependent type theory builds on work on the computational interpretation of modal logic with *simple* types. Some of this work involves a standard notion of context; most relevantly to this paper, the calculus for Intuitionistic K of Bellin et al. (2001), which employs *explicit substitutions* in terms. Departing from standard contexts, Fitch-style calculi were introduced independently by Borghuis (1994) and Martini and Masini (1996). Recent work by Clouston (2018) argued that Fitch-style calculus can be extended to a variety of different modal logics and gave a sound categorical interpretation by modelling the modality as a right adjoint. Another non-standard notion of context are the *dual contexts* introduced by Davies and Pfenning (2001) for the modal logic Intuitionistic S4 of comonads. Here, a context $\Delta; \Gamma$ is understood as meaning $\Box \Delta \wedge \Gamma$, so the structure in the context is modelled by the modality itself, not its left adjoint. Recent work by Kavvos (2017) has extended this approach to a variety of modal logics, including Intuitionistic K.

There exists recent work employing variants of dual contexts for modal *dependent* type theory, all involving (co)monads rather than the more basic logic of this paper. Spatial type theory (Shulman 2018), designed for applications in homotopy type theory (see also (Licata et al. 2018; Wellen 2017)), extends the Davies–Pfenning calculus for a comonad with both dependent types and a second modality, a monad right adjoint to the comonad. Second, the calculus for parametricity of Nuyts et al. (2017) uses *three* zones to extend Davies–Pfenning with a monad *left* adjoint to the comonad. They focus on Π - and Σ -types with modalised arguments, but a more standard modality can be extracted by taking the second argument of a modalised Σ -type to be the unit type. In both the above works, the leftmost modality is intended to itself be a right adjoint, so they potentially could also be captured by a Fitch-style calculus. Third, de Paiva and Ritter (2016) suggest a generalisation of Davies–Pfenning with some unusual properties, as \Box types carry an auxiliary typed variable and Π -types may only draw their argument from the modal context. We finally note that the dual contexts approach has inspired the mode theories of Licata et al. (2017), but this line of work as yet does not support a term calculus.

We, however, do not know how to apply the dual context approach to modal logics where the modality is not a (co)monad. For example, it is not obvious how to extend Kavvos’s simply typed calculus for Intuitionistic K. This should be compared to the ease of extending the simply typed Fitch-style calculus with dependent types. We hope that Fitch-style calculi continue to provide a relatively simple setting for modal dependent type theory as we explore the extensions discussed in the next subsection.

We are not aware of any successful extensions of the explicit substitution approach to dependent types; our own experiments with this while developing Guarded Dependent Type Theory (Clouston et al. 2015) suggest that this is probably possible but becomes unwieldy with real examples. Far more successful was the cloTT (Bahr et al. 2017) discussed in Section 5, which can now be seen to have rediscovered the Fitch-style framework, albeit with the innovation of named locks to control fixed-point unfoldings. That work provides the inspiration for the more foundational developments of this paper.

7.2 Future work

We wish to develop operational semantics for dependent Fitch-style calculi, and conjecture that standard techniques for sound normalisation and canonicity can be extended, as was possible for simply typed Fitch-style calculi (Borghuis 1994; Clouston 2018), and for cloTT (Bahr et al. 2017). Such results should then lead to practical implementation.

The modal axiom Intuitionistic K was used in this paper, because it provides a basic notion of modal necessity and holds of many useful models. Nonetheless, for particular applications, we will want to develop Fitch-style calculi corresponding to more particular logics. There can be no algorithm for converting additional axioms to well-behaved calculi, but we know that Fitch-style calculi are extremely versatile in the simply typed case (Clouston 2018), and cloTT provides one example of this with dependent types. In particular, we are interested in Fitch-style calculi with multiple interacting modalities, each of which is assigned its own lock; we hope to develop guarded type theory with both \triangleright and \Box modalities in this style.

The notion of CwF with a weak CwF endomorphism (Definition 14) is more general than our CwF+A, as it does not require the existence of a left adjoint. Because a weak CwF endomorphism must preserve products, it appears to be a rival candidate for a model of dependent type theory with the K axiom. However, we do not know how to capture this class of models in syntax. Understanding this would be valuable because *truncation* (Awodey and Bauer 2004), considered as an endofunctor, for example, on sets, defines such a morphism but is not a right adjoint. Truncation allows one to move between general types and propositions. For example, combining it with guarded types would allow us to formalise the work in this field that makes that distinction (Birkedal et al. 2012; Clouston et al. 2015).

Notes

- 1 For an introduction to modal logic, see, for example, Blackburn *et al.* (2002).
- 2 For ‘Kripke’, not to be confused with Streicher’s K (Streicher 1993).
- 3 This should not be confused with models where there are adjoint functors between different categories that can be composed to define a monad or comonad.
- 4 This capability was achieved by Bizjak *et al.* (2016) via *delayed substitutions*, but this construction does not straightforwardly support an operational semantics (Bahr *et al.* 2017).

Acknowledgements. This work was supported in part by the following research grants: Guarded Homotopy Type Theory (grant number 12386) from VILLUM FONDEN; Type theories for reactive programming (grant number 13156) from VILLUM FONDEN; Guarded recursive types in the foundations of programming from the Danish Council for Independent Research (FNU); and Air Force Office of Scientific Research project “Homotopy Type Theory and Probabilistic Computation” (grant number 12595060).

References

- Atkey, R. and McBride, C. (2013) Productive coprogramming with guarded recursion. In: *18th ACM SIGPLAN International Conference on Functional Programming (ICFP 2013)*.
- Awodey, S. and Bauer, A. (2004). Propositions as types. *Journal of Logic and Computation* **14** (4), 447–471.
- Bahr, P., Grathwohl, H. B. and Mögelberg, R. E. (2017). The clocks are ticking: No more delays! In: *2017 32nd Annual ACM/IEEE Symposium on Logic in Computer Science (LICS)*, IEEE, 1–12.
- Bellin, G., De Paiva, V. and Ritter, E. (2001). Extended curry-howard correspondence for a basic constructive modal logic. In: *Proceedings of Methods for Modalities*.
- Bierman, G. M. and de Paiva, V. C. V. (2000) On an intuitionistic modal logic. *Studia Logica* **65** (3), 383–416.
- Birkedal, L., Bizjak, A., Clouston, R., Grathwohl, H. B., Spitters, B. and Vezzosi, A. (2018). Guarded cubical type theory. *Journal of Automated Reasoning*.
- Birkedal, L. and Mögelberg, R. E. (2013). Intensional type theory with guarded recursive types qua fixed points on universes. In: *Proceedings of the 2013 28th Annual ACM/IEEE Symposium on Logic in Computer Science*, IEEE Computer Society, 213–222.
- Birkedal, L., Mögelberg, R. E., Schwinghammer, J. and Støvring, K. (2012). First steps in synthetic guarded domain theory: Step-indexing in the topos of trees. *LMCS* **8**(4), 1–45.
- Bizjak, A., Grathwohl, H. B., Clouston, R., Mögelberg, R. E. and Birkedal, L. (2016). Guarded dependent type theory with coinductive types. In: *International Conference on Foundations of Software Science and Computation Structures*, Springer, 20–35.
- Bizjak, A. and Mögelberg, R. (2018). Denotational semantics for guarded dependent type theory, *arXiv:1802.03744*.
- Blackburn, P., De Rijke, M. and Venema, Y. (2002). *Modal Logic*, Vol. 53. *Cambridge Tracts in Theoretical Computer Science*, Cambridge University Press.
- Borghuis, V. A. J. (1994). *Coming to terms with modal logic: on the interpretation of modalities in typed lambda-calculus*, PhD thesis, Technische Universiteit Eindhoven.
- Castellan, S. (2014). *Dependent type theory as the initial category with families*, Technical report, Chalmers University of Technology. Internship Report. <http://iso.mor.phis.me/archives/2011-2012/stage-2012-goteburg/report.pdf>
- Castellan, S., Clairambault, P. and Dybjer, P. (2017). Undecidability of equality in the free locally cartesian closed category (extended version). *LMCS* **13**(4).
- Clairambault, P. and Dybjer, P. (2014). The biequivalence of locally cartesian closed categories and Martin-Löf type theories. *Mathematical Structures in Computer Science* **24**(6).
- Clouston, R. (2018). Fitch-style modal lambda calculi. In: *International Conference on Foundations of Software Science and Computation Structures*, Springer, 258–275.
- Clouston, R., Bizjak, A., Grathwohl, H. B. and Birkedal, L. (2015). Programming and reasoning with guarded recursion for coinductive types. In: *International Conference on Foundations of Software Science and Computation Structures*, Springer, 407–421.
- Cohen, C., Coquand, T., Huber, S. and Mörtberg, A. (2018). Cubical type theory: A constructive interpretation of the univalence axiom. In: Uustalu, T. (ed.) *21st International Conference on Types for Proofs and Programs (TYPES 2015)*, Vol. 69. Leibniz International Proceedings in Informatics (LIPIcs), Schloss Dagstuhl–Leibniz-Zentrum fuer Informatik, Dagstuhl, Germany, 5:1–5:34.
- Coquand, T. (2012). Presheaf model of type theory. Unpublished note. www.cse.chalmers.se/~coquand/presheaf.pdf
- Davies, R. and Pfenning, F. (2001). A modal analysis of staged computation. *JACM* **48** (3), 555–604.
- de Paiva, V. and Ritter, E. (2016). Fibrational modal type theory. *Electronic Notes in Theoretical Computer Science* **323**, 143–161.

- Dybjer, P. (1995). Internal type theory. In: *International Workshop on Types for Proofs and Programs*, Springer, 120–134.
- Fitch, F. B. (1952). *Symbolic Logic, An Introduction*, Ronald Press Co.
- Giraud, J. (1965). Cohomologie non abélienne, *C. R. Acad. Sci. Paris* **260**, 2666–2668.
- Hofmann, M. (1994). On the interpretation of type theory in locally cartesian closed categories. In: *International Workshop on Computer Science Logic*, Springer, 427–441.
- Hofmann, M. (1997). Syntax and semantics of dependent types. In: *Extensional Constructs in Intensional Type Theory*, Springer, 13–54.
- Johnstone, P. (2002). *Sketches of An Elephant: A Topos Theory Compendium*, Oxford University Press.
- Kapulkin, C. and Lumsdaine, P. L. (2018). The simplicial model of univalent foundations (after Voevodsky). *Journal of the European Mathematical Society*. To appear.
- Kavvos, G. (2017). Dual-context calculi for modal logic. In *2017 32nd Annual ACM/IEEE Symposium on Logic in Computer Science (LICS)*, IEEE, 1–12.
- Krishnaswami, N. R., Pdrad, P. and Benton, N. (2015). Integrating linear and dependent types. In: *ACM SIGPLAN Notices*, Vol. 50, ACM, 17–30.
- Licata, D. R., Orton, I., Pitts, A. M. and Spitters, B. (2018). Internal universes in models of homotopy type theory. In: *3rd International Conference on Formal Structures for Computation and Deduction, FSCD 2018*, July 9–12, 2018, Oxford, UK.
- Licata, D. R., Shulman, M. and Riley, M. (2017). A fibrational framework for substructural and modal logics. In: *LIPICs-Leibniz International Proceedings in Informatics*, Vol. 84, Schloss Dagstuhl-Leibniz-Zentrum fuer Informatik.
- Lumsdaine, P. L. and Warren, M. A. (2015). The local universes model: An overlooked coherence construction for dependent type theories. *ACM Trans. Comput. Logic* **16** (3), 23:1–23:31.
- Luo, Z. (1989). ECC, an extended calculus of constructions. In: *Proceedings of the Fourth Annual Symposium on Logic in Computer Science (LICS 1989)*, Pacific Grove, California, USA, June 5–8, 1989, 386–395.
- Mannaa, B. and Møgelberg, R. E. (2018). The clocks they are adjunctions denotational semantics for clocked type theory. In: *3rd International Conference on Formal Structures for Computation and Deduction, FSCD 2018*, July 9–12, 2018, Oxford, UK, 23:1–23:17.
- Martini, S. and Masini, A. (1996). A computational interpretation of modal proofs. In: *Proof theory of modal logic*, Springer, 213–241.
- Møgelberg, R. E. (2014). A type theory for productive coprogramming via guarded recursion. In *Joint Meeting of the Twenty-Third EACSL Annual Conference on Computer Science Logic (CSL) and the Twenty-Ninth Annual ACM/IEEE Symposium on Logic in Computer Science (LICS)*, CSL-LICS 2014, Vienna, Austria, July 14 – 18, 2014, 71:1–71:10.
- Nakano, H. (2000). A modality for recursion. In: *15th Annual IEEE Symposium on Logic in Computer Science*, Santa Barbara, California, USA, June 26–29, 2000, 255–266.
- Norell, U. (2007). *Towards a practical programming language based on dependent type theory*, PhD thesis, Chalmers University of Technology.
- Nuyts, A., Vezzosi, A. and Devriese, D. (2017). Parametric quantifiers for dependent type theory. *PACMPL* **1** (ICFP), 32:1–32:29.
- Pitts, A. M. (2013). *Nominal Sets: Names and Symmetry in Computer Science*, Vol. 57. *Cambridge Tracts in Theoretical Computer Science*, Cambridge University Press.
- Pitts, A. M., Matthiesen, J. and Derikx, J. (2015). A dependent type theory with abstractable names. *Electronic Notes in Theoretical Computer Science* **312**, 19–50.
- Rijke, E., Shulman, M. and Spitters, B. (2018). Modalities in homotopy type theory. *LMCS*.
- Shulman, M. (2018). Brouwer’s fixed-point theorem in real-cohesive homotopy type theory. *Mathematical Structures in Computer Science* **28** (6), 856–941.
- Streicher, T. (1993). *Investigations into intensional type theory*, Habilitation thesis, Ludwig Maximilian Universität.
- Vákár, M. (2017). *In search of effectful dependent types*, PhD thesis, University of Oxford.
- Voevodsky, V. (2014). A C-system defined by a universe category, *arXiv:1409.7925*.
- Wellen, F. (2017). *Formalizing Cartan Geometry in Modal Homotopy Type Theory*, PhD thesis, Karlsruher Institut für Technologie.