

# On the decidability of priced timed games

T. Brihaye<sup>1</sup>, G. Geeraerts<sup>2</sup>, S. Krishna<sup>3</sup>, L. Manasa<sup>3</sup> and A. Trivedi<sup>3</sup>

<sup>1</sup> Université de Mons – Belgium

<sup>2</sup> Université Libre de Bruxelles – Belgium

<sup>3</sup> Indian Institute of Technology Bombay – India


HIGHLIGHTS of Logic, Games and Automata – Paris

# Outline of the talk

- 1 Priced Timed Games and Optimal Strategies
- 2 Existing results
- 3 Going further...

# Classical game (qualitative, zero-sum, turn-based)

A game  $\mathcal{G} = ((V, E), V_0, V_1, \mathbf{G})$ . Pl. 0 aims at reaching  $\mathbf{G}$ .



A winning strategy for Pl. 0 from A


## Winning strategy for Pl. 0

A strategy  $\lambda_0 : V^* V_0 \rightarrow V$  is winning for Pl. 0 from  $v$  iff

$$\forall \lambda_1 : V^* V_1 \rightarrow V \quad \text{Out}(v, \lambda_0, \lambda_1) \text{ visits } \mathbf{G}.$$

# Priced Game

Pl. 0 aims at reaching **G** while minimising the cost.



A strategy for Pl. 0 ensuring 11 from A


## Strategy for Pl. 0 ensuring a cost $K$

A strategy  $\lambda_0$  ensures a cost  $K$  for Pl. 0 from  $v$  iff

$$\sup_{\lambda_1} \text{Cost}(\text{Out}(v, \lambda_0, \lambda_1)) \leq K.$$

# Priced game

Pl. 0 aims at reaching **G** while minimising the cost.



Optimal strategy for Pl. 0 from A (ensuring 9)


## Optimal Strategy for Pl. 0

A strategy  $\lambda_0^*$  is optimal for Pl. 0 from  $v$  iff

$$\sup_{\lambda_1} \text{Cost}(\text{Out}(v, \lambda_0^*, \lambda_1)) = \inf_{\lambda_0} \sup_{\lambda_1} \text{Cost}(\text{Out}(v, \lambda_0, \lambda_1)).$$

# Timed Game

Pl. 0 aims at reaching **G** while minimising the time.



A strategy for Pl. 0 ensuring 10 t.u. from  $(A, 0, 0)$

## Strategy for Pl. 0 ensuring a time $T$

A strategy  $\lambda_0$  ensures a time  $T$  for Pl. 0 from  $(\ell, 0)$  iff

$$\sup_{\lambda_1} \text{Time}(\text{Out}((\ell, 0), \lambda_0, \lambda_1)) \leq T.$$


# About optimal strategies in Timed Game

## Optimal Strategy for Pl. 0

A strategy  $\lambda_0^*$  is optimal for Pl. 0 from  $(\ell, 0)$  iff

$$\sup_{\lambda_1} \text{Time}(\text{Out}((\ell, 0), \lambda_0^*, \lambda_1)) = \inf_{\lambda_0} \sup_{\lambda_1} \text{Time}(\text{Out}((\ell, 0), \lambda_0, \lambda_1)).$$

Optimal strategies do not always exist in timed game!!!



Pl. 0 can choose to wait any  $t > 0$  before reaching **G**.

$$\inf_{\lambda_0} \sup_{\lambda_1} \text{Time}(\text{Out}((A, 0), \lambda_0, \lambda_1)) = \inf_{t > 0} t = 0.$$

However, there is no strategy  $\lambda_0^*$  such that


$$\sup_{\lambda_1} \text{Time}(\text{Out}((A, 0), \lambda_0^*, \lambda_1)) = 0.$$

# About optimal strategies in Timed Game (continued)

## $\epsilon$ -optimal Strategy for Pl. 0

Given  $\epsilon > 0$ , a strategy  $\lambda_0^*$  is  $\epsilon$ -optimal for Pl. 0 from  $(\ell, 0)$  iff

$$\sup_{\lambda_1} \text{Time}(\text{Out}((\ell, 0), \lambda_0^*, \lambda_1)) \leq \inf_{\lambda_0} \sup_{\lambda_1} \text{Time}(\text{Out}((\ell, 0), \lambda_0, \lambda_1)) + \epsilon.$$



There is no optimal strategy for Pl. 0 from  $(A, 0)$ .

But for all  $\epsilon > 0$ , there is an  $\epsilon$ -optimal strategy for Pl. 0 from  $(A, 0)$ .




# About optimal strategies in Timed Game (continued)

## $\epsilon$ -optimal Strategy for Pl. 0

Given  $\epsilon > 0$ , a strategy  $\lambda_0^*$  is  $\epsilon$ -optimal for Pl. 0 from  $(\ell, 0)$  iff

$$\sup_{\lambda_1} \text{Time}(\text{Out}((\ell, 0), \lambda_0^*, \lambda_1)) \leq \inf_{\lambda_0} \sup_{\lambda_1} \text{Time}(\text{Out}((\ell, 0), \lambda_0, \lambda_1)) + \epsilon.$$



There is no optimal strategy for Pl. 0 from  $(A, 0)$ .


But for all  $\epsilon > 0$ , there is an  $\epsilon$ -optimal strategy for Pl. 0 from  $(A, 0)$ .

## Remark

The classical *region* of timed automata is a right tool to solve timed game.

# Priced Timed Game

Pl. 0 aims at reaching **G** while minimising the cost.



A strategy for Pl. 0 ensuring 52 from  $(A, 0, 0)$

## Strategy for Pl. 0 ensuring a cost $K$

A strategy  $\lambda_0$  ensures a cost  $K$  for Pl. 0 from  $(\ell, 0)$  iff

$$\sup_{\lambda_1} \text{Cost}(\text{Out}((\ell, 0), \lambda_0, \lambda_1)) \leq K.$$

# About optimal strategies in Priced Timed Game


## Remark

Clearly optimal strategies do not always exist in Priced Timed Game

# About optimal strategies in Priced Timed Game

## Remark

Clearly optimal strategies do not always exist in Priced Timed Game




$$\inf_{\lambda_0} \sup_{\lambda_1} \text{Cost}(\text{Out}((A, 0), \lambda_0, \lambda_1)) = \inf_{0 \leq t \leq 1} \max\{5t + 7(1 - t), 5t + 3(2 - t)\}$$

# About optimal strategies in Priced Timed Game

## Remark

Clearly optimal strategies do not always exist in Priced Timed Game



$$\begin{aligned} \inf_{\lambda_0} \sup_{\lambda_1} \text{Cost}(\text{Out}((A, 0), \lambda_0, \lambda_1)) &= \inf_{0 \leq t \leq 1} \max\{5t + 7(1 - t), 5t + 3(2 - t)\} \\ &= \inf_{0 \leq t \leq 1} \max\{-2t + 7, 2t + 6\} = 6.5 \end{aligned}$$

The optimal strategy for Pl. 0 asks to take the transition after  $\frac{1}{4}$  t.u.

# Outline of the talk

- 1 Priced Timed Games and Optimal Strategies
- 2 Existing results
- 3 Going further...

# The $K$ -bounded problem

## The $K$ -bounded Problem

Given  $\mathcal{A}$  a PTG and  $K \in \mathbb{N}$ , decide whether there exists  $\lambda_0^*$  such that

$$\sup_{\lambda_1} \text{Cost}(\text{Out}(\ell_0, 0), \lambda_0^*, \lambda_1)) \leq K.$$

# Decidability results

The  $K$ -bounded problem is **decidable** on

- **Timed Games**

E. Asarin and O. Maler. As soon as possible : Time optimal control for timed automata. 1999

- **Priced Timed Games** under strong non-Zenoness of the cost

R. Alur, M. Bernadsky, and P. Madhusudan. Optimal reachability for weighted timed games. 2004

P. Bouyer, F. Cassez, E. Fleury, and K. G. Larsen. Optimal strategies in priced timed game automata. 2004

- **Priced Timed Games** with one clock

P. Bouyer, K. Larsen, N. Markey, and J. Rasmussen. Almost optimal strategies in one clock priced timed games. 2006

The value  $\inf_{\lambda_0} \sup_{\lambda_1} \text{Cost}(\text{Out}((\ell_0, 0), \lambda_0, \lambda_1))$  can be computed.



# Undecidability results

The  $K$ -bounded Problem is **undecidable** on

- **Priced Timed Games** with 6 clocks and non-negative prices.

T. Brihaye, V. Bruyère, J.-F. Raskin. On optimal timed strategies. 2005.

- **Priced Timed Games** with 3 clocks and non-negative prices.

P. Bouyer, T. Brihaye, N. Markey. Improved Undecidability Results on Weighted Timed Automata. 2006

# Outline of the talk

- 1 Priced Timed Games and Optimal Strategies
- 2 Existing results
- 3 Going further...

## Questions still open...

- What about the variants of the  $K$ -bounded Problem?

### The $K$ -bounded Problem (strict version)

Given  $\mathcal{A}$  a PTG and  $K \in \mathbb{N}$ , decide whether there exists  $\lambda_0^*$  such that

$$\sup_{\lambda_1} \text{Cost}(\text{Out}((\ell_0, 0), \lambda_0^*, \lambda_1)) < K.$$

### The $K$ -bounded Problem ( $\epsilon$ -version)

Given  $\mathcal{A}$  a PTG and  $K \in \mathbb{N}$ , decide whether for all  $\epsilon > 0$ , there exists  $\lambda_0^\epsilon$  such that

$$\sup_{\lambda_1} \text{Cost}(\text{Out}((\ell_0, 0), \lambda_0^\epsilon, \lambda_1)) \leq K + \epsilon.$$

- What happens if we consider concurrent games, positive costs, ...?

# The time-bounded framework

**Undecidable** problems become **decidable** when considering their **time-bounded** version :

- Time-bounded language inclusion for Timed Automata

J. Ouaknine, A. Rabinovich, and J. Worrell. Time-bounded verification. 2009

- Reachability for Hybrid Automata

T. Brihaye, L. Doyen, G. Geeraerts, J. Ouaknine, J.-F. Raskin, J. Worrell : On Reachability for Hybrid Automata over Bounded Time. 2011

# The time-bounded framework

**Undecidable** problems become **decidable** when considering their **time-bounded** version :

- Time-bounded language inclusion for Timed Automata

J. Ouaknine, A. Rabinovich, and J. Worrell. Time-bounded verification. 2009

- Reachability for Hybrid Automata

T. Brihaye, L. Doyen, G. Geeraerts, J. Ouaknine, J.-F. Raskin, J. Worrell : On Reachability for Hybrid Automata over Bounded Time. 2011

Would this work for the  $K$ -bounded Problem on PTG?

## New (undecidability) results

- The **time-bounded**,  $K$ -bounded Pbm is **undecidable** on **PTG**.
- The **strict version of** the  $K$ -bounded Pbm is **undecidable** on **PTG**.
- The  $\epsilon$ -**version of** the  $K$ -bounded Pbm is **undecidable** on **PTG**.
- The  $K$ -bounded Pbm is **undecidable** on **concurrent PTG with 2 clocks**.

## New (undecidability) results

- The **time-bounded**,  $K$ -bounded Pbm is **undecidable** on **PTG**.
- The **strict version of** the  $K$ -bounded Pbm is **undecidable** on **PTG**.
- The  $\epsilon$ -**version of** the  $K$ -bounded Pbm is **undecidable** on **PTG**.
- The  $K$ -bounded Pbm is **undecidable** on **concurrent PTG with 2 clocks**.

We hope to obtain :

- a precise characterisation of the decidability border,
- a new **decidability result** (with positive cost, few clocks, and ???)

Thank you !!!