

Greibach Normal Form for ω -Algebraic Systems and Weighted Simple ω -Pushdown Automata

Manfred Droste^a, Sven Dziadek^{a,1}, Werner Kuich^{b,2}

^a*Institut für Informatik, Universität Leipzig, Germany*

^b*Institut für Diskrete Mathematik und Geometrie, Technische Universität Wien, Austria*

Abstract

In weighted automata theory, many classical results on formal languages have been extended into a quantitative setting. Here, we investigate weighted context-free languages of infinite words, a generalization of ω -context-free languages (Cohen, Gold 1977) and an extension of weighted context-free languages of finite words (Chomsky, Schützenberger 1963). As in the theory of formal grammars, these weighted context-free languages, or ω -algebraic series, can be represented as solutions of mixed ω -algebraic systems of equations and by weighted ω -pushdown automata.

In our first main result, we show that (mixed) ω -algebraic systems can be transformed into *Greibach normal form*. We use the Greibach normal form in our second main result to prove that *simple ω -reset pushdown automata* recognize all ω -algebraic series. Simple ω -reset automata do not use ϵ -transitions and can change the stack only by at most one symbol. These results generalize fundamental properties of context-free languages to weighted context-free languages.

Keywords: Greibach normal form, weighted automata, omega-pushdown automata, omega context-free languages

1. Introduction

Context-free languages provide a fundamental concept for programming languages in computer science. In order to model quantitative properties, already in 1963, Chomsky and Schützenberger [3] introduced weighted context-free languages. The theory of weighted pushdown automata has been extensively studied; for background, we refer the reader to the survey [21] and the books [22, 18, 11]. In 1977, Cohen and Gold [4] investigated context-free languages of infinite words. Weighted ω -context-free languages, i.e., ω -algebraic series were studied by Ésik and Kuich [16].

The goal of this paper is the investigation of weighted context-free languages and weighted pushdown automata on infinite words. As in [18], the weighted context-free languages of infinite words are described by solutions of ω -algebraic systems and mixed ω -algebraic systems of equations. In our first main result, we show that these systems can be transformed into a Greibach normal form. In the literature, Greibach normal forms, central for the theory of context-free languages of finite words, have been established for ω -context-free languages (of infinite words), see [4], and also for algebraic systems of equations for series over finite words [22, 18]; this latter result is employed in our proof. Hence here we extend these classical results to a weighted version for infinite words.

Recently, Droste, Ésik and Kuich introduced weighted ω -pushdown automata in [9, 10]. In our second main result, we consider weighted *simple* ω -pushdown automata that we call simple ω -reset pushdown automata here. These automata do not use ϵ -transitions and utilize only three simple stack commands: popping a symbol, pushing a symbol or leaving the stack unaltered; moreover, it is only possible to read the topmost stack symbol by popping it. Observe that together with the restriction of not allowing ϵ -transitions, restrictions for the actions on the stack are

Email addresses: droste@informatik.uni-leipzig.de (Manfred Droste), dziadek@informatik.uni-leipzig.de (Sven Dziadek), werner.kuich@tuwien.ac.at (Werner Kuich)

¹Supported by Deutsche Forschungsgemeinschaft (DFG), Graduiertenkolleg 1763 (QuantLA)

²Partially supported by Austrian Science Fund (FWF): grant no. I1661 N25

non-trivial. In our second main result we show that these *simple* ω -reset pushdown automata recognize all weighted ω -context-free languages. For our proof, we use that ω -algebraic systems can be brought into Greibach normal form by our present first main result. Our construction of simple ω -reset pushdown automata is deduced from the construction used in a recent corresponding result [6] that states that simple reset pushdown automata on finite words recognize all algebraic series.

We believe the model of simple ω -reset pushdown automata to be very natural. Similar expressivity equivalence results in the unweighted case hold for context-free languages of finite words, hidden in a proof by Blass and Gurevich [1], and also for ω -context-free languages, see [8].

After the Preliminaries, in Section 3, we characterize ω -algebraic series by a series of equivalent statements.

In Section 4, we define mixed ω -algebraic systems and their canonical solutions; l^{th} canonical solutions are by definition unique. This allows us to perform the following proof method in Section 7: The proof that each of two expressions is the m^{th} component of the l^{th} canonical solution implies the equality of these two expressions. (Compare this with the proof method in continuous semirings: The proof that each of two expressions is the m^{th} component of the least solution of an algebraic system implies the equality of these two expressions.) The main result of Section 4 states that each ω -algebraic series is a component of a canonical solution of a mixed ω -algebraic system in Greibach normal form.

In Section 5 we specialize the main result of Section 4: now each ω -algebraic series is a component of a canonical solution of an ω -algebraic system in Greibach normal form.

We consider simple reset pushdown automata in Section 6 and recall the result of [6] that for each algebraic series r there exists a simple reset pushdown automaton with behavior r .

Simple ω -reset pushdown automata are introduced in Section 7. The main result of this section and of the whole paper is that for each ω -algebraic series r it is possible to construct a simple ω -reset pushdown automaton with behavior r . The proof of this main result is performed by our proof method described above. We consider an ω -algebraic series that is the m^{th} component of the l^{th} canonical solution of an ω -algebraic system in Greibach normal form and construct a simple ω -reset pushdown automaton whose moves depend only on the coefficients of this Greibach normal form. We prove that the behavior of this simple ω -reset pushdown automaton equals the m^{th} component of the l^{th} canonical solution of this Greibach normal form. Hence, we conclude that for each ω -algebraic series r we can construct a simple ω -reset pushdown automaton with behavior r .

A preliminary version of this paper appeared in [7]. In this version, we strengthen the first main result by proving that already ω -algebraic systems can be transformed into Greibach normal form. In [7], we only showed the existence of the Greibach normal form for mixed ω -algebraic systems. The stronger result in this work allows us to generalize the second main result: weighted simple ω -pushdown recognize all ω -algebraic series. For this, we needed to adapt the construction such that our simple ω -reset pushdown automata behave exactly like the canonical solutions of ω -algebraic systems. Furthermore, we add a result (see Theorem 2) describing ω -powers of matrices considering Büchi-acceptance. We give complete arguments and further examples for our results.

2. Preliminaries

For the convenience of the reader, we recall definitions and results from Ésik, Kuich [18].

A monoid $\langle S, +, 0 \rangle$ is called *complete* if it is equipped with sum operations \sum_I for all families $(a_i \mid i \in I)$ of elements of S , where I is an arbitrary index set, such that the following conditions are satisfied (see Conway [5], Eilenberg [14], Kuich [21]):

- (i) $\sum_{i \in \emptyset} a_i = 0$, $\sum_{i \in \{j\}} a_i = a_j$, $\sum_{i \in \{j,k\}} a_i = a_j + a_k$ for $j \neq k$,
- (ii) $\sum_{j \in J} (\sum_{i \in I_j} a_i) = \sum_{i \in I} a_i$, if $\bigcup_{j \in J} I_j = I$ and $I_j \cap I_{j'} = \emptyset$ for $j \neq j'$.

Furthermore, a semiring $\langle S, +, \cdot, 0, 1 \rangle$ is called *complete* if $\langle S, +, 0 \rangle$ is a complete monoid and if we additionally have

- (iii) $\sum_{i \in I} (c \cdot a_i) = c \cdot (\sum_{i \in I} a_i)$, $\sum_{i \in I} (a_i \cdot c) = (\sum_{i \in I} a_i) \cdot c$.

This means that a semiring S is complete if it has “infinite sums” (i) that are an extension of the finite sums, (ii) that are associative and commutative and (iii) that satisfy the distributivity laws.

A semiring S equipped with an additional unary star operation $*$: $S \rightarrow S$ is called a *starsemiring*. In complete semirings for each element a , the *star* a^* of a is defined by

$$a^* = \sum_{j \geq 0} a^j.$$

Hence, each complete semiring is a starsemiring, called a *complete starsemiring*.


Starsemirings allow us to generalize the star operation to matrices. Let $M \in S^{n \times n}$, then we define $M^* \in S^{n \times n}$ inductively as in Ésik, Kuich [18], pp. 14–15 as follows. For $n = 1$ and $M = (a)$, for $a \in S$, we let $M^* = (a^*)$. Now, for $n > 1$, we partition M into submatrices, called *blocks*,

$$M = \begin{pmatrix} a & b \\ c & d \end{pmatrix}, \quad (1)$$

with $a \in S^{1 \times 1}$, $b \in S^{1 \times (n-1)}$, $c \in S^{(n-1) \times 1}$, $d \in S^{(n-1) \times (n-1)}$, and we define

$$M^* = \begin{pmatrix} (a + bd^*c)^* & (a + bd^*c)^*bd^* \\ (d + ca^*b)^*ca^* & (d + ca^*b)^* \end{pmatrix}. \quad (2)$$

Whenever we use a matrix M as defined in (1), the corresponding automaton can be illustrated as follows:



A semiring is called *continuous* if it is ordered, each directed subset has a least upper bound and addition and multiplication preserve the least upper bound of directed sets. Any continuous semiring is complete. See Ésik, Kuich [18] for background.

Suppose that S is a semiring and V is a commutative monoid written additively. We call V a (left) S -semimodule if V is equipped with a (left) action

$$\begin{aligned} S \times V &\rightarrow V \\ (s, v) &\mapsto sv \end{aligned}$$

subject to the following rules:

$$\begin{aligned} s(s'v) &= (ss')v, & (s + s')v &= sv + s'v, & s(v + v') &= sv + sv', \\ 1v &= v, & 0v &= 0, & s0 &= 0, \end{aligned}$$

for all $s, s' \in S$ and $v, v' \in V$. If V is an S -semimodule, we call (S, V) a *semiring-semimodule pair*.

Suppose that (S, V) is a semiring-semimodule pair such that S is a starsemiring and S and V are equipped with an omega operation $^\omega : S \rightarrow V$. Then we call (S, V) a *starsemiring-omegasemimodule pair*.

Ésik, Kuich [19] define a *complete semiring-semimodule pair* to be a semiring-semimodule pair (S, V) such that S is a complete semiring and V is a complete monoid with

$$s\left(\sum_{i \in I} v_i\right) = \sum_{i \in I} sv_i \quad \text{and} \quad \left(\sum_{i \in I} s_i\right)v = \sum_{i \in I} s_iv,$$

for all $s \in S$, $v \in V$, and for all families $(s_i)_{i \in I}$ over S and $(v_i)_{i \in I}$ over V ; moreover, it is required that an *infinite product operation*

$$S^\omega \ni (s_1, s_2, \dots) \mapsto \prod_{j \geq 1} s_j \in V$$

is given mapping infinite sequences over S to V subject to the following three conditions:

$$\begin{aligned} \text{(i)} \quad & \prod_{i \geq 1} s_i = \prod_{i \geq 1} (s_{n_{i-1}+1} \cdots s_{n_i}), \\ \text{(ii)} \quad & s_1 \cdot \prod_{i \geq 1} s_{i+1} = \prod_{i \geq 1} s_i, \\ \text{(iii)} \quad & \prod_{j \geq 1} \sum_{i_j \in I_j} s_{i_j} = \sum_{(i_1, i_2, \dots) \in I_1 \times I_2 \times \dots} \prod_{j \geq 1} s_{i_j}, \end{aligned}$$

where in the first equation $0 = n_0 \leq n_1 \leq n_2 \leq \dots$ and I_1, I_2, \dots are arbitrary index sets. This means that the left action of the semimodule is distributive and it is required that it has “infinite products” mapping infinite sequences over S to V such that the product (i) can be partitioned (an infinite form of associativity), (ii) can be extended from the left and (iii) satisfies an infinite distributivity law.

Suppose that (S, V) is complete. Then we define

$$s^* = \sum_{i \geq 0} s^i \quad \text{and} \quad s^\omega = \prod_{i \geq 1} s,$$

for all $s \in S$. This turns (S, V) into a starsemiring-omegasemimodule pair. Observe that, if (S, V) is a complete semiring-semimodule pair, then $0^\omega = 0$.

A *star-omega semiring* is a semiring S equipped with unary operations $*$ and $^\omega : S \rightarrow S$. A star-omega semiring S is called *complete* if (S, S) is a complete semiring-semimodule pair, i.e., if S is complete and is equipped with an infinite product operation that satisfies the three conditions stated above. A complete star-omega semiring S is called *continuous* if the semiring S is continuous.

Example 1. Formal languages are covered by our model. Let $\langle \mathbb{B}, +, \cdot, 0, 1 \rangle$ be the Boolean semiring. Then let $0^* = 1^* = 1$ and take infima as infinite products. This makes \mathbb{B} a continuous star-omega and commutative semiring. It then follows that $\mathbb{B}\langle\langle \Sigma^* \rangle\rangle \times \mathbb{B}\langle\langle \Sigma^\omega \rangle\rangle$ is isomorphic to formal languages of finite and infinite words with the usual operations.

The semiring $\langle \mathbb{N}^\infty, +, \cdot, 0, 1 \rangle$ with $\mathbb{N}^\infty = \mathbb{N} \cup \{\infty\}$ and the natural infinite product operation of numbers is a continuous star-omega and commutative semiring.

The tropical semiring $\langle \mathbb{N}^\infty, \min, +, \infty, 0 \rangle$ with the usual infinite sum operation as infinite product is a commutative semiring and a continuous star-omega semiring.

Analogously, the arctic semiring $\langle \bar{\mathbb{N}}, \max, +, -\infty, 0 \rangle$ with $\bar{\mathbb{N}} = \mathbb{N} \cup \{-\infty, \infty\}$ and the infinite sum operation as infinite product is a commutative semiring and a continuous star-omega semiring.

A *Conway semiring* (see Conway [5], Bloom, Ésik [2]) is a starsemiring S satisfying the *sum star identity*

$$(a + b)^* = a^*(ba^*)^*$$

and the *product star identity*

$$(ab)^* = 1 + a(ba)^*b$$

for all $a, b \in S$. Observe that by Ésik, Kuich [18], Theorem 1.2.24, each complete starsemiring is a Conway semiring.

Note that from the identities in Conway semirings, it follows

$$\begin{aligned} a^* &= 1 + aa^* = 1 + a^*a, \\ a(ba)^* &= (ab)^*a, \end{aligned} \tag{3}$$

for all $a, b \in S$.

If S is a Conway semiring then so is $S^{n \times n}$. Let $M \in S^{n \times n}$. Assume that $n > 1$ and write M as in (1). Applying the identities of Conway semirings, we get an equivalent definition (cf. Conway [5], pp. 27–28) to (2):

$$M^* = \begin{pmatrix} (a + bd^*c)^* & a^*b(d + ca^*b)^* \\ d^*c(a + bd^*c)^* & (d + ca^*b)^* \end{pmatrix}. \tag{4}$$

Following Bloom, Ésik [2], we call a starsemiring-omegasemimodule pair (S, V) a *Conway semiring-semimodule pair* if S is a Conway semiring and if the omega operation satisfies the *sum omega identity* and the *product omega identity*:

$$(a + b)^\omega = (a^*b)^\omega + (a^*b)^*a^\omega \quad \text{and} \quad (ab)^\omega = a(ba)^\omega,$$

for all $a, b \in S$. By Ésik, Kuich [19] each complete semiring-semimodule pair is a Conway semiring-semimodule pair.

Observe that the *omega fixed-point equation* holds, i.e.

$$aa^\omega = a^\omega,$$

for all $a \in S$.

Consider a starsemiring-omegasemimodule pair (S, V) . Following Bloom, Ésik [2], we define a matrix operation $\omega: S^{n \times n} \rightarrow V^{n \times 1}$ on a starsemiring-omegasemimodule pair (S, V) as follows. If $n = 0$, M^ω is the unique element of V^0 , and if $n = 1$, so that $M = (a)$, for some $a \in S$, $M^\omega = (a^\omega)$. Assume now that $n > 1$ and write M as in (1). Then

$$M^\omega = \begin{pmatrix} (a + bd^*c)^\omega + (a + bd^*c)^*bd^\omega \\ (d + ca^*b)^\omega + (d + ca^*b)^*ca^\omega \end{pmatrix}.$$

Additionally, the *matrix star identity* is valid for Conway semirings and states that the star of a matrix is independent of the partitioning of the matrix. The *matrix omega identity* is valid for Conway semiring-semimodule pairs and states that the operation $^\omega$ is independent of the partitioning of the matrix, i.e., the blocks of (1) can have arbitrary sizes: $a \in S^{n_1 \times n_1}$, $b \in S^{n_1 \times n_2}$, $c \in S^{n_2 \times n_1}$, $d \in S^{n_2 \times n_2}$ for $n_1 + n_2 = n$. If (S, V) is a Conway semiring-semimodule pair, then so is $(S^{n \times n}, V^n)$. See also Ésik, Kuich [18], page 106.

Following Ésik, Kuich [17], we define matrix operations $\omega, t: S^{n \times n} \rightarrow V^{n \times 1}$ for $0 \leq t \leq n$ as follows. Assume that $M \in S^{n \times n}$ is decomposed into blocks a, b, c, d as in (1), but with a of dimension $t \times t$ and d of dimension $(n-t) \times (n-t)$. Then

$$M^{\omega, t} = \begin{pmatrix} (a + bd^*c)^\omega \\ d^*c(a + bd^*c)^\omega \end{pmatrix}. \quad (5)$$

Observe that $M^{\omega, 0} = 0$ and $M^{\omega, n} = M^\omega$. Intuitively, M can be interpreted as an adjacency matrix and $M^{\omega, t}$ are infinite paths where the first t states are repeated states, i.e., states that are Büchi-accepting.

The next theorem states that, in case of a Conway semiring, $M^{\omega, t}$, for $0 \leq t \leq n$, can be computed also in a way different from its definition and, with certain limits, is independent of the partitioning of the matrix M .

Theorem 2. *Let S be a Conway semiring and $0 \leq t \leq k \leq n$. Assume $M \in S^{n \times n}$ is decomposed into blocks*

$$M = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$$

with block a being of dimension $k \times k$ and block d of dimension $(n-k) \times (n-k)$.

Then we have,

$$M^{\omega, t} = \begin{pmatrix} (a + bd^*c)^{\omega, t} \\ d^*c(a + bd^*c)^{\omega, t} \end{pmatrix}. \quad (6)$$

Proof. The proof resembles the proof of the matrix omega identity (cf. [18], Theorem 5.3.13). Assume $M \in S^{n \times n}$ is decomposed into nine blocks

$$M = \begin{pmatrix} f & g & h \\ i & a & b \\ j & c & d \end{pmatrix}$$

with dimensions $f \in S^{t \times t}$, $a \in S^{(k-t) \times (k-t)}$ and $d \in S^{(n-k) \times (n-k)}$. Consider the following two partitionings:

$$M = \left(\begin{array}{c|cc} f & g & h \\ \hline i & a & b \\ \hline j & c & d \end{array} \right) \quad M' = \left(\begin{array}{cc|c} f & g & h \\ \hline i & a & b \\ \hline j & c & d \end{array} \right)$$

Now we need to show that $M^{\omega,t}$, calculated as in (5)

$$M^{\omega,t} = \left(\begin{array}{c} \alpha \\ \left(\begin{array}{cc} a & b \\ c & d \end{array} \right)^* \begin{pmatrix} i \\ j \end{pmatrix} \alpha \end{array} \right),$$

where

$$\alpha = \left(f + \begin{pmatrix} g & h \end{pmatrix} \begin{pmatrix} a & b \\ c & d \end{pmatrix}^* \begin{pmatrix} i \\ j \end{pmatrix} \right)^\omega$$

is equal to $M'^{\omega,t}$, calculated as in (6)

$$M'^{\omega,t} = \left(d^* \begin{pmatrix} \mu \\ j & c \end{pmatrix} \mu \right),$$

where

$$\mu = \left(\begin{pmatrix} f & g \\ i & a \end{pmatrix} + \begin{pmatrix} h \\ b \end{pmatrix} d^* \begin{pmatrix} j & c \end{pmatrix} \right)^{\omega,t}.$$

In the case $t = k$, we have

$$M = M' = \begin{pmatrix} f & h \\ j & d \end{pmatrix}.$$

It follows that

$$\begin{aligned} \alpha &= (f + hd^*j)^\omega \\ &= (f + hd^*j)^{\omega,t} = \mu, \end{aligned}$$

where the second equality is due to t being the full dimension of $f + hd^*j$. The second components of $M^{\omega,t}$ and $M'^{\omega,t}$ then both reduce to $d^*j(f + hd^*j)^\omega$.

If $k = n$, we have

$$M = M' = \begin{pmatrix} f & g \\ i & a \end{pmatrix}.$$

Now, the second component of $M'^{\omega,t}$ and the second summand of μ have dimension 0 and thus

$$M'^{\omega,t} = \begin{pmatrix} f & g \\ i & a \end{pmatrix}^{\omega,t} = M^{\omega,t}$$

Hence, in the following, we can assume $t < k < n$.

First, we compute $M^{\omega,t}$. We denote the blocks of $M^{\omega,t}$ by $(M^{\omega,t})_i$ for $1 \leq i \leq 3$. Then we have

$$\begin{aligned} (M^{\omega,t})_1 &= \alpha = \left(f + \begin{pmatrix} g & h \end{pmatrix} \begin{pmatrix} a & b \\ c & d \end{pmatrix}^* \begin{pmatrix} i \\ j \end{pmatrix} \right)^\omega \\ &= \left(f + \begin{pmatrix} g & h \end{pmatrix} \begin{pmatrix} (a + bd^*c)^* & a^*b(d + ca^*b)^* \\ d^*c(a + bd^*c)^* & (d + ca^*b)^* \end{pmatrix} \begin{pmatrix} i \\ j \end{pmatrix} \right)^\omega \\ &= \left(f + \begin{pmatrix} g & h \end{pmatrix} \begin{pmatrix} (a + bd^*c)^*i + a^*b(d + ca^*b)^*j \\ d^*c(a + bd^*c)^*i + (d + ca^*b)^*j \end{pmatrix} \right)^\omega \\ &= (f + g(a + bd^*c)^*i + ga^*b(d + ca^*b)^*j \\ &\quad + hd^*c(a + bd^*c)^*i + h(d + ca^*b)^*j)^\omega. \end{aligned}$$

Here, we used the star of a matrix in the form shown in (4). We will now compute the other two blocks by using the star of a matrix as in (2):

$$\begin{aligned}
\begin{pmatrix} (M^{\omega,t})_2 \\ (M^{\omega,t})_3 \end{pmatrix} &= \begin{pmatrix} a & b \\ c & d \end{pmatrix}^* \begin{pmatrix} i \\ j \end{pmatrix} \alpha \\
&= \begin{pmatrix} (a + bd^*c)^* & (a + bd^*c)^*bd^* \\ (d + ca^*b)^*ca^* & (d + ca^*b)^* \end{pmatrix} \begin{pmatrix} i \\ j \end{pmatrix} \alpha \\
&= \begin{pmatrix} (a + bd^*c)^*i + (a + bd^*c)^*bd^*j \\ (d + ca^*b)^*ca^*i + (d + ca^*b)^*j \end{pmatrix} \alpha \\
&= \begin{pmatrix} ((a + bd^*c)^*i + (a + bd^*c)^*bd^*j)\alpha \\ ((d + ca^*b)^*ca^*i + (d + ca^*b)^*j)\alpha \end{pmatrix}
\end{aligned}$$

Now, we compute $M'^{\omega,t}$. We denote the blocks of $M'^{\omega,t}$ by $(M'^{\omega,t})_i$ for $1 \leq i \leq 3$. Then we have

$$\begin{aligned}
\begin{pmatrix} (M'^{\omega,t})_1 \\ (M'^{\omega,t})_2 \end{pmatrix} &= \mu = \left(\begin{pmatrix} f & g \\ i & a \end{pmatrix} + \begin{pmatrix} h \\ b \end{pmatrix} d^* \begin{pmatrix} j & c \end{pmatrix} \right)^{\omega,t} \\
&= \left(\begin{pmatrix} f & g \\ i & a \end{pmatrix} + \begin{pmatrix} hd^*j & hd^*c \\ bd^*j & bd^*c \end{pmatrix} \right)^{\omega,t} \\
&= \begin{pmatrix} f + hd^*j & g + hd^*c \\ i + bd^*j & a + bd^*c \end{pmatrix}^{\omega,t} \\
&= \begin{pmatrix} \delta \\ (a + bd^*c)^*(i + bd^*j)\delta \end{pmatrix},
\end{aligned}$$

where

$$\delta = (f + hd^*j + (g + hd^*c)(a + bd^*c)^*(i + bd^*j))^\omega.$$

It remains to calculate

$$\begin{aligned}
(M'^{\omega,t})_3 &= d^* \begin{pmatrix} j & c \end{pmatrix} \mu \\
&= d^*(j + c(a + bd^*c)^*(i + bd^*j))\delta.
\end{aligned}$$

The last step is to verify the three equalities $(M^{\omega,t})_i = (M'^{\omega,t})_i$ for $1 \leq i \leq 3$. The first equality follows basically from Lemma 1.2.16 of [18]. We will mark the use of Lemma 1.2.16 by \diamond and obtain

$$\begin{aligned}
(M^{\omega,t})_1 &= \alpha = (f + g(a + bd^*c)^*i + ga^*b(d + ca^*b)^*j \\
&\quad + hd^*c(a + bd^*c)^*i + h(d + ca^*b)^*j)^\omega \\
&\stackrel{\diamond}{=} (f + hd^*j + g(a + bd^*c)^*i + g(a + bd^*c)^*bd^*j \\
&\quad + hd^*c(a + bd^*c)^*i + hd^*c(a + bd^*c)^*bd^*j)^\omega \\
&= (f + hd^*j + g(a + bd^*c)^*(i + bd^*j) + hd^*c(a + bd^*c)^*(i + bd^*j))^\omega \\
&= (f + hd^*j + (g + hd^*c)(a + bd^*c)^*(i + bd^*j))^\omega \\
&= \delta = (M'^{\omega,t})_1
\end{aligned}$$

For the second equality, we have

$$\begin{aligned}
(M^{\omega,t})_2 &= ((a + bd^*c)^*i + (a + bd^*c)^*bd^*j)\alpha \\
&= ((a + bd^*c)^*(i + bd^*j))\delta \\
&= (M'^{\omega,t})_2.
\end{aligned}$$

Now, for the third equality, it suffices to prove

$$(d + ca^*b)^*ca^*i + (d + ca^*b)^*j = d^*(j + c(a + bd^*c)^*(i + bd^*j)).$$

We have

$$\begin{aligned} d^*(j + c(a + bd^*c)^*(i + bd^*j)) &= d^*j + d^*c(a + bd^*c)^*(i + bd^*j) \\ &= d^*j + d^*c(a + bd^*c)^*i + d^*c(a + bd^*c)^*bd^*j \\ &= d^*j + d^*c(a^*bd^*c)^*a^*i + d^*c(a^*bd^*c)^*a^*bd^*j \\ &= d^*j + (d^*ca^*b)^*d^*ca^*i + (d^*ca^*b)^*d^*ca^*bd^*j \\ &= (d^*ca^*b)^*d^*ca^*i + (d^*ca^*b)^*d^*ca^*bd^*j + d^*j \\ &= (d^*ca^*b)^*d^*ca^*i + ((d^*ca^*b)^*d^*ca^*b + 1)d^*j \\ &= (d^*ca^*b)^*d^*ca^*i + (d^*ca^*b)^*d^*j \\ &= (d + ca^*b)^*ca^*i + (d + ca^*b)^*j. \end{aligned}$$

Note that for this calculation, we rely heavily on commutativity of addition, distributivity and the sum star identity and the product star identity of Conway semirings together with their derived identities (3). This completes the proof. \square

For a complete definition of quemirings, we refer the reader to [18], page 110. Here we note that a quemiring T is isomorphic to a quemiring $S \times V$ determined by the semiring-semimodule pair (S, V) ; it follows that we can identify every element t of a quemiring T by a pair (s, v) of a semiring-semimodule pair (S, V) . A quemiring is an algebraic structure with an addition given componentwise, i.e.,

$$(s, v) + (s', v') = (s + s', v + v'),$$

a semidirect product type multiplication (using that S acts on V), i.e.,

$$(s, v) \cdot (s', v') = (ss', v + sv'),$$

and two constants $0 = (0, 0)$ and $1 = (1, 0)$ (and a unary operation \mathbb{I} , but we will not use it here). A quemiring $S \times V$ satisfies a set of axioms inherited from semiring-semimodule pairs; those axioms make a quemiring *quasi a semiring* (cf. Elgot [15], Ésik, Kuich [18], page 109; in fact, a quemiring is not necessarily distributive from the left and 0 only behaves like a zero from the left). Also, one can define a natural star operation on $S \times V$, i.e.,

$$(s, v)^\otimes = (s^*, s^\omega + s^*v),$$

making it a *generalized starquemiring*, see [18].

For an alphabet Σ , we call mappings r of Σ^* into S *series*. The collection of all such series r is denoted by $S\langle\langle\Sigma^*\rangle\rangle$. We call the set $\text{supp}(r) = \{w \mid (r, w) \neq 0\}$ the *support* of a series r . The set of series with finite support $S\langle\Sigma^*\rangle = \{s \in S\langle\langle\Sigma^*\rangle\rangle \mid \text{supp}(s) \text{ is finite}\}$ is called the set of *polynomials*. We denote by $S\langle\Sigma\rangle$, $S\langle\{\epsilon\}\rangle$ and $S\langle\Sigma \cup \{\epsilon\}\rangle$ the series with support in Σ , $\{\epsilon\}$ and $\Sigma \cup \{\epsilon\}$, respectively. Series s with $|\text{supp}(s)| \leq 1$ are called monomials. Note that polynomials are finite sums of monomials.

Mappings of Σ^ω into S are called ω -series and their collection is denoted by $S\langle\langle\Sigma^\omega\rangle\rangle$. See [22, 18] for more information. Examples of monomials in $S\langle\Sigma^*\rangle$ for a semiring $\langle S, +, \cdot, 0, 1 \rangle$ are 0 , w , sw for $s \in S$ and $w \in \Sigma^*$, defined by

$$\begin{aligned} (0, w) &= 0 \text{ for all } w, \\ (w, w) &= 1 \text{ and } (w, w') = 0 \text{ for } w \neq w', \\ (sw, w) &= s \text{ and } (sw, w') = 0 \text{ for } w \neq w'. \end{aligned}$$

3. ω -Algebraic Systems

This and the next two sections describe the Greibach normal form for (mixed) ω -algebraic series.

For this section and the next two sections, Sections 3, 4 and 5, S is a continuous, and therefore complete, star-omega semiring. If we consider $S\langle\langle\Sigma^*\rangle\rangle$ or $S\langle\langle\Sigma^\omega\rangle\rangle$, then we assume additionally that the underlying semiring S is commutative. Let further Σ denote an alphabet.

By Theorem 5.5.5 of Ésik, Kuich [18], $(S\langle\langle\Sigma^*\rangle\rangle, S\langle\langle\Sigma^\omega\rangle\rangle)$ is a complete semiring-semimodule pair, hence a Conway semiring-semimodule pair, satisfying $\epsilon^\omega = 0$. Hence, $S\langle\langle\Sigma^*\rangle\rangle \times S\langle\langle\Sigma^\omega\rangle\rangle$ is a generalized starquemiring.

In the sequel, x, y and z denote vectors of dimension n , i.e., $x = (x_1, \dots, x_n)$, $y = (y_1, \dots, y_n)$ and $z = (z_1, \dots, z_n)$. Later, we will also use z of dimension m . It is clear by the context whether they are used as row or as column vectors. Similar conventions hold for vectors p, σ, ω and τ . Moreover, X denotes the set of variables $\{x_1, \dots, x_n\}$ for $S\langle\langle\Sigma^*\rangle\rangle$, while $\{z_1, \dots, z_n\}$ is the set of variables for $S\langle\langle\Sigma^\omega\rangle\rangle$. The set Y denotes the set of variables $\{y_1, \dots, y_n\}$ for the quemiring $S\langle\langle\Sigma^*\rangle\rangle \times S\langle\langle\Sigma^\omega\rangle\rangle$. We will be working with two different generalizations of ω -context-free grammars, the ω -algebraic systems and the mixed ω -algebraic systems.

An ω -algebraic system over the quemiring $S\langle\langle\Sigma^*\rangle\rangle \times S\langle\langle\Sigma^\omega\rangle\rangle$ consists of an algebraic system over $S\langle\langle\Sigma^*\rangle\rangle \times S\langle\langle\Sigma^\omega\rangle\rangle$

$$y = p(y), \quad p \in (S\langle\langle(\Sigma \cup Y)^*\rangle\rangle)^{n \times 1}.$$

The vector of quemiring elements $\tau \in (S\langle\langle\Sigma^*\rangle\rangle \times S\langle\langle\Sigma^\omega\rangle\rangle)^n$ is a solution of the ω -algebraic system

$$y = p(y),$$

if

$$\tau = p(\tau).$$

Note that every p_i is a polynomial, i.e., a finite sum of monomials in $S\langle\langle(\Sigma \cup Y)^*\rangle\rangle$. Let $y_i = (x_i, z_i)$, for $1 \leq i \leq n$. Now, we can apply the quemiring addition and multiplication to p .

Consider a monomial

$$t(y_1, \dots, y_n) = s w_0 y_{i_1} w_1 \dots w_{k-1} y_{i_k} w_k,$$

where $s \in S$ and $w_i \in \Sigma^*$ for $1 \leq i \leq k$. Note that from the quemiring operations, we have

$$t((x_1, z_1), \dots, (x_n, z_n)) = (s w_0 x_{i_1} w_1 \dots w_{k-1} x_{i_k} w_k, s w_0 z_{i_1} + s w_0 x_{i_1} w_1 z_{i_2} + \dots + s w_0 x_{i_1} w_1 \dots w_{k-2} x_{i_{k-1}} w_{k-1} z_{i_k}).$$

Therefore, following Ésik, Kuich [18], p. 138, we define

$$t_x(x_1, \dots, x_n, z_1, \dots, z_n) = s w_0 z_{i_1} + s w_0 x_{i_1} w_1 z_{i_2} + \dots + s w_0 x_{i_1} w_1 \dots w_{k-2} x_{i_{k-1}} w_{k-1} z_{i_k},$$

and for a polynomial $p(y_1, \dots, y_n) = \sum_{1 \leq j \leq m} t_j(y_1, \dots, y_n)$, we let

$$p_x(x_1, \dots, x_n, z_1, \dots, z_n) = \sum_{1 \leq j \leq m} (t_j)_x(x_1, \dots, x_n, z_1, \dots, z_n).$$

For an ω -algebraic system $y = p(y)$ over $S\langle\langle\Sigma^*\rangle\rangle \times S\langle\langle\Sigma^\omega\rangle\rangle$, we call $x = p(x)$, $z = p_x(x, z)$ the mixed ω -algebraic system over $S\langle\langle\Sigma^*\rangle\rangle \times S\langle\langle\Sigma^\omega\rangle\rangle$ induced by $y = p(y)$.

In general, a mixed ω -algebraic system over the quemiring $S\langle\langle\Sigma^*\rangle\rangle \times S\langle\langle\Sigma^\omega\rangle\rangle$ consists of an algebraic system over $S\langle\langle\Sigma^*\rangle\rangle$

$$x = p(x), \quad p \in (S\langle\langle(\Sigma \cup X)^*\rangle\rangle)^{n \times 1}$$

and a linear system over $S\langle\langle\Sigma^\omega\rangle\rangle$

$$z = \varrho(x)z, \quad \varrho \in (S\langle\langle(\Sigma \cup X)^*\rangle\rangle)^{m \times m}.$$

The pair $(\sigma, \omega) \in (S\langle\langle\Sigma^*\rangle\rangle)^n \times (S\langle\langle\Sigma^\omega\rangle\rangle)^m$ is a solution of the mixed ω -algebraic system

$$x = p(x), \quad z = \varrho(x)z,$$

if

$$\sigma = p(\sigma), \quad \omega = \varrho(\sigma)\omega.$$

Observe that, by Theorem 5.5.1 of Ésik, Kuich [18], $\omega^{(k)} = \varrho(\sigma)^{\omega, k}$ for each $1 \leq k \leq n$, is solution for the linear system

$$z = \varrho(\sigma)z.$$

A solution $(\sigma_1, \dots, \sigma_n)$ of the algebraic system $x = p(x)$ is termed *least solution* if

$$\sigma_i \leq \tau_i, \quad \text{for each } 1 \leq i \leq n,$$

for all solutions (τ_1, \dots, τ_n) of $x = p(x)$.

If σ is the least solution of $x = p(x)$, then $z = \varrho(\sigma)z$ is an $S^{\text{alg}}\langle\langle\Sigma^*\rangle\rangle$ -linear system and $(\sigma, \omega^{(k)}) = (\sigma, \varrho(\sigma)^{\omega, k})$, where $k \in \{0, 1, \dots, m\}$, is called k^{th} *canonical solution* of $x = p(x)$, $z = \varrho(x)z$. Observe that the k^{th} canonical solution is unique by definition. A solution (σ, ω) is called *canonical*, if there exists a k such that (σ, ω) is the k^{th} canonical solution. The k^{th} canonical solution of an ω -algebraic system $y = p(y)$ is defined to be the k^{th} canonical solution of the mixed ω -algebraic system $x = p(x)$, $z = p_x(x, z)$ induced by $y = p(y)$.

Recall that $S^{\text{alg}}\langle\langle\Sigma^*\rangle\rangle$ is the collection of all components of least solutions of algebraic systems

$$x_i = p_i \quad \text{where } p_i \in S\langle\langle\Sigma \cup X\rangle\rangle^* \text{ for } 1 \leq i \leq n.$$

We define $S^{\text{alg}}\langle\langle\Sigma^\omega\rangle\rangle$ to be the collection of all components of vectors $M^{\omega, k}$, where $M \in (S^{\text{alg}}\langle\langle\Sigma^*\rangle\rangle)^{n \times n}$, $n \geq 1$, and $k \in \{1, \dots, n\}$. Moreover, we define $\omega\text{-Rat}(S^{\text{alg}}\langle\langle\Sigma^*\rangle\rangle)$ to be the ω -Kleene closure of (i.e., the generalized starquemiring generated by) $S^{\text{alg}}\langle\langle\Sigma^*\rangle\rangle$.

Example 3. We consider the following ω -algebraic system over the quemiring $\mathbb{B}\langle\langle\Sigma^*\rangle\rangle \times \mathbb{B}\langle\langle\Sigma^\omega\rangle\rangle$ for the Boolean semiring $\langle\mathbb{B}, +, \cdot, 0, 1\rangle$

$$\begin{aligned} y_1 &= y_2 y_1 + \epsilon \\ y_2 &= a y_2 b + \epsilon, \end{aligned}$$

where $a, b \in \Sigma$. This induces the following mixed ω -algebraic system

$$\begin{aligned} x_1 &= x_2 x_1 + \epsilon & z_1 &= z_2 + x_2 z_1 \\ x_2 &= a x_2 b + \epsilon & z_2 &= a z_2. \end{aligned}$$

Then for the algebraic system $x = p(x)$ over $\mathbb{B}\langle\langle\Sigma^*\rangle\rangle$, we get the least solution $\sigma_2 = \sum_{n \geq 0} a^n b^n$ and therefore $\sigma_1 = (\sum_{n \geq 0} a^n b^n)^*$. For the semimodule part, we can consider the first canonical solution where only z_1 is Büchi-accepting and the second canonical solution where both z_1 and z_2 are Büchi-accepting. The first canonical solution of the mixed ω -algebraic system $x = p(x)$, $z = p_x(x, z)$ over $\mathbb{B}\langle\langle\Sigma^*\rangle\rangle \times \mathbb{B}\langle\langle\Sigma^\omega\rangle\rangle$ is then $(\sigma_1, \sigma_2; (\sum_{n \geq 0} a^n b^n)^\omega, 0)$. The second canonical solution would be $(\sigma_1, \sigma_2; (\sum_{n \geq 0} a^n b^n)^\omega + (\sum_{n \geq 0} a^n b^n)^* a^\omega, a^\omega)$.

Example 4. We consider the following mixed ω -algebraic system over the quemiring $\mathbb{N}^\infty\langle\langle\Sigma^*\rangle\rangle \times \mathbb{N}^\infty\langle\langle\Sigma^\omega\rangle\rangle$ for the tropical semiring $\langle\mathbb{N}^\infty, \min, +, \infty, 0\rangle$

$$\begin{aligned} x_1 &= 1 a x_1 b + 1 a b & z_1 &= c z_1 \\ & & z_2 &= x_1 z_1 + z_1 \end{aligned}$$

where $a, b, c \in \Sigma$ and using the natural number 1.

Then for the algebraic system $x = p(x)$ over $\mathbb{N}^\infty\langle\langle\Sigma^*\rangle\rangle$, we get the least solution $\sigma = a^n b^n \mapsto n$. The first canonical solution of the mixed ω -algebraic system $x = p(x)$, $z = \varrho(x)z$ over $\mathbb{N}^\infty\langle\langle\Sigma^*\rangle\rangle \times \mathbb{N}^\infty\langle\langle\Sigma^\omega\rangle\rangle$ is then $(\sigma, c^\omega \mapsto 0, a^n b^n c^\omega \mapsto n)$. Hence the series $a^n b^n c^\omega \mapsto n$ is ω -algebraic but it is clearly not recognizable by a weighted automaton without stack.

Now we have the following characterization of ω -algebraic series.

Theorem 5. *Let S be a continuous complete star-omega semiring with the underlying semiring S being commutative and let Σ be an alphabet. Then the following statements are equivalent for $(s, v) \in S\langle\langle\Sigma^*\rangle\rangle \times S\langle\langle\Sigma^\omega\rangle\rangle$:*

- (i) $(s, v) \in S^{\text{alg}}\langle\langle\Sigma^*\rangle\rangle \times S^{\text{alg}}\langle\langle\Sigma^\omega\rangle\rangle$,
- (ii) $(s, v) \in \omega\text{-Rat}(S^{\text{alg}}\langle\langle\Sigma^*\rangle\rangle)$,
- (iii) $(s, v) = \|\mathfrak{A}\|$, where \mathfrak{A} is a finite $S^{\text{alg}}\langle\langle\Sigma^*\rangle\rangle$ -automaton over $S\langle\langle\Sigma^*\rangle\rangle \times S\langle\langle\Sigma^\omega\rangle\rangle$,
- (iv) $s \in S^{\text{alg}}\langle\langle\Sigma^*\rangle\rangle$ and $v = \sum_{1 \leq j \leq l} s_j t_j^\omega$ for some $l \geq 0$, where $s_j, t_j \in S^{\text{alg}}\langle\langle\Sigma^*\rangle\rangle$,
- (v) (s, v) is a component of a canonical solution of a mixed ω -algebraic system over $S\langle\langle\Sigma^*\rangle\rangle \times S\langle\langle\Sigma^\omega\rangle\rangle$.

Proof. The statements (ii), (iii) and (iv) are equivalent by Theorem 5.4.9 (see also Theorem 5.6.6) of Ésik, Kuich [18].

(iii) \Rightarrow (v): Assume that $(s, v) = \|\mathfrak{A}\|$, where $\mathfrak{A} = (n, I, M, P, k)$ is a finite $S^{\text{alg}}\langle\langle\Sigma^*\rangle\rangle$ -automaton. Without loss of generality \mathfrak{A} is normalized by Theorem 5.4.2 of Ésik, Kuich [18]; i.e., $I = e_i$ for some i . Hence, $(s, v) = ((M^*P)_i, (M^{\omega,k})_i)$ is a component of the k^{th} canonical solution of the mixed ω -algebraic system

$$x = Mx + P, \quad z = Mz.$$

(v) \Rightarrow (i): Assume there exists a mixed ω -algebraic system $x = p(x), z = \varrho(x)z$, with canonical solution $(\sigma, \varrho(\sigma)^{\omega,k})$ such that $(s, v) = (\sigma_i, (\varrho(\sigma)^{\omega,k})_j)$ for some i and j . Since the entries of σ and $\varrho(\sigma)$ are in $S^{\text{alg}}\langle\langle\Sigma^*\rangle\rangle$, (s, v) is in $S^{\text{alg}}\langle\langle\Sigma^*\rangle\rangle \times S^{\text{alg}}\langle\langle\Sigma^\omega\rangle\rangle$.

(i) \Rightarrow (iv): Now assume $s \in S^{\text{alg}}\langle\langle\Sigma^*\rangle\rangle$ and $v = (M^{\omega,k})_i$ for some $M \in (S^{\text{alg}}\langle\langle\Sigma^*\rangle\rangle)^{n \times n}$, $n \geq 1$, and $i, k \in \{1, \dots, n\}$. By the definition of $M^{\omega,k}$, each entry of $M^{\omega,k}$ is of the form $\sum_{1 \leq j \leq l} s_j t_j^\omega$ for some $l \geq 0$, where $s_j, t_j \in S^{\text{alg}}\langle\langle\Sigma^*\rangle\rangle$ for $1 \leq j \leq l$. \square

4. Greibach Normal Form for Mixed ω -Algebraic Systems

In this section we show that for any element (s, v) of $S^{\text{alg}}\langle\langle\Sigma^*\rangle\rangle \times S^{\text{alg}}\langle\langle\Sigma^\omega\rangle\rangle$ there exists a mixed ω -algebraic system in Greibach normal form such that (s, v) is a component of a solution of this ω -algebraic system.

Similar to the definition for algebraic systems on finite words (cf. also Greibach [20]), a mixed ω -algebraic system

$$x = p(x), \quad z = \varrho(x)z$$

is in *Greibach normal form* if

$$\begin{aligned} \text{supp}(p_i(x)) &\subseteq \{\epsilon\} \cup \Sigma \cup \Sigma X \cup \Sigma XX, & \text{for all } 1 \leq i \leq n, & \quad \text{and} \\ \text{supp}(\varrho_{ij}(x)) &\subseteq \Sigma \cup \Sigma X, & \text{for all } 1 \leq i, j \leq m. \end{aligned}$$

For the construction of the Greibach normal form we need a corollary to Theorem 5.

Corollary 6. *The following statement for $(s, v) \in S\langle\langle\Sigma^*\rangle\rangle \times S\langle\langle\Sigma^\omega\rangle\rangle$ is equivalent to the statements (i) to (v) of Theorem 5:*

$s \in S^{\text{alg}}\langle\langle\Sigma^\rangle\rangle$ and $v = \sum_{1 \leq j \leq l} s_j t_j^\omega$ for some $l \geq 0$, where $s_j, t_j \in S^{\text{alg}}\langle\langle\Sigma^*\rangle\rangle$ with $(t_j, \epsilon) = 0$; moreover $(s_j, \epsilon) = 0$ or $s_j = (s_j, \epsilon)\epsilon$.*

Proof. Assume $(s_j, \epsilon) \neq 0$. Then $s_j = (s_j, \epsilon)\epsilon + s'_j$ where $(s'_j, \epsilon) = 0$, and $s_j t_j^\omega = (s_j, \epsilon)t_j^\omega + s'_j t_j^\omega$.

Assume $(t_j, \epsilon) \neq 0$. Then $t_j = (t_j, \epsilon)\epsilon + t'_j$, where $(t'_j, \epsilon) = 0$. Since $(S\langle\langle\Sigma^*\rangle\rangle, S\langle\langle\Sigma^\omega\rangle\rangle)$ is a Conway semiring-semimodule pair satisfying $\epsilon^\omega = 0$, we obtain $t_j^\omega = ((t_j, \epsilon)^* \epsilon^* t'_j)^\omega$ with $(t_j, \epsilon)^* \epsilon^* t'_j \in S^{\text{alg}}\langle\langle\Sigma^*\rangle\rangle$, since $((t_j, \epsilon)\epsilon)^\omega = (t_j, \epsilon)^\omega \epsilon^\omega = 0$. \square

We now assume that $(s, v) \in S^{\text{alg}}\langle\langle\Sigma^*\rangle\rangle \times S\langle\langle\Sigma^\omega\rangle\rangle$ is given in the form of Corollary 6 with $l = 1$. By Theorem 2.4.10 of Ésik, Kuich [18], there exist algebraic systems in Greibach normal form whose first component of their least solutions equals s_1, t_1 .

Firstly, we deal with the case $(s_1, \epsilon) = 0$. Let

$$x_i = p_i(x) + \sum_{1 \leq j \leq n} p_{ij}(x)x_j, \quad \text{for each } 1 \leq i \leq n, \quad (*)$$

where $\text{supp}(p_i(x)) \subseteq \Sigma \cup \Sigma X$, $\text{supp}(p_{ij}(x)) \subseteq \Sigma X$, be the algebraic system in Greibach normal form for s_1 and

$$x'_i = p'_i(x') + \sum_{1 \leq j \leq m} p'_{ij}(x')x'_j, \quad \text{for each } 1 \leq i \leq m, \quad (**)$$

where $\text{supp}(p'_i(x')) \subseteq \Sigma \cup \Sigma X'$, $\text{supp}(p'_{ij}(x')) \subseteq \Sigma X'$, be the algebraic system in Greibach normal form for t_1 . Let σ and σ' with $\sigma_1 = s_1$ and $\sigma'_1 = t_1$ be the least solutions of $(*)$ and $(**)$, respectively.

Consider now the mixed ω -algebraic system consisting of the algebraic system $(*)$, $(**)$ over $S\langle\langle\Sigma^*\rangle\rangle$ and the linear system over $S\langle\langle\Sigma^\omega\rangle\rangle$

$$\begin{aligned} z'' &= p'_1(x')z'' + \sum_{1 \leq j \leq m} p'_{1j}(x')z'_j, \\ z'_i &= p'_i(x')z'' + \sum_{1 \leq j \leq m} p'_{ij}(x')z'_j, \quad \text{for } 1 \leq i \leq m, \\ z_i &= p_i(x)z'' + \sum_{1 \leq j \leq n} p_{ij}(x)z_j, \quad \text{for } 1 \leq i \leq n. \end{aligned} \quad (***)$$

Observe that the mixed ω -algebraic system is in Greibach normal form. We then order the variables of the mixed ω -algebraic system $(*)$, $(**)$, $(***)$ as $x_1, \dots, x_n; x'_1, \dots, x'_m; z'', z'_1, \dots, z'_m; z_1, \dots, z_n$. After an example, we will prove that

$$(\sigma_1, \dots, \sigma_n; \sigma'_1, \dots, \sigma'_m; \sigma'_1 \sigma_1^\omega; \sigma'_1 \sigma_1^\omega, \dots, \sigma'_m \sigma_1^\omega; \sigma_1 \sigma_1^\omega, \dots, \sigma_n \sigma_1^\omega) \quad (7)$$

is a solution of $(*)$, $(**)$, $(***)$. Observe that $\sigma'_1 \sigma_1^\omega = \sigma_1^\omega$.

Example 7. Consider the quemiring $\mathbb{N}^\infty\langle\langle\Sigma^*\rangle\rangle \times \mathbb{N}^\infty\langle\langle\Sigma^\omega\rangle\rangle$ for the tropical semiring $\langle\mathbb{N}^\infty, \min, +, \infty, 0\rangle$. Note that subsequently, 1 stands for the natural number 1 and the neutral element of the semiring multiplication is $\mathbb{1} = 0$.

We now define algebraic systems in Greibach normal form for $s = a^n b^n \mapsto n$ and $t = ((dd)^* c) \mapsto 0$. Let

$$\begin{aligned} x_1 &= 1ax_2 + 1ax_1x_2 & x'_1 &= c + dx'_2x'_1 \\ x_2 &= b & x'_2 &= d \end{aligned}$$

Here, x_1 is the start variable for s and x'_1 is the start variable for t . In the proof, these two systems are called $(*)$ and $(**)$. Now, we construct a mixed ω -algebraic system:

$$\begin{aligned} z'' &= cz'' + dx'_2z'_1 \\ z'_1 &= cz'' + dx'_2z'_1 & z'_2 &= dz'' \\ z_1 &= 1ax_2z'' + 1ax_1z_2 & z_2 &= bz'' \end{aligned}$$

In the new system (corresponding to $(***)$), variable z'' is Büchi-accepting and variable z_1 acts as the start variable, i.e., we consider the forth component (with the ordering $z'', z'_1, z'_2, z_1, z_2$) of the first canonical solution. The semimodule part of the solution is $st^\omega = a^n b^n ((dd)^* c)^\omega \mapsto n$. Note that the equation for z'' is needed in this example because z'_1 is not allowed to be Büchi-accepting to prevent $(dd)^\omega$ as part of the canonical solution.

Lemma 8. The mixed ω -algebraic system $(*)$, $(**)$, $(***)$ has solution

$$(\sigma_1, \dots, \sigma_n; \sigma'_1, \dots, \sigma'_m; \sigma'_1 \sigma_1^\omega; \sigma'_1 \sigma_1^\omega, \dots, \sigma'_m \sigma_1^\omega; \sigma_1 \sigma_1^\omega, \dots, \sigma_n \sigma_1^\omega)$$

Proof. Again, observe that $\sigma_1^{\omega} = \sigma_1' \sigma_1'^{\omega}$. We obtain, for the first equation,

$$\begin{aligned} p_1'(\sigma') \sigma_1'^{\omega} + \sum_{1 \leq j \leq m} p_{1j}'(\sigma') \sigma_j' \sigma_1'^{\omega} &= (p_1'(\sigma') + \sum_{1 \leq j \leq m} p_{1j}'(\sigma') \sigma_j') \sigma_1'^{\omega} \\ &= \sigma_1' \sigma_1'^{\omega}, \end{aligned}$$

then, for $1 \leq i \leq m$, and the second equation,

$$\begin{aligned} p_i'(\sigma') \sigma_1'^{\omega} + \sum_{1 \leq j \leq m} p_{ij}'(\sigma') \sigma_j' \sigma_1'^{\omega} &= (p_i'(\sigma') + \sum_{1 \leq j \leq m} p_{ij}'(\sigma') \sigma_j') \sigma_1'^{\omega} \\ &= \sigma_i' \sigma_1'^{\omega}, \end{aligned}$$

and, for $1 \leq i \leq n$, and the third equation,

$$\begin{aligned} p_i(\sigma) \sigma_1^{\omega} + \sum_{1 \leq j \leq n} p_{ij}(\sigma) \sigma_j \sigma_1^{\omega} &= (p_i(\sigma) + \sum_{1 \leq j \leq n} p_{ij}(\sigma) \sigma_j) \sigma_1^{\omega} \\ &= \sigma_i \sigma_1^{\omega}. \end{aligned}$$

□

But we need more: We will prove that this solution is the first canonical solution of (*), (**), (***) .

Lemma 9. *The solution (7) is the first canonical solution of the mixed ω -algebraic system (*), (**), (***) .*

Proof. Let

$$\begin{aligned} P_{1m}'(x') &= (p_{11}'(x') \quad \cdots \quad p_{1m}'(x')), \\ P_{m1}'(x') &= \begin{pmatrix} p_1'(x') \\ \vdots \\ p_m'(x') \end{pmatrix}, & P_{mm}'(x') &= \begin{pmatrix} p_{11}'(x') & \cdots & p_{1m}'(x') \\ \vdots & & \vdots \\ p_{m1}'(x') & \cdots & p_{mm}'(x') \end{pmatrix}, \\ P_{n1}(x) &= \begin{pmatrix} p_1(x) \\ \vdots \\ p_n(x) \end{pmatrix}, & P_{nn}(x) &= \begin{pmatrix} p_{11}(x) & \cdots & p_{1n}(x) \\ \vdots & & \vdots \\ p_{n1}(x) & \cdots & p_{nn}(x) \end{pmatrix}, \\ z &= \begin{pmatrix} z_1 \\ \vdots \\ z_n \end{pmatrix}, & z' &= \begin{pmatrix} z_1' \\ \vdots \\ z_m' \end{pmatrix}, \end{aligned}$$

and

$$M(x, x') = \begin{pmatrix} p_1'(x') & P_{1m}'(x') & 0 \\ P_{m1}'(x') & P_{mm}'(x') & 0 \\ P_{n1}(x) & 0 & P_{nn}(x) \end{pmatrix}.$$

Then the linear system (***) can be written in the form

$$\begin{pmatrix} z'' \\ z' \\ z \end{pmatrix} = M(x, x') \begin{pmatrix} z'' \\ z' \\ z \end{pmatrix}.$$

Hence, the first canonical solution of (*), (**), (***) is $(\sigma, \sigma', M(\sigma, \sigma')^{\omega, 1})$. Before we prove our lemma, we prove three identities.

The system (*) can be written in the form

$$x = P_{n1}(x) + P_{nn}(x)x, \quad \text{for } x = (x_1, \dots, x_n)^{\top}.$$

By the diagonal identity (see Proposition 2.2.11 of Ésik, Kuich [18]) the system

$$x = P_{n1}(\sigma) + P_{nm}(\sigma)x$$

has the same least solution as (*). Hence,

$$\sigma = P_{nm}(\sigma)^* P_{n1}(\sigma). \quad (8)$$

The system (**) can be written in the form

$$x' = P'_{m1}(x') + P'_{mm}(x')x', \quad \text{for } x' = (x'_1, \dots, x'_m)^\top.$$

Again, by the diagonal identity (see Proposition 2.2.11 of Ésik, Kuich [18]) the system

$$x' = P'_{m1}(\sigma') + P'_{mm}(\sigma')x'$$

has the same solution. Hence

$$\sigma' = P'_{mm}(\sigma')^* P'_{m1}(\sigma'). \quad (9)$$

It follows for the first component

$$\begin{aligned} \sigma'_1 &= (P'_{mm}(\sigma')^* P'_{m1}(\sigma'))_1 \\ &= (P'_{m1}(\sigma') + P'_{mm}(\sigma')^+ P'_{m1}(\sigma'))_1 \\ &= (P'_{m1}(\sigma') + P'_{mm}(\sigma') P'_{mm}(\sigma')^* P'_{m1}(\sigma'))_1 \\ &= p'_1(\sigma') + P'_{1m}(\sigma') P'_{mm}(\sigma')^* P'_{m1}(\sigma'). \end{aligned} \quad (10)$$

We now compute

$$\begin{aligned} (M^{\omega,1}(\sigma, \sigma'))_{z''} &= \left[p'_1(\sigma') + (P'_{1m}(\sigma') \quad 0) \begin{pmatrix} P'_{mm}(\sigma') & 0 \\ 0 & P_{nm}(\sigma) \end{pmatrix}^* \begin{pmatrix} P'_{m1}(\sigma') \\ P_{n1}(\sigma) \end{pmatrix} \right]^\omega \\ &= \left[p'_1(\sigma') + (P'_{1m}(\sigma') \quad 0) \begin{pmatrix} P'_{mm}(\sigma')^* & 0 \\ 0 & P_{nm}(\sigma)^* \end{pmatrix} \begin{pmatrix} P'_{m1}(\sigma') \\ P_{n1}(\sigma) \end{pmatrix} \right]^\omega \\ &= [p'_1(\sigma') + P'_{1m}(\sigma') P'_{mm}(\sigma')^* P'_{m1}(\sigma')]^\omega \\ &= \sigma'_1{}^\omega. \end{aligned}$$

The last equality is by (10).

When starting with another variable z_i or z'_j for $1 \leq i \leq n$ and $1 \leq j \leq m$, we get

$$\begin{aligned} (M^{\omega,1}(\sigma, \sigma'))_{(z', z)} &= \begin{pmatrix} P'_{mm}(\sigma') & 0 \\ 0 & P_{nm}(\sigma) \end{pmatrix}^* \begin{pmatrix} P'_{m1}(\sigma') \\ P_{n1}(\sigma) \end{pmatrix} (M^{\omega,1}(\sigma, \sigma'))_{z''} \\ &= \begin{pmatrix} P'_{mm}(\sigma')^* & 0 \\ 0 & P_{nm}(\sigma)^* \end{pmatrix} \begin{pmatrix} P'_{m1}(\sigma') \\ P_{n1}(\sigma) \end{pmatrix} \sigma'_1{}^\omega \\ &= \begin{pmatrix} P'_{mm}(\sigma')^* P'_{m1}(\sigma') \\ P_{nm}(\sigma)^* P_{n1}(\sigma) \end{pmatrix} \sigma'_1{}^\omega \end{aligned}$$

Thus, by (9), we have, for $1 \leq i \leq m$,

$$(M^{\omega,1}(\sigma, \sigma'))_{z'_i} = [P'_{mm}(\sigma')^* P'_{m1}(\sigma')]_i \sigma'_1{}^\omega = \sigma'_i \sigma'_1{}^\omega,$$

and, by (8), we have, for $1 \leq i \leq n$,

$$(M^{\omega,1}(\sigma, \sigma'))_{z_i} = [P_{nm}(\sigma)^* P_{n1}(\sigma)]_i \sigma'_1{}^\omega = \sigma_i \sigma'_1{}^\omega.$$

This completes the proof. \square

Secondly, we deal with the case $s_1 = (s_1, \epsilon)\epsilon$. Consider now the mixed ω -algebraic system consisting of (**) and the linear system over $S \langle\langle \Sigma^\omega \rangle\rangle$

$$\begin{aligned} z'' &= p'_1(x')z'' + \sum_{1 \leq j \leq m} p'_{1j}(x')z'_j, \\ z'_i &= p'_i(x')z'' + \sum_{1 \leq j \leq m} p'_{ij}(x')z'_j, \quad 1 \leq i \leq m, \\ z_1 &= (s_1, \epsilon)p'_1(x')z'' + (s_1, \epsilon) \sum_{1 \leq j \leq m} p'_{1j}(x')z'_j. \end{aligned} \tag{****}$$

Lemma 10. *The mixed ω -algebraic system (**), (****) has solution*

$$(\sigma'_1, \dots, \sigma'_m; \sigma'_1 \sigma_1'^\omega; \sigma'_1 \sigma_1'^\omega, \dots, \sigma'_m \sigma_1'^\omega; (s_1, \epsilon) \sigma_1'^\omega). \tag{11}$$

Proof. As in the proof of Lemma 8, we obtain that $(\sigma'_1 \sigma_1'^\omega; \sigma'_1 \sigma_1'^\omega, \dots, \sigma'_m \sigma_1'^\omega)$ is solution of the z'' - and z'_i -equations, $1 \leq i \leq m$. For the right side of the z_1 -equation, we obtain

$$(s_1, \epsilon) \left(p'_1(\sigma') \sigma_1'^\omega + \sum_{1 \leq j \leq m} p'_{1j}(\sigma') \sigma'_j \sigma_1'^\omega \right) = (s_1, \epsilon) \sigma'_1 \sigma_1'^\omega = (s_1, \epsilon) \sigma_1'^\omega. \quad \square$$

But again we need more: We will prove that this solution is the first canonical solution of (**), (****).

Lemma 11. *The solution (11) is the first canonical solution of the mixed algebraic system (**), (****).*

Proof. Let

$$M_\epsilon(x') = \begin{pmatrix} p'_1(x') & P'_{1m}(x') & 0 \\ P'_{m1}(x') & P'_{mm}(x') & 0 \\ (s_1, \epsilon)p'_1(x') & (s_1, \epsilon)P'_{1m}(x') & 0 \end{pmatrix}.$$

Then the linear system (****) can be written in the form

$$\begin{pmatrix} z'' \\ z' \\ z_1 \end{pmatrix} = M_\epsilon(x') \begin{pmatrix} z'' \\ z' \\ z_1 \end{pmatrix}.$$

Hence, the first canonical solution of (**), (****) is $(\sigma', M_\epsilon(\sigma')^{\omega,1})$. We now compute

$$\begin{aligned} (M_\epsilon^{\omega,1}(\sigma'))_{z''} &= \left[p'_1(\sigma') + \begin{pmatrix} P'_{1m}(\sigma') & 0 \end{pmatrix} \begin{pmatrix} P'_{mm}(\sigma') & 0 \\ (s_1, \epsilon)P'_{1m}(\sigma') & 0 \end{pmatrix}^* \begin{pmatrix} P'_{m1}(\sigma') \\ (s_1, \epsilon)p'_1(\sigma') \end{pmatrix} \right]^\omega \\ &= \left[p'_1(\sigma') + \begin{pmatrix} P'_{1m}(\sigma') & 0 \end{pmatrix} \begin{pmatrix} P'_{mm}(\sigma')^* & 0 \\ (s_1, \epsilon)P'_{1m}(\sigma')P'_{mm}(\sigma')^* & 1 \end{pmatrix} \begin{pmatrix} P'_{m1}(\sigma') \\ (s_1, \epsilon)p'_1(\sigma') \end{pmatrix} \right]^\omega \\ &= \left[p'_1(\sigma') + P'_{1m}(\sigma')P'_{mm}(\sigma')^*P'_{m1}(\sigma') \right]^\omega \\ &= \sigma_1'^\omega. \end{aligned}$$

The last equality is by (10).

When starting with another variable z'_i or z_1 for $1 \leq i \leq m$, we get

$$\begin{aligned} (M_\epsilon^{\omega,1}(\sigma'))_{(z', z_1)} &= \begin{pmatrix} P'_{mm}(\sigma') & 0 \\ (s_1, \epsilon)P'_{1m}(\sigma') & 0 \end{pmatrix}^* \begin{pmatrix} P'_{m1}(\sigma') \\ (s_1, \epsilon)p'_1(\sigma') \end{pmatrix} (M_\epsilon^{\omega,1}(\sigma'))_{z''} \\ &= \begin{pmatrix} P'_{mm}(\sigma')^* & 0 \\ (s_1, \epsilon)P'_{1m}(\sigma')P'_{mm}(\sigma')^* & 1 \end{pmatrix} \begin{pmatrix} P'_{m1}(\sigma') \\ (s_1, \epsilon)p'_1(\sigma') \end{pmatrix} \sigma_1'^\omega \\ &= \begin{pmatrix} P'_{mm}(\sigma')^*P'_{m1}(\sigma') \\ (s_1, \epsilon)P'_{1m}(\sigma')P'_{mm}(\sigma')^*P'_{m1}(\sigma') + (s_1, \epsilon)p'_1(\sigma') \end{pmatrix} \sigma_1'^\omega \end{aligned}$$

Thus, by (9), we have, for $1 \leq i \leq m$,

$$(M_{\epsilon}^{\omega,1}(\sigma'))_{z_i} = [P'_{mm}(\sigma')^* P'_{m1}(\sigma')]_i \sigma'_1{}^{\omega} = \sigma'_i \sigma'_1{}^{\omega},$$

and, by (10), we have

$$\begin{aligned} (M_{\epsilon}^{\omega,1}(\sigma'))_{z_1} &= ((s_1, \epsilon) P'_{1m}(\sigma') P'_{mm}(\sigma')^* P'_{m1}(\sigma') + (s_1, \epsilon) p'_1(\sigma')) \sigma'_1{}^{\omega} \\ &= (s_1, \epsilon) (P'_{1m}(\sigma') P'_{mm}(\sigma')^* P'_{m1}(\sigma') + p'_1(\sigma')) \sigma'_1{}^{\omega} \\ &= (s_1, \epsilon) \sigma'_1 \sigma'_1{}^{\omega} = (s_1, \epsilon) \sigma'_1{}^{\omega}. \end{aligned}$$

□

We now consider general sums of series of the above form. The next lemma shows how to construct a mixed ω -algebraic system whose canonical solution is the sum of the canonical solutions of multiple mixed ω -algebraic systems as given in Lemmas 9 and 11.

Lemma 12. *Let $(s, v) \in S^{alg}\langle\langle\Sigma^*\rangle\rangle \times S\langle\langle\Sigma^{\omega}\rangle\rangle$ be given in the form of Corollary 6. Then there exists a mixed ω -algebraic system in Greibach normal form such that v is a component of its l^{th} canonical solution.*

Proof. Let $v = \sum_{1 \leq i \leq l} s_i t_i^{\omega}$ as in the statement of Corollary 6 and let $l \geq 1$. By Lemmas 9 and 11, for $1 \leq i \leq l$, there exist mixed ω -algebraic systems

$$\begin{aligned} x_i &= p_i(x_i), \\ \begin{pmatrix} z_i \\ \bar{z}_i \end{pmatrix} &= M_i(x_i) \begin{pmatrix} z_i \\ \bar{z}_i \end{pmatrix}, \end{aligned} \tag{\#}$$

in Greibach normal form with

$$M_i(x_i) = \begin{pmatrix} a_i & b_i \\ c_i & d_i \end{pmatrix},$$

where

$$\begin{aligned} a_i &\in (S\langle\langle\Sigma \cup X\rangle\rangle)^{1 \times 1}, \\ b_i &\in (S\langle\langle\Sigma \cup X\rangle\rangle)^{1 \times (n_i-1)}, \\ c_i &\in (S\langle\langle\Sigma \cup X\rangle\rangle)^{(n_i-1) \times 1}, \\ d_i &\in (S\langle\langle\Sigma \cup X\rangle\rangle)^{(n_i-1) \times (n_i-1)}, \end{aligned}$$

such that $s_i t_i^{\omega}$ is a component of the first canonical solution of the i^{th} system. We will assume without loss of generality that $s_i t_i^{\omega}$ is the first component of variable \bar{z}_i , i.e.,

$$s_i t_i^{\omega} = [(M_i^{\omega,1})_{\bar{z}_i}]_1 = [(d_i^* c_i)(a_i + b_i d_i^* c_i)^{\omega}]_1. \tag{12}$$

Similarly to the case of summation in Theorem 5.4.4 of Ésik, Kuich [18], we consider now the mixed ω -algebraic system consisting of the algebraic systems (\#) over $S\langle\langle\Sigma^*\rangle\rangle$ and the linear system over $S\langle\langle\Sigma^{\omega}\rangle\rangle$

$$\hat{z} = M \hat{z}, \tag{\#\#}$$

with

$$M = \begin{pmatrix} \begin{pmatrix} a_1 & & \\ & \ddots & \\ & & a_l \end{pmatrix} & \begin{pmatrix} b_1 & & \\ & \ddots & \\ & & b_l \end{pmatrix} & 0 \\ \begin{pmatrix} c_1 & & \\ & \ddots & \\ & & c_l \end{pmatrix} & \begin{pmatrix} d_1 & & \\ & \ddots & \\ & & d_l \end{pmatrix} & 0 \\ \begin{pmatrix} c_1 & \cdots & c_l \end{pmatrix} & \begin{pmatrix} d_1 & \cdots & d_l \end{pmatrix} & 0 \end{pmatrix}, \quad \hat{z} = \begin{pmatrix} z_1 \\ \vdots \\ z_l \\ \bar{z}_1 \\ \vdots \\ \bar{z}_l \\ z' \end{pmatrix}.$$

Note that this system $\#\#$ is still in Greibach normal form.

We order the variables of the mixed ω -algebraic system $(\#)$, $(\#\#)$ as $z_1, \dots, z_l; \bar{z}_1, \dots, \bar{z}_l; z'$. We now compute the l^{th} canonical solution, starting with variable $z = (z_1, \dots, z_l)^\top$. Then

$$\begin{aligned}
(M^{\omega,l})_z &= \left[\begin{pmatrix} a_1 & & \\ & \ddots & \\ & & a_l \end{pmatrix} + \begin{pmatrix} b_1 & & \\ & \ddots & \\ & & b_l \end{pmatrix} 0 \begin{pmatrix} d_1 & & \\ & \ddots & \\ & & d_l \end{pmatrix} 0 \begin{pmatrix} c_1 & & \\ & \ddots & \\ & & c_l \end{pmatrix} \right]^\omega \\
&= \left[\begin{pmatrix} a_1 & & \\ & \ddots & \\ & & a_l \end{pmatrix} + \begin{pmatrix} b_1 & & \\ & \ddots & \\ & & b_l \end{pmatrix} 0 \begin{pmatrix} d_1 & & \\ & \ddots & \\ & & d_l \end{pmatrix}^* \begin{pmatrix} c_1 & & \\ & \ddots & \\ & & c_l \end{pmatrix} \right]^\omega \\
&= \left[\begin{pmatrix} a_1 & & \\ & \ddots & \\ & & a_l \end{pmatrix} + \begin{pmatrix} b_1 & & \\ & \ddots & \\ & & b_l \end{pmatrix} \begin{pmatrix} d_1 & & \\ & \ddots & \\ & & d_l \end{pmatrix}^* \begin{pmatrix} c_1 & & \\ & \ddots & \\ & & c_l \end{pmatrix} \right]^\omega \\
&= \begin{pmatrix} (a_1 + b_1 d_1^* c_1)^\omega \\ \vdots \\ (a_l + b_l d_l^* c_l)^\omega \end{pmatrix}.
\end{aligned}$$

When starting with the new variable z' , we get a sum of the original solutions:

$$\begin{aligned}
(M^{\omega,l})_{z'} &= \left[\begin{pmatrix} d_1 & & \\ & \ddots & \\ & & d_l \end{pmatrix}^* \begin{pmatrix} c_1 & & \\ & \ddots & \\ & & c_l \end{pmatrix} (M^{\omega,l})_z \right]_{l+1} \\
&= \left[\begin{pmatrix} d_1 & & \\ & \ddots & \\ & & d_l \end{pmatrix}^* 0 \begin{pmatrix} c_1 & & \\ & \ddots & \\ & & c_l \end{pmatrix} (M^{\omega,l})_z \right]_{l+1} \\
&= \left[\begin{pmatrix} d_1 & & \\ & \ddots & \\ & & d_l \end{pmatrix}^* \begin{pmatrix} c_1 & & \\ & \ddots & \\ & & c_l \end{pmatrix} + \begin{pmatrix} c_1 & & \\ & \ddots & \\ & & c_l \end{pmatrix} \right]_{l+1} (M^{\omega,l})_z \\
&= \left[\begin{pmatrix} d_1^* c_1 & & \\ & \ddots & \\ & & d_l^* c_l \end{pmatrix} \begin{pmatrix} (a_1 + b_1 d_1^* c_1)^\omega \\ \vdots \\ (a_l + b_l d_l^* c_l)^\omega \end{pmatrix} \right]_{l+1}
\end{aligned}$$

$$\begin{aligned}
&= \begin{pmatrix} d_1^* c_1 (a_1 + b_1 d_1^* c_1)^\omega \\ \vdots \\ d_l^* c_l (a_l + b_l d_l^* c_l)^\omega \\ \sum_{1 \leq i \leq l} (d_i d_i^* c_i + c_i)(a_i + b_i d_i^* c_i)^\omega \end{pmatrix}_{l+1} \\
&= \sum_{1 \leq i \leq l} (d_i d_i^* c_i + c_i)(a_i + b_i d_i^* c_i)^\omega \\
&= \sum_{1 \leq i \leq l} (d_i^* c_i)(a_i + b_i d_i^* c_i)^\omega
\end{aligned}$$

Thus, the first component is (by identity (12))

$$\begin{aligned}
[(M^{\omega, l})_{z'}]_1 &= \left[\sum_{1 \leq i \leq l} (d_i^* c_i)(a_i + b_i d_i^* c_i)^\omega \right]_1 \\
&= \sum_{1 \leq i \leq l} [(d_i^* c_i)(a_i + b_i d_i^* c_i)^\omega]_1 = \sum_{1 \leq i \leq l} s_i t_i^\omega = v.
\end{aligned}$$

□

We can now conclude the following theorem.

Theorem 13. *The following statement for $(s, v) \in S\langle\langle\Sigma^*\rangle\rangle \times S\langle\langle\Sigma^\omega\rangle\rangle$ is equivalent to the statements of Theorem 5: (s, v) is component of a canonical solution of a mixed ω -algebraic system over $S\langle\langle\Sigma^*\rangle\rangle \times S\langle\langle\Sigma^\omega\rangle\rangle$ in Greibach normal form.*

Proof. The above statement trivially implies statement (v) of Theorem 5. By Corollary 6 and Lemma 12, the statements of Theorem 5 imply the above statement. □

5. Greibach Normal Form for ω -Algebraic Systems

For the following sections, we need the Greibach normal form not only for mixed ω -algebraic systems but also for ω -algebraic systems. So we show in this section a specialization of Theorem 13 for ω -algebraic systems.

Similar to the definition for mixed ω -algebraic systems, an ω -algebraic system

$$y = p(y)$$

where $\{y_1, \dots, y_n\}$ is a set of variables for the quemiring $S^{\text{alg}}\langle\langle\Sigma^*\rangle\rangle \times S^{\text{alg}}\langle\langle\Sigma^\omega\rangle\rangle$, is in *Greibach normal form* if

$$\text{supp}(p_i(y)) \subseteq \{\epsilon\} \cup \Sigma \cup \Sigma Y \cup \Sigma Y Y, \quad \text{for all } 1 \leq i \leq n.$$

Our first main result is the following.

Theorem 14. *The following statement for $(s, v) \in S\langle\langle\Sigma^*\rangle\rangle \times S\langle\langle\Sigma^\omega\rangle\rangle$ is equivalent to the statements of Theorem 5: (s, v) is component of a canonical solution of an ω -algebraic system over $S\langle\langle\Sigma^*\rangle\rangle \times S\langle\langle\Sigma^\omega\rangle\rangle$ in Greibach normal form.*

Proof. By Theorem 13, we can assume that (s, v) is component of the t^{th} canonical solution of a mixed ω -algebraic system over $S\langle\langle\Sigma^*\rangle\rangle \times S\langle\langle\Sigma^\omega\rangle\rangle$ in Greibach normal form for a $t \in \mathbb{N}$. Let the mixed ω -algebraic system be given in the following form:

$$x_i = p_i + \sum_{1 \leq j \leq n} (p_{ij}x + q_{ij})x_j, \quad \text{for } 1 \leq i \leq n, \quad (*)$$

$$z_i = \sum_{1 \leq j \leq m} (p'_{ij}x + q'_{ij})z_j, \quad \text{for } 1 \leq i \leq m, \quad (**)$$

where

$$\begin{aligned}
p_{ij} &\in S\langle\Sigma\rangle^{1 \times n}, & \text{for } 1 \leq i, j \leq n, \\
p'_{ij} &\in S\langle\Sigma\rangle^{1 \times n}, & \text{for } 1 \leq i, j \leq m,
\end{aligned}$$

and

$$\begin{aligned} \text{supp}(p_i) &\subseteq \{\epsilon\} \cup \Sigma, & \text{supp}(p_{ij}x) &\subseteq \Sigma X, & \text{supp}(q_{ij}) &\subseteq \Sigma, \\ \text{supp}(p'_{ij}x) &\subseteq \Sigma X, & \text{supp}(q'_{ij}) &\subseteq \Sigma. \end{aligned}$$

Note that

$$p_{ij}x = \sum_{1 \leq k \leq n} (p_{ij})_k x_k;$$

we decided for this notation because of brevity, important especially in matrices.

For the remainder of the proof, consider integers k and l to be fixed such that the t^{th} canonical solution of $(*)$, $(**)$ is (σ, ω) with $\sigma_k = s$ and $\omega_l = v$.

We will later need a simple implication: We can write the linear system $(**)$ as

$$z = P'_{mm}(x)z,$$

where

$$P'_{mm}(x) = \begin{pmatrix} p'_{11}x + q'_{11} & \cdots & p'_{1m}x + q'_{1m} \\ \vdots & \ddots & \vdots \\ p'_{m1}x + q'_{m1} & \cdots & p'_{mm}x + q'_{mm} \end{pmatrix}.$$

Note that $t \leq m$. It follows that

$$\omega = P'_{mm}(\sigma)^{\omega, t}. \quad (13)$$

Now, we construct from $(*)$, $(**)$ an ω -algebraic system $(***)$ where the variables x are substituted by \bar{y} and z by \hat{y} . Additionally, we add a new equation and a new variable \dot{y} to combine the k^{th} component of the semiring part and the l^{th} component of the semimodule part:

$$\begin{aligned} \hat{y}_i &= \sum_{1 \leq j \leq m} (p'_{ij}\bar{y} + q'_{ij})\hat{y}_j, & \text{for } 1 \leq i \leq m, \\ \bar{y}_i &= p_i + \sum_{1 \leq j \leq n} (p_{ij}\bar{y} + q_{ij})\bar{y}_j, & \text{for } 1 \leq i \leq n, \\ \dot{y} &= p_k + \sum_{1 \leq j \leq n} (p_{kj}\bar{y} + q_{kj})\bar{y}_j + \sum_{1 \leq j \leq m} (p'_{lj}\bar{y} + q'_{lj})\hat{y}_j. \end{aligned} \quad (***)$$

Note that $(***)$ is in Greibach normal form. Moreover, note that we order the equations such that the first equations are those corresponding to the old equations of variables z_i . This ensures that the t^{th} canonical solution still considers the correct variables as Büchi-accepting.

Claim: The $(m+n+1)^{\text{th}}$ component of the t^{th} canonical solution of $(***)$ is $(\sigma_k, \omega_l) = (s, v)$.

We now compute this solution. The t^{th} canonical solution of the ω -algebraic system $(***)$ is defined to be the t^{th} canonical solution of the mixed ω -algebraic system induced by $(***)$. The corresponding induced mixed ω -algebraic system is given by the algebraic system over $S^{\text{alg}}(\langle \Sigma^* \rangle)$

$$\begin{aligned} \hat{x}_i &= \sum_{1 \leq j \leq m} (p'_{ij}\bar{x} + q'_{ij})\hat{x}_j, & \text{for } 1 \leq i \leq m, \\ \bar{x}_i &= p_i + \sum_{1 \leq j \leq n} (p_{ij}\bar{x} + q_{ij})\bar{x}_j, & \text{for } 1 \leq i \leq n, \\ \dot{x} &= p_k + \sum_{1 \leq j \leq n} (p_{kj}\bar{x} + q_{kj})\bar{x}_j + \sum_{1 \leq j \leq m} (p'_{lj}\bar{x} + q'_{lj})\hat{x}_j, \end{aligned} \quad (\#)$$

and the linear system over $S^{\text{alg}}\langle\langle\Sigma^\omega\rangle\rangle$

$$\begin{aligned}\hat{z}_i &= \sum_{1 \leq j \leq m} (p'_{ij}\bar{x} + q'_{ij})\hat{z}_j + p'_{ij}\bar{z}, & \text{for } 1 \leq i \leq m, \\ \bar{z}_i &= \sum_{1 \leq j \leq n} (p_{ij}\bar{x} + q_{ij})\bar{z}_j + p_{ij}\bar{z}, & \text{for } 1 \leq i \leq n, \\ \hat{z} &= \sum_{1 \leq j \leq n} (p_{kj}\bar{x} + q_{kj})\bar{z}_j + p_{kj}\bar{z} + \sum_{1 \leq j \leq m} (p'_{lj}\bar{x} + q'_{lj})\hat{z}_j + p'_{lj}\bar{z}.\end{aligned}\tag{##}$$

Claim: $(0, \dots, 0; \sigma; \sigma_k)$ is the least solution of (#).

First, we prove that it is a solution by plugging it into the right sides of the equations. We have for the first m equations, and for $1 \leq i \leq m$,

$$\sum_{1 \leq j \leq m} (p'_{ij}\sigma + q'_{ij})0 = 0.$$

Then for the second set of equations and $1 \leq i \leq n$,

$$p_i + \sum_{1 \leq j \leq n} (p_{ij}\sigma + q_{ij})\sigma_j = \sigma_i;$$

because σ is a solution of (*). Finally, we obtain by the same reason, for the last equation,

$$\begin{aligned}p_k + \sum_{1 \leq j \leq n} (p_{kj}\sigma + q_{kj})\sigma_j + \sum_{1 \leq j \leq m} (p'_{lj}\sigma + q'_{lj})0_j &= p_k + \sum_{1 \leq j \leq n} (p_{kj}\sigma + q_{kj})\sigma_j + 0 \\ &= \sigma_k.\end{aligned}$$

The algebraic system (#) is strict and therefore has a unique solution. This means that $(0, \dots, 0; \sigma; \sigma_k)$ is also the least solution. This proves the claim.

Now consider the linear system (##). Let $P'_{mn}(\bar{x})$ be defined as above and let further

$$\begin{aligned}P_{nn}(\bar{x}) &= \begin{pmatrix} p_{11}\bar{x} + q_{11} & \cdots & p_{1n}\bar{x} + q_{1n} \\ \vdots & \ddots & \vdots \\ p_{n1}\bar{x} + q_{n1} & \cdots & p_{nn}\bar{x} + q_{nn} \end{pmatrix}, \\ R_{nn} &= \begin{pmatrix} \sum_{1 \leq j \leq n} (p_{1j})_1 & \cdots & \sum_{1 \leq j \leq n} (p_{1j})_n \\ \vdots & \ddots & \vdots \\ \sum_{1 \leq j \leq n} (p_{nj})_1 & \cdots & \sum_{1 \leq j \leq n} (p_{nj})_n \end{pmatrix}, \\ R'_{mn} &= \begin{pmatrix} \sum_{1 \leq j \leq m} (p'_{1j})_1 & \cdots & \sum_{1 \leq j \leq m} (p'_{1j})_n \\ \vdots & \ddots & \vdots \\ \sum_{1 \leq j \leq m} (p'_{mj})_1 & \cdots & \sum_{1 \leq j \leq m} (p'_{mj})_n \end{pmatrix}.\end{aligned}$$

Note that for (##) and for $1 \leq i \leq m$, we have

$$\begin{aligned}\sum_{1 \leq j \leq m} p'_{ij}\bar{z} &= \sum_{1 \leq j \leq m} \sum_{1 \leq k \leq n} (p'_{ij})_k \bar{z}_k \\ &= \sum_{1 \leq k \leq n} \sum_{1 \leq j \leq m} (p'_{ij})_k \bar{z}_k \\ &= \left(\sum_{1 \leq j \leq m} (p'_{ij})_1, \dots, \sum_{1 \leq j \leq m} (p'_{ij})_n \right) \bar{z} \\ &= (R'_{mn})_i \bar{z}.\end{aligned}$$

Analogously, we can prove $\sum_{1 \leq j \leq n} p_{ij} \bar{z} = (R_{nn})_i \bar{z}$. We let

$$M(\hat{x}, \bar{x}, x) = \begin{pmatrix} P'_{mm}(\bar{x}) & R'_{mn} & 0 \\ 0 & P_{nn}(\bar{x}) + R_{nn} & 0 \\ (P'_{mm}(\bar{x}))_l & (P_{nn}(\bar{x}))_k + (R_{nn})_k + (R'_{mn})_l & 0 \end{pmatrix},$$

then the linear system (##) can be written as

$$\begin{pmatrix} \hat{z} \\ \bar{z} \\ z \end{pmatrix} = M(\hat{x}, \bar{x}, x) \begin{pmatrix} \hat{z} \\ \bar{z} \\ z \end{pmatrix}.$$

Now, we can plug the semiring part $(0, \sigma, \sigma_k)$ of the solution into M . By Theorem 2, the semimodule part of the canonical solution of (#), (##) is

$$M(0, \sigma, \sigma_k)^{\omega, t} = \begin{pmatrix} P'_{mm}(\sigma) + R'_{mn} & 0 \\ \chi & 0 \end{pmatrix}^* \begin{pmatrix} 0 \\ (P'_{mm}(\sigma))_l \end{pmatrix} \xi^{\omega, t}$$

with

$$\chi = (P_{nn}(\sigma))_k + (R_{nn})_k + (R'_{mn})_l$$

and

$$\begin{aligned} \xi &= P'_{mm}(\sigma) + \begin{pmatrix} R'_{mn} & 0 \end{pmatrix} \begin{pmatrix} P_{nn}(\sigma) + R_{nn} & 0 \\ \chi & 0 \end{pmatrix}^* \begin{pmatrix} 0 \\ (P'_{mm}(\sigma))_l \end{pmatrix} \\ &= P'_{mm}(\sigma) + \begin{pmatrix} R'_{mn} & 0 \end{pmatrix} \begin{pmatrix} (P_{nn}(\sigma) + R_{nn})^* & 0 \\ \chi(P_{nn}(\sigma) + R_{nn})^* & 1 \end{pmatrix} \begin{pmatrix} 0 \\ (P'_{mm}(\sigma))_l \end{pmatrix} \\ &= P'_{mm}(\sigma) + \begin{pmatrix} R'_{mn}(P_{nn}(\sigma) + R_{nn})^* & 0 \end{pmatrix} \begin{pmatrix} 0 \\ (P'_{mm}(\sigma))_l \end{pmatrix} \\ &= P'_{mm}(\sigma) + 0 = P'_{mm}(\sigma). \end{aligned}$$

It follows that

$$\begin{aligned} M(0, \sigma, \sigma_k)^{\omega, t} &= \begin{pmatrix} P'_{mm}(\sigma)^{\omega, t} & 0 \\ \begin{pmatrix} P_{nn}(\sigma) + R_{nn} & 0 \\ \chi & 0 \end{pmatrix}^* \begin{pmatrix} 0 \\ (P'_{mm}(\sigma))_l \end{pmatrix} P'_{mm}(\sigma)^{\omega, t} \end{pmatrix} \\ &= \begin{pmatrix} P'_{mm}(\sigma)^{\omega, t} & 0 \\ \begin{pmatrix} (P_{nn}(\sigma) + R_{nn})^* & 0 \\ \chi(P_{nn}(\sigma) + R_{nn})^* & 1 \end{pmatrix} \begin{pmatrix} 0 \\ (P'_{mm}(\sigma))_l \end{pmatrix} P'_{mm}(\sigma)^{\omega, t} \end{pmatrix} \\ &= \begin{pmatrix} P'_{mm}(\sigma)^{\omega, t} & 0 \\ \begin{pmatrix} 0 \\ (P'_{mm}(\sigma))_l \end{pmatrix} P'_{mm}(\sigma)^{\omega, t} \end{pmatrix} \\ &= \begin{pmatrix} P'_{mm}(\sigma)^{\omega, t} & 0 \\ \begin{pmatrix} P'_{mm}(\sigma) & 0 \\ (P'_{mm}(\sigma))_l & P'_{mm}(\sigma)^{\omega, t} \end{pmatrix} \end{pmatrix}. \end{aligned}$$

Now, we have for the last component

$$\begin{aligned} (M(0, \sigma, \sigma_k)^{\omega, t})_{m+n+1} &= (P'_{mm}(\sigma))_l P'_{mm}(\sigma)^{\omega, t} \\ &= (P'_{mm}(\sigma) P'_{mm}(\sigma)^{\omega, t})_l \\ &= (P'_{mm}(\sigma)^{\omega, t})_l = \omega_l, \end{aligned}$$

where the third equality is by Theorem 5.5.1 of [18] and the last equality is by (13). In summary, the $(n + m + 1)^{\text{th}}$ component of the t^{th} canonical solution of $(\#)$, $(\#\#)$ is $(\sigma_k, \omega_l) = (s, v)$. As defined for ω -algebraic systems, it then follows that also the t^{th} canonical solution of $(\ast \ast \ast)$ is (s, v) . \square

As the mixed ω -algebraic system in the preceding proof does not depend on the previous discussion and since we proved that we can construct the Greibach normal form when needed, we infer the following.

Corollary 15. *Let (s, v) be a component of a canonical solution of a mixed ω -algebraic system over $S \langle \langle \Sigma^* \rangle \rangle \times S \langle \langle \Sigma^\omega \rangle \rangle$. Then we can construct an ω -algebraic system over $S \langle \langle \Sigma^* \rangle \rangle \times S \langle \langle \Sigma^\omega \rangle \rangle$ (in Greibach normal form) where (s, v) is a component of a canonical solution.*

6. Simple Reset Pushdown Automata

Now that we have proved the existence of the Greibach normal form for every ω -algebraic system and every mixed ω -algebraic system, we want to use it in the second part of the paper to show that each ω -algebraic series can be represented as the behavior of a simple ω -reset pushdown automaton. The next section will prove that result. For the proof, we will need the corresponding result for finite words as an intermediate step. We have shown in [6], that for every algebraic series r (of finite words), there exists a simple reset pushdown automaton with behavior r . We recall the construction of the simple reset pushdown automata here for the convenience of the reader, as variants of these automata will be used in Section 7 for ω -algebraic series.

Following Kuich, Salomaa [22] and Kuich [21], we introduce pushdown transitions matrices. These matrices can be considered as adjacency matrices of graphs representing automata. A special form, the reset pushdown matrices, is used for pushdown automata starting with an empty stack and allowing the automaton to push onto the empty stack. Here, we are interested in simple reset pushdown matrices, introduced in [6]. This simple form allows the automaton only to push one symbol, to pop one symbol or to ignore the stack. The corresponding automata, the simple reset pushdown automata are a generalization of the unweighted automata used in [8]. They do not use ϵ -transitions and do not allow the inspection of the topmost stack symbol.

A matrix $M \in (S^{n \times n})^{\Gamma^* \times \Gamma^*}$ is called *row-finite* if $\{\pi' \mid M_{\pi, \pi'} \neq 0\}$ is finite for all $\pi \in \Gamma^*$.

Let Γ be an alphabet, called *pushdown alphabet* and let $n \geq 1$. A matrix $\bar{M} \in (S^{n \times n})^{\Gamma^* \times \Gamma^*}$ is called a *pushdown matrix* (with *pushdown alphabet* Γ and *stateset* $\{1, \dots, n\}$) if

- (i) \bar{M} is row-finite;
- (ii) for all $\pi_1, \pi_2 \in \Gamma^*$,

$$\bar{M}_{\pi_1, \pi_2} = \begin{cases} \bar{M}_{p, \pi}, & \text{if there exist } p \in \Gamma, \pi, \pi' \in \Gamma^* \text{ with } \pi_1 = p\pi' \text{ and } \pi_2 = \pi\pi', \\ 0, & \text{otherwise.} \end{cases}$$

Intuitively, here (ii) means that the infinite pushdown matrix \bar{M} is fully represented already by the blocks $\bar{M}_{p, \pi}$ where $p \in \Gamma, \pi \in \Gamma^*$, and (i) means that only finitely many such blocks are nonzero.

Let Γ be a pushdown alphabet and $\{1, \dots, n\}, n \geq 1$, be a set of states. A *reset matrix* $M_R \in (S^{n \times n})^{\Gamma^* \times \Gamma^*}$ is a row-finite matrix such that

$$(M_R)_{\pi_1, \pi_2} = 0 \quad \text{for } \pi_1, \pi_2 \in \Gamma^* \text{ with } \pi_1 \neq \epsilon.$$

A *reset pushdown matrix* $M \in (S^{n \times n})^{\Gamma^* \times \Gamma^*}$ is the sum of a reset matrix M_R and a pushdown matrix \bar{M} ,

$$M = M_R + \bar{M}.$$

Intuitively, a reset pushdown matrix is similar to a pushdown matrix with the additional possibility to push onto the empty stack, i.e., $M_{\epsilon, \pi}$ is allowed to be nonzero. Note that the entries of reset pushdown matrices are determined by finitely many values because it is row-finite and property (ii) of pushdown matrices ensures that the value of $M_{p\pi', \pi\pi'}$ is equal to (and therefore can be derived from) $M_{p, \pi}$.

A reset pushdown matrix M is called *simple* if, $M \in ((S\langle\Sigma\rangle)^{n \times n})^{\Gamma^* \times \Gamma^*}$ for some $n \geq 1$, and for all $p, p_1 \in \Gamma$,

$$M_{p,\epsilon}, M_{p,p} = M_{\epsilon,\epsilon} \text{ and } M_{p,p_1 p} = M_{\epsilon,p_1},$$

are the only blocks $M_{\pi,\pi'}$, where $\pi \in \{\epsilon, p\}$ and $\pi' \in \Gamma^*$, that may be unequal to the zero matrix 0.

Hence, a simple reset pushdown matrix M is defined by its blocks $M_{\epsilon,\epsilon}$ and $M_{p,\epsilon}, M_{\epsilon,p}$ ($p \in \Gamma$). Intuitively, the automata will only be allowed to ignore the stack (modeled by $M_{\epsilon,\epsilon}$), pop one symbol ($M_{p,\epsilon}$) or push one symbol ($M_{\epsilon,p}$). Note also that the matrix $M \in ((S\langle\Sigma\rangle)^{n \times n})^{\Gamma^* \times \Gamma^*}$ forbids ϵ -transitions. Moreover, the equalities $M_{p,p} = M_{\epsilon,\epsilon}$ and $M_{p,p_1 p} = M_{\epsilon,p_1}$ imply that the next transition does not depend on the topmost symbol of the stack except when popping it (modeled by $M_{p,\epsilon}$).

A *reset pushdown automaton* (with input alphabet Σ) $\mathfrak{A} = (n, \Gamma, I, M, P)$ is given by

- a set of states $\{1, \dots, n\}$, $n \geq 1$,
- a pushdown alphabet Γ ,
- a reset pushdown matrix $M \in ((S\langle\Sigma \cup \{\epsilon\}\rangle)^{n \times n})^{\Gamma^* \times \Gamma^*}$ called *transition matrix*,
- a row vector $I \in (S\langle\{\epsilon\}\rangle)^{1 \times n}$, called *initial state vector*,
- a column vector $P \in (S\langle\{\epsilon\}\rangle)^{n \times 1}$, called *final state vector*.

The *behavior* $\|\mathfrak{A}\|$ of a reset pushdown automaton \mathfrak{A} is defined by

$$\|\mathfrak{A}\| = I(M^*)_{\epsilon,\epsilon} P.$$

A reset pushdown automaton $\mathfrak{A} = (n, \Gamma, I, M, P)$ is called *simple* if M is a simple reset pushdown matrix.

Example 18 will show a simple reset pushdown automaton and the corresponding simple reset pushdown matrix.

Given a series $r \in S^{\text{alg}}\langle\langle\Sigma^*\rangle\rangle$, we want to construct a simple reset pushdown automaton with behavior r . By Theorems 5.10 and 5.4 of [21], r is a component of the unique solution of a strict algebraic system in Greibach normal form.

We only consider the algebraic series r with $(r, \epsilon) = 0$; cf. [6] for the other case. So we assume without loss of generality that r is the x_1 -component of the unique solution of the algebraic system (14) with variables x_1, \dots, x_n

$$x_i = p_i, \quad 1 \leq i \leq n,$$

of the form

$$\begin{aligned} x_i = & \sum_{1 \leq j, k \leq n} \sum_{a \in \Sigma} (p_i, ax_j x_k) ax_j x_k + \\ & \sum_{1 \leq j \leq n} \sum_{a \in \Sigma} (p_i, ax_j) ax_j + \\ & \sum_{a \in \Sigma} (p_i, a) a. \end{aligned} \tag{14}$$

As in [6], we now construct the simple reset pushdown automaton $\mathfrak{A}_m = (n+1, \Gamma, I_m, M, P)$, $1 \leq m \leq n$, with $r = \|\mathfrak{A}_1\|$ as follows:

We let $\Gamma = \{x_1, \dots, x_n\}$; we also denote the state $n+1$ by f ; the entries of M of the form $(M_{x_k, x_k})_{i,j}, (M_{x_k, \epsilon})_{i,j}, (M_{\epsilon, x_k})_{i,j}, (M_{\epsilon, \epsilon})_{i,j}, (M_{\epsilon, \epsilon})_{i,f}$, $1 \leq i, j, k \leq n$, that may be unequal to 0 are

$$\begin{aligned} (M_{\epsilon, x_k})_{i,j} &= \sum_{a \in \Sigma} (p_i, ax_j x_k) a, \\ (M_{x_k, x_k})_{i,j} &= (M_{\epsilon, \epsilon})_{i,j} = \sum_{a \in \Sigma} (p_i, ax_j) a, \\ (M_{x_k, \epsilon})_{i,k} &= (M_{x_k, x_k})_{i,f} = (M_{\epsilon, \epsilon})_{i,f} = \sum_{a \in \Sigma} (p_i, a) a; \end{aligned}$$




Figure 1: Example 18: Simple reset pushdown automaton, where $\downarrow X$ means push symbol X , $\uparrow X$ means pop X , and $\#$ leaves the stack unaltered. All shown transitions have a weight equal to the natural number 0 except the two transitions reading letter a and pushing B onto the stack that have weight 1. All other possible transitions have weight $-\infty$.

we further put $(I_m)_m = \epsilon$, $(I_m)_i = 0$ for $1 \leq i \leq m-1$ and $m+1 \leq i \leq n+1$; finally let $P_f = \epsilon$ and $P_j = 0$ for $1 \leq j \leq n$;

The following motivation will be essential for our later construction for ω -pushdown automata. Intuitively, the variables in the algebraic system are simulated by states in the simple reset pushdown automaton \mathfrak{A}_m . By the Greibach normal form, only two variables on the right-hand side are allowed. The first is modeled directly by changing the state, the second is pushed to the pushdown tape and the state is changed to it later when the variable is popped again. The special final state f will only be used as the last state.

Note that $(M_{x_k, x_l})_{i, f}$ allows the automaton to change to the final state with a non-empty pushdown tape. This is an artificial addition to fit the definition of simple reset pushdown matrices. If the simple reset automaton is not popping a symbol from the pushdown tape, it cannot distinguish between different pushdown states. Even though the automaton can enter the final state too early, it can not continue from there as it is a sink.

Observe that $\|\mathfrak{A}_m\| = ((M^*)_{\epsilon, \epsilon})_{s, f}$ for all $1 \leq m \leq n$.

This simple reset pushdown matrix M is called the simple pushdown matrix *induced* by the Greibach normal form (14). The simple reset pushdown automata \mathfrak{A}_m , $1 \leq m \leq n$, are called the simple reset pushdown automata *induced* by the Greibach normal form (14).

The following (main) theorem of [6] states that the behavior of the simple reset pushdown automata induced by the Greibach normal form (14) is the unique solution of the original algebraic system (14).

Theorem 16 (Theorem 11 of [6]). *The unique solution of the algebraic system (14) is*

$$(\|\mathfrak{A}_1\|, \dots, \|\mathfrak{A}_n\|) = (((M^*)_{\epsilon, \epsilon})_{1, f}, \dots, ((M^*)_{\epsilon, \epsilon})_{n, f}).$$

Corollary 17 (Corollary 12 of [6]). *Let $r \in S^{alg}(\langle \Sigma^* \rangle)$. Then there exists a simple reset pushdown automaton with behavior r .*

Example 18. Consider the semiring $\bar{\mathbb{N}}(\langle \Sigma^* \rangle)$ for the arctic semiring $\langle \bar{\mathbb{N}}, \max, +, -\infty, 0 \rangle$ with $\bar{\mathbb{N}} = \mathbb{N} \cup \{-\infty, \infty\}$. Analogously to Example 7, we let $\mathbb{0} = -\infty$ and $\mathbb{1} = 0$ and we note that in the following, 1 stands for the natural number 1.

We define the algebraic system

$$\begin{aligned} S &= aMS + 1aNT + 1aN & T &= aMT + aM \\ M &= b + aMB & N &= b + 1aNB \\ B &= b \end{aligned}$$

with the variables S, T, M, N, B . These variables facilitate reading the equations, but for comparison with equation (14), consider the variable mapping $x_1 = T, x_2 = S, x_3 = N, x_4 = M, x_5 = B$.

Now, the variable M derives a string $a^n b^{n+1}$ for $n \in \mathbb{N}$. The variable N does the same but at the same time produces the weight n . The variables S and T add another a .

Let $L = \{a^n b^n \mid n \geq 1\}$. In total, the second component (i.e., with S being the start variable) of the least solution is u with $(u, a^{n_1} b^{n_1} a^{n_2} b^{n_2} \dots a^{n_k} b^{n_k}) = \max n_i$ for $k \geq 1$ and $(u, w) = -\infty$ for $w \notin L^+$.

From this, we can construct a simple reset pushdown automaton $\mathfrak{A}_2 = (n, \Gamma, I, M, P)$ as shown in Figure 1. Thus, we have $n = 6$, $\Gamma = \{T, S, N, M, B\}$. The initial state vector is $I_2 = \epsilon$ and $I_i = 0$ for $i \neq 2$. The final state vector is $P_6 = \epsilon$ and $P_i = 0$ for $i \neq 6$. The simple reset pushdown matrix is defined as

$$M = \begin{pmatrix} M_{\epsilon, \epsilon} & M_{\epsilon, T} & M_{\epsilon, S} & M_{\epsilon, N} & M_{\epsilon, M} & M_{\epsilon, B} & \dots \\ M_{T, \epsilon} & M_{\epsilon, \epsilon} & 0 & 0 & 0 & 0 & \dots \\ M_{S, \epsilon} & 0 & M_{\epsilon, \epsilon} & 0 & 0 & 0 & \dots \\ M_{N, \epsilon} & 0 & 0 & M_{\epsilon, \epsilon} & 0 & 0 & \dots \\ M_{M, \epsilon} & 0 & 0 & 0 & M_{\epsilon, \epsilon} & 0 & \dots \\ M_{B, \epsilon} & 0 & 0 & 0 & 0 & M_{\epsilon, \epsilon} & \dots \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \ddots \end{pmatrix},$$

with, for instance

$$M_{\epsilon, \epsilon} = \begin{pmatrix} 0 & 0 & 0 & a & 0 & 0 \\ 0 & 0 & 1a & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & b \\ 0 & 0 & 0 & 0 & 0 & b \\ 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix} \text{ and } M_{\epsilon, B} = \begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1a & 0 & 0 & 0 \\ 0 & 0 & 0 & a & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix}.$$

The rest of the matrix M can be inferred by the rules of pushdown matrices. The behavior $\|\mathfrak{A}_2\|$ is equal to the second component of the least solution of the algebraic system above.

7. Simple ω -Reset Pushdown Automata

In this section, we will prove that for every ω -algebraic series r , there exists a simple ω -reset pushdown automaton with behavior r . We first prove some results for infinite applications of simple reset pushdown matrices. Then we introduce simple ω -reset pushdown automata, and the main theorem will show that they can recognize all ω -algebraic series.

In the sequel, (S, V) is a complete semiring-semimodule pair.

We will use sets P_l comprising infinite sequences over $\{1, \dots, n\}$ as defined in [9]:

$$P_l = \{(j_1, j_2, \dots) \in \{1, \dots, n\}^\omega \mid j_t \leq l \text{ for infinitely many } t \geq 1\}.$$

We obtain, for a reset pushdown matrix $M \in (S^{n \times n})^{\Gamma^* \times \Gamma^*}$, $\pi \in \Gamma^+$ and for $1 \leq j \leq n$,

$$((M^{\omega, l})_\pi)_j = \sum_{\pi_1, \pi_2, \dots \in \Gamma^*} \sum_{(j_1, j_2, \dots) \in P_l} (M_{\pi, \pi_1})_{j, j_1} (M_{\pi_1, \pi_2})_{j_1, j_2} (M_{\pi_2, \pi_3})_{j_2, j_3} \dots \quad (15)$$

Observe the following summation identity: Assume that M_1, M_2, \dots are matrices in $S^{n \times n}$. Then for $0 \leq l \leq n$, $1 \leq j \leq n$, and $m \geq 1$, we have

$$\sum_{(j_1, j_2, \dots) \in P_l} (M_1)_{j, j_1} (M_2)_{j_1, j_2} \dots = \sum_{1 \leq j_1, \dots, j_m \leq n} (M_1)_{j, j_1} \dots (M_m)_{j_{m-1}, j_m} \sum_{(j_{m+1}, j_{m+2}, \dots) \in P_l} (M_{m+1})_{j_m, j_{m+1}} \dots$$

By Theorem 5.5.1 of Ésik, Kuich [18] we obtain, for a finite matrix M and for $0 \leq l \leq n$, the equality $MM^{\omega, l} = M^{\omega, l}$. By Theorem 6 of Droste, Ésik, Kuich [9], we have a similar result for pushdown matrices. We will now show the same equality for a reset pushdown matrix M .

Theorem 19. Let (S, V) be a complete semiring-semimodule pair and let further $M \in (S^{n \times n})^{\Gamma^* \times \Gamma^*}$ be a reset pushdown transition matrix. Then, for $0 \leq l \leq n$,

$$M^{\omega, l} = MM^{\omega, l}.$$

Proof. We obtain for $\pi_0 \in \Gamma^*$ and $1 \leq j_0 \leq n$,

$$\begin{aligned} ((MM^{\omega, l})_{\pi_0})_{j_0} &= \sum_{\pi \in \Gamma^*} \sum_{1 \leq j \leq n} (M_{\pi_0, \pi})_{j_0, j} \sum_{\pi_1, \pi_2, \dots \in \Gamma^*} \sum_{(j_1, j_2, \dots) \in P_l} (M_{\pi, \pi_1})_{j, j_1} (M_{\pi_1, \pi_2})_{j_1, j_2} \cdots \\ &= \sum_{\pi, \pi_1, \pi_2, \dots \in \Gamma^*} \sum_{(j, j_1, j_2, \dots) \in P_l} (M_{\pi_0, \pi})_{j_0, j} (M_{\pi, \pi_1})_{j, j_1} (M_{\pi_1, \pi_2})_{j_1, j_2} \cdots \\ &= ((M^{\omega, l})_{\pi_0})_{j_0}. \end{aligned}$$

□

The following two theorems compare reset pushdown matrices to pushdown matrices in the course of an infinite application. They state that either the topmost stack symbol p is popped or the reset pushdown matrix behaves similar to pushdown matrices.

Theorem 20. Let (S, V) be a complete semiring-semimodule pair. Let M be a reset pushdown matrix. Then

$$(M^{\omega})_p = (\bar{M}^{\omega})_p + (\bar{M}^*)_{p, \epsilon} (M^{\omega})_{\epsilon}, \text{ for any } p \in \Gamma.$$

Proof. We obtain, for $p \in \Gamma$,

$$\begin{aligned} (M^{\omega})_p &= \sum_{\pi_1, \pi_2, \dots \in \Gamma^+} M_{p, \pi_1} M_{\pi_1, \pi_2} \cdots + \sum_{l \geq 1} \sum_{\pi_1, \dots, \pi_{l-1} \in \Gamma^+} M_{p, \pi_1} \cdots M_{\pi_{l-1}, \epsilon} (M^{\omega})_{\epsilon} \\ &= (\bar{M}^{\omega})_p + \sum_{l \geq 1} (\bar{M}^l)_{p, \epsilon} (M^{\omega})_{\epsilon} \\ &= (\bar{M}^{\omega})_p + (\bar{M}^*)_{p, \epsilon} (M^{\omega})_{\epsilon}. \end{aligned}$$

□

Theorem 21. Let (S, V) be a complete semiring-semimodule pair. Let M be a reset pushdown matrix and $0 \leq l \leq n$. Then

$$(M^{\omega, l})_p = (\bar{M}^{\omega, l})_p + (\bar{M}^*)_{p, \epsilon} (M^{\omega, l})_{\epsilon}, \text{ for any } p \in \Gamma.$$

Proof. We use the proof of Theorem 20. We obtain, for $p \in \Gamma$, $0 \leq l \leq n$ and $1 \leq j \leq n$,

$$\begin{aligned} ((M^{\omega, l})_p)_j &= \sum_{\pi_1, \pi_2, \dots \in \Gamma^*} \sum_{(j_1, j_2, \dots) \in P_l} (M_{p, \pi_1})_{j, j_1} (M_{\pi_1, \pi_2})_{j_1, j_2} \cdots \\ &= \sum_{\pi_1, \pi_2, \dots \in \Gamma^+} \sum_{(j_1, j_2, \dots) \in P_l} (M_{p, \pi_1})_{j, j_1} (M_{\pi_1, \pi_2})_{j_1, j_2} \cdots \\ &\quad + \sum_{l \geq 1} \sum_{\pi_1, \dots, \pi_{l-1} \in \Gamma^+} \sum_{\substack{(j_1, j_2, \dots) \in P_l \\ \pi_{l+1}, \pi_{l+2}, \dots \in \Gamma^*}} (M_{p, \pi_1})_{j, j_1} \cdots (M_{\pi_{l-1}, \epsilon})_{j_{l-1}, j_l} \cdot (M_{\epsilon, \pi_{l+1}})_{j_l, j_{l+1}} (M_{\pi_{l+1}, \pi_{l+2}})_{j_{l+1}, j_{l+2}} \cdots \\ &= ((\bar{M}^{\omega, l})_p)_j + \sum_{l \geq 1} \sum_{\pi_1, \dots, \pi_{l-1} \in \Gamma^+} \sum_{1 \leq j_1, \dots, j_l \leq n} (M_{p, \pi_1})_{j, j_1} \cdots (M_{\pi_{l-1}, \epsilon})_{j_{l-1}, j_l} \cdot \\ &\quad \sum_{\pi_{l+1}, \pi_{l+2}, \dots \in \Gamma^*} \sum_{(j_{l+1}, j_{l+2}, \dots) \in P_l} (M_{\epsilon, \pi_{l+1}})_{j_l, j_{l+1}} (M_{\pi_{l+1}, \pi_{l+2}})_{j_{l+1}, j_{l+2}} \cdots \\ &= ((\bar{M}^{\omega, l})_p)_j + \sum_{l \geq 1} \sum_{1 \leq j' \leq n} ((\bar{M}^l)_{p, \epsilon})_{j, j'} ((M^{\omega, l})_{\epsilon})_{j'} \\ &= ((\bar{M}^{\omega, l})_p)_j + (\bar{M}^*)_{p, \epsilon} (M^{\omega, l})_{\epsilon}{}_j. \end{aligned}$$

□

For simple reset pushdown matrices, the following two lemmas state that infinite paths starting with symbol p on the pushdown tape can either ignore that symbol or pop it and then continue with an infinite path from the empty tape.

Lemma 22. Let (S, V) be a complete semiring-semimodule pair. Let M be a simple reset pushdown matrix. Then, for $p \in \Gamma$,

$$(M^\omega)_p = (M^\omega)_\epsilon + (M^*)_{\epsilon, \epsilon} M_{p, \epsilon} (M^\omega)_\epsilon.$$

Proof. We obtain, for $p \in \Gamma$,

$$\begin{aligned} (M^\omega)_p &= \sum_{\pi_1, \pi_2, \dots \in \Gamma^*} M_{p, \pi_1} M_{\pi_1, \pi_2} \cdots \\ &= \sum_{\pi_1, \pi_2, \dots \in \Gamma^*} M_{p, \pi_1 p} M_{\pi_1 p, \pi_2 p} \cdots + \sum_{t \geq 0} \sum_{\pi_1, \dots, \pi_{t-1} \in \Gamma^*} M_{p, \pi_1 p} \cdots M_{\pi_{t-1} p, p} M_{p, \epsilon} (M^\omega)_\epsilon \\ &= (M^\omega)_\epsilon + \left(\sum_{t \geq 0} \sum_{\pi_1, \dots, \pi_{t-1} \in \Gamma^*} M_{\epsilon, \pi_1} \cdots M_{\pi_{t-1}, \epsilon} \right) M_{p, \epsilon} (M^\omega)_\epsilon \\ &= (M^\omega)_\epsilon + \sum_{t \geq 0} (M^t)_{\epsilon, \epsilon} M_{p, \epsilon} (M^\omega)_\epsilon \\ &= (M^\omega)_\epsilon + (M^*)_{\epsilon, \epsilon} M_{p, \epsilon} (M^\omega)_\epsilon. \end{aligned}$$

□

Lemma 23. Let (S, V) be a complete semiring-semimodule pair. Let M be a simple reset pushdown matrix. Then, for $p \in \Gamma$ and $0 \leq l \leq n$,

$$(M^{\omega, l})_p = (M^{\omega, l})_\epsilon + (M^*)_{\epsilon, \epsilon} M_{p, \epsilon} (M^{\omega, l})_\epsilon.$$

Proof. We use the proof of Lemma 22. We obtain, for $p \in \Gamma$, $0 \leq l \leq n$ and $1 \leq j \leq n$,

$$\begin{aligned} ((M^{\omega, l})_p)_j &= \sum_{\pi_1, \pi_2, \dots \in \Gamma^*} \sum_{(j_1, j_2, \dots) \in P_l} (M_{p, \pi_1})_{j, j_1} (M_{\pi_1, \pi_2})_{j_1, j_2} \cdots \\ &= \sum_{\pi_1, \pi_2, \dots \in \Gamma^*} \sum_{(j_1, j_2, \dots) \in P_l} (M_{p, \pi_1 p})_{j, j_1} (M_{\pi_1 p, \pi_2 p})_{j_1, j_2} \cdots \\ &\quad + \sum_{t \geq 0} \sum_{\pi_1, \pi_2, \dots \in \Gamma^*} \sum_{(j_1, j_2, \dots) \in P_l} (M_{p, \pi_1 p})_{j, j_1} \cdots (M_{\pi_{t-1} p, p})_{j_{t-1}, j_t} \cdot \\ &\quad (M_{p, \epsilon})_{j_t, j_{t+1}} (M_{\epsilon, \pi_{t+2}})_{j_{t+1}, j_{t+2}} (M_{\pi_{t+2}, \pi_{t+3}})_{j_{t+2}, j_{t+3}} \cdots \\ &= ((M^{\omega, l})_\epsilon)_j \\ &\quad + \sum_{t \geq 0} \sum_{\pi_1, \dots, \pi_{t-1} \in \Gamma^*} \sum_{1 \leq j_1, \dots, j_{t+1} \leq n} (M_{p, \pi_1 p})_{j, j_1} \cdots (M_{\pi_{t-1} p, p})_{j_{t-1}, j_t} (M_{p, \epsilon})_{j_t, j_{t+1}} \cdot \\ &\quad \sum_{\pi_{t+2}, \pi_{t+3}, \dots \in \Gamma^*} \sum_{(j_{t+2}, j_{t+3}, \dots) \in P_l} (M_{\epsilon, \pi_{t+2}})_{j_{t+1}, j_{t+2}} (M_{\pi_{t+2}, \pi_{t+3}})_{j_{t+2}, j_{t+3}} \cdots \\ &= ((M^{\omega, l})_\epsilon)_j + \sum_{t \geq 0} \sum_{\pi_1, \dots, \pi_{t-1} \in \Gamma^*} \sum_{1 \leq j_1, \dots, j_{t+1} \leq n} (M_{\epsilon, \pi_1})_{j, j_1} \cdots (M_{\pi_{t-1}, \epsilon})_{j_{t-1}, j_t} \cdot \\ &\quad (M_{p, \epsilon})_{j_t, j_{t+1}} ((M^{\omega, l})_\epsilon)_{j_{t+1}} \\ &= ((M^\omega)_\epsilon)_j + \sum_{t \geq 0} \sum_{1 \leq j', j'' \leq n} ((M^t)_{\epsilon, \epsilon})_{j, j'} (M_{p, \epsilon})_{j', j''} ((M^{\omega, l})_\epsilon)_{j''} \\ &= ((M^\omega)_\epsilon + (M^*)_{\epsilon, \epsilon} M_{p, \epsilon} (M^{\omega, l})_\epsilon)_j. \end{aligned}$$

□

Next, an ω -reset pushdown automaton

$$\mathfrak{A} = (n, \Gamma, I, M, P, l)$$

is given by a reset pushdown automaton (n, Γ, I, M, P) and an integer l with $0 \leq l \leq n$, which indicates that $1, \dots, l$ are the repeated states of \mathfrak{A} . The behavior $\|\mathfrak{A}\|$ of this ω -reset pushdown automaton \mathfrak{A} is defined by

$$\|\mathfrak{A}\| = I(M^*)_{\epsilon, \epsilon} P + I(M^{\omega, l})_\epsilon.$$

The ω -reset pushdown automaton $\mathfrak{A} = (n, \Gamma, I, M, P, l)$ is called *simple* if M is a simple reset pushdown matrix.




Figure 2: Example 24: Simple ω -reset pushdown automaton, where (\downarrow, X) means push symbol X , (\uparrow, X) means pop X , and $\#$ leaves the stack unaltered. All transitions shown have a weight equal to the natural number 0 except the two transitions reading letter a and pushing a symbol onto the stack that have weight 1. All other possible transitions have weight ∞ .

Example 24. Figure 2 shows a simple ω -reset pushdown automaton $\mathcal{A} = (4, \Gamma, I, M, P, 1)$ over the quemiring $\mathbb{N}^\infty \langle \langle \Sigma^* \rangle \rangle \times \mathbb{N}^\infty \langle \langle \Sigma^\omega \rangle \rangle$ for the tropical semiring $\langle \mathbb{N}^\infty, \min, +, 0 = \infty, 1 = 0 \rangle$ with $\Sigma = \{a, b, c\}$, $\Gamma = \{Z_0, X\}$, $I_2 = 0$, $I_i = \infty$ for $i \neq 2$ and $P_i = \infty$ for all $1 \leq i \leq 4$. Then the adjacency matrix M of the automaton shown in Figure 2 is a simple reset pushdown matrix. As an indication, M is defined with $(M_{\epsilon, \epsilon})_{1,1} = 0c$, $(M_{\epsilon, \epsilon})_{2,1} = 0c$, $(M_{\epsilon, Z_0})_{2,3} = 1a$, etc., resulting in e.g.,

$$M_{\epsilon, \epsilon} = \begin{pmatrix} 0c & 0 & 0 & 0 \\ 0c & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix} \text{ and finally } M = \begin{pmatrix} M_{\epsilon, \epsilon} & M_{\epsilon, Z_0} & M_{\epsilon, X} & \cdots \\ M_{Z_0, \epsilon} & M_{\epsilon, \epsilon} & 0 & \cdots \\ M_{X, \epsilon} & 0 & M_{\epsilon, \epsilon} & \cdots \\ \vdots & \vdots & \vdots & \ddots \end{pmatrix},$$

where the excluded part of M can be derived from the rules of pushdown and simple reset pushdown matrices. The automaton \mathcal{A} has the behavior $a^n b^n c^\omega \mapsto n$, similar to the mixed ω -algebraic system in Example 4.

Example 25. Reconsider Example 18. We define the simple ω -reset pushdown automaton $\mathcal{A}_2 = (6, \Gamma, I, M, P, 1)$ where we define the state ordering T, S, N, M, B, F to make state T Büchi-accepting. The behavior in the semiring part is equal to before; the behavior in the semimodule part is u with $(u, a^{n_1} b^{n_1} a^{n_2} b^{n_2} \dots) = \max n_i$ and $(u, w) = -\infty$ for $w \notin \{a^n b^n \mid n \geq 1\}^\omega$.

Example 26. Consider the ω -algebraic system

$$\begin{aligned} y_1 &= a + cy_1 \\ y_2 &= ay_1 y_2 + ay_1. \end{aligned} \tag{16}$$

We will consider the second component of the first canonical solution, i.e., variable y_1 is Büchi-accepting and variable y_2 is the start variable.

The ω -algebraic system induces the following mixed ω -algebraic system

$$\begin{aligned} x_1 &= a + cx_1 & z_1 &= cz_1 \\ x_2 &= ax_1 x_2 + ax_1 & z_2 &= az_1 + ax_1 z_2. \end{aligned} \tag{17}$$

The least solution of $x = p(x)$ is

$$\sigma = \begin{pmatrix} c^* a \\ (ac^* a)^+ \end{pmatrix}.$$

Now, we write the linear system $z = \varrho(\sigma)z$ in the matrix form and compute the first canonical solution.


$$\begin{aligned} \varrho(\sigma)^{\omega, 1} &= \begin{pmatrix} c & 0 \\ a & a(c^* a) \end{pmatrix}^{\omega, 1} \\ &= \begin{pmatrix} (c + 0(aa)^* a)^\omega \\ (ac^* a)^* a(c + 0(aa)^* a)^\omega \end{pmatrix} \\ &= \begin{pmatrix} c^\omega \\ (ac^* a)^* ac^\omega \end{pmatrix} =: \begin{pmatrix} \omega_1^{(1)} \\ \omega_2^{(1)} \end{pmatrix} = \omega^{(1)} \end{aligned}$$

Note that the second component, $\omega_2^{(1)}$, does not contain the ω -words $(ac^*a)^\omega$ even though for an unweighted ω -context-free grammar corresponding to (16), the derivation

$$y_2 \rightarrow ay_1y_2 \rightarrow (aa)y_2 \rightarrow (aa)ay_1y_2 \rightarrow (aa)^2y_2 \rightarrow^\omega a^\omega$$

would be successful even with only y_1 Büchi-accepting. The difference is due to the fact that y_1 is not significant in the ω -algebraic system above because it is exchanged by x_1 in the derivation (for more information, see [18] pp. 140 ff.).

Now, we look at the simple ω -reset pushdown automaton induced by ω -algebraic system (16):




The behavior of this automaton is

$$\begin{aligned} & (((M^*)_{\epsilon, \epsilon})_{1, f}, ((M^*)_{\epsilon, \epsilon})_{2, f}; ((M^{\omega, 1})_{\epsilon})_1, ((M^{\omega, 1})_{\epsilon})_2) \\ = & (c^*a, (ac^*a)^+; c^\omega, (ac^*a)^*ac^\omega + (ac^*a)^\omega) \end{aligned}$$

Here, the first two components are equal to σ , as desired. But the last component differs from $\omega_2^{(1)}$; the last component is however equal to the behavior of unweighted ω -context-free grammars.

Note that the desired component $\omega_2^{(1)} = (ac^*a)^*ac^\omega$ is not recognized by this automaton, even when changing the Büchi-accepting states. If no states are Büchi-accepting, the behavior is 0, if all of them are Büchi-accepting, we have the same behavior as above. If only state 2 is Büchi-accepting (can be achieved by renaming), we only recognize $(ac^*a)^\omega$.

We now propose a different construction; this new construction models exactly the canonical solutions of mixed ω -algebraic systems. The following is the simple ω -reset pushdown automaton induced by the mixed ω -algebraic system (17); this new construction will be defined after the example. Basically, the construction is similar to the old construction but it differentiates between variables x and z ; it therefore uses the states $x_1, \dots, x_n, z_1, \dots, z_n$:



This simple ω -reset pushdown automaton has exactly the behavior $(\sigma, \omega^{(1)})$. This means, if only z_1 is Büchi-accepting, then the automaton does not allow the run $(ac^*a)^\omega$.

The rest of the paper will show that in general, the l^{th} canonical solution of a mixed ω -algebraic system $x = p(x), z = \varrho(x)z$ is exactly the behavior of the simple ω -reset pushdown automaton induced by $x = p(x), z = \varrho(x)z$.

Given a series $r \in S^{\text{alg}}\langle\langle\Sigma^*\rangle\rangle \times S^{\text{alg}}\langle\langle\Sigma^\omega\rangle\rangle$, we want to construct a simple ω -reset pushdown automaton with behavior r . By Theorem 14 and Theorem 5, r is a component of a canonical solution of an ω -algebraic system (18)

(compare this to the algebraic system (14)) in Greibach normal form over the quemiring $S\langle\langle\Sigma^*\rangle\rangle \times S\langle\langle\Sigma^\omega\rangle\rangle$,

$$\begin{aligned} y_i = & \sum_{1 \leq j, k \leq n} \sum_{a \in \Sigma} (p_i, ay_j y_k) ay_j y_k + \\ & \sum_{1 \leq j \leq n} \sum_{a \in \Sigma} (p_i, ay_j) ay_j + \\ & \sum_{a \in \Sigma} (p_i, a) a. \end{aligned} \quad (18)$$

The variables of this system are y_i , ($1 \leq i \leq n$); they are variables for $S\langle\langle\Sigma^*\rangle\rangle \times S\langle\langle\Sigma^\omega\rangle\rangle$. The system (18) induces the following mixed ω -algebraic system:

$$\begin{aligned} x_i = & \sum_{1 \leq j, k \leq n} \sum_{a \in \Sigma} (p_i, ay_j y_k) ax_j x_k + \\ & \sum_{1 \leq j \leq n} \sum_{a \in \Sigma} (p_i, ay_j) ax_j + \\ & \sum_{a \in \Sigma} (p_i, a) a. \end{aligned} \quad (19)$$

and

$$z_i = \sum_{1 \leq j, k \leq n} \sum_{a \in \Sigma} (p_i, ay_j y_k) a(z_j + x_j z_k) + \sum_{1 \leq j \leq n} \sum_{a \in \Sigma} (p_i, ay_j) az_j \quad (20)$$

But this system hides information, for instance, $y_j y_k$ will never be derived by two consecutive variables $z_j z_k$ of $S\langle\langle\Sigma^\omega\rangle\rangle$. Our new construction is therefore based on the following mixed ω -algebraic system that can be gained from the last system (14), (19) by renaming:

$$\begin{aligned} x_i = & \sum_{1 \leq j, k \leq n} \sum_{a \in \Sigma} (p_i, ax_j x_k) ax_j x_k + \\ & \sum_{1 \leq j \leq n} \sum_{a \in \Sigma} (p_i, ax_j) ax_j + \\ & \sum_{a \in \Sigma} (p_i, a) a. \end{aligned} \quad (21)$$

and

$$z_i = \sum_{1 \leq j, k \leq n} \sum_{a \in \Sigma} (p_i, ax_j z_k) ax_j z_k + \sum_{1 \leq j \leq n} \sum_{a \in \Sigma} (p_i, az_j) az_j \quad (22)$$

We now want to construct a simple ω -reset pushdown automaton. Here, we introduce our new construction. Let $\mathfrak{A}_m^l = (2n+1, \Gamma, I_m, M, P, l)$, $1 \leq m \leq n$, $0 \leq l \leq n$, be defined as follows:

We let $\Gamma = \{X_1, \dots, X_n, Z_1, \dots, Z_n\}$; we denote the states $1, \dots, 2n+1$ by $z_1, \dots, z_n, x_1, \dots, x_n, f$; the entries of M of the form $(M_{\pi, \pi'})_{v, v'}$ for $1 \leq v, v' \leq 2n+1$ and for $\pi, \pi' \in \Gamma^*$ with $|\pi|, |\pi'| \leq 1$ that may be unequal to 0 are

$$\begin{aligned} (M_{\epsilon, X_k})_{x_i, x_j} &= \sum_{a \in \Sigma} (p_i, ax_j x_k) a, \\ (M_{Z_k, Z_k})_{x_i, x_j} &= (M_{X_k, X_k})_{x_i, x_j} = (M_{\epsilon, \epsilon})_{x_i, x_j} = \sum_{a \in \Sigma} (p_i, ax_j) a, \\ (M_{Z_k, \epsilon})_{x_i, z_k} &= (M_{X_k, \epsilon})_{x_i, x_k} = (M_{Z_k, Z_k})_{x_i, f} = (M_{X_k, X_k})_{x_i, f} = (M_{\epsilon, \epsilon})_{x_i, f} = \sum_{a \in \Sigma} (p_i, a) a, \\ (M_{Z_k, Z_k})_{z_i, z_j} &= (M_{X_k, X_k})_{z_i, z_j} = (M_{\epsilon, \epsilon})_{z_i, z_j} = \sum_{a \in \Sigma} (p_i, az_j) a, \\ (M_{\epsilon, Z_k})_{z_i, x_j} &= \sum_{a \in \Sigma} (p_i, ax_j z_k) a, \end{aligned}$$

for $1 \leq i, j, k \leq n$; we further put $(I_m)_{x_m} = (I_m)_{z_m} = \epsilon$, and $(I_m)_{x_i} = (I_m)_{z_i} = 0$ for $1 \leq i \leq m-1$ and $m+1 \leq i \leq n$ and $(I_m)_f = 0$; finally let $P_f = \epsilon$ and $P_j = 0$ for $1 \leq j \leq 2n$;

In the following, we assume that $r \in S^{\text{alg}}\langle\langle\Sigma^*\rangle\rangle \times S^{\text{alg}}\langle\langle\Sigma^\omega\rangle\rangle$ is the m^{th} component of the l^{th} canonical solution of (18). We want to show that for the l^{th} canonical solution $\tau = (\sigma, \omega)$ of (14), (20), and therefore also of (18), we have $\tau_m = \sigma_m + \omega_m = \|\mathfrak{A}_m^l\|$.

This simple reset pushdown matrix M is called the simple reset pushdown matrix *induced* by the Greibach normal form (14), (20). The simple ω -reset pushdown automata \mathfrak{A}_m^l ($1 \leq m \leq n$, $0 \leq l \leq n$) are called the simple ω -reset pushdown automata *induced* by the Greibach normal form (14), (20).

For the rest of the paper, we will use the following notation (cf. [22], page 179). Note that $M \in (S^{k \times k})^{\Gamma^* \times \Gamma^*}$ for $k = 2n+1$. By isomorphism, we can transform this into $\widehat{M} \in (S^{\Gamma^* \times \Gamma^*})^{k \times k}$. We then have $(M_{\pi, \pi'})_{v, v'} = (\widehat{M}_{v, v'})_{\pi, \pi'}$ for $\pi, \pi' \in \Gamma^*$ and $1 \leq v, v' \leq 2n+1$. (By the notation $1 \leq v \leq 2n+1$, we mean v can be any of the states $z_1, \dots, z_n, x_1, \dots, x_n, f$.)

Example 27. This notation allows us to add up matrices with suitable pushdown indexes while still keeping the information of the states. For instance, note that

$$\sum_{1 \leq k \leq n} \sum_{\pi \in \Gamma^*} (\widehat{M}_{z_i, x_k})_{\epsilon, \pi} (\widehat{M}_{x_k, z_j})_{\pi, \epsilon} = \sum_{1 \leq k \leq n} (\widehat{M}_{z_i, x_k} \widehat{M}_{x_k, z_j})_{\epsilon, \epsilon}.$$

Now consider the term

$$\sum_{1 \leq k \leq n} \sum_{\pi \in \Gamma^*} (M_{\epsilon, \pi})_{z_i, x_k} (M_{\pi, \epsilon})_{x_k, z_j},$$

which cannot be simplified because $\sum_{\pi \in \Gamma^*} (M_{\epsilon, \pi} M_{\pi, \epsilon})_{z_i, z_j}$ does no longer hold the information that the path passes only through states x_i , i.e., it contains also the path $(M_{\epsilon, \pi})_{z_i, z_k} (M_{\pi, \epsilon})_{z_k, z_j}$ (for all $1 \leq k \leq n$). In the proofs below, we will specifically need to distinguish paths that pass through states x_i and those that pass through states z_i as in the mixed ω -algebraic system, we also distinguish between variables x_i for finite derivations and variables z_i for infinite derivations.

Lemma 28. Let $M \in (S^{k \times k})^{\Gamma^* \times \Gamma^*}$ be a reset pushdown matrix. Then,

$$\widehat{M}^* = \widehat{M}^*.$$

Proof. For $1 \leq v, v' \leq k$ and for $\pi, \pi' \in \Gamma^*$, we obtain

$$\begin{aligned} ((\widehat{M}^*)_{v, v'})_{\pi, \pi'} &= ((M^*)_{\pi, \pi'})_{v, v'} \\ &= \sum_{n \geq 0} ((M^n)_{\pi, \pi'})_{v, v'} \\ &= \sum_{n \geq 0} ((\widehat{M})_{v, v'})_{\pi, \pi'} \\ &= ((\widehat{M})_{v, v'})_{\pi, \pi'}. \end{aligned}$$

□

Similarly, we need the above result for another operator.

Lemma 29. Let $M \in (S^{k \times k})^{\Gamma^* \times \Gamma^*}$ be a reset pushdown matrix. Then, for $1 \leq l \leq k$,

$$\widehat{M}^{\omega, l} = \widehat{M}^{\omega, l}.$$

Proof. For $1 \leq v \leq k$ and for $\pi \in \Gamma^*$, we obtain

$$\begin{aligned} ((\widehat{M}^{\omega, l})_v)_\pi &= ((M^{\omega, l})_\pi)_v \\ &= \sum_{\pi_1, \pi_2, \dots \in \Gamma^*} \sum_{(v_1, v_2, \dots) \in P_l} (M_{\pi, \pi_1})_{v, v_1} (M_{\pi_1, \pi_2})_{v_1, v_2} (M_{\pi_2, \pi_3})_{v_2, v_3} \cdots \\ &= \sum_{\pi_1, \pi_2, \dots \in \Gamma^*} \sum_{(v_1, v_2, \dots) \in P_l} (\widehat{M}_{v, v_1})_{\pi, \pi_1} (\widehat{M}_{v_1, v_2})_{\pi_1, \pi_2} (\widehat{M}_{v_2, v_3})_{\pi_2, \pi_3} \cdots \\ &= ((\widehat{M})_v)_\pi. \end{aligned}$$

□

Let M be a simple reset pushdown matrix induced by the Greibach normal form (14), (20). We define some blocks of the matrix \widehat{M} to make the following argumentation easier. We take the idea of the above-mentioned isomorphism and divide \widehat{M} like

$$\widehat{M} = \begin{pmatrix} \widehat{M}_{z,z} & \widehat{M}_{z,x} & 0 \\ \widehat{M}_{x,z} & \widehat{M}_{x,x} & \widehat{M}_{x,f} \\ 0 & 0 & 0 \end{pmatrix}, \quad (21)$$

where the respective blocks are defined as

$$\begin{aligned} \widehat{M}_{z,z} &= \begin{pmatrix} \widehat{M}_{z_1,z_1} & \cdots & \widehat{M}_{z_1,z_n} \\ \vdots & \ddots & \vdots \\ \widehat{M}_{z_n,z_1} & \cdots & \widehat{M}_{z_n,z_n} \end{pmatrix}, & \widehat{M}_{z,x} &= \begin{pmatrix} \widehat{M}_{z_1,x_1} & \cdots & \widehat{M}_{z_1,x_n} \\ \vdots & \ddots & \vdots \\ \widehat{M}_{z_n,x_1} & \cdots & \widehat{M}_{z_n,x_n} \end{pmatrix}, \\ \widehat{M}_{x,z} &= \begin{pmatrix} \widehat{M}_{x_1,z_1} & \cdots & \widehat{M}_{x_1,z_n} \\ \vdots & \ddots & \vdots \\ \widehat{M}_{x_n,z_1} & \cdots & \widehat{M}_{x_n,z_n} \end{pmatrix}, & \widehat{M}_{x,x} &= \begin{pmatrix} \widehat{M}_{x_1,x_1} & \cdots & \widehat{M}_{x_1,x_n} \\ \vdots & \ddots & \vdots \\ \widehat{M}_{x_n,x_1} & \cdots & \widehat{M}_{x_n,x_n} \end{pmatrix}, & \widehat{M}_{x,f} &= \begin{pmatrix} \widehat{M}_{x_1,f} \\ \vdots \\ \widehat{M}_{x_n,f} \end{pmatrix}, \end{aligned}$$

and where each $\widehat{M}_{v,v'} \in S^{\Gamma^* \times \Gamma^*}$ for $1 \leq v, v' \leq 2n+1$. For notational convenience, we also set

$$\widehat{M}_{z_i,x} = \begin{pmatrix} \widehat{M}_{z_i,x_1} & \cdots & \widehat{M}_{z_i,x_n} \end{pmatrix}, \quad \widehat{M}_{x,z_i} = \begin{pmatrix} \widehat{M}_{x_1,z_i} \\ \vdots \\ \widehat{M}_{x_n,z_i} \end{pmatrix}.$$

Note that we have not defined the blocks $\widehat{M}_{z,f}$, $\widehat{M}_{f,z}$, $\widehat{M}_{f,x}$ and $\widehat{M}_{f,f}$ as they would all be zero by our construction for simple reset pushdown matrices induced by the Greibach normal form (14), (20).

Analogously, let $M_{z,z}, M_{z,x}, M_{x,z}, M_{x,x}, M_{x,f} \in (S^{(2n+1) \times (2n+1)})^{\Gamma^* \times \Gamma^*}$ be the isomorphic copy of $\widehat{M}_{z,z}, \widehat{M}_{z,x}, \widehat{M}_{x,z}, \widehat{M}_{x,x}, \widehat{M}_{x,f}$, respectively. Then, for $u, v \in \{x, z\}$ and for $\pi, \pi' \in \Gamma^*$, the matrix $(M_{u,v})_{\pi, \pi'}$ is $M_{\pi, \pi'}$ restricted to the variables u_i, v_j (for $1 \leq i, j \leq n$). Similarly, $M_{x,f}$ is M restricted to variables x_i, f (for $1 \leq i \leq n$). For instance, $(\widehat{M}_{x,x})^*$ and equally $(M_{x,x})^*$ consider only paths passing through states x_i and no paths through y_i or f (for $1 \leq i \leq n$). Their only difference is the order of indexes.

The following theorem computes the behavior of induced simple ω -reset pushdown automata.

Theorem 30. *Let M be a simple reset pushdown matrix induced by the Greibach normal form (14), (20). Then, for all $1 \leq i \leq n$ and $0 \leq l \leq n$,*

$$((M^{\omega,l})_\epsilon)_{x_i} = ((M^{\omega,l})_\epsilon)_f = 0,$$

and

$$((M^{\omega,l})_\epsilon)_{z_i} = (((M_{z,z} + M_{z,x}(M^*)_{x,x}M_{x,z})^{\omega,l})_\epsilon)_i.$$

Proof. For the matrix M , we have, by above notation (21),

$$\widehat{M} = \begin{pmatrix} \widehat{M}_{z,z} & \widehat{M}_{z,x} & 0 \\ \widehat{M}_{x,z} & \widehat{M}_{x,x} & \widehat{M}_{x,f} \\ 0 & 0 & 0 \end{pmatrix} = \left(\begin{array}{c|cc} \widehat{M}_{z,z} & \widehat{M}_{z,x} & 0 \\ \hline \widehat{M}_{x,z} & \widehat{M}_{x,x} & \widehat{M}_{x,f} \\ 0 & 0 & 0 \end{array} \right).$$

Thus, by Theorem 2 and Lemma 29, we obtain

$$M^{\omega,l} = \begin{pmatrix} \alpha^{\omega,l} \\ \left(\begin{pmatrix} M_{x,x} & M_{x,f} \\ 0 & 0 \end{pmatrix}^* \begin{pmatrix} M_{x,z} \\ 0 \end{pmatrix} \right) \alpha^{\omega,l} \end{pmatrix},$$

where

$$\begin{aligned}
\alpha &= M_{z,z} + \begin{pmatrix} M_{z,x} & 0 \\ 0 & 0 \end{pmatrix} \begin{pmatrix} M_{x,x} & M_{x,f} \\ 0 & 0 \end{pmatrix}^* \begin{pmatrix} M_{x,z} \\ 0 \end{pmatrix} \\
&= M_{z,z} + \begin{pmatrix} M_{z,x} & 0 \\ 0 & 0 \end{pmatrix} \begin{pmatrix} (M_{x,x})^* & (M_{x,x})^* M_{x,f} \\ 0 & 1 \end{pmatrix} \begin{pmatrix} M_{x,z} \\ 0 \end{pmatrix} \\
&= M_{z,z} + \begin{pmatrix} M_{z,x}(M_{x,x})^* & M_{z,x}(M_{x,x})^* M_{x,f} \\ 0 & 0 \end{pmatrix} \begin{pmatrix} M_{x,z} \\ 0 \end{pmatrix} \\
&= M_{z,z} + M_{z,x}(M_{x,x})^* M_{x,z}.
\end{aligned} \tag{22}$$

Now, we continue with the term from before and get

$$\begin{aligned}
M^{\omega,l} &= \begin{pmatrix} \alpha^{\omega,l} \\ \begin{pmatrix} M_{x,x} & M_{x,f} \\ 0 & 0 \end{pmatrix}^* \begin{pmatrix} M_{x,z} \\ 0 \end{pmatrix} \alpha^{\omega,l} \end{pmatrix} \\
&= \begin{pmatrix} (M_{z,z} + M_{z,x}(M_{x,x})^* M_{x,z})^{\omega,l} \\ \begin{pmatrix} M_{x,x} & M_{x,f} \\ 0 & 0 \end{pmatrix}^* \begin{pmatrix} M_{x,z} \\ 0 \end{pmatrix} (M_{z,z} + M_{z,x}(M_{x,x})^* M_{x,z})^{\omega,l} \end{pmatrix} \\
&= \begin{pmatrix} (M_{z,z} + M_{z,x}(M_{x,x})^* M_{x,z})^{\omega,l} \\ \begin{pmatrix} (M_{x,x})^* & (M_{x,x})^* M_{x,f} \\ 0 & 1 \end{pmatrix} \begin{pmatrix} M_{x,z} \\ 0 \end{pmatrix} (M_{z,z} + M_{z,x}(M_{x,x})^* M_{x,z})^{\omega,l} \end{pmatrix} \\
&= \begin{pmatrix} (M_{z,z} + M_{z,x}(M_{x,x})^* M_{x,z})^{\omega,l} \\ \begin{pmatrix} (M_{x,x})^* M_{x,z} \\ 0 \end{pmatrix} (M_{z,z} + M_{z,x}(M_{x,x})^* M_{x,z})^{\omega,l} \end{pmatrix} \\
&= \begin{pmatrix} (M_{z,z} + M_{z,x}(M_{x,x})^* M_{x,z})^{\omega,l} \\ ((M_{x,x})^* M_{x,z})(M_{z,z} + M_{z,x}(M_{x,x})^* M_{x,z})^{\omega,l} \\ 0 \end{pmatrix}.
\end{aligned}$$

Then, we start the run of the automaton with an empty stack and get

$$\begin{aligned}
(M^{\omega,l})_\epsilon &= \begin{pmatrix} (M_{z,z} + M_{z,x}(M_{x,x})^* M_{x,z})^{\omega,l} \\ ((M_{x,x})^* M_{x,z})(M_{z,z} + M_{z,x}(M_{x,x})^* M_{x,z})^{\omega,l} \\ 0 \end{pmatrix}_\epsilon \\
&= \begin{pmatrix} ((M_{z,z} + M_{z,x}(M_{x,x})^* M_{x,z})^{\omega,l})_\epsilon \\ (((M_{x,x})^* M_{x,z})(M_{z,z} + M_{z,x}(M_{x,x})^* M_{x,z})^{\omega,l})_\epsilon \\ 0 \end{pmatrix}_\epsilon \\
&= \begin{pmatrix} ((M_{z,z} + M_{z,x}(M_{x,x})^* M_{x,z})^{\omega,l})_\epsilon \\ \sum_{\pi \in \Gamma^*} ((M_{x,x})^* M_{x,z})_{\epsilon,\pi} ((M_{z,z} + M_{z,x}(M_{x,x})^* M_{x,z})^{\omega,l})_\pi \\ 0 \end{pmatrix} \\
&\stackrel{4}{=} \begin{pmatrix} ((M_{z,z} + M_{z,x}(M_{x,x})^* M_{x,z})^{\omega,l})_\epsilon \\ \sum_{\pi \in \Gamma^*} 0 ((M_{z,z} + M_{z,x}(M_{x,x})^* M_{x,z})^{\omega,l})_\pi \\ 0 \end{pmatrix} \\
&= \begin{pmatrix} ((M_{z,z} + M_{z,x}(M_{x,x})^* M_{x,z})^{\omega,l})_\epsilon \\ 0 \\ 0 \end{pmatrix}
\end{aligned}$$

where the forth equality uses the fact that $((M_{x,x})^* M_{x,z})_{\epsilon,\pi} = 0$ which is because $(M_{x,z})_{\pi,\pi'} = 0$ for all $\pi \neq Z_k \pi''$ ($1 \leq k \leq n$ and $\pi'' \in \Gamma^*$) and at the same time, $((M_{x,x})^*)_{\epsilon, Z_k \pi''} = 0$ because only $(M_{z,x})_{\epsilon, Z_k} \neq 0$ by construction.

The vector $(M^{\omega,l})_\epsilon$ is indexed by $z_1, \dots, z_n, x_1, \dots, x_n, f$, thus completing the proof. \square

We want to apply the results from Section 6. The following three lemmas investigate the star operation applied to simple reset pushdown matrices M induced by the Greibach normal form (14), (20). The lemmas state that in a computation $(M^*)_{\epsilon, \epsilon}$, the new states y_k are never reached when starting in a state x_i and therefore, these computations are equivalent to the computations $(M'^*)_{\epsilon, \epsilon}$ for M' being induced by the Greibach normal form (14), i.e., for M' built by the old construction.

Lemma 31. *Let M be a simple reset pushdown matrix induced by the Greibach normal form (14), (20). Then, for all $1 \leq i, k \leq n$,*

$$((M^*)_{\epsilon, \epsilon})_{x_k, x_i} = (((M_{x, x})^*)_{\epsilon, \epsilon})_{x_k, x_i}.$$

Proof. Let $\Delta = \{X_1, \dots, X_n\}$. We have

$$\begin{aligned} ((M^*)_{\epsilon, \epsilon})_{x_k, x_i} &= \sum_{t \geq 0} ((M^t)_{\epsilon, \epsilon})_{x_k, x_i} \\ &= \sum_{t \geq 0} \sum_{\pi_1, \dots, \pi_{t-1} \in \Gamma^*} (M_{\epsilon, \pi_1} M_{\pi_1, \pi_2} \cdots M_{\pi_{t-1}, \epsilon})_{x_k, x_i} \\ &= \sum_{t \geq 0} \sum_{\substack{\pi_1 \in \Delta^* \\ \pi_2, \dots, \pi_{t-1} \in \Gamma^*}} \sum_{1 \leq j_1 \leq n} (M_{\epsilon, \pi_1})_{x_k, x_{j_1}} (M_{\pi_1, \pi_2} \cdots M_{\pi_{t-1}, \epsilon})_{x_{j_1}, x_i} \\ &= \sum_{t \geq 0} \sum_{\pi_1, \dots, \pi_{t-1} \in \Delta^*} \sum_{1 \leq j_1, \dots, j_{t-1} \leq n} (M_{\epsilon, \pi_1})_{x_k, x_{j_1}} (M_{\pi_1, \pi_2})_{x_{j_1}, x_{j_2}} \cdots (M_{\pi_{t-1}, \epsilon})_{x_{j_{t-1}}, x_i} \\ &= ((\sum_{t \geq 0} (M_{x, x})^t)_{\epsilon, \epsilon})_{x_k, x_i} = (((M_{x, x})^*)_{\epsilon, \epsilon})_{x_k, x_i}, \end{aligned}$$

where the third equality (and similarly the forth equality) is by definition of induced pushdown matrices; the blocks $(M_{\epsilon, X_k})_{x_i, x_j}$, $(M_{X_k, X_k})_{x_i, x_j}$ and $(M_{\epsilon, \epsilon})_{x_i, x_j}$ are the only non-null blocks that describe a step in the matrix starting from a state x_i and having ϵ or X_k as the topmost stack symbol. \square

Lemma 32. *Let M be a simple reset pushdown matrix induced by the Greibach normal form (14), (20). Then, we have*

$$((M_{x, x} + M_{x, z}(M_{z, z})^* M_{z, x})_{\epsilon, \epsilon})_{\epsilon, \epsilon} = ((M_{x, x})^*)_{\epsilon, \epsilon}.$$

Proof. Let $\Delta = \{X_1, \dots, X_n\}$. In some sense similar to the proof of Lemma 31, we have

$$\begin{aligned} &((M_{x, x} + M_{x, z}(M_{z, z})^* M_{z, x})_{\epsilon, \epsilon})_{\epsilon, \epsilon} \\ &= \left(\sum_{t \geq 0} (M_{x, x} + M_{x, z}(M_{z, z})^* M_{z, x})^t \right)_{\epsilon, \epsilon} \\ &= \sum_{t \geq 0} \sum_{\pi_1, \dots, \pi_{t-1} \in \Gamma^*} (M_{x, x} + M_{x, z}(M_{z, z})^* M_{z, x})_{\epsilon, \pi_1} \cdots (M_{x, x} + M_{x, z}(M_{z, z})^* M_{z, x})_{\pi_{t-1}, \epsilon} \\ &= \sum_{t \geq 0} \sum_{\pi_1, \dots, \pi_{t-1} \in \Gamma^*} \left((M_{x, x})_{\epsilon, \pi_1} + \left(\sum_{\pi, \pi' \in \Gamma^*} (M_{x, z})_{\epsilon, \pi} ((M_{z, z})^*)_{\pi, \pi'} (M_{z, x})_{\pi, \pi_1} \right) \right) \cdots (M_{x, x} + M_{x, z}(M_{z, z})^* M_{z, x})_{\pi_{t-1}, \epsilon} \\ &= \sum_{t \geq 0} \sum_{\substack{\pi_1 \in \Delta^* \\ \pi_2, \dots, \pi_{t-1} \in \Gamma^*}} (M_{x, x})_{\epsilon, \pi_1} (M_{x, x} + M_{x, z}(M_{z, z})^* M_{z, x})_{\pi_1, \pi_2} \cdots (M_{x, x} + M_{x, z}(M_{z, z})^* M_{z, x})_{\pi_{t-1}, \epsilon} \\ &= \sum_{t \geq 0} \sum_{\substack{\pi_1 \in \Delta^* \\ \pi_2, \dots, \pi_{t-1} \in \Gamma^*}} (M_{x, x})_{\epsilon, \pi_1} \left((M_{x, x})_{\pi_1, \pi_2} + \left(\sum_{\pi, \pi' \in \Gamma^*} (M_{x, z})_{\pi_1, \pi} ((M_{z, z})^*)_{\pi, \pi'} (M_{z, x})_{\pi, \pi_2} \right) \right) \cdots (M_{x, x} + M_{x, z}(M_{z, z})^* M_{z, x})_{\pi_{t-1}, \epsilon} \\ &= \sum_{t \geq 0} \sum_{\pi_1, \dots, \pi_{t-1} \in \Delta^*} (M_{x, x})_{\epsilon, \pi_1} (M_{x, x})_{\pi_1, \pi_2} \cdots (M_{x, x})_{\pi_{t-1}, \epsilon} = ((M_{x, x})^*)_{\epsilon, \epsilon}, \end{aligned}$$

where the forth equality is because $(M_{x, z})_{\epsilon, \pi} = 0$ for all $\pi \in \Gamma^*$. Similarly, for the sixth equality, we use the fact that $(M_{x, z})_{\pi_i, \pi} = 0$ for all $\pi_i \in \Delta^*$ (and $\pi \in \Gamma^*$). \square

Lemma 33. Let M be a simple reset pushdown matrix induced by the Greibach normal form (14), (20) and M' be induced by the Greibach normal form (14). Then, for all $1 \leq i \leq n$,

$$((M^*)_{\epsilon, \epsilon})_{x_i, f} = ((M'^*)_{\epsilon, \epsilon})_{i, f}.$$

Proof. Note that by construction, we have

$$\widehat{M}' = \begin{pmatrix} \widehat{M}_{x,x} & \widehat{M}_{x,f} \\ 0 & 0 \end{pmatrix}.$$

By applying Lemma 28, we infer

$$M'^* = \begin{pmatrix} (M_{x,x})^* & (M_{x,x})^* M_{x,f} \\ 0 & 1 \end{pmatrix},$$

and we get

$$((M'^*)_{\epsilon, \epsilon})_{i, f} = (((M_{x,x})^* M_{x,f})_{\epsilon, \epsilon})_i. \quad (23)$$

At the same time, we have

$$\begin{aligned} \widehat{M} &= \begin{pmatrix} \widehat{M}_{z,z} & \widehat{M}_{z,x} & 0 \\ \widehat{M}_{x,z} & \widehat{M}_{x,x} & \widehat{M}_{x,f} \\ 0 & 0 & 0 \end{pmatrix} \\ &= \left(\begin{array}{c|cc} \widehat{M}_{z,z} & \widehat{M}_{z,x} & 0 \\ \hline \widehat{M}_{x,z} & \widehat{M}_{x,x} & \widehat{M}_{x,f} \\ 0 & 0 & 0 \end{array} \right). \end{aligned}$$

By Lemma 28, we obtain

$$M^* = \begin{pmatrix} \alpha^* & \alpha^* (M_{z,x} \ 0) \\ \beta^* \begin{pmatrix} M_{x,z} \\ 0 \end{pmatrix} & \beta^* \end{pmatrix},$$

with

$$\begin{aligned} \alpha &= M_{z,z} + \begin{pmatrix} M_{z,x} & 0 \end{pmatrix} \begin{pmatrix} M_{x,x} & M_{x,f} \\ 0 & 0 \end{pmatrix}^* \begin{pmatrix} M_{x,z} \\ 0 \end{pmatrix} \\ &= M_{z,z} + M_{z,x} (M_{x,x})^* M_{x,z}, \end{aligned}$$

by (22) in the proof of Theorem 30 and

$$\begin{aligned} \beta^* &= \left(\begin{pmatrix} \widehat{M}_{x,x} & \widehat{M}_{x,f} \\ 0 & 0 \end{pmatrix} + \begin{pmatrix} M_{x,z} \\ 0 \end{pmatrix} (M_{z,z})^* (M_{z,x} \ 0) \right)^* \\ &= \left(\begin{pmatrix} M_{x,x} & M_{x,f} \\ 0 & 0 \end{pmatrix} + \begin{pmatrix} M_{x,z} (M_{z,z})^* M_{z,x} & 0 \\ 0 & 0 \end{pmatrix} \right)^* \\ &= \begin{pmatrix} M_{x,x} + M_{x,z} (M_{z,z})^* M_{z,x} & M_{x,f} \\ 0 & 0 \end{pmatrix}^* \\ &= \begin{pmatrix} (M_{x,x} + M_{x,z} (M_{z,z})^* M_{z,x})^* & (M_{x,x} + M_{x,z} (M_{z,z})^* M_{z,x})^* M_{x,f} \\ 0 & 1 \end{pmatrix}. \end{aligned} \quad (24)$$

We deduce that

$$\begin{aligned}
((M^*)_{\epsilon, \epsilon})_{x_i, f} &= (((M_{x, x} + M_{x, z}(M_{z, z})^* M_{z, x})^* M_{x, f})_{\epsilon, \epsilon})_i \\
&= (((M_{x, x} + M_{x, z}(M_{z, z})^* M_{z, x})^*)_{\epsilon, \epsilon} (M_{x, f})_{\epsilon, \epsilon})_i \\
&= (((M_{x, x})^*)_{\epsilon, \epsilon} (M_{x, f})_{\epsilon, \epsilon})_i \\
&= (((M_{x, x})^* M_{x, f})_{\epsilon, \epsilon})_i \\
&= ((M')_{\epsilon, \epsilon})_{i, f},
\end{aligned}$$

where the third equality is by Lemma 32 and the last equality is by (23). This concludes the proof. \square

The following lemma investigates the final state f in infinite paths. It states that a finite run of induced simple ω -reset pushdown automata is equivalent to another path only through states x and with symbol Z_j initially on the pushdown tape and ending in state z_j with an empty pushdown tape.

Lemma 34. *Let M be a simple reset pushdown matrix induced by the Greibach normal form (14), (20). Then, for all $1 \leq j, k \leq n$,*

$$((M^*)_{\epsilon, \epsilon})_{x_k, f} = (((M_{x, x})^*)_{Z_j, Z_j} M_{Z_j, \epsilon})_{x_k, z_j}.$$

Proof. The beginning of the proof is similar to the proof of Lemma 10 of [6]. We obtain

$$\begin{aligned}
((M^*)_{\epsilon, \epsilon})_{x_k, f} &= ((M^+)_{\epsilon, \epsilon})_{x_k, f} = ((M^* M)_{\epsilon, \epsilon})_{x_k, f} \\
&= \sum_{1 \leq v_1 \leq 2n+1} ((M^*)_{\epsilon, \epsilon})_{x_k, v_1} (M_{\epsilon, \epsilon})_{v_1, f} + \sum_{1 \leq v_1 \leq 2n+1} \sum_{P \in \Gamma} ((M^*)_{\epsilon, P})_{x_k, v_1} (M_{P, \epsilon})_{v_1, f} \\
&\stackrel{4}{=} \sum_{1 \leq v_1 \leq 2n+1} ((M^*)_{\epsilon, \epsilon})_{x_k, v_1} (M_{\epsilon, \epsilon})_{v_1, f} \\
&\stackrel{5}{=} \sum_{1 \leq i \leq n} ((M^*)_{\epsilon, \epsilon})_{x_k, x_i} (M_{\epsilon, \epsilon})_{x_i, f} \\
&\stackrel{6}{=} \sum_{1 \leq i \leq n} (((M_{x, x})^*)_{\epsilon, \epsilon})_{x_k, x_i} (M_{\epsilon, \epsilon})_{x_i, f} \\
&\stackrel{7}{=} \sum_{1 \leq i \leq n} (((M_{x, x})^*)_{Z_j, Z_j})_{x_k, x_i} (M_{Z_j, \epsilon})_{x_i, z_j} = (((M_{x, x})^*)_{Z_j, Z_j} M_{Z_j, \epsilon})_{x_k, z_j},
\end{aligned}$$

where the forth equality is since $(M_{P, \epsilon})_{v_1, f} = 0$ for all $1 \leq v_1 \leq 2n+1$ and $P \in \Gamma$ by our construction. In the fifth equality, we use the fact that $(M_{\epsilon, \epsilon})_{v_1, f} = 0$ for $v_1 \neq x_i$ ($1 \leq i \leq n$). The sixth equality is by Lemma 31. The seventh equality is also by construction and by the definition of pushdown matrices. \square

The following lemma treats a case similar to the previous lemma. It states that an infinite path starting with symbol Z_k on the pushdown tape is equivalent to a finite run starting with an empty pushdown tape and ending in state f followed by an infinite run that starts in state z_k with an empty pushdown tape.

Lemma 35. *Let M be a simple reset pushdown matrix induced by the Greibach normal form (14), (20). Then, for all $1 \leq j, k \leq n$ and $0 \leq l \leq n$,*

$$((M^{\omega, l})_{Z_k})_{x_j} = ((M^*)_{\epsilon, \epsilon})_{x_j, f} ((M^{\omega, l})_{\epsilon})_{z_k}.$$

Proof. By Lemma 23, we have

$$\begin{aligned}
((M^{\omega, l})_{Z_k})_{x_j} &= [(M^{\omega, l})_{\epsilon} + (M^*)_{\epsilon, \epsilon} M_{Z_k, \epsilon} (M^{\omega, l})_{\epsilon}]_{x_j} \\
&= ((M^{\omega, l})_{\epsilon})_{x_j} + ((M^*)_{\epsilon, \epsilon} M_{Z_k, \epsilon} (M^{\omega, l})_{\epsilon})_{x_j}.
\end{aligned}$$

Consider the first summand. By Theorem 30, we know that

$$((M^{\omega, l})_{\epsilon})_{x_j} = 0.$$

Now, we consider the second summand. For $1 \leq j, k \leq n$, we have

$$\begin{aligned}
((M^*)_{\epsilon, \epsilon} M_{Z_k, \epsilon} (M^{\omega, l})_{\epsilon})_{x_j} &= \sum_{1 \leq v_1, v_2 \leq 2n+1} ((M^*)_{\epsilon, \epsilon})_{x_j, v_1} (M_{Z_k, \epsilon})_{v_1, v_2} ((M^{\omega, l})_{\epsilon})_{v_2} \\
&= \sum_{1 \leq v_1 \leq 2n+1} ((M^*)_{\epsilon, \epsilon})_{x_j, v_1} (M_{Z_k, \epsilon})_{v_1, z_k} ((M^{\omega, l})_{\epsilon})_{z_k} \\
&= \sum_{1 \leq v_1 \leq 2n+1} ((M^*)_{\epsilon, \epsilon})_{x_j, v_1} (M_{\epsilon, \epsilon})_{v_1, f} ((M^{\omega, l})_{\epsilon})_{z_k} \\
&= ((M^*)_{\epsilon, \epsilon} M_{\epsilon, \epsilon})_{x_j, f} ((M^{\omega, l})_{\epsilon})_{z_k} \\
&= ((M^*)_{\epsilon, \epsilon})_{x_j, f} ((M^{\omega, l})_{\epsilon})_{z_k},
\end{aligned}$$

where the second equality holds because we defined $(M_{Z_k, \epsilon})_{v_1, v_2} = 0$ for $v_2 \neq z_k$ and the third equality is because we have $(M_{Z_k, \epsilon})_{v_1, z_k} = (M_{\epsilon, \epsilon})_{v_1, f}$ for induced simple pushdown matrices. The result follows. \square

We now discuss the behaviors of our constructed simple ω -reset pushdown automata.

Lemma 36. *Let the simple ω -reset pushdown automata $\mathfrak{A}_m^l = (2n+1, \Gamma, I_m, M, P, l)$, for $1 \leq m \leq n$ and $0 \leq l \leq n$, be induced by the Greibach normal form (14), (20). We then have*

$$\|\mathfrak{A}_m^l\| = ((M^*)_{\epsilon, \epsilon})_{x_m, f} + ((M^{\omega, l})_{\epsilon})_{z_m}.$$

Proof. Let $1 \leq m \leq n$ and $0 \leq l \leq n$. We obtain

$$\begin{aligned}
\|\mathfrak{A}_m^l\| &= I(M^*)_{\epsilon, \epsilon} P + I(M^{\omega, l})_{\epsilon} \\
&= ((M^*)_{\epsilon, \epsilon})_{x_m, f} + ((M^*)_{\epsilon, \epsilon})_{z_m, f} + ((M^{\omega, l})_{\epsilon})_{x_m} + ((M^{\omega, l})_{\epsilon})_{z_m}, \\
&= ((M^*)_{\epsilon, \epsilon})_{x_m, f} + ((M^*)_{\epsilon, \epsilon})_{z_m, f} + ((M^{\omega, l})_{\epsilon})_{z_m}.
\end{aligned}$$

where the last equality is by Theorem 30.

It remains to show that $((M^*)_{\epsilon, \epsilon})_{z_m, f} = 0$. We have

$$\widehat{M} = \left(\begin{array}{c|cc} \widehat{M}_{z, z} & \widehat{M}_{z, x} & 0 \\ \hline \widehat{M}_{x, z} & \widehat{M}_{x, x} & \widehat{M}_{x, f} \\ 0 & 0 & 0 \end{array} \right).$$

Now let

$$M^* = \begin{pmatrix} \alpha & \beta \\ \gamma & \delta \end{pmatrix},$$

where we are only interested in the second component of β . By lemma 28 and by (24) in the proof of Lemma 33, we have

$$\begin{aligned}
\beta &= (M_{z, z})^* \begin{pmatrix} M_{z, x} & 0 \end{pmatrix} \left[\begin{pmatrix} M_{x, x} & M_{x, f} \\ 0 & 0 \end{pmatrix} + \begin{pmatrix} M_{x, z} \\ 0 \end{pmatrix} (M_{z, z})^* \begin{pmatrix} M_{z, x} & 0 \end{pmatrix} \right]^* \\
&= \begin{pmatrix} (M_{z, z})^* M_{z, x} & 0 \end{pmatrix} \begin{pmatrix} (M_{x, x} + M_{x, z} (M_{z, z})^* M_{z, x})^* & (M_{x, x} + M_{x, z} (M_{z, z})^* M_{z, x})^* M_{x, f} \\ 0 & 1 \end{pmatrix} \\
&= \begin{pmatrix} (M_{z, z})^* M_{z, x} (M_{x, x} + M_{x, z} (M_{z, z})^* M_{z, x})^*, & (M_{z, z})^* M_{z, x} (M_{x, x} + M_{x, z} (M_{z, z})^* M_{z, x})^* M_{x, f} \end{pmatrix}.
\end{aligned}$$

Now, we obtain

$$\begin{aligned}
((M^*)_{\epsilon, \epsilon})_{z_m, f} &= \left(((M_{z, z})^* M_{z, x} (M_{x, x} + M_{x, z} (M_{z, z})^* M_{z, x})^* M_{x, f})_{\epsilon, \epsilon} \right)_m \\
&= \left(((M_{z, z})^* M_{z, x} (M_{x, x} + M_{x, z} (M_{z, z})^* M_{z, x})^*)_{\epsilon, \epsilon} (M_{x, f})_{\epsilon, \epsilon} \right)_m \\
&= \left(((M_{z, z})^*)_{\epsilon, \epsilon} (M_{z, x} (M_{x, x} + M_{x, z} (M_{z, z})^* M_{z, x})^*)_{\epsilon, \epsilon} (M_{x, f})_{\epsilon, \epsilon} \right)_m \\
&= \sum_{1 \leq i \leq n} \left(((M_{z, z})^*)_{\epsilon, \epsilon} (M_{z, x})_{\epsilon, Z_i} ((M_{x, x} + M_{x, z} (M_{z, z})^* M_{z, x})^*)_{Z_i, \epsilon} (M_{x, f})_{\epsilon, \epsilon} \right)_m, \tag{25}
\end{aligned}$$

where in the second equality, we have $(M_{x,f})_{\pi,\epsilon} = 0$ for $\pi \neq \epsilon$. The third equality uses that $(M_{z,z})_{\epsilon,\pi}^* = 0$ for $\pi \neq \epsilon$. In the forth equality, we have $(M_{z,x})_{\epsilon,\pi} = 0$ for $\pi \notin \{Z_i \mid 1 \leq i \leq n\}$.

We concentrate on the factor in the center, where we have

$$\begin{aligned} & ((M_{x,x} + M_{x,z}(M_{z,z})^* M_{z,x})^*)_{Z_i,\epsilon} \\ &= \sum_{t \geq 0} \sum_{\pi_1, \dots, \pi_{t-1} \in \Gamma^*} (M_{x,x} + M_{x,z}(M_{z,z})^* M_{z,x})_{Z_i,\pi_1} \cdots (M_{x,x} + M_{x,z}(M_{z,z})^* M_{z,x})_{\pi_{t-1},\epsilon} \\ &= \sum_{t \geq 0} \sum_{\pi_1, \dots, \pi_{t-2} \in \Gamma^*} (M_{x,x} + M_{x,z}(M_{z,z})^* M_{z,x})_{Z_i,\pi_1} \cdots (M_{x,x} + M_{x,z}(M_{z,z})^* M_{z,x})_{\pi_{t-2},\epsilon} (M_{x,x})_{\epsilon,\epsilon} \\ &= \sum_{t \geq 0} (M_{x,x})_{Z_i,\epsilon} \cdots (M_{x,x})_{\epsilon,\epsilon} (M_{x,x})_{\epsilon,\epsilon} = 0, \end{aligned}$$

where in the second (and similarly in the third) equality we have $(M_{x,x} + M_{x,z}(M_{z,z})^* M_{z,x})_{\pi_{t-1},\epsilon} = (M_{x,x})_{\epsilon,\epsilon}$ because $(M_{x,x})_{\pi_{t-1},\epsilon} = 0$ for $\pi_{t-1} \neq \epsilon$ and because

$$(M_{x,z}(M_{z,z})^* M_{z,x})_{\pi_{t-1},\epsilon} = \sum_{\pi, \pi' \in \Gamma^*} (M_{x,z})_{\pi_{t-1},\pi} ((M_{z,z})^*)_{\pi,\pi'} (M_{z,x})_{\pi',\epsilon} = 0$$

as $(M_{z,x})_{\pi',\epsilon} = 0$ for all π' . In the last equality, $(M_{x,x})_{Z_i,\epsilon} = 0$.

We now plug this into (25) and obtain

$$\begin{aligned} ((M^*)_{\epsilon,\epsilon})_{z_m,f} &= \sum_{1 \leq i \leq n} \left(((M_{z,z})^*)_{\epsilon,\epsilon} (M_{z,x})_{\epsilon,Z_i} ((M_{x,x} + M_{x,z}(M_{z,z})^* M_{z,x})^*)_{Z_i,\epsilon} (M_{x,f})_{\epsilon,\epsilon} \right)_m \\ &= \sum_{1 \leq i \leq n} \left(((M_{z,z})^*)_{\epsilon,\epsilon} (M_{z,x})_{\epsilon,Z_i} 0 (M_{x,f})_{\epsilon,\epsilon} \right)_m = 0. \end{aligned}$$

This completes the proof. \square

The following theorem compares the behavior of induced simple ω -reset pushdown automata with the solutions of system (18).

Theorem 37. *Let (S, V) be a complete semiring-semimodule pair. Let the simple ω -reset pushdown automata \mathfrak{A}_m^l , for $1 \leq m \leq n$ and $0 \leq l \leq n$, be induced by the Greibach normal form (14), (20).*

Then, for $0 \leq l \leq n$,

$$(\|\mathfrak{A}_1^l\|, \dots, \|\mathfrak{A}_n^l\|) = (((M^*)_{\epsilon,\epsilon})_{x_1,f}, ((M^{\omega,l})_{\epsilon})_{z_1}, \dots, ((M^*)_{\epsilon,\epsilon})_{x_n,f}, ((M^{\omega,l})_{\epsilon})_{z_n})$$

is a solution of (18).

Proof. We show that

$$(((M^*)_{\epsilon,\epsilon})_{x_1,f}, \dots, ((M^*)_{\epsilon,\epsilon})_{x_n,f}) \quad \text{and} \quad (((M^{\omega,l})_{\epsilon})_{z_1}, \dots, ((M^{\omega,l})_{\epsilon})_{z_n})$$

are solutions of the mixed ω -algebraic system (14), (20).

Let now M' be induced by the Greibach normal form (14). By Theorem 16, $((M')_{\epsilon,\epsilon})_{1,f}, \dots, ((M')_{\epsilon,\epsilon})_{n,f}$ is a solution of (14). By Lemma 33, we deduce that $((M^*)_{\epsilon,\epsilon})_{x_1,f}, \dots, ((M^*)_{\epsilon,\epsilon})_{x_n,f}$ is also a solution of (14).

We now show that $((M^{\omega,l})_{\epsilon})_{z_1}, \dots, ((M^{\omega,l})_{\epsilon})_{z_n}$ is a solution of (20) and substitute it into the right sides of (20):

$$\begin{aligned} & \sum_{1 \leq j,k \leq n} \sum_{a \in \Sigma} (p_i, a x_j z_k) a \sigma_j \omega_k + \sum_{1 \leq j \leq n} \sum_{a \in \Sigma} (p_i, a z_j) a \omega_j \\ &= \sum_{1 \leq j,k \leq n} (M_{\epsilon,Z_k})_{z_i,x_j} ((M^*)_{\epsilon,\epsilon})_{x_j,f} ((M^{\omega,l})_{\epsilon})_{z_k} + \sum_{1 \leq j \leq n} (M_{\epsilon,\epsilon})_{z_i,z_j} ((M^{\omega,l})_{\epsilon})_{z_j} \\ &= \sum_{1 \leq j,k \leq n} (M_{\epsilon,Z_k})_{z_i,x_j} ((M^{\omega,l})_{\epsilon})_{z_k} + \sum_{1 \leq j \leq n} (M_{\epsilon,\epsilon})_{z_i,z_j} ((M^{\omega,l})_{\epsilon})_{z_j} \\ &= \sum_{1 \leq k \leq n} (M_{\epsilon,Z_k} (M^{\omega,l})_{Z_k})_{z_i} + (M_{\epsilon,\epsilon} (M^{\omega,l})_{\epsilon})_{z_i} \\ &= (M M^{\omega,l})_{\epsilon})_{z_i} = ((M^{\omega,l})_{\epsilon})_{z_i}, \quad \text{for each } 1 \leq i \leq n, \end{aligned}$$

where the second equality is by Lemma 35, the last equality is by Theorem 19. \square

The theorem above, Theorem 37, is not sufficient for our main result. The following theorem extends the previous theorem by stating that $(\|\mathfrak{A}_1^l\|, \dots, \|\mathfrak{A}_n^l\|)$ is a canonical solution of (18).

Theorem 38. *Let (S, V) be a complete semiring-semimodule pair. Let the simple ω -reset pushdown automata \mathfrak{A}_m^l , for $1 \leq m \leq n$ and $0 \leq l \leq n$, be induced by the Greibach normal form (14), (20).*

Then, for $0 \leq l \leq n$,

$$(\|\mathfrak{A}_1^l\|, \dots, \|\mathfrak{A}_n^l\|) = (((M^*)_{\epsilon, \epsilon})_{x_1, f} + ((M^{\omega, l})_{\epsilon})_{z_1}, \dots, ((M^*)_{\epsilon, \epsilon})_{x_n, f} + ((M^{\omega, l})_{\epsilon})_{z_n})$$

is the l^{th} canonical solution of (18).

Proof. We show that

$$(((M^*)_{\epsilon, \epsilon})_{x_1, f}, \dots, ((M^*)_{\epsilon, \epsilon})_{x_n, f}) \quad \text{and} \quad (((M^{\omega, l})_{\epsilon})_{z_1}, \dots, ((M^{\omega, l})_{\epsilon})_{z_n})$$

is the l^{th} canonical solution of the mixed ω -algebraic system (14), (20).

Let M' be induced by the Greibach normal form (14). Then, by Theorem 16, $((M')_{\epsilon, \epsilon})_{1, f}, \dots, ((M')_{\epsilon, \epsilon})_{n, f}$ is the unique (and therefore least) solution of (14). By Lemma 33, we can conclude that $\sigma = (((M^*)_{\epsilon, \epsilon})_{x_1, f}, \dots, ((M^*)_{\epsilon, \epsilon})_{x_n, f})$ is also the least solution of (14).

Fix l with $1 \leq l \leq n$ for the remainder of the proof. It remains to show that for the system (20), written as $z = \varrho(x)z$, we have

$$\varrho(\sigma)^{\omega, l} = (((M^{\omega, l})_{\epsilon})_{z_1}, \dots, ((M^{\omega, l})_{\epsilon})_{z_n})$$

We start with the right side of equation (20). We have, for $1 \leq i \leq n$,

$$\begin{aligned} \varrho(\sigma)_i z &= \sum_{1 \leq j, k \leq n} \sum_{a \in \Sigma} (p_i, ax_j z_k) a \sigma_j z_k + \sum_{1 \leq j \leq n} \sum_{a \in \Sigma} (p_i, az_j) a z_j \\ &= \sum_{1 \leq j, k \leq n} \sum_{a \in \Sigma} (p_i, ax_k z_j) a \sigma_k z_j + \sum_{1 \leq j \leq n} \sum_{a \in \Sigma} (p_i, az_j) a z_j \\ &= \sum_{1 \leq j \leq n} \left(\sum_{1 \leq k \leq n} \sum_{a \in \Sigma} (p_i, ax_k z_j) a \sigma_k + \sum_{a \in \Sigma} (p_i, az_j) a \right) z_j \\ &= \sum_{1 \leq j \leq n} \left(\sum_{1 \leq k \leq n} (M_{\epsilon, Z_j})_{z_i, x_k} ((M^*)_{\epsilon, \epsilon})_{x_k, f} + (M_{\epsilon, \epsilon})_{z_i, z_j} \right) z_j \\ &\stackrel{5}{=} \sum_{1 \leq j \leq n} \left(\sum_{1 \leq k \leq n} (M_{\epsilon, Z_j})_{z_i, x_k} (((M_{x, x})^*)_{Z_j, Z_j} M_{Z_j, \epsilon})_{x_k, z_j} + (M_{\epsilon, \epsilon})_{z_i, z_j} \right) z_j \\ &= \sum_{1 \leq j \leq n} \left(\sum_{1 \leq k, k' \leq n} (M_{\epsilon, Z_j})_{z_i, x_k} (((M_{x, x})^*)_{Z_j, Z_j})_{x_k, x_{k'}} (M_{Z_j, \epsilon})_{x_{k'}, z_j} + (M_{\epsilon, \epsilon})_{z_i, z_j} \right) z_j \\ &= \sum_{1 \leq j \leq n} \left(\sum_{1 \leq k, k' \leq n} (\widehat{M}_{z_i, x_k})_{\epsilon, Z_j} (((\widehat{M}_{x, x})^*)_{x_k, x_{k'}})_{Z_j, Z_j} (\widehat{M}_{x_{k'}, z_j})_{Z_j, \epsilon} + (\widehat{M}_{z_i, z_j})_{\epsilon, \epsilon} \right) z_j \\ &\stackrel{8}{=} \sum_{1 \leq j \leq n} \left(\sum_{1 \leq k, k' \leq n} \sum_{P \in \Gamma} (\widehat{M}_{z_i, x_k})_{\epsilon, P} (((\widehat{M}_{x, x})^*)_{x_k, x_{k'}})_{P, P} (\widehat{M}_{x_{k'}, z_j})_{P, \epsilon} + (\widehat{M}_{z_i, z_j})_{\epsilon, \epsilon} \right) z_j \\ &= \sum_{1 \leq j \leq n} \left(\sum_{1 \leq k, k' \leq n} (\widehat{M}_{z_i, x_k} ((\widehat{M}_{x, x})^*)_{x_k, x_{k'}} \widehat{M}_{x_{k'}, z_j})_{\epsilon, \epsilon} + (\widehat{M}_{z_i, z_j})_{\epsilon, \epsilon} \right) z_j \\ &= \sum_{1 \leq j \leq n} \left(\widehat{M}_{z_i, x} (\widehat{M}_{x, x})^* \widehat{M}_{x, z_j} + \widehat{M}_{z_i, z_j} \right)_{\epsilon, \epsilon} z_j, \end{aligned}$$

where the fifth equality is by Lemma 34. The eighth equality is because for $P \neq Z_j$, we have $(\widehat{M}_{x_{k'}, z_j})_{P, \epsilon} = 0$.

Now for ϱ of the system $z = \varrho(x)z$, we obtain

$$\begin{aligned}\varrho(\sigma) &= \begin{pmatrix} (\widehat{M}_{z_1,x}(\widehat{M}_{x,x})^* \widehat{M}_{x,z_1} + \widehat{M}_{z_1,z_1})_{\epsilon,\epsilon} & \cdots & (\widehat{M}_{z_1,x}(\widehat{M}_{x,x})^* \widehat{M}_{x,z_n} + \widehat{M}_{z_1,z_n})_{\epsilon,\epsilon} \\ \vdots & \ddots & \vdots \\ (\widehat{M}_{z_n,x}(\widehat{M}_{x,x})^* \widehat{M}_{x,z_1} + \widehat{M}_{z_n,z_1})_{\epsilon,\epsilon} & \cdots & (\widehat{M}_{z_n,x}(\widehat{M}_{x,x})^* \widehat{M}_{x,z_n} + \widehat{M}_{z_n,z_n})_{\epsilon,\epsilon} \end{pmatrix} \\ &= (M_{z,x}(M_{x,x})^* M_{x,z} + M_{z,z})_{\epsilon,\epsilon}.\end{aligned}$$

Then, we have

$$\begin{aligned}(\varrho(\sigma)^{\omega,l})_j &= (((M_{z,x}(M_{x,x})^* M_{x,z} + M_{z,z})_{\epsilon,\epsilon})^{\omega,l})_j \\ &= \sum_{(j_1, j_2, \dots) \in P_l} (\widehat{M}_{z_j,x}(\widehat{M}_{x,x})^* \widehat{M}_{x,z_{j_1}} + \widehat{M}_{z_j,z_{j_1}})_{\epsilon,\epsilon} (\widehat{M}_{z_{j_1},x}(\widehat{M}_{x,x})^* \widehat{M}_{x,z_{j_2}} + \widehat{M}_{z_{j_1},z_{j_2}})_{\epsilon,\epsilon} \cdots \\ &\stackrel{4}{=} \sum_{(j_1, j_2, \dots) \in P_l} \sum_{\pi_1, \pi_2, \dots \in \Gamma^*} (\widehat{M}_{z_j,x}(\widehat{M}_{x,x})^* \widehat{M}_{x,z_{j_1}} + \widehat{M}_{z_j,z_{j_1}})_{\epsilon,\pi_1} (\widehat{M}_{z_{j_1},x}(\widehat{M}_{x,x})^* \widehat{M}_{x,z_{j_2}} + \widehat{M}_{z_{j_1},z_{j_2}})_{\pi_1,\pi_2} \cdots \\ &= (((M_{z,x}(M_{x,x})^* M_{x,z} + M_{z,z})_{\epsilon,\epsilon})^{\omega,l})_j,\end{aligned}\tag{26}$$

where the forth equality uses the fact that $(\widehat{M}_{z_i,x}(\widehat{M}_{x,x})^* \widehat{M}_{x,z_j} + \widehat{M}_{z_i,z_j})_{\epsilon,\pi} = 0$ for $\pi \neq \epsilon$ which is because $(\widehat{M}_{z_i,z_j})_{\epsilon,\pi} = 0$ for $\pi \neq \epsilon$ by definition and because, by our construction, we have

$$M_{z,x}(M_{x,x})^* M_{x,z} = \sum_{1 \leq j \leq n} (M_{z,x})_{\epsilon,Z_j} ((M_{x,x})^*)_{Z_j,Z_j} (M_{x,z})_{Z_j,\epsilon}.$$

Inductively, the above argument can be applied to all factors $(\widehat{M}_{z_{j_i},x}(\widehat{M}_{x,x})^* \widehat{M}_{x,z_{j_{i+1}}} + \widehat{M}_{z_{j_i},z_{j_{i+1}}})_{\pi_i,\pi_{i+1}}$ because we learn from the preceding factor that $\pi_i = \epsilon$.

Now, we proceed from the other direction. From Theorem 30, we know that for the simple ω -reset pushdown automaton \mathfrak{A}_m^l and a variable z_j , we have

$$\begin{aligned}((M^{\omega,l})_{\epsilon})_{z_j} &= (((M_{z,z} + M_{z,x}(M_{x,x})^* M_{x,z})_{\epsilon,\epsilon})^{\omega,l})_j \\ &= \varrho(\sigma)_j^{\omega,l},\end{aligned}$$

where the last equality is by (26). This completes the proof. \square

We now combine our previous discussion and Theorem 38 to get our second main result.

Corollary 39. *Let S be a continuous star-omega semiring with the underlying semiring S being commutative and let $r \in S^{\text{alg}}\langle\langle\Sigma^*\rangle\rangle \times S^{\text{alg}}\langle\langle\Sigma^\omega\rangle\rangle$.*

Then there exists a simple ω -reset pushdown automaton with behavior r .

Proof. Let $r \in S^{\text{alg}}\langle\langle\Sigma^*\rangle\rangle \times S^{\text{alg}}\langle\langle\Sigma^\omega\rangle\rangle$. As discussed on page 29, by Theorem 14 (and Theorem 5), r is a component of a canonical solution of an ω -algebraic system in Greibach normal form over $S\langle\langle\Sigma^*\rangle\rangle \times S\langle\langle\Sigma^\omega\rangle\rangle$. Let (18) be such a system and assume that the m^{th} component of the l^{th} canonical solution of (18) is r , i.e., assume $\tau_m = r$ for the l^{th} canonical solution τ .

Now, we can construct the simple ω -reset pushdown automata \mathfrak{A}_m^l induced by the Greibach normal form (14), (20), for which, by Theorem 38, $(\|\mathfrak{A}_1^l\|, \dots, \|\mathfrak{A}_n^l\|)$ is the l^{th} canonical solution of (18). As the l^{th} canonical solution is unique, we can conclude that

$$\|\mathfrak{A}_m^l\| = \tau_m = r.$$

\square

8. Discussion

We have extended the characterization of ω -algebraic series so that we can use the ω -Kleene closure to transfer the property of Greibach normal form from algebraic systems to mixed ω -algebraic systems. This generalizes a fundamental property from context-free languages.

We believe that the same technique can be used to transfer other properties of algebraic systems to infinite words. Cohen, Gold [4] use this technique also for the elimination of chain rules, for the Chomsky normal form and for effective decision methods of emptiness, finiteness and infiniteness.

The second part of this paper applies the Greibach normal form for the construction of ω -pushdown automata. Simple ω -reset pushdown automata do not use ϵ -transitions; in the literature, this is also called a *realtime* pushdown automaton. Realtime pushdown automata read a symbol of the input word in every transition - exactly like context-free grammars in Greibach normal form generate a letter in every derivation step. Additionally, each derivation step of context-free grammars in Greibach normal form increases the number of non-terminals in the sentential form by at most one. We showed that for realtime pushdown automata it suffices to handle at most one stack symbol per transition. Here the Greibach normal form provides exactly the properties needed to construct simple ω -reset pushdown automata.

The model of simple ω -reset pushdown automata seems to be very natural. They occur when applying general homomorphisms to nested-word automata [1, 8]. Their unweighted counterparts have been used for a Büchi-type logical characterization of timed pushdown languages of finite words [12] and ω -context-free languages [8]. Also in the weighted setting, simple reset pushdown automata of finite words have been used in [13]. A Büchi-type logical characterization for weighted ω -context-free languages is currently in development and uses the simple ω -reset pushdown automata introduced here.

References

- [1] Blass, A., & Gurevich, Y. (2006). A note on nested words. *Microsoft Research*, URL: <https://www.microsoft.com/en-us/research/publication/180-a-note-on-nested-words/>.
- [2] Bloom, S. L., & Ésik, Z. (1993). *Iteration Theories*. EATCS Monographs on Theoretical Computer Science. Springer. doi:10.1007/978-3-642-78034-9.
- [3] Chomsky, N., & Schützenberger, M. P. (1963). The algebraic theory of context-free languages. In *Studies in Logic and the Foundations of Mathematics* (pp. 118–161). Elsevier volume 35: Computer Programming and Formal Systems. doi:10.1016/S0049-237X(08)72023-8.
- [4] Cohen, R. S., & Gold, A. Y. (1977). Theory of ω -languages I: Characterizations of ω -context-free languages. *Journal of Computer and System Sciences*, 15, 169–184. doi:10.1016/S0022-0000(77)80004-4.
- [5] Conway, J. H. (1971). *Regular Algebra and Finite Machines*. Chapman and Hall.
- [6] Droste, M., Dziadek, S., & Kuich, W. (2019). Weighted simple reset pushdown automata. *Theoretical Computer Science*, 777, 252–259. doi:10.1016/j.tcs.2019.01.016.
- [7] Droste, M., Dziadek, S., & Kuich, W. (2019). Greibach normal form for ω -algebraic systems and weighted simple ω -pushdown automata. In *Foundations of Software Technology and Theoretical Computer Science (FSTTCS 2019)* (pp. 38:1–38:14). volume 150 of *LIPIcs*. doi:10.4230/LIPIcs.FSTTCS.2019.38.
- [8] Droste, M., Dziadek, S., & Kuich, W. (2020). Logic for ω -pushdown automata. *Information and Computation*, . Special issue on "Weighted Automata", Accepted for publication.
- [9] Droste, M., Ésik, Z., & Kuich, W. (2017). The triple-pair construction for weighted ω -pushdown automata. In *Conference on Automata and Formal Languages (AFL 2017)* (pp. 101–113). volume 252 of *Electronic Proceedings in Theoretical Computer Science*. doi:10.4204/EPTCS.252.12.
- [10] Droste, M., & Kuich, W. (2017). A Kleene theorem for weighted ω -pushdown automata. *Acta Cybernetica*, 23, 43–59. doi:10.14232/actacyb.23.1.2017.4.
- [11] Droste, M., Kuich, W., & Vogler, H. (Eds.) (2009). *Handbook of Weighted Automata*. EATCS Monographs in Theoretical Computer Science. Springer. doi:10.1007/978-3-642-01492-5.
- [12] Droste, M., & Perevoshchikov, V. (2015). A logical characterization of timed pushdown languages. In *Computer Science Symposium in Russia (CSR 2015)* (pp. 189–203). Springer volume 9139 of *LNCS*. doi:10.1007/978-3-319-20297-6_13.
- [13] Droste, M., & Perevoshchikov, V. (2015). Logics for weighted timed pushdown automata. In *Fields of Logic and Computation II* (pp. 153–173). Springer. doi:10.1007/978-3-319-23534-9_9.
- [14] Eilenberg, S. (1974). *Automata, Languages, and Machines* volume 59, Part A of *Pure and Applied Mathematics*. Elsevier. doi:10.1016/S0079-8169(08)60880-6.
- [15] Elgot, C. C. (1976). Matricial theories. *Journal of Algebra*, 42, 391–421. doi:10.1016/0021-8693(76)90106-X.
- [16] Ésik, Z., & Kuich, W. (2004). A semiring-semimodule generalization of ω -context-free languages. In *Theory Is Forever* (pp. 68–80). Springer volume 3113 of *LNCS*. doi:10.1007/978-3-540-27812-2_7.
- [17] Ésik, Z., & Kuich, W. (2005). A semiring-semimodule generalization of ω -regular languages II. *Journal of Automata, Languages and Combinatorics*, 10, 243–264. doi:10.25596/jalc-2005-243.

- [18] Ésik, Z., & Kuich, W. (2007). *Modern Automata Theory*. URL: <http://www.dmg.tuwien.ac.at/kuich>.
- [19] Ésik, Z., & Kuich, W. (2007). On iteration semiring-semimodule pairs. *Semigroup Forum*, 75, 129–159. doi:10.1007/s00233-007-0709-7.
- [20] Greibach, S. A. (1965). A new normal-form theorem for context-free phrase structure grammars. *Journal of the ACM*, 12, 42–52. doi:10.1145/321250.321254.
- [21] Kuich, W. (1997). Semirings and formal power series: Their relevance to formal languages and automata. In *Handbook of Formal Languages* chapter 9. (pp. 609–677). Springer volume 1: Word, Language, Grammar. doi:10.1007/978-3-642-59136-5_9.
- [22] Kuich, W., & Salomaa, A. (1986). *Semirings, Automata, Languages* volume 5 of *EATCS Monographs on Theoretical Computer Science*. Springer. doi:10.1007/978-3-642-69959-7.