

The lattice of subsemilattices of a semilattice

LEONID LIBKIN* AND ILYA MUCHNIK

This note makes two observations about lattices of subsemilattices. First, we establish relationship between direct decompositions of such lattices and ordinal sum decompositions of semilattices. Then we give a characterization of the subsemilattice-lattices.

Let us recall some terminology. L will always stand for a semilattice, whose operation will be denoted by \circ . The ordering on L is given by letting $l_1 \leq l_2$ iff $l_1 \circ l_2 = l_2$, i.e. L is always a join-semilattice. Subsemilattices of L , ordered by inclusion, form a subsemilattice-lattice denoted by $\text{Sub } L$. In $\text{Sub } L$ the meet operation is intersection, and the join operation is defined as follows: $L_1 \vee L_2 = L_1 \cup L_2 \cup \{l_1 \circ l_2 \mid l_1 \in L_1, l_2 \in L_2\}$. An element a of an arbitrary lattice \mathcal{L} is called *neutral* if $m(a, x, y) = M(a, x, y)$ for all $x, y \in \mathcal{L}$, where $m(a, x, y) = (a \wedge x) \vee (a \wedge y) \vee (x \wedge y)$ and $M(a, x, y) = (a \vee x) \wedge (a \vee y) \wedge (x \vee y)$. Notice that $m(a, x, y) \leq M(a, x, y)$ holds in any lattice.

LEMMA 1. *Let L be a semilattice and L_0 its subsemilattice. Then L_0 is a neutral element of $\text{Sub } L$ iff $L - L_0$ is a subsemilattice of L and every element of L_0 is comparable with every element of $L - L_0$.*

Proof. Let L_0 be a subsemilattice of L such that $L - L_0$ is a subsemilattice of L as well and every element of L_0 is comparable with every element of $L - L_0$. We must prove that, for any $L_1, L_2 \in \text{Sub } L$, $M(L_0, L_1, L_2) \subseteq m(L_0, L_1, L_2)$. Let $x \in M(L_0, L_1, L_2)$. Since $L_0 \vee L_i = L_0 \cup L_i$, $i = 1, 2$, there are 12 cases, but only one of them is nontrivial: $x \in L_0$ and $x = l_1 \circ l_2$, where $l_1 \in L_1$, $l_2 \in L_2$. If l_1 and l_2 are comparable, then either $x \in L_1$ or $x \in L_2$; hence $x \in m(L_0, L_1, L_2)$. If l_1 and l_2 are not comparable, then $l_1, l_2 \in L_0$ and $x \in (L_0 \wedge L_1) \vee (L_0 \wedge L_2) \subseteq m(L_0, L_1, L_2)$. Conversely, if $L_0 \in \text{Sub } L$ and $L - L_0$ is not a subsemilattice of L , then there exist

Presented by R. Freese.

Received November 3, 1989; accepted in final form May 19, 1992.

*While working on the final version, this author was supported by NSF Grants IRI-86-10617 and CCR-90-57570 and ONR Grant N00014-88-K0634.

$l_1, l_2 \notin L_0$ such that $l_1 \circ l_2 \in L_0$. But then $m(L_0, \{l_1\}, \{l_2\}) \neq M(L_0, \{l_1\}, \{l_2\})$. If $L - L_0$ is a subsemilattice of L and there exist incomparable $l_1 \in L_0$, $l_2 \notin L_0$ and $l = l_1 \circ l_2$, then $m(L_0, L - L_0, \{l_2\}) \neq M(L_0, L - L_0, \{l_2\})$ if $l \notin L_0$ and $m(L_0, \{l_1\}, \{l_2\}) \neq M(L_0, \{l_1\}, \{l_2\})$ if $l \in L_0$. Hence, L_0 is not neutral.

LEMMA 2. *Sub $L \simeq \mathcal{L}_1 \times \mathcal{L}_2$ iff there exists a neutral element L_0 of Sub L and that $\mathcal{L}_1 \simeq \text{Sub } L_0$ and $\mathcal{L}_2 \simeq \text{Sub } L - L_0$.*

Proof. By Theorem 1 of [2, p. 152], the direct decompositions of Sub L into two factors are of form $\text{Sub } L \simeq (L_0) \times [L_0)$, where L_0 is neutral. By Lemma 1, $\varphi : \text{Sub } L - L_0 \rightarrow [L_0)$ defined by $\varphi(L') = L' \cup L_0$ is a lattice isomorphism if L_0 is neutral. The lemma follows now from the fact that $\text{Sub } L_0 \simeq (L_0]$. \square

COROLLARY 1. *An arbitrary semilattice L can not be represented as an ordinal sum of its proper subsemilattices iff Sub L is directly indecomposable.* \square

COROLLARY 2. *If L is finite, then Sub L is directly indecomposable iff it is subdirectly irreducible.*

Proof. One direction is obvious. To prove that a directly indecomposable Sub L is subdirectly irreducible, assume that $|L| \geq 2$, since Sub L for a one-element L is a two-element chain and, therefore, subdirectly irreducible. Let $\mathbf{1}$ be the greatest element of L . We will show that $\Theta(\emptyset, \{\mathbf{1}\})$ is a unique atom of the congruence lattice of Sub L . Since one-element subsemilattices are exactly the atoms of Sub L , it is enough to show that $\Theta(\emptyset, \{\mathbf{1}\}) \leq \Theta(\emptyset, \{l\})$ for each $l \in L$, $l \neq \mathbf{1}$ or, equivalently, that $\{\mathbf{1}\}/\emptyset \approx_\omega \{l\}/\emptyset$. Notice that if $l_1 \circ l_2 = l$ in L , then $\{l\}/\emptyset \sim_\omega \{l_1, l_2, l\}/\{l_2\} \sim_\omega \{l_1\}/\emptyset$ in Sub L .

Since Sub L is directly indecomposable, by Corollary 1 for any element $l \in L$, $l \neq \mathbf{1}$, there exists $l' \in L$ incomparable with l , i.e. $l \circ l' > l$. Since L is finite, for any $l \neq \mathbf{1}$ there is a finite sequence l_0, l_1, \dots, l_{2n} , where $l_0 = l$, $l_{2n} = \mathbf{1}$, l_{2i} and l_{2i+1} are incomparable and $l_{2i+2} = l_{2i} \circ l_{2i+1}$, $i = 0, \dots, n-1$. The existence of such a sequence and the observation made above immediately imply $\{\mathbf{1}\}/\emptyset \approx_\omega \{l\}/\emptyset$. \square

Notice that any neutral element of Sub L is complemented. Neutral complemented elements of any lattice \mathcal{L} form a Boolean sublattice of \mathcal{L} denoted by $\text{Cen}(\mathcal{L})$ [2]. It follows from Lemma 1 that intersection of an arbitrary family of neutral elements of Sub L is neutral. Hence, $\text{Cen}(\text{Sub } L)$ is a complete lattice. Moreover, intersection of all neutral elements containing $l \in L$ is an atom of $\text{Cen}(\text{Sub } L)$. Therefore, $\text{Cen}(\text{Sub } L)$ is an atomic Boolean lattice whose atoms are exactly ordinally indecomposable subsemilattices of L . From this we conclude

THEOREM 1. *Let L be an arbitrary semilattice. Then $\text{Sub } L$ can be represented as a direct product of directly indecomposable lattices, $\text{Sub } L \simeq \prod_{i \in I} \text{Sub } L_i$, where $L = \bigoplus_{i \in I} L_i$ is a representation of L as an ordinal sum of ordinally indecomposable subsemilattices.* \square

In the finite case the structure of $\text{Cen}(\text{Sub } L)$ allows us to list all the direct decompositions of $\text{Sub } L$. If $L = \bigoplus_{i \in I} L_i$, where each L_i is ordinally indecomposable and $\text{Sub } L \simeq \mathcal{L}_1 \times \cdots \times \mathcal{L}_m$, then there exist disjoint sets $I_1, \dots, I_m \subseteq I$ such that $I_1 \cup \cdots \cup I_m = I$, $L'_j = \bigoplus_{i \in I_j} L_i$ and $\mathcal{L}_j \simeq \text{Sub } L'_j$ for $j = 1, \dots, m$.

We conclude the paper by characterizing the subsemilattice-lattices. An atomistic lattice \mathcal{L} is called *biatomic* [1] if for any two non-zero $x, y \in \mathcal{L}$ and an atom $z \leq x \vee y$ there exist atoms $x' \leq x, y' \leq y$ such that $z \leq x' \vee y'$. We say that a biatomic lattice \mathcal{L} satisfies property (S_n) if for any ideal V generated by n atoms $a_1, \dots, a_n \in \mathcal{L}$ there exists a finite semilattice L_V such that $V \simeq \text{Sub } L_V$, and the natural embedding of ideals $V \rightarrow W$ induces the embedding of semilattices $L_V \rightarrow L_W$.

THEOREM 2. *A lattice \mathcal{L} is isomorphic to $\text{Sub } L$ for some semilattice L iff it is algebraic, biatomic and satisfies (S_3) .*

Proof. The ‘only if’ part is obvious. To prove the ‘if’ part, denote the set of atoms of \mathcal{L} by $A(\mathcal{L})$ and the set of atoms under $x \in \mathcal{L}$ by $A(x)$. Notice that (S_3) implies that $(*)$ for every $X \subseteq A(\mathcal{L})$ with $|X| \leq 3$ there exists a semilattice operation \circ_X on $A(\bigvee X)$ such that $(\bigvee X) \simeq \text{Sub } \langle A(\bigvee X), \circ_X \rangle$ and $\circ_Y = \circ_X \upharpoonright_{A(\bigvee Y)}$ for every $Y \subseteq A(\bigvee X)$ with $|Y| \leq 3$.

Define a binary operation \circ on $A(\mathcal{L})$ by $a_1 \circ a_2 = a_1 \circ_{\{a_1, a_2\}} a_2$. Clearly, \circ is idempotent and commutative. That \circ is associative follows from $(*)$. Thus, \circ is a semilattice operation on $A(\mathcal{L})$. Define $\varphi : \mathcal{L} \rightarrow \text{Sub } \langle A(\mathcal{L}), \circ \rangle$ by $\varphi(y) = \{x \in A(\mathcal{L}) \mid x \leq y\}$. That φ is well-defined follows from $(*)$. The remaining properties of \mathcal{L} guarantee that φ is an isomorphism. Thus, $\mathcal{L} \simeq \text{Sub } \langle A(\mathcal{L}), \circ \rangle$. \square

Remark. Neutral elements of a lattice $\text{Sub } L$ were characterized in Lemma 1. One can easily check that a weaker condition characterizes distributive and standard elements. In fact, L_0 is a distributive element of $\text{Sub } L$ iff it is standard iff for all $l_1 \in L_0, l_2 \notin L_0$ either $l_1 \leq l_2$ or $l_1 \circ l_2 \in L_0$.

Acknowledgements

We would like to thank E. T. Schmidt and W. Dziobiak for their helpful comments.

REFERENCES

- [1] Bennett, M. K., *Biatomic lattices*, Algebra Universalis 24 (1987), 60–73.
- [2] Grätzer, G. *General Lattice Theory*, Springer Verlag, Berlin, 1978.

*Department of Computer and Information Science
University of Pennsylvania
Philadelphia, Pennsylvania
U.S.A.*

*24 Chestnut St.
Waltham, Massachusetts
U.S.A.*