



Contents lists available at ScienceDirect

Journal of Algebra

www.elsevier.com/locate/jalgebra



Products of ideals and jet schemes

Gleb Pogudin¹

Institute for Algebra, Johannes Kepler University, 4040 Linz, Austria



ARTICLE INFO

Article history:

Received 30 January 2017

Available online 2 February 2018

Communicated by Leonard L. Scott, Jr.

MSC:

12H20

12H05

14B10

Keywords:

Jets schemes

Algebraic differential equations

Differential polynomials

ABSTRACT

In the present paper, we give a full description of the jet schemes of the polynomial ideal $(x_1 \cdots x_n) \in k[x_1, \dots, x_n]$ over a field of zero characteristic. We use this description to answer questions about products and intersections of ideals emerged recently in algorithmic studies of algebraic differential equations.

© 2018 Elsevier Inc. All rights reserved.

1. Introduction

Properties of ideals in rings of differential polynomials are essential for the algorithmic study of algebraic differential equations [3,15]. The following fact is an important part of the recent first bound for the effective differential elimination ([15]). If h_1, \dots, h_n are natural numbers, then there exists d such that for every ideals I_1, \dots, I_n in a commutative associative differential algebra over a field of zero characteristic

E-mail address: pogudin.gleb@gmail.com.

¹ Current address: Courant Institute of Mathematical Sciences, New York University, USA.

$$\left(I_1^{(h_1)} \cdots I_n^{(h_n)}\right)^d \subset (I_1 \cdots I_n)^{(h_1 + \cdots + h_n)}, \quad (1.1)$$

where, for an ideal I , we denote the ideal generated by derivatives of elements of I of order at most h by $I^{(h)}$. Such an inclusion allows us to reduce a problem about an arbitrary ideal to the problem about an ideal with additional useful properties (for example, to prime ideals in [15]). This kind of reduction is expected to have many potential applications to the algorithmic problems about algebraic differential equations. The value of such d was not needed for obtaining the bound in [15], but we expect that it can be used for refining this and related bounds similarly to the way the Noether exponent was used in [3, Lemma 3.1] and [15, Lemma 5.5]. The problem of determining the number d in (1.1) as well as many other questions about products of ideals in differential algebras can be reduced to questions about the jet ideal of the polynomial ideal generated by $x_1 \cdots x_n \in k[x_1, \dots, x_n]$ (see [15, Lemma 6.2]).

Recently, jet schemes were successfully applied to study singularities of algebraic varieties (for example, [14]). In [1,16], jet ideals of the ideal generated by $x_1 \cdots x_n$ were studied, minimal primes and their multiplicities were found. In the context of differential algebra, jet schemes and related objects proved to be one of main tools in differential algebraic geometry (see [12,8]).

In the present paper, we generalize results of [1,16] giving a full description of jet schemes of the ideal generated by $x_1 \cdots x_n$ (Section 4), where an unexpected connection to Shubert calculus occurs (see Proposition 3.10). Then we use the obtained information in order to solve the original problem about differential algebraic equations, namely find the minimal possible number d in (1.1) (see Corollary 5.2 and Corollary 5.4).

We also note that our problem is connected to a classical membership problem for the differential ideal generated by $x_1 \cdots x_n$ (see [9,5,6]). In particular, if a differential polynomial belongs to any of the jet ideals of the polynomial ideal generated by $x_1 \cdots x_n$, then it belongs to the corresponding differential ideal, and if a differential polynomial does not belong to any of these jet ideals, it does not belong to the differential ideal. Our structural results give an effective way for deciding if any of these situations takes place (see Remark 5.5).

2. Preliminaries

2.1. Differential algebra

Throughout the paper, all fields are assumed to be of characteristic zero. Unless otherwise stated, all algebras are commutative, associative, and with unity.

Let R be a ring. A map $D: R \rightarrow R$ satisfying $D(a+b) = D(a) + D(b)$ and $D(ab) = aD(b) + D(a)b$ for all $a, b \in R$ is called a *derivation*. A *differential ring* R is a ring with a specified derivation. In this case, we will denote $D(x)$ by x' and $D^n(x)$ by $x^{(n)}$. A differential ring that is a field will be called a *differential field*.

A differential ring A is said to be a *differential K -algebra* over a differential field K if A is a K -algebra and the restriction of the derivation of A on K coincides with the derivation on K . An ideal I of a differential ring R is said to be a *differential ideal* if $a' \in I$ for all $a \in I$. The differential ideal generated by $a_1, \dots, a_n \in I$ will be denoted by $[a_1, \dots, a_n]$.

Let A be a differential K -algebra. We consider the polynomial ring

$$A[x, x', \dots, x^{(n)}, \dots],$$

where x, x', x'', \dots are algebraically independent variables. Extending the derivation from A to $A[x, x', \dots]$ by $D(x^{(n)}) = x^{(n+1)}$ we obtain a differential algebra. This algebra is called the *algebra of differential polynomials* in x over A and we denote it by $A\{x\}$. Iterating this construction, we define the algebra of differential polynomials in variables x_1, \dots, x_n over A and denote it by $A\{x_1, \dots, x_n\}$.

2.2. Jet schemes

For general scheme-theoretic treatment, we refer reader to [14] (the same object is called the arc space in [12]). We will need only coordinate description of the jet ideal of the polynomial ideal.

Let $I = (g_1, \dots, g_s)$ be an ideal in the polynomial ring $k[x_1, \dots, x_n]$ and $m \in \mathbb{Z}_{\geq 0} \cup \{\infty\}$. By $J_m(I)$ we will denote the m -th jet ideal of I which is an ideal in the polynomial ring

$$R_m = k[x_1^{(0)}, x_1^{(1)}, \dots, x_1^{(m)}, x_2^{(0)}, \dots, x_n^{(m)}]$$

generated by coefficients of the following truncated power series

$$g_i(x_1^{(0)} + x_1^{(1)}t + \dots + x_1^{(m)}t^m, \dots, x_n^{(0)} + x_n^{(1)}t + \dots + x_n^{(m)}t^m) \in k[[t]]/(t^{m+1}),$$

where $i = 1, \dots, s$.

For example, the 0-th jet ideal of I coincides with I after renaming $x_i^{(0)}$ into x_i for all i .

Remark 2.1. Let us consider the infinite jet ring R_∞ over a differential field K . It can be endowed with the structure of differential K -algebra by $(x_i^{(j)})' = (j+1)x_i^{(j+1)}$. Moreover, the differential algebra obtained by this procedure is differentially isomorphic to the algebra of differential polynomials $K\{y_1, \dots, y_n\}$ via $\varphi(x_i^{(j)}) = \frac{y_i^{(j)}}{j!}$. Then if coefficients of g_1, \dots, g_s are constants in K , the infinite jet ideal $J_\infty(I)$ corresponds to a differential ideal generated by $g_1(y_1, \dots, y_n), \dots, g_s(y_1, \dots, y_n)$ in $K\{y_1, \dots, y_n\}$. This correspondence will allow us to apply results about jet schemes to problems about algebraic differential equations.

3. Facts about symmetric polynomials

In this section we collect facts about the ideal generated by symmetric polynomials with zero constant term, which plays a central role in Shubert calculus (see [10,11]). We also introduce algebras $A_\lambda(k)$ (Definition 3.3) which locally describe the jet schemes of the ideal $(x_1 \cdots x_n)$ (Section 4).

Consider the algebra $k[z_1, \dots, z_\ell]$ of polynomials over k . By \mathbf{z} we denote the tuple (z_1, \dots, z_ℓ) . By $h_d(\mathbf{z})$ we denote the complete symmetric polynomial of degree d which is a sum of all monomials of degree d in z_1, \dots, z_ℓ :

$$h_d(\mathbf{z}) = \sum_{i_1 \leq i_2 \leq \dots \leq i_d} z_{i_1} z_{i_2} \cdots z_{i_d}.$$

By \mathbf{z}_k we denote the tuple (z_k, \dots, z_ℓ) . For example, $\mathbf{z}_1 = \mathbf{z}$, $\mathbf{z}_\ell = (z_\ell)$, and $h_d(\mathbf{z}_\ell) = z_\ell^d$. By $I_S \subset k[z_1, \dots, z_\ell]$ we denote the ideal generated by all symmetric polynomials with respect to z_1, \dots, z_ℓ with zero constant term. We will use the following lemma which follows from Proposition 2.1 and Remark 2.1 from [13].

Lemma 3.1. *Polynomials $h_1(\mathbf{z}_1), h_2(\mathbf{z}_2), \dots, h_\ell(\mathbf{z}_\ell)$ constitute a Gröbner basis of I_S with respect to lexicographical ordering $z_1 > z_2 > \dots > z_\ell$.*

Also, for every $j \geq i$, the polynomial $h_j(\mathbf{z}_i)$ lies in I_S .

Throughout the rest of the section, we will refer to the basis from Lemma 3.1 as the basis of I_S .

Lemma 3.2. *Let $p(\mathbf{z}) \in k[z_1, \dots, z_\ell]$, and $d \geq 1$. By \tilde{p} we denote the reduction of p with respect to the Gröbner basis of I_S with respect to lexicographic ordering with $z_1 > z_2 > \dots > z_\ell$. Then*

$$p(z_1, \dots, z_s, z_{s+1}, \dots, z_\ell) = p(z_1, \dots, z_{s+1}, z_s, \dots, z_\ell)$$

implies

$$\tilde{p}(z_1, \dots, z_s, z_{s+1}, \dots, z_\ell) = \tilde{p}(z_1, \dots, z_{s+1}, z_s, \dots, z_\ell).$$

Proof. By g_i we denote the i -th element of the Gröbner basis of I_S . Lemma 3.1 implies that there are ℓ of them, and $g_i = h_i(\mathbf{z}_i)$. We will prove that if p is not reduced with respect to g_1, \dots, g_ℓ , then there exists $q(\mathbf{z})$ such that the leading monomial of q is lower than the leading monomial of p , q is also symmetric with respect to z_s and z_{s+1} , and $p - q \in I_S$. We can not apply this procedure infinitely many times, so at some point we will obtain q such that q is symmetric with respect to z_s and z_{s+1} and reduced with respect to g_1, \dots, g_ℓ . The uniqueness of the reduction with respect to a Gröbner basis implies that $q = \tilde{p}$.

So, assume that p is not reduced with respect to I_S . Since the reduction with respect to a Gröbner basis is always equal to the lead-reduction, the leading monomial of p is not reduced with respect to g_i for some i . Let the leading monomial of p be $M_1 = z_1^{d_1} \cdots z_\ell^{d_\ell}$. Due to the symmetry, the monomial $M_2 = z_1^{d_1} \cdots z_s^{d_{s+1}} z_{s+1}^{d_s} \cdots z_\ell^{d_\ell}$ occurs in p with the same coefficient, say c . Since M_1 is greater or equal then M_2 , $d_s \geq d_{s+1}$. We consider three cases

1. $i \neq s$ and $i \neq s+1$. Let us consider

$$q = \begin{cases} p - c \frac{M_1 + M_2}{z_i^{d_i}} g_i, & \text{if } d_s > d_{s+1}, \\ p - c \frac{M_1}{z_i^{d_i}} g_i, & \text{if } d_s = d_{s+1}. \end{cases}$$

The terms cM_1 cancel, so the leading monomial of q is less then M_1 . Moreover, q is still symmetric with respect to z_s and z_{s+1} .

2. $i = s$. Since $d_s \geq s$, Lemma 3.1 implies that $h_{d_s}(\mathbf{z}_s), h_{d_s+i}(\mathbf{z}_{s+1}) \in I_S$ for every $i > 0$. By N we denote $\frac{M_1}{z_s^{d_s} z_{s+1}^{d_{s+1}}}$. Consider

$$q = p - cN \left(z_{s+1}^{d_{s+1}} h_{d_s}(\mathbf{z}_s) - \sum_{i=1}^{d_{s+1}} z_s^{d_{s+1}-i} h_{d_s+i}(\mathbf{z}_{s+1}) \right).$$

We denote $z_{s+1}^{d_{s+1}} h_{d_s}(\mathbf{z}_s)$ and $\sum_{i=1}^{d_{s+1}} z_s^{d_{s+1}-i} h_{d_s+i}(\mathbf{z}_{s+1})$ by f_1 and f_2 , respectively. Since $d_s \geq d_{s+1}$, the leading monomial of $f_1 - f_2$ is $z_s^{d_s} z_{s+1}^{d_{s+1}}$. Hence, the leading monomial of q is less than the leading monomial of p . Since N does not involve neither z_s , nor z_{s+1} , it is sufficient to prove that $f_1 - f_2$ is symmetric with respect to z_s and z_{s+1} . The polynomial f_1 is a sum of all monomials S in \mathbf{z}_s of the total degree $d_s + d_{s+1}$ with $\deg_{z_{s+1}} S \geq d_{s+1}$. The polynomial f_2 is a sum of all monomials S in \mathbf{z}_s of the total degree $d_s + d_{s+1}$ with $\deg_{z_s} S < d_{s+1}$. Let $S_1 = z_1^{e_1} \cdots z_\ell^{e_\ell}$ be an arbitrary monomial of the total degree $d_s + d_{s+1}$, and $S_2 = z_1^{e_1} \cdots z_s^{e_{s+1}} z_{s+1}^{e_s} \cdots z_\ell^{e_\ell}$ is the monomial symmetric to S_1 with respect to z_s and z_{s+1} . We also assume that $e_s > e_{s+1}$. It is sufficient to prove that S_1 and S_2 occur in $f_1 - f_2$ with the same coefficient. There are three cases:

- (a) $e_s > e_{s+1} \geq d_{s+1}$. Then S_1 and S_2 occur in f_1 , but none of them occurs in f_2 .
 - (b) $d_{s+1} > e_s > e_{s+1}$. Then S_1 and S_2 occur in f_2 , but none of them occurs in f_1 .
 - (c) $e_s \geq d_{s+1} > e_{s+1}$. Then S_2 occurs both in f_1 and f_2 , so $f_1 - f_2$ does not involve S_2 . Also, S_1 does not occur in f_1 and f_2 .
3. $i = s+1$. Since $d_s \geq d_{s+1}$, p is also not reduced with respect to g_s , so this case can be reduced to the previous. \square

Definition 3.3. Let us remind that by I_S we denote the ideal of $k[z_1, \dots, z_\ell]$ generated by symmetric polynomials with respect to z_1, \dots, z_ℓ without constant term. For the

rest of the section, we fix a positive integer n . For the vector $\lambda = (\lambda_1, \dots, \lambda_n) \in \mathbb{Z}_{\geq 0}^n$, by Λ_i we denote the partial sum $\lambda_1 + \dots + \lambda_{i-1}$, assuming also $\Lambda_1 = 0$. Assume that $\Lambda_{n+1} = \lambda_1 + \dots + \lambda_n = \ell$. We will say that a polynomial $p \in k[z_1, \dots, z_\ell]$ is λ -symmetric if it is symmetric with respect to $z_{\Lambda_i+1}, \dots, z_{\Lambda_{i+1}}$ for every $1 \leq i \leq n$. The notion of λ -symmetric polynomials generalizes the notion of block-symmetric polynomials [2].

For Lemmas 3.4 and 3.5, we fix some $\lambda = (\lambda_1, \dots, \lambda_n) \in \mathbb{Z}_{\geq 0}^n$ such that $\lambda_1 + \dots + \lambda_n = \ell$. We denote the elementary symmetric polynomial of degree d in t_1, \dots, t_r by $\sigma_d(t_1, \dots, t_r)$ for any variables t_1, \dots, t_r and $1 \leq d \leq r$.

Lemma 3.4. *Let $p \in k[z_1, \dots, z_\ell]$ be a λ -symmetric polynomial belonging to I_S . Then there exist λ -symmetric polynomials $a_1, \dots, a_\ell \in k[z_1, \dots, z_\ell]$ such that*

$$p = a_1 \sigma_1(z_1, \dots, z_\ell) + \dots + a_\ell \sigma_\ell(z_1, \dots, z_\ell).$$

Proof. Since $p \in I_S$, there exist $b_1, \dots, b_\ell \in k[z_1, \dots, z_\ell]$ such that

$$p = b_1 \sigma_1(z_1, \dots, z_\ell) + \dots + b_\ell \sigma_\ell(z_1, \dots, z_\ell). \quad (3.1)$$

Let $G := S_{\lambda_1} \times \dots \times S_{\lambda_n} \subset S_\ell$ be a subgroup in the group of all permutations of z_1, \dots, z_ℓ . We consider a linear operator Sym_λ on $k[z_1, \dots, z_m]$ defined by

$$\text{Sym}_\lambda(q) := \frac{1}{|G|} \sum_{\sigma \in G} \sigma(q), \text{ for } q \in k[z_1, \dots, z_\ell].$$

Then we have $\text{Sym}_\lambda(ab) = a \text{Sym}_\lambda(b)$ for every λ -symmetric a and every $b \in k[z_1, \dots, z_\ell]$. Moreover $\text{Sym}_\lambda(b)$ is λ -symmetric for every $b \in k[z_1, \dots, z_\ell]$. Applying Sym_λ to both sides of (3.1), we obtain

$$p = \text{Sym}_\lambda(b_1) \sigma_1(z_1, \dots, z_\ell) + \dots + \text{Sym}_\lambda(b_\ell) \sigma_\ell(z_1, \dots, z_\ell).$$

Since every polynomial in the image of Sym_λ is λ -symmetric, we set $a_i := \text{Sym}_\lambda(b_i)$ for every $1 \leq i \leq \ell$. \square

Lemma 3.5. *Let $p \in k[z_1, \dots, z_\ell]$ be a λ -symmetric polynomial. Then p belongs to the subalgebra generated by*

$$\{\sigma_j(z_{\Lambda_i+1}, \dots, z_{\Lambda_{i+1}}) \mid 1 \leq i \leq n, 1 \leq j \leq \lambda_i\}.$$

Proof. We will prove the statement of the lemma by induction on n . The base case $n = 1$ is exactly the fundamental theorem on symmetric polynomials. Let $n > 1$ and consider a λ -symmetric polynomial $p \in k[z_1, \dots, z_\ell]$. We write p in the form $\sum_{i=1}^N a_i m_i$, where $a_i \in k[z_1, \dots, z_{\Lambda_{n-1}}]$ and m_i are distinct monomials in $z_{\Lambda_{n-1}+1}, \dots, z_{\Lambda_n}$. Since p

is λ -symmetric, $a_i = a_j$ for every $1 \leq i, j \leq N$ such that m_i and m_j can be obtained from each other by permuting the variables. Grouping together m_i 's with equal a_i 's, we obtain a representation $p = \sum_{i=1}^M b_i s_i$, where $b_i \in k[z_1, \dots, z_{\Lambda_{n-1}}]$ and s_i is a symmetric polynomial in $z_{\Lambda_{n-1}+1}, \dots, z_{\Lambda_n}$. The fundamental theorem on symmetric polynomials imply that each s_i is contained in the subalgebra generated by $\sigma_1(z_{\Lambda_{n-1}+1}, \dots, z_{\Lambda_n}), \dots, \sigma_{\lambda_n}(z_{\Lambda_{n-1}+1}, \dots, z_{\Lambda_n})$. Since p is λ -symmetric, b_1, \dots, b_M are $(\lambda_1, \dots, \lambda_{n-1})$ -symmetric, so they belong to the subalgebra generated by

$$\{\sigma_j(z_{\Lambda_i+1}, \dots, z_{\Lambda_{i+1}}) \mid 1 \leq i \leq n-1, 1 \leq j \leq \lambda_i\}$$

due to the induction hypothesis. Thus, the lemma is proved. \square

Definition 3.6. By $A_\lambda(k)$ we denote the subalgebra of $k[z_1, \dots, z_\ell]/I_S$ consisting of images of all λ -symmetric polynomials.

Lemma 3.7. For all $\ell \in \mathbb{Z}_{>0}$ and $\lambda \in \mathbb{Z}_{\geq 0}^n$ with $\lambda_1 + \dots + \lambda_n = \ell$,

$$\dim_k A_\lambda(k) = \frac{\ell!}{\prod_{i=1}^n \lambda_i!}.$$

Proof. Let us identify every element of $A_\lambda(k)$ with its normal form with respect to the Gröbner basis $\{g_1, \dots, g_\ell\}$ from Lemma 3.1. Lemma 3.2 implies that $A_\lambda(k)$ consists of polynomials p such that p is reduced with respect to $\{g_1, \dots, g_\ell\}$, and p is λ -symmetric. By G we denote the subgroup of S_ℓ such that for every $\sigma \in G$ and $\Lambda_i + 1 \leq j \leq \Lambda_{i+1}$, $\sigma(j)$ satisfies $\Lambda_i + 1 \leq \sigma(j) \leq \Lambda_{i+1}$. Then $G \cong S_{\lambda_1} \times S_{\lambda_2} \times \dots \times S_{\lambda_n}$.

By B we denote the set of vectors $\mathbf{d} = (d_1, \dots, d_\ell) \in \mathbb{Z}_{\geq 0}^\ell$ such that $d_{\Lambda_i+1} \leq \Lambda_i$ and $d_{\Lambda_i+1} \geq d_{\Lambda_i+2} \geq \dots \geq d_{\Lambda_{i+1}}$ for every $i \leq n$. For every such vector, we associate the polynomial

$$f_{\mathbf{d}} = \sum_{\sigma \in G} z_{\sigma(1)}^{d_1} z_{\sigma(2)}^{d_2} \cdots z_{\sigma(\ell)}^{d_\ell}.$$

We claim that these polynomials constitute a k -basis of $A_\lambda(k)$. The leading monomial of $f_{\mathbf{d}}$ is $z_1^{d_1} \cdots z_\ell^{d_\ell}$, so polynomials $f_{\mathbf{d}}$ have distinct leading monomials. Hence, they are linearly independent. It is sufficient to prove that for every $p \in A_\lambda(k)$, the leading monomial of p coincides with the leading monomial of $f_{\mathbf{d}}$ for some \mathbf{d} . If $M = z_1^{e_1} \cdots z_\ell^{e_\ell}$ is the leading monomial of p , then $e_{\Lambda_i+1} \leq \Lambda_i$ for all i since p is reduced with respect to the Gröbner basis. Assume also that for some k and i such that $\Lambda_i + 1 \leq k < \Lambda_{i+1}$, we have $e_k < e_{k+1}$. Due to the symmetry, the monomial $M \frac{z_k^{e_{k+1}} z_{k+1}^{e_k}}{z_k^{e_k} z_{k+1}^{e_{k+1}}}$, which is larger than M , occurs in p . Hence, the exponent of M lies in B .

Thus, we need to compute the cardinality of B . For every fixed i , the number of tuples $(d_{\Lambda_i+1}, \dots, d_{\Lambda_{i+1}})$ such that $\Lambda_i \geq d_{\Lambda_i+1} \geq \dots \geq d_{\Lambda_{i+1}}$ is equal to the number of multisets

of size $\Lambda_{i+1} - \Lambda_i = \lambda_i$ whose elements are chosen from $\{0, 1, \dots, \Lambda_i\}$. The number of such multisets is $\binom{\Lambda_i + \lambda_i}{\lambda_i} = \binom{\Lambda_{i+1}}{\lambda_i}$. Hence, the total number of elements in B is

$$\prod_{i=1}^n \binom{\Lambda_{i+1}}{\lambda_i} = \frac{\ell!}{\prod_{i=1}^n \lambda_i!}. \quad \square$$

Definition 3.8. For every $i \leq n$, by $A_\lambda^i(k)$ we denote the subalgebra of $A_\lambda(k)$ consisting of images of symmetric polynomials from $k[z_{\Lambda_i+1}, \dots, z_{\Lambda_{i+1}}]$.

For every nonzero $a \in A_\lambda(k)$, by $\nu(a)$ we denote the minimal total degree of a monomial in the reduction of a with respect to the Gröbner basis of I_S . Note that since I_S is homogeneous, $\nu(a)$ does not depend on the ordering of variables. We also observe that if $ab \neq 0$, then $\nu(ab) = \nu(a) + \nu(b)$.

The following [Propositions 3.9 and 3.10](#) are important for applications of our results to differential algebra (see the proof of [Theorem 5.1](#)).

Proposition 3.9. *Let $a \in A_\lambda^i(k)$ be an element such that $\nu(a) = r > 0$. Then $a^d = 0$ for all $d > \frac{\lambda_i}{r}(\ell - \lambda_i)$.*

Proof. Reordering variables if necessary, we can assume that $i = 1$. Then a is an image of $p(z_1, \dots, z_{\lambda_1})$, where p is a symmetric polynomial in $z_1, \dots, z_{\lambda_1}$ over k without constant term. Let g_1, \dots, g_ℓ be the Gröbner basis of I_S from [Lemma 3.1](#). [Lemma 3.2](#) implies, that for every d , the reduction of $p(z_1, \dots, z_{\lambda_1})^d$ with respect to this basis is a polynomial symmetric with respect to $z_1, \dots, z_{\lambda_1}$ and with respect to all other variables. Let us denote this reduction by q . Since $\deg_{z_1} q < \deg_{z_1} g_1 = 1$, q does not involve z_1 . Symmetry implies that q does not involve also $z_2, \dots, z_{\lambda_1}$. Analogously, for $i > \lambda_1$ we have $\deg_{z_i} q < \deg_{z_{\lambda_1+1}} g_{\lambda_1+1} = \lambda_1 + 1$. Hence, the total degree of q does not exceed $\lambda_1(\ell - \lambda_1)$. On the other hand, I_S is a homogeneous ideal, so the total degree of q can not be less than $\nu(p^d) = dr$. Thus, for all $d > \frac{\lambda_1}{r}(\ell - \lambda_1)$, q is zero. \square

Proposition 3.10. *Let $a = x_{\Lambda_i+1} + \dots + x_{\Lambda_{i+1}} \in A_\lambda^i(k)$. Then $a^{\lambda_i(\ell - \lambda_i)} \neq 0$.*

Proof. Our proof will use some facts about Shubert polynomials, in particular, Monk's formula. For details and profound account, we refer reader to the original paper [\[11\]](#) and the monograph [\[10\]](#). We will briefly recall the most important facts. Shubert polynomials are polynomials in z_1, \dots, z_ℓ indexed by permutations. By \mathfrak{S}_ω we will denote the Shubert polynomial corresponding to a permutation $\omega \in S_\ell$. We will use several properties of these polynomials:

1. The set of all Shubert polynomials in z_1, \dots, z_ℓ constitute a k -basis of $k[z_1, \dots, z_\ell]/I_S$ (see [\[10, Proposition 2.5.3\]](#)).

2. By s_i we denote the transposition $(i, i + 1) \in S_\ell$. Then $\mathfrak{S}_{s_i} = z_1 + \dots + z_i$ (see [10, §2.7.1]).
3. (Monk's formula) For every permutation $\omega \in S_\ell$:

$$\mathfrak{S}_{s_r} \cdot \mathfrak{S}_\omega \equiv \sum_{\substack{j \leq r < k \\ l(\omega(jk)) = l(\omega) + 1}} \mathfrak{S}_{\omega(jk)} \pmod{I_S},$$

where $l(\omega)$ is the number of inversions in ω (see [10, Theorem 2.7.1] and [11, Theorem 3]).

Let us return to the proof of the proposition. Reordering variables if necessary, we may assume that $i = 1$. Using the language of Shubert polynomials, we want to prove that $\mathfrak{S}_{s_{\lambda_1}}^{\lambda_1(\ell - \lambda_1)} = \mathfrak{S}_{s_{\lambda_1}}^{\lambda_1(\ell - \lambda_1)} \mathfrak{S}_e \neq 0 \pmod{I_S}$, where e is the identity in S_ℓ . The expression $\mathfrak{S}_{s_{\lambda_1}}^{\lambda_1(\ell - \lambda_1)}$ can be represented as a sum of Shubert polynomials with nonnegative integer coefficients by applying Monk's formula $\lambda_1(\ell - \lambda_1) - 1$ times. Due to Monk's formula, every summand in this representation corresponds to some permutation ω such that $l(\omega) = \lambda_1(\ell - \lambda_1)$ and ω can be written as a product of $\lambda_1(\ell - \lambda_1)$ transpositions of the form (jk) , where $j \leq \lambda_1 < k$. Showing that there exists at least one such permutation would imply that there is at least one such summand, so $\mathfrak{S}_{s_{\lambda_1}}^{\lambda_1(\ell - \lambda_1)} \neq 0 \pmod{I_S}$. We claim that the following permutation satisfies both properties

$$\omega = \begin{pmatrix} 1 & 2 & \dots & \lambda_1 & \lambda_1 + 1 & \lambda_1 + 2 & \dots & \ell \\ \ell - \lambda_1 + 1 & \ell - \lambda_1 + 2 & \dots & \ell & 1 & 2 & \dots & \ell - \lambda_1 \end{pmatrix}.$$

Direct computation shows that the number of inversions is exactly $\lambda_1(\ell - \lambda_1)$. The second condition is satisfied, because ω admits the following representation

$$\omega = \prod_{i=1}^{\lambda_1} (i\ell)(i\ell - 1) \cdots (i\lambda_1 + 1). \quad \square$$

Remark 3.11. It can be shown, that ω in the proof above is the only permutation such that $l(\omega) = \lambda_1(\ell - \lambda_1)$ and ω can be written as a product of $\lambda_1(\ell - \lambda_1)$ transpositions of the form (jk) , where $j \leq \lambda_1 < k$. Then we will obtain the congruence $\mathfrak{S}_{s_{\lambda_1}} \equiv N \mathfrak{S}_\omega \pmod{I_S}$, where N is the number of representations of ω as a product of $\lambda_1(\ell - \lambda_1)$ transpositions of the form (jk) , where $j \leq \lambda_1 < k$.

For the special case $\lambda = (2, 1, \dots, 1)$, there is a one-to-one correspondence between the set of such representations and the set of expressions containing $\ell - 2$ pair of parenthesis, which are correctly matched, where transpositions of the form $(1i)$ correspond to open parenthesis, and transpositions of the form $(2i)$ correspond to close parenthesis (it can be proved using the combinatorial discussion after Theorem 3 in [11]). Thus, in this case $N = C_{\ell-2}$, the $\ell - 2$ -th Catalan number. Moreover, since ω is a Grassmanian permutation, [10, Proposition 2.6.8] implies that $\mathfrak{S}_\omega = z_3^2 \cdots z_\ell^2$. Hence, we obtain the

following curious congruence

$$(z_1 + z_2)^{2(\ell-2)} \equiv C_{\ell-2} z_3^2 \cdots z_\ell^2 \pmod{I_S}.$$

4. Structure of the jet schemes of $x_1 \cdots x_n$

The following theorem describes the structure of the m -th jet ideal of $(x_1 \cdots x_n)$ in terms of algebras $A_\lambda(k)$ introduced in Definition 3.3. For an arbitrary $\lambda = (\lambda_1, \dots, \lambda_n) \in \mathbb{Z}_{\geq 0}^n$ such that $\lambda_1 + \dots + \lambda_n = m + 1$, by \mathfrak{p}_λ we denote the prime ideal in $R = k[x_i^{(j)} \mid 1 \leq i \leq n, 0 \leq j \leq m]$ generated by polynomials $x_i^{(j)}$ with $1 \leq i \leq n$ and $0 \leq j < \lambda_i$. In [1, Theorem 2.2] it is proved that minimal primes of $J_m(x_1 \cdots x_n)$ are exactly all ideals of the form \mathfrak{p}_λ .

Theorem 4.1. *Let n, m be positive integers. Consider R and \mathfrak{p}_λ defined above. If we denote the subfield $k(x_i^{(j)} \mid 1 \leq i \leq n, \lambda_i \leq j \leq m)$ of the field of fractions of R by K_λ , then*

$$(R/J_m(x_1 \cdots x_n))_{\mathfrak{p}_\lambda} \cong A_\lambda(K_\lambda),$$

where the isomorphism is an isomorphism of K_λ -algebras.

Lemma 3.7 and Theorem 4.1 imply the following corollary, which gives an alternative proof of the main result of [16].

Corollary 4.2. *For every $\lambda = (\lambda_1, \dots, \lambda_n)$ such that $\lambda_1 + \dots + \lambda_n = m + 1$, the multiplicity of the ideal $J_m(x_1 \cdots x_n)$ along \mathfrak{p}_λ is equal to $\frac{(m+1)!}{\lambda_1! \cdots \lambda_n!}$.*

Proof of Theorem 4.1. For every $i \leq n$, we set $\Lambda_i = \lambda_1 + \dots + \lambda_{i-1}$ (and $\Lambda_0 = 0$). We will denote $J_m(x_1 \cdots x_n)$ by J_m .

Assume that $\lambda_i = m + 1$ for some i , say for $i = 1$. Then $\lambda = (m + 1, 0, \dots, 0)$. The ideal generated by J_m in the localization $R_{\mathfrak{p}_\lambda}$ is $(x_1^{(0)}, \dots, x_1^{(m)})$, so $(R/J_m)_{\mathfrak{p}_\lambda} \cong K_\lambda$. Since $A_\lambda(K_\lambda) \cong K_\lambda$, in what follows we can assume that $\lambda_i \leq m$ for all i .

By Q_λ we denote the primary component of J_m corresponding to prime ideal \mathfrak{p}_λ . Let $B = (R/J_m)_{\mathfrak{p}_\lambda} = (R/Q_\lambda)_{\mathfrak{p}_\lambda}$. Then B is a local K_λ -algebra with maximal ideal $\mathfrak{p}_\lambda B$. For every element $a \in R_{\mathfrak{p}_\lambda}$, by \bar{a} we will denote its image in B .

We also introduce a local algebra $C_\lambda(k)$ defined by

$$C_\lambda(k) = k[y_{i,j} \mid 1 \leq i \leq n, 0 \leq j < \lambda_i] / I,$$

where I is generated by nonleading coefficients of the polynomial

$$\prod_{i=1}^n (y_{i,0} + y_{i,1}t + \dots + y_{i,\lambda_i-1}t^{\lambda_i-1} + t^{\lambda_i}).$$

We denote the coefficient of the monomial t^k in the above polynomial by f_k . Thus, $I = (f_0, \dots, f_m)$. It will turn out that $A_\lambda(K_\lambda)$, B , and $C_\lambda(K_\lambda)$ are isomorphic.

Lemma 4.3. *For every field k , local k -algebras $A_\lambda(k)$ and $C_\lambda(k)$ are isomorphic.*

Proof. Consider two polynomial algebras $S_1 = k[y_{i,j} \mid 1 \leq i \leq n, 0 \leq j < \lambda_i]$ and $S_2 = k[z_1, \dots, z_{m+1}]$. Then $C_\lambda(k) = S_1/I$, where I is defined above, and $A_\lambda(k) \subset A(k) = S_2/I_S$, where I_S is the ideal generated by all polynomials symmetric with respect to z_1, \dots, z_{m+1} without constant terms. Furthermore,

$$I_S = (\sigma_1(z_1, \dots, z_{m+1}), \dots, \sigma_{m+1}(z_1, \dots, z_{m+1})),$$

where $\sigma_d(z_1, \dots, z_{m+1})$ denotes the elementary symmetric polynomial in z_1, \dots, z_{m+1} of degree $1 \leq d \leq m+1$. We define a k -algebra homomorphism $\alpha: S_1 \rightarrow S_2$ by

$$\alpha(y_{i,j}) = \sigma_{\lambda_i-j}(z_{\lambda_i+1}, \dots, z_{\lambda_i+1}). \quad (4.1)$$

Since elementary symmetric polynomials are algebraically independent, α is injective. Extending α to $S_1[t] \rightarrow S_2[t]$ by $\alpha(t) = t$, we obtain

$$\begin{aligned} \alpha \left(\prod_{i=1}^n (y_{i,0} + y_{i,1}t + \dots + y_{i,\lambda_i-1}t^{\lambda_i-1} + t^{\lambda_i}) \right) &= \\ \prod_{i=1}^n (\alpha(y_{i,0}) + \alpha(y_{i,1})t + \dots + \alpha(y_{i,\lambda_i-1})t^{\lambda_i-1} + t^{\lambda_i}) &= \\ \prod_{i=1}^{m+1} (t + z_i) = \sum_{i=0}^{m+1} \sigma_{m+1-i}(z_1, \dots, z_{m+1})t^i. \end{aligned}$$

Hence, $\alpha(f_k) = \sigma_{m+1-k}(z_1, \dots, z_{m+1})$ for $0 \leq k \leq m$, so $\alpha(I) \subset \alpha(S_1) \cap I_S$. We claim that $\alpha(I) = \alpha(S_1) \cap I_S$. Let $\alpha(p) \in \alpha(S_1) \cap I_S$, where $p \in S_1$. Since $\alpha(p)$ is λ -symmetric, Lemma 3.4 implies that there exist λ -symmetric $a_1, \dots, a_{m+1} \in S_2$ such that $\alpha(p) = \sum_{j=1}^{m+1} a_j \sigma_j(z_1, \dots, z_{m+1})$. Lemma 3.5 implies that there exist $q_1, \dots, q_{m+1} \in S_1$ such that

$a_j = \alpha(q_j)$ for all $1 \leq j \leq m+1$. Hence, $\alpha(p) = \alpha \left(\sum_{j=1}^{m+1} q_j f_{m+1-j} \right)$, so the injectivity of α implies that $p \in I$.

Since $\alpha(I) = \alpha(S_1) \cap I_S$, α defines an injective homomorphism $\beta: S_1/I \rightarrow S_2/I_S$, which gives us the embedding $\beta: C_\lambda(k) \rightarrow A(k)$. Lemma 3.5 together with (4.1) imply that the image $\beta(C_\lambda(k))$ coincides with the set of all λ -symmetric polynomials, so β gives a desired isomorphism. \square

Lemma 4.4. *There exists a surjective homomorphism $\varphi_\lambda: C_\lambda(K_\lambda) \rightarrow B$.*

Proof. We set $X_i(t) = x_i^{(0)} + x_i^{(1)}t + \dots + x_i^{(m)}t^m \in R_{\mathfrak{p}_\lambda}[t]/(t^{m+1})$. For fixed $i \leq n$, we will find $b_0, \dots, b_{m-\lambda_i} \in R_{\mathfrak{p}_\lambda}$ such that the polynomial $X_i(t)P_i(t)$, where $P_i(t) = b_0 + b_1t + \dots + b_{m-\lambda_i}t^{m-\lambda_i}$, is equal to some monic polynomial of degree λ_i modulo t^{m+1} . Then elements $b_0, \dots, b_{m-\lambda_i}$ should satisfy the following linear system

$$\begin{pmatrix} x_i^{(\lambda_i)} & \dots & x_i^{(m-1)} & x_i^{(m)} \\ \vdots & \ddots & \vdots & \vdots \\ \dots & \dots & x_i^{(\lambda_i)} & x_i^{(\lambda_i+1)} \\ \dots & \dots & x_i^{(\lambda_i-1)} & x_i^{(\lambda_i)} \end{pmatrix} \begin{pmatrix} b_{m-\lambda_i} \\ \vdots \\ b_1 \\ b_0 \end{pmatrix} = \begin{pmatrix} 0 \\ \vdots \\ 0 \\ 1 \end{pmatrix}.$$

The matrix M of this system is upper-triangular modulo ideal $\mathfrak{p}_\lambda R_{\mathfrak{p}_\lambda}$ with $x_i^{(\lambda_i)}$ on the diagonal, so there is a unique vector $(b_{m-\lambda_i}, \dots, b_1, b_0) \in R_{\mathfrak{p}_\lambda}^{m-\lambda_i+1}$ satisfying this system. Moreover, $b_0 \notin \mathfrak{p}_\lambda R_{\mathfrak{p}_\lambda}$.

We denote the product $X_i(t)P_i(t)$ reduced modulo t^{m+1} by $Q_i(t) = a_{i,0} + a_{i,1}t + \dots + a_{i,\lambda_i-1}t^{\lambda_i-1} + t^{\lambda_i}$. Since $x_i^{(0)}, \dots, x_i^{(\lambda_i-1)} \in \mathfrak{p}_\lambda R_{\mathfrak{p}_\lambda}$, elements $a_{i,0}, \dots, a_{i,\lambda_i-1}$ also lie in $\mathfrak{p}_\lambda R_{\mathfrak{p}_\lambda}$, so their images in B are nilpotents. We also note that

$$\prod_{i=1}^n Q_i(t) \equiv \left(\prod_{i=1}^n X_i(t) \right) \cdot \left(\prod_{i=1}^n P_i(t) \right) \pmod{t^{m+1}}. \quad (4.2)$$

In what follows, the image of element $a \in R_{\mathfrak{p}_\lambda}$ in B will be denoted by \bar{a} . By S we denote the k -subalgebra of B generated by $\bar{a}_0, \dots, \bar{a}_{\lambda_i-1}, \bar{x}_i^{(\lambda_i)}, \dots, \bar{x}_i^{(m)}$, and the inverse of $\bar{x}_i^{(\lambda_i)}$, we set also $\mathfrak{p} = \mathfrak{p}_\lambda B \cap S$. We claim that $\bar{x}_i^{(0)}, \dots, \bar{x}_i^{(\lambda_i-1)} \in S$. For every j , \bar{b}_j can be written as $\frac{c_j}{c}$, where $c = \det \bar{M}$ and c_j is a polynomial in $\bar{x}_i^{(\lambda_i)}, \dots, \bar{x}_i^{(m)}$ over k . Then the following matrix equality

$$\begin{pmatrix} c_0 & c_1 & \dots & c_{\lambda_i-1} \\ 0 & c_0 & \dots & c_{\lambda_i-1} \\ \vdots & \ddots & \ddots & \vdots \\ 0 & \dots & 0 & c_0 \end{pmatrix} \begin{pmatrix} \bar{x}_i^{(\lambda_i-1)} \\ \bar{x}_i^{(\lambda_i-2)} \\ \vdots \\ \bar{x}_i^{(0)} \end{pmatrix} = \begin{pmatrix} c\bar{a}_{i,\lambda_i-1} \\ c\bar{a}_{i,\lambda_i-2} \\ \vdots \\ c\bar{a}_{i,0} \end{pmatrix} \quad (4.3)$$

can be considered as a system of polynomial equations in $\bar{x}_i^{(0)}, \dots, \bar{x}_i^{(\lambda_i-1)}$ over S . We denote the matrix in the left-hand side of (4.3) by N . Vector $(0, \dots, 0) \in S^{\lambda_i}$ gives a solution of this system modulo \mathfrak{p} . The Jacobian matrix of the system (4.3) at $(0, \dots, 0)$ is equal to N modulo \mathfrak{p} . The determinant $\det N = c_0^{\lambda_i}$ at $(0, \dots, 0)$ is equal to $\left(x_i^{(\lambda_i)}\right)^{\lambda_i(m-\lambda_i)}$ modulo \mathfrak{p} , so it is an invertible element of S . Since \mathfrak{p} is a nilpotent ideal, S is complete with respect to it. Thus, Hensel's Lemma ([7, Sect. 4]) implies that there exists a unique solution of (4.3) in S^{λ_i} , which is equal to $(0, \dots, 0)$ modulo \mathfrak{p} . Repeating the same argument for algebra B and ideal $\mathfrak{p}_\lambda B$, we have that (4.3) has a unique solution in $B \supset S$. Hence, these solutions coincide, and are both equal to $(\bar{x}_i^{(0)}, \dots, \bar{x}_i^{(\lambda_i-1)})$.

So, $\bar{x}_i^{(0)}, \dots, \bar{x}_i^{(\lambda_i-1)} \in S$, and B is generated by $\{a_{i,j} \mid 1 \leq i \leq n, 0 \leq j < \lambda_i\}$ as a K_λ -algebra.

Consider $\varphi: K_\lambda[y_{i,j} \mid 1 \leq i \leq n, 0 \leq j < \lambda_i] \rightarrow B$ defined by $\varphi(y_{i,j}) = \bar{a}_{i,j}$. Since the coefficients of $\prod_i^n X_i(t)$ generate $J_m(x_1 \cdots x_n)$, equality (4.2) implies that $\text{Ker } \varphi \supset I$, so we obtain a homomorphism $\varphi_\lambda: C_\lambda(K_\lambda) \rightarrow B$. Since $a_{i,j}$ generate B over K_λ , φ_λ is surjective, so we are done. \square

Let us return to the proof of Theorem 4.1. Lemma 4.3 and Lemma 4.4 imply that for every $\lambda \in \mathbb{Z}_{\geq 0}^n$ such that $\lambda_1 + \dots + \lambda_n = m + 1$ inequality

$$\dim_{K_\lambda} (R/J_m(x_1 \cdots x_n))_{\mathfrak{p}_\lambda} \leq \dim_{K_\lambda} A_\lambda(K_\lambda) \quad (4.4)$$

holds. Moreover, the equality in (4.4) would imply that φ_λ is an isomorphism.

The left-hand side of (4.4) is the multiplicity of J_m along \mathfrak{p}_λ . Every \mathfrak{p}_λ defines an affine subspace, so has degree one. Since J_m is an ideal of codimension $m + 1$ generated by $m + 1$ polynomials, Bézout theorem ([4, Theorem 1.7.7]) implies that the sum of multiplicities of these prime components is equal to the product of degrees of generating polynomials. The latter is equal to n^{m+1} . Summing up inequality (4.4) for all λ such that $\lambda_1 + \dots + \lambda_n = m + 1$, we obtain

$$n^{m+1} \leq \sum_{\lambda} \dim A_\lambda(K_\lambda) = (\text{Lemma 3.7}) = \sum_{\lambda_1 + \dots + \lambda_n = m+1} \frac{(m+1)!}{\lambda_1! \cdots \lambda_n!} = n^{m+1}.$$

Thus, all inequalities of the form (4.4) are equalities, so all φ_λ are isomorphisms. \square

Corollary 4.5. *Let us consider the isomorphism*

$$\psi_\lambda: (R/J_m(x_1 \cdots x_n))_{\mathfrak{p}_\lambda} \rightarrow A_\lambda(K_\lambda)$$

from Theorem 4.1. For every $1 \leq i \leq n$ and $0 \leq j < \lambda_i$, the image $\psi_\lambda(x_i^{(j)})$ lies in $A_\lambda^i(K_\lambda)$.

Moreover, $\nu(\psi_\lambda(x_i^{(j)})) = \lambda_i - j$ (see Definition 3.3).

Proof. In what follows we will use the notation from the proof of Theorem 4.1. Let us fix i and j such that $1 \leq i \leq n$ and $0 \leq j < \lambda_i$. The proof of Lemma 4.4 implies that the image of $x_i^{(j)}$ in the algebra $C_\lambda(K_\lambda)$ lies in the subalgebra generated by $y_{i,0}, \dots, y_{i,\lambda_i-1}$. The construction of the isomorphism between $C_\lambda(K_\lambda)$ and $A_\lambda(K_\lambda)$ (see (4.1)) implies that the images of $y_{i,0}, \dots, y_{i,\lambda_i-1}$ lie in $A_\lambda^i(K_\lambda)$.

In order to prove the second claim we need to introduce some notation. Due to the proof of Theorem 4.1, the isomorphism ψ_λ can be factored as $\psi_\lambda = \beta \circ \gamma$, where $\beta: C_\lambda(K_\lambda) \rightarrow A_\lambda(K_\lambda)$ is the isomorphism constructed in Lemma 4.3, and

$$\gamma: B_\lambda = (R/J_m(x_1 \cdots x_n))_{\mathfrak{p}_\lambda} \rightarrow C_\lambda(K_\lambda)$$

is the inverse to the isomorphism φ_λ constructed in [Lemma 4.4](#) (its injectivity proved in [Theorem 4.1](#)). For elements $b \in B_\lambda$ and $c \in C_\lambda(K_\lambda)$, we can define function ν by $\nu(c) = \nu(\beta(c))$ and $\nu(b) = \nu(\gamma(b))$. Our goal can be reformulated as $\nu(x_i^{(j)}) \geq \lambda_i - j$. Definitions of β and γ imply that $\nu(\bar{a}_{i,j}) = \nu(y_{i,j}) = \lambda_i - j$, and for every invertible element z from $A_\lambda(K_\lambda)$, B_λ , or $C_\lambda(K_\lambda)$ we have $\nu(z) = 0$.

We will prove the second claim of the corollary by induction on j . For the case $j = 0$, we consider the last equation in [\(4.3\)](#). Since c and c_0 are invertible, we have $\nu(\bar{x}_i^{(0)}) = \nu(\bar{a}_{i,0}) = \lambda_i$. For an arbitrary j , we consider $(\lambda_i - j)$ -th equation in [\(4.3\)](#). Again, due to invertibility of c , ν of the right-hand side is $\lambda_i - j$. The value of ν of all summands on the left-hand side except $c_0 \bar{x}_i^{(j)}$ is larger than $\lambda_i - j$ due to the induction hypothesis. Since c_0 is invertible, $\nu(\bar{x}_i^{(j)}) = \lambda_i - j$. \square

5. Applications to differential algebra

Theorem 5.1. *Let K be a differential field, $(h_1, \dots, h_n) \in \mathbb{Z}_{\geq 0}^n$, and $H = h_1 + \dots + h_n$. By I_h we denote the ideal*

$$(y_1 \cdots y_n, (y_1 \cdots y_n)', \dots, (y_1 \cdots y_n)^{(h)}) \subset K\{y_1, \dots, y_n\}.$$

Then

1. The differential polynomial $y_1^{(h_1)} \cdots y_n^{(h_n)}$ does not lie in the radical of I_{H-1} .
2. The minimal d such that $(y_1^{(h_1)} \cdots y_n^{(h_n)})^d \in I_H$ is equal to $\max_{1 \leq i \leq n} (h_i + 1)(H - h_i) + 1$.

Proof. The first claim can be deduced from the result of [\[1\]](#), but we prefer to give a more straightforward argument. Let us consider a (not differential) homomorphism $\varphi: K\{y_1, \dots, y_n\} \rightarrow K$ defined by

$$\varphi(y_i^{(j)}) = \begin{cases} 0, & \text{if } j < h_i, \\ 1, & \text{otherwise.} \end{cases}$$

For every tuple $(i_1, \dots, i_n) \in \mathbb{Z}_{\geq 0}^n$ such that $i_1 + \dots + i_n = h \leq H - 1$, there exists j such that $i_j < h_j$. Hence, $\varphi(y_1^{(i_1)} \cdots y_n^{(i_n)}) = 0$, so $\varphi((y_1 \cdots y_n)^{(h)}) = 0$. On the other hand, $\varphi(y_1^{(h_1)} \cdots y_n^{(h_n)}) = 1$. Thus, the first claim is proved.

By D we denote the number $\max_{1 \leq i \leq n} (h_i + 1)(H - h_i)$. We want to formulate our problem using the language of jet ideals. Since the ideal $\tilde{I}_H = I_H \cap K[y_i^{(j)} \mid 1 \leq i \leq n, j \leq H]$ is also generated by $y_1 \cdots y_n, (y_1 \cdots y_n)', \dots, (y_1 \cdots y_n)^{(H)}$, it is sufficient to consider \tilde{I}_H . Similarly to [Remark 2.1](#), the isomorphism

$$\psi: K[y_i^{(j)} \mid 1 \leq i \leq n, j \leq H] \rightarrow R_H = K[x_i^{(j)} \mid 1 \leq i \leq n, j \leq H]$$

defined by $\psi(y_i^{(j)}) = j!x_i^{(j)}$ sends \tilde{I}_H to the H -th jet ideal of $x_1 \cdots x_n$. We denote the latter by J_H . Since $\psi\left(y_1^{(h_1)} \cdots y_n^{(h_n)}\right) = \left(\prod_{i=1}^n h_i!\right) x_1^{(h_1)} \cdots x_n^{(h_n)}$, it is sufficient to prove that the minimal d such that $\left(x_1^{(h_1)} \cdots x_n^{(h_n)}\right)^d \in J_H$ is equal to $D + 1$.

First, we prove that for every $d > D$

$$\left(x_1^{(h_1)} \cdots x_n^{(h_n)}\right)^d \in J_H.$$

Let us consider a minimal prime of J_H . Due to the discussion before [Theorem 4.1](#), it is of the form \mathfrak{p}_λ , where $\lambda = (\lambda_1, \dots, \lambda_n) \in \mathbb{Z}_{\geq 0}^n$ such that $\lambda_1 + \dots + \lambda_n = H + 1$. [Theorem 4.1](#) also implies that

$$B_\lambda = (K\{x_1, \dots, x_n\}/J_H)_{\mathfrak{p}_\lambda} \cong A_\lambda(L)$$

for some field L . It is sufficient to show that for all λ the image of $\left(x_1^{(h_1)} \cdots x_n^{(h_n)}\right)^d$ in this localization is zero. Since $\lambda_1 + \dots + \lambda_n = H + 1 > h_1 + \dots + h_n$, there exists i such that $h_i < \lambda_i$. Due to [Corollary 4.5](#), $\nu\left(\psi_\lambda\left(x_i^{(h_i)}\right)\right) = \lambda_i - h_i$, so [Proposition 3.9](#) implies, that in B_λ

$$\left(x_i^{(h_i)}\right)^{d_0} = 0, \text{ if } d_0 > \frac{\lambda_i}{\lambda_i - h_i}(H + 1 - \lambda_i).$$

Since $H - h_i \geq H + 1 - \lambda_i$ and $(\lambda_i - h_i)(h_i + 1) \geq \lambda_i$, so we obtain

$$D \geq (h_i + 1)(H - h_i) \geq \frac{\lambda_i}{\lambda_i - h_i}(H + 1 - \lambda_i).$$

Hence, $\left(x_i^{(h_i)}\right)^d = 0$ in B_λ .

Now it is sufficient to prove that $\left(x_1^{(h_1)} \cdots x_n^{(h_n)}\right)^D \notin J_H$. Without loss of generality, we can assume that $h_1 \geq h_2 \geq \dots \geq h_n$, so $D = (h_1 + 1)(H - h_1)$. Set $\lambda = (h_1 + 1, h_2, \dots, h_n)$ and

$$H_i = (h_1 + 1) + h_2 + \dots + h_{i-1} = \lambda_1 + \dots + \lambda_{i-1}$$

for all $i \leq n$. We define a homomorphism $\psi: R_H \rightarrow A_\lambda(K) \subset K[z_1, \dots, z_{H+1}]/I_S$ by

$$\psi\left(x_i^{(j)}\right) = \begin{cases} \sigma_{\lambda_i-j}\left(z_{H_i+1}, \dots, z_{H_i+1}\right), & \text{if } j < \lambda_i, \\ 1, & \text{if } j = \lambda_i, \\ 0, & \text{otherwise.} \end{cases}$$

We claim that $J_H \subset \text{Ker } \psi$. Note that coefficients of $1, t, \dots, t^H$ in the product

$$\prod_{i=1}^n \left(\psi \left(x_i^{(0)} \right) + \psi \left(x_i^{(1)} \right) t + \dots + \psi \left(x_i^{(H)} \right) t^H \right) \in A_\lambda(K)[t]$$

are exactly images of generators of J_H . Similarly to the proof of [Lemma 4.3](#), we obtain

$$\prod_{i=1}^n \left(\psi \left(x_i^{(0)} \right) + \psi \left(x_i^{(1)} \right) t + \dots + \psi \left(x_i^{(H)} \right) t^H \right) = \prod_{i=1}^{H+1} (t + z_i) = \sum_{i=1}^{H+1} \sigma_i(z_1, \dots, z_{H+1}) t^{H+1-i} + t^{H+1} = t^{H+1}.$$

Hence, $J_H \subset \text{Ker } \psi$. On the other hand, $\psi \left(x_1^{(h_1)} \dots x_n^{(h_n)} \right) = z_1 + \dots + z_{h_1+1}$. [Proposition 3.10](#) implies that $\psi \left(\left(x_1^{(h_1)} \dots x_n^{(h_n)} \right)^D \right) \neq 0$, so $\left(x_1^{(h_1)} \dots x_n^{(h_n)} \right)^D \notin J_H$. \square

Corollary 5.2. *Let n be a positive integer, and h_1, \dots, h_n be non negative integers with the sum $H := h_1 + \dots + h_n$. Then for ideals (not necessarily differential) I_1, \dots, I_n in a differential K -algebra A the following inclusion holds*

$$\left(I_1^{(h_1)} \dots I_n^{(h_n)} \right)^{\max(h_i+1)(H-h_i)+1} \subset (I_1 \dots I_n)^{(h_1+\dots+h_n)}.$$

Proof. In Section 5 of [\[15\]](#) (see Corollary 6.2 and Lemma 6.3) it was proved that if for some s

$$\left(x_1^{(h_1)} \dots x_n^{(h_n)} \right)^s \in \left(x_1 \dots x_n, (x_1 \dots x_n)', \dots, (x_1 \dots x_n)^{(h_1+\dots+h_n)} \right) \subset K\{x_1, \dots, x_n\},$$

then for the same s the inclusion

$$\left(I_1^{(h_1)} \dots I_n^{(h_n)} \right)^s \subset (I_1 \dots I_n)^{(h_1+\dots+h_n)} \quad (5.1)$$

holds for any ideals in any differential polynomial ring $K\{y_1, \dots, y_N\}$.

We claim that the same inclusion also holds in any commutative associative differential K -algebra A . Let I_1, \dots, I_n be ideals in A . Let a be any element of $\left(I_1^{(h_1)} \dots I_n^{(h_n)} \right)^s$. Then there are finitely generated ideals $J_1 \subset I_1, \dots, J_n \subset I_n$ such that $a \in \left(J_1^{(h_1)} \dots J_n^{(h_n)} \right)^s$. Let a_1, \dots, a_N be the union of the sets of generators of J_1, \dots, J_n . Consider a homomorphism of differential K -algebras $\varphi: K\{y_1, \dots, y_N\} \rightarrow A$ defined by $\varphi(y_i) = a_i$. Then [\(5.1\)](#) implies that every $b \in \varphi^{-1}(a)$ belongs to $(\varphi^{-1}(J_1) \dots \varphi^{-1}(J_n))^{(h_1+\dots+h_m)}$. Applying φ , we conclude that

$$a \in (J_1 \dots J_n)^{(h_1+\dots+h_m)} \subset (I_1 \dots I_n)^{(h_1+\dots+h_m)},$$

so the claim is proved.

Now the corollary follows directly from [Theorem 5.1](#). \square

Remark 5.3. [Theorem 5.1](#) also shows that the degree in [Corollary 5.2](#) can not be lower. As an example we can take principal ideals $I_1 = (x_1)$, $I_2 = (x_2)$, ..., $I_n = (x_n)$ in $K\{x_1, \dots, x_n\}$.

Analogously to [\[15, Corollary 6.3\]](#) one can use the inclusions

$$(J_1 \cap \dots \cap J_n)^n \subset J_1 \cdots J_n \subset J_1 \cap \dots \cap J_n$$

valid for arbitrary ideals in any K -algebra to deduce the following result from [Corollary 5.2](#).

Corollary 5.4. *Let n be a positive integer, and h_1, \dots, h_n be non negative integers with the sum $H := h_1 + \dots + h_n$. Then for ideals (not necessarily differential) I_1, \dots, I_n in a differential K -algebra A , the following inclusion holds*

$$\left(I_1^{(h_1)} \cap \dots \cap I_n^{(h_n)} \right)^{n \max(h_i+1)(H-h_i)+n} \subset (I_1 \cap \dots \cap I_n)^{(h_1+\dots+h_n)}.$$

Remark 5.5. [Theorem 5.1](#) also gives a partial solution for classical membership problem for differential ideal $[x_1 \dots x_n] \in K\{x_1, \dots, x_n\}$. More precisely, it implies that if for every $h_1, \dots, h_n \in \mathbb{Z}_{\geq 0}$ and $H = h_1 + \dots + h_n$

$$\left(x_1^{(h_1)} \cdots x_n^{(h_n)} \right)^d \in [x_1 \cdots x_n], \quad (5.2)$$

where $d = \max_{1 \leq i \leq n} (h_i + 1)(H - h_i) + 1$. Possibly, the result of such form can be also deduced using techniques of [\[6\]](#), but it is not straightforward.

Let us also note, that unlike [Theorem 5.1](#) the degree d in (5.2) is not optimal in general. In other words, taking more prolongations, it is possible to lower the degree d . For example, in the case $n = 2$ and $h_1 = h_2 = 2$ the result of Levi ([\[9, Theorem 3.2\]](#)) implies that $\left(x_1^{(2)} x_2^{(2)} \right)^5 \in [x_1 x_2]$.

Acknowledgments

This work was supported by the Austrian Science Fund FWF grant Y464-N18 and NSF grants CCF-1563942, CCF-0952591, DMS-1606334.

The author is grateful to Alexey Ovchinnikov and Thieu Vo Ngoc for useful discussions and to the referee for numerous comments and suggestions, which helped to improve the paper.

References

- [1] R. Goward Jr., K. Smith, The jet scheme of a monomial scheme, *Comm. Algebra* 35 (5) (2006) 1591–1598, <https://doi.org/10.1080/00927870500454927>.
- [2] Frederic Green, Daniel Kreymer, Emanuele Viola, Block-symmetric polynomials correlate with parity better than symmetric, *Comput. Complexity* 26 (2) (2017) 323–364, <https://doi.org/10.1007/s00037-017-0153-3>.
- [3] R. Gustavson, M. Kondratieva, A. Ovchinnikov, New effective differential nullstellensatz, *Adv. Math.* 290 (2016) 1138–1158, <https://doi.org/10.1016/j.aim.2015.12.021>.
- [4] R. Hartshorne, *Algebraic Geometry*, *Grad. Texts in Math.*, vol. 52, Springer, New York, 1977.
- [5] A.P. Hillman, D.G. Mead, K.B. O’Keefe, E.S. O’Keefe, Ideals generated by products, *Proc. Amer. Math. Soc.* 17 (3) (1966) 717–719, <http://www.jstor.org/stable/2035398>.
- [6] A.P. Hillman, D.G. Mead, K.B. O’Keefe, E.S. O’Keefe, A dynamic programming generalization of xy to n variables, *Proc. Amer. Math. Soc.* 17 (3) (1966) 720–723, <http://www.jstor.org/stable/2035399>.
- [7] F.-V. Kuhlmann, Valuation theoretic and model theoretic aspects of local uniformization, in: *Resolution of Singularities: A Research Textbook in Tribute to Oskar Zariski*, in: *Progr. Math.*, vol. 181, Birkhäuser, 1997.
- [8] O. Léon Sanchez, On the model companion of partial differential fields with an automorphism, *Israel J. Math.* 212 (1) (2016) 419–442, <https://doi.org/10.1007/s11856-016-1292-y>.
- [9] H. Levi, On the structure of differential polynomials and on their theory of ideals, *Trans. Amer. Math. Soc.* 51 (1942) 532–568, <http://www.jstor.org/stable/1990077>.
- [10] L. Manivel, *Symmetric Functions, Shubert Polynomials and Degeneracy Loci*, *SMF/AMS Texts Monogr.*, vol. 6, 2001.
- [11] D. Monk, The geometry of flag manifolds, *Proc. Lond. Math. Soc.* (3) 9 (1959) 253–268.
- [12] R. Moosa, T. Scanlon, Jet and prolongation spaces, *J. Ints. Math. Jussieu* 9 (2) (2010) 391–430, <https://doi.org/10.1017/S1474748010000010>.
- [13] T. Mora, M. Sala, On the Gröbner bases of some symmetric systems and their application to coding theory, *J. Symbolic Comput.* 35 (2) (2003) 177–194, [https://doi.org/10.1016/S0747-7171\(02\)00131-1](https://doi.org/10.1016/S0747-7171(02)00131-1).
- [14] M. Mustata, Jet schemes of locally complete intersection canonical singularities, *Invent. Math.* 145 (3) (2001) 397–424, [https://doi.org/10.1016/S0747-7171\(02\)00131-1](https://doi.org/10.1016/S0747-7171(02)00131-1).
- [15] Alexey Ovchinnikov, Gleb Pogudin, N. Thieu Vo, Upper bound for effective differential elimination, [arXiv:1610.04022](https://arxiv.org/abs/1610.04022), 2016.
- [16] C. Yeun, The multiplicity of jet schemes of a simple normal crossing divisor, *Comm. Algebra* 35 (12) (2007) 3909–3911, <https://doi.org/10.1080/00927870701512168>.