

# Bounds for D-finite Closure Properties

Manuel Kauers

- D-finite?

- D-finite
  - Solutions of linear recurrences / differential equations with polynomial coefficients.

- D-finite
  - Solutions of linear recurrences / differential equations with polynomial coefficients.
- Closure properties?

- D-finite
  - Solutions of linear recurrences / differential equations with polynomial coefficients.
- Closure properties
  - If  $f$  and  $g$  are D-finite, then also  $f + g$  and  $fg$  are D-finite (and some others, too).

- D-finite
  - Solutions of linear recurrences / differential equations with polynomial coefficients.
- Closure properties
  - If  $f$  and  $g$  are D-finite, then also  $f + g$  and  $fg$  are D-finite (and some others, too).
- Bounds?

- D-finite
  - Solutions of linear recurrences / differential equations with polynomial coefficients.
- Closure properties
  - If  $f$  and  $g$  are D-finite, then also  $f + g$  and  $fg$  are D-finite (and some others, too).
- Bounds
  - Want to know the sizes of the resulting recurrences / differential equations in advance.

- D-finite
  - Solutions of linear recurrences / differential equations with polynomial coefficients.
- Closure properties
  - If  $f$  and  $g$  are D-finite, then also  $f + g$  and  $fg$  are D-finite (and some others, too).
- Bounds
  - Want to know the sizes of the resulting recurrences / differential equations in advance.



A sequence  $(a_n)_{n=0}^{\infty}$  in a field  $K$  is called **D-finite** if there exist polynomials  $p_0, \dots, p_r$ , not all zero, such that

$$p_r(n)a_{n+r} + \dots + p_1(n)a_{n+1} + p_0(n)a_n = 0.$$

A sequence  $(a_n)_{n=0}^{\infty}$  in a field  $K$  is called **D-finite** if there exist polynomials  $p_0, \dots, p_r$ , not all zero, such that

$$p_r(n)a_{n+r} + \dots + p_1(n)a_{n+1} + p_0(n)a_n = 0.$$

**Examples:**

A sequence  $(a_n)_{n=0}^{\infty}$  in a field  $K$  is called **D-finite** if there exist polynomials  $p_0, \dots, p_r$ , not all zero, such that

$$p_r(n)a_{n+r} + \dots + p_1(n)a_{n+1} + p_0(n)a_n = 0.$$

**Examples:**

- $2^n$ :

A sequence  $(a_n)_{n=0}^{\infty}$  in a field  $K$  is called **D-finite** if there exist polynomials  $p_0, \dots, p_r$ , not all zero, such that

$$p_r(n)a_{n+r} + \dots + p_1(n)a_{n+1} + p_0(n)a_n = 0.$$

**Examples:**

- $2^n$ :  $a_{n+1} - 2a_n = 0$

A sequence  $(a_n)_{n=0}^{\infty}$  in a field  $K$  is called **D-finite** if there exist polynomials  $p_0, \dots, p_r$ , not all zero, such that

$$p_r(n)a_{n+r} + \dots + p_1(n)a_{n+1} + p_0(n)a_n = 0.$$

**Examples:**

- $2^n$ :  $a_{n+1} - 2a_n = 0$
- $n!$ :

A sequence  $(a_n)_{n=0}^{\infty}$  in a field  $K$  is called **D-finite** if there exist polynomials  $p_0, \dots, p_r$ , not all zero, such that

$$p_r(n)a_{n+r} + \dots + p_1(n)a_{n+1} + p_0(n)a_n = 0.$$

**Examples:**

- $2^n$ :  $a_{n+1} - 2a_n = 0$
- $n!$ :  $a_{n+1} - (n+1)a_n = 0$

A sequence  $(a_n)_{n=0}^{\infty}$  in a field  $K$  is called **D-finite** if there exist polynomials  $p_0, \dots, p_r$ , not all zero, such that

$$p_r(n)a_{n+r} + \dots + p_1(n)a_{n+1} + p_0(n)a_n = 0.$$

**Examples:**

- $2^n$ :  $a_{n+1} - 2a_n = 0$
- $n!$ :  $a_{n+1} - (n+1)a_n = 0$
- $\sum_{k=0}^n \frac{(-1)^k}{k!}$ :

A sequence  $(a_n)_{n=0}^{\infty}$  in a field  $K$  is called **D-finite** if there exist polynomials  $p_0, \dots, p_r$ , not all zero, such that

$$p_r(n)a_{n+r} + \dots + p_1(n)a_{n+1} + p_0(n)a_n = 0.$$

**Examples:**

- $2^n$ :  $a_{n+1} - 2a_n = 0$
- $n!$ :  $a_{n+1} - (n+1)a_n = 0$
- $\sum_{k=0}^n \frac{(-1)^k}{k!}$ :  $(n+2)a_{n+2} - (n+1)a_{n+1} - a_n = 0$



A sequence  $(a_n)_{n=0}^{\infty}$  in a field  $K$  is called **D-finite** if there exist polynomials  $p_0, \dots, p_r$ , not all zero, such that

$$p_r(n)a_{n+r} + \dots + p_1(n)a_{n+1} + p_0(n)a_n = 0.$$

**Examples:**

- $2^n$ :  $a_{n+1} - 2a_n = 0$
- $n!$ :  $a_{n+1} - (n+1)a_n = 0$
- $\sum_{k=0}^n \frac{(-1)^k}{k!}$ :  $(n+2)a_{n+2} - (n+1)a_{n+1} - a_n = 0$
- Fibonacci numbers, Harmonic numbers, Perrin numbers, diagonal Delannoy numbers, Motzkin numbers, Catalan numbers, Apéry numbers, Schröder numbers, . . .

A sequence  $(a_n)_{n=0}^{\infty}$  in a field  $K$  is called **D-finite** if there exist polynomials  $p_0, \dots, p_r$ , not all zero, such that

$$p_r(n)a_{n+r} + \dots + p_1(n)a_{n+1} + p_0(n)a_n = 0.$$

**Examples:**

- $2^n$ :  $a_{n+1} - 2a_n = 0$
- $n!$ :  $a_{n+1} - (n+1)a_n = 0$
- $\sum_{k=0}^n \frac{(-1)^k}{k!}$ :  $(n+2)a_{n+2} - (n+1)a_{n+1} - a_n = 0$
- Fibonacci numbers, Harmonic numbers, Perrin numbers, diagonal Delannoy numbers, Motzkin numbers, Catalan numbers, Apéry numbers, Schröder numbers, . . .
- Many sequences which have no name and no closed form.

A formal power series  $f(x) \in K[[x]]$  is called **D-finite** if there exist polynomials  $p_0, \dots, p_r$ , not all zero, such that

$$p_r(x)f^{(r)}(x) + \dots + p_1(x)f'(x) + p_0(x)f(x) = 0.$$

A formal power series  $f(x) \in K[[x]]$  is called **D-finite** if there exist polynomials  $p_0, \dots, p_r$ , not all zero, such that

$$p_r(x)f^{(r)}(x) + \dots + p_1(x)f'(x) + p_0(x)f(x) = 0.$$

**Examples:**

A formal power series  $f(x) \in K[[x]]$  is called **D-finite** if there exist polynomials  $p_0, \dots, p_r$ , not all zero, such that

$$p_r(x)f^{(r)}(x) + \dots + p_1(x)f'(x) + p_0(x)f(x) = 0.$$

**Examples:**

- $\exp(x)$ :

A formal power series  $f(x) \in K[[x]]$  is called **D-finite** if there exist polynomials  $p_0, \dots, p_r$ , not all zero, such that

$$p_r(x)f^{(r)}(x) + \dots + p_1(x)f'(x) + p_0(x)f(x) = 0.$$

**Examples:**

- $\exp(x)$ :  $f'(x) - f(x) = 0$
- $\log(1 - x)$ :

A formal power series  $f(x) \in K[[x]]$  is called **D-finite** if there exist polynomials  $p_0, \dots, p_r$ , not all zero, such that

$$p_r(x)f^{(r)}(x) + \dots + p_1(x)f'(x) + p_0(x)f(x) = 0.$$

**Examples:**

- $\exp(x)$ :  $f'(x) - f(x) = 0$
- $\log(1 - x)$ :  $(x - 1)f''(x) - f'(x) = 0$
- $\frac{1}{1 + \sqrt{1 - x^2}}$ :

A formal power series  $f(x) \in K[[x]]$  is called **D-finite** if there exist polynomials  $p_0, \dots, p_r$ , not all zero, such that

$$p_r(x)f^{(r)}(x) + \dots + p_1(x)f'(x) + p_0(x)f(x) = 0.$$

**Examples:**

- $\exp(x)$ :  $f'(x) - f(x) = 0$
- $\log(1 - x)$ :  $(x - 1)f''(x) - f'(x) = 0$
- $\frac{1}{1+\sqrt{1-x^2}}$ :  $(x^3 - x)f''(x) + (4x^2 - 3)f'(x) + 2xf(x) = 0$



A formal power series  $f(x) \in K[[x]]$  is called **D-finite** if there exist polynomials  $p_0, \dots, p_r$ , not all zero, such that

$$p_r(x)f^{(r)}(x) + \dots + p_1(x)f'(x) + p_0(x)f(x) = 0.$$

**Examples:**

- $\exp(x)$ :  $f'(x) - f(x) = 0$
- $\log(1 - x)$ :  $(x - 1)f''(x) - f'(x) = 0$
- $\frac{1}{1+\sqrt{1-x^2}}$ :  $(x^3 - x)f''(x) + (4x^2 - 3)f'(x) + 2xf(x) = 0$
- Bessel functions, Hankel functions, Struve functions, Airy functions, Polylogarithms, Elliptic integrals, the Error function, Kelvin functions, Mathieu functions, ...

A formal power series  $f(x) \in K[[x]]$  is called **D-finite** if there exist polynomials  $p_0, \dots, p_r$ , not all zero, such that

$$p_r(x)f^{(r)}(x) + \dots + p_1(x)f'(x) + p_0(x)f(x) = 0.$$

### Examples:

- $\exp(x)$ :  $f'(x) - f(x) = 0$
- $\log(1 - x)$ :  $(x - 1)f''(x) - f'(x) = 0$
- $\frac{1}{1+\sqrt{1-x^2}}$ :  $(x^3 - x)f''(x) + (4x^2 - 3)f'(x) + 2xf(x) = 0$
- Bessel functions, Hankel functions, Struve functions, Airy functions, Polylogarithms, Elliptic integrals, the Error function, Kelvin functions, Mathieu functions, ...
- Many functions which have no name and no closed form.

Write equations in terms of operators:

Write equations in terms of operators:

$$p_r(x)f^{(r)}(x) + \cdots + p_1(x)f'(x) + p_0(x)f(x) = 0$$

Write equations in terms of operators:

$$(p_r(x)D^r + \cdots + p_1(x)D + p_0(x)) \cdot f(x) = 0$$

Write equations in terms of operators:

$$(p_r(x)D^r + \cdots + p_1(x)D + p_0(x)) \cdot f(x) = 0$$

$$p_r(n)a_{n+r} + \cdots + p_1(n)a_{n+1} + p_0(n)a_n = 0$$

Write equations in terms of operators:

$$(p_r(x)D^r + \cdots + p_1(x)D + p_0(x)) \cdot f(x) = 0$$

$$(p_r(n)S^r + \cdots + p_1(n)S + p_0(n)) \cdot a_n = 0$$

Write equations in terms of operators:

$$(p_r(x)D^r + \cdots + p_1(x)D + p_0(x)) \cdot f(x) = 0$$

$$(p_r(n)S^r + \cdots + p_1(n)S + p_0(n)) \cdot a_n = 0$$

These operators belong to the **Ore algebras**  $C[x][D]$  and  $C[n][S]$ , respectively.



Write equations in terms of operators:

$$(p_r(x)D^r + \cdots + p_1(x)D + p_0(x)) \cdot f(x) = 0$$

$$(p_r(n)S^r + \cdots + p_1(n)S + p_0(n)) \cdot a_n = 0$$

These operators belong to the **Ore algebras**  $C[x][D]$  and  $C[n][S]$ , respectively.

Ore algebras are non-commutative rings, with a multiplication defined in such a way that  $(L_1 L_2) \cdot f = L_1 \cdot (L_2 \cdot f)$ .

Write equations in terms of operators:

$$(p_r(x)D^r + \cdots + p_1(x)D + p_0(x)) \cdot f(x) = 0$$

$$(p_r(n)S^r + \cdots + p_1(n)S + p_0(n)) \cdot a_n = 0$$

These operators belong to the **Ore algebras**  $C[x][D]$  and  $C[n][S]$ , respectively.

Ore algebras are non-commutative rings, with a multiplication defined in such a way that  $(L_1 L_2) \cdot f = L_1 \cdot (L_2 \cdot f)$ .

Examples:  $Dx = xD + 1$ ,  $Sn = (n + 1)S$ .

In general:  $A = C[x][\partial]$  with commutation rule  $\partial x = \sigma(x)\partial + \delta(x)$ ,  
with suitable maps  $\sigma, \delta: C[x] \rightarrow C[x]$ .

In general:  $A = C[x][\partial]$  with commutation rule  $\partial x = \sigma(x)\partial + \delta(x)$ , with suitable maps  $\sigma, \delta: C[x] \rightarrow C[x]$ .

If there is an action  $A \times F \rightarrow F$  of  $A$  on a  $C[x]$ -module  $F$  (a “function space”), then  $f \in F$  is called **D-finite** (w.r.t. this action), if there exists  $L \in A \setminus \{0\}$  with  $L \cdot f = 0$ .

In general:  $A = C[x][\partial]$  with commutation rule  $\partial x = \sigma(x)\partial + \delta(x)$ , with suitable maps  $\sigma, \delta: C[x] \rightarrow C[x]$ .

If there is an action  $A \times F \rightarrow F$  of  $A$  on a  $C[x]$ -module  $F$  (a “function space”), then  $f \in F$  is called **D-finite** (w.r.t. this action), if there exists  $L \in A \setminus \{0\}$  with  $L \cdot f = 0$ .

### Examples:

- Differential operators:  $\sigma = \text{id}$ ,  $\delta = \frac{d}{dx}$ ,  $F = C[[x]]$
- Recurrence operators:  $\sigma(x) = x + 1$ ,  $\delta = 0$ ,  $F = C^{\mathbb{N}}$

- D-finite
  - Solutions of linear recurrences / differential equations with polynomial coefficients.
- Closure properties
  - If  $f$  and  $g$  are D-finite, then also  $f + g$  and  $fg$  are D-finite (and some others, too).
- Bounds
  - Want to know the sizes of the resulting recurrences / differential equations in advance.

- D-finite

- Solutions of linear recurrences / differential equations with polynomial coefficients.

- Closure properties

- If  $f$  and  $g$  are D-finite, then also  $f + g$  and  $fg$  are D-finite (and some others, too).

- Bounds

- Want to know the sizes of the resulting recurrences / differential equations in advance.

Closure properties are useful for proving identities.



Closure properties are useful for proving identities.

Example:

$$\sum_{k=0}^n \frac{2k+1}{k+1} P_k^{(1,-1)}(x) - \frac{1}{1-x} (2 - P_n(x) - P_{n+1}(x)) = 0$$

Closure properties are useful for proving identities.

Example:

$$\sum_{k=0}^n \underbrace{\frac{2k+1}{k+1}}_{\text{recurrence of order 1}} \underbrace{P_k^{(1,-1)}(x)}_{\text{recurrence of order 2}} - \frac{1}{1-x} \left( 2 - \underbrace{P_n(x)}_{\text{recurrence of order 2}} - \underbrace{P_{n+1}(x)}_{\text{recurrence of order 2}} \right) = 0$$

recurrence of order 2
recurrence of order 4

recurrence of order 5
recurrence of order 3

recurrence of order 7

What makes this work?

What makes this work?

Suppose  $L \cdot f = 0$  for some operator  $L$  of order  $r$ .

What makes this work?

Suppose  $L \cdot f = 0$  for some operator  $L$  of order  $r$ .

Then  $\partial^r f$  is a  $C(x)$ -linear combination of  $f, \dots, \partial^{r-1} f$ .

What makes this work?

Suppose  $L \cdot f = 0$  for some operator  $L$  of order  $r$ .

Then  $\partial^r f$  is a  $C(x)$ -linear combination of  $f, \dots, \partial^{r-1} f$ .

Same with  $\partial^i f$  for every  $i \in \mathbb{N}$ .

What makes this work?

Suppose  $L \cdot f = 0$  for some operator  $L$  of order  $r$ .

Then  $\partial^r f$  is a  $C(x)$ -linear combination of  $f, \dots, \partial^{r-1} f$ .

Same with  $\partial^i f$  for every  $i \in \mathbb{N}$ .

So they all live in some  $C(x)$ -vector space of dimension at most  $r$ .

What makes this work?

Suppose  $L \cdot f = 0$  for some operator  $L$  of order  $r$ .

Then  $\partial^r f$  is a  $C(x)$ -linear combination of  $f, \dots, \partial^{r-1} f$ .

Same with  $\partial^i f$  for every  $i \in \mathbb{N}$ .

So they all live in some  $C(x)$ -vector space of dimension at most  $r$ .

$$\dim_{C(x)}(C(x)[\partial] \cdot f) = \dim_{C(x)}(C(x)[\partial]/\langle L \rangle) \leq r.$$



What makes this work?

Suppose  $L \cdot f = 0$  for some operator  $L$  of order  $r$ .

Then  $\partial^r f$  is a  $C(x)$ -linear combination of  $f, \dots, \partial^{r-1} f$ .

Same with  $\partial^i f$  for every  $i \in \mathbb{N}$ .

So they all live in some  $C(x)$ -vector space of dimension at most  $r$ .

$$\dim_{C(x)}(C(x)[\partial] \cdot f) = \dim_{C(x)}(C(x)[\partial]/\langle L \rangle) \leq r.$$

(Compare  $\dim_{\mathbb{Q}} \mathbb{Q}[\sqrt{2}] = \dim_{\mathbb{Q}} \mathbb{Q}[x]/\langle x^2 - 2 \rangle = 2$ .)

What makes this work?

Suppose  $L \cdot f = 0$  for some operator  $L$  of order  $r$ .

Suppose  $M \cdot g = 0$  for some operator  $M$  of order  $s$ .

What makes this work?

Suppose  $L \cdot f = 0$  for some operator  $L$  of order  $r$ .

Suppose  $M \cdot g = 0$  for some operator  $M$  of order  $s$ .

Then all  $\partial^i g$  live in some  $C(x)$ -vector space of dimension at most  $s$ .

What makes this work?

Suppose  $L \cdot \mathbf{f} = 0$  for some operator  $L$  of order  $r$ .

Suppose  $M \cdot \mathbf{g} = 0$  for some operator  $M$  of order  $s$ .

Then all  $\partial^i \mathbf{g}$  live in some  $C(x)$ -vector space of dimension at most  $s$ .

$$\dim_{C(x)}(C(x)[\partial] \cdot \mathbf{g}) = \dim_{C(x)}(C(x)[\partial]/\langle M \rangle) \leq r.$$

What makes this work?

Suppose  $L \cdot f = 0$  for some operator  $L$  of order  $r$ .

Suppose  $M \cdot g = 0$  for some operator  $M$  of order  $s$ .

$$\dim_{C(x)}(C(x)[\partial] \cdot (f + g))$$

What makes this work?

Suppose  $L \cdot f = 0$  for some operator  $L$  of order  $r$ .

Suppose  $M \cdot g = 0$  for some operator  $M$  of order  $s$ .

$$\begin{aligned} & \dim_{C(x)} (C(x)[\partial] \cdot (f + g)) \\ & \leq \dim_{C(x)} (C(x)[\partial] \cdot f + C(x)[\partial] \cdot g) \end{aligned}$$

What makes this work?

Suppose  $L \cdot f = 0$  for some operator  $L$  of order  $r$ .

Suppose  $M \cdot g = 0$  for some operator  $M$  of order  $s$ .

$$\begin{aligned} & \dim_{C(x)} (C(x)[\partial] \cdot (f + g)) \\ & \leq \dim_{C(x)} (C(x)[\partial] \cdot f + C(x)[\partial] \cdot g) \\ & \leq \dim_{C(x)} (C(x)[\partial] \cdot f) + \dim_{C(x)} (C(x)[\partial] \cdot g) \end{aligned}$$

What makes this work?

Suppose  $L \cdot f = 0$  for some operator  $L$  of order  $r$ .

Suppose  $M \cdot g = 0$  for some operator  $M$  of order  $s$ .

$$\begin{aligned} & \dim_{C(x)}(C(x)[\partial] \cdot (f + g)) \\ & \leq \dim_{C(x)}(C(x)[\partial] \cdot f + C(x)[\partial] \cdot g) \\ & \leq \dim_{C(x)}(C(x)[\partial] \cdot f) + \dim_{C(x)}(C(x)[\partial] \cdot g) \\ & \leq r + s. \end{aligned}$$



What makes this work?

Suppose  $L \cdot f = 0$  for some operator  $L$  of order  $r$ .

Suppose  $M \cdot g = 0$  for some operator  $M$  of order  $s$ .

$$\begin{aligned} & \dim_{C(x)}(C(x)[\partial] \cdot (f + g)) \\ & \leq \dim_{C(x)}(C(x)[\partial] \cdot f + C(x)[\partial] \cdot g) \\ & \leq \dim_{C(x)}(C(x)[\partial] \cdot f) + \dim_{C(x)}(C(x)[\partial] \cdot g) \\ & \leq r + s. \end{aligned}$$

Any  $r+s+1$  elements of  $C(x)[\partial] \cdot (f+g)$  are  $C(x)$ -linearly dependent.

What makes this work?

Suppose  $L \cdot \mathbf{f} = 0$  for some operator  $L$  of order  $r$ .

Suppose  $M \cdot \mathbf{g} = 0$  for some operator  $M$  of order  $s$ .

In particular, there exists a nontrivial relation

$$p_{s+r}(x) \partial^{s+r}(\mathbf{f}+\mathbf{g}) + \cdots + p_1(x) \partial(\mathbf{f}+\mathbf{g}) + p_0(x) (\mathbf{f}+\mathbf{g}) = 0.$$

What makes this work?

Suppose  $L \cdot f = 0$  for some operator  $L$  of order  $r$ .

Suppose  $M \cdot g = 0$  for some operator  $M$  of order  $s$ .

In particular, there exists a nontrivial relation

$$p_{s+r}(x) \partial^{s+r}(f+g) + \cdots + p_1(x) \partial(f+g) + p_0(x) (f+g) = 0.$$

This shows that  $f + g$  is D-finite.

What makes this work?

Suppose  $L \cdot f = 0$  for some operator  $L$  of order  $r$ .

Suppose  $M \cdot g = 0$  for some operator  $M$  of order  $s$ .

In particular, there exists a nontrivial relation

$$p_{s+r}(x) \partial^{s+r}(f+g) + \cdots + p_1(x) \partial(f+g) + p_0(x) (f+g) = 0.$$

This shows that  $f + g$  is D-finite.

A similar reasoning shows that the product  $fg$  is D-finite.

Example  $f_n = 2^n + n!$

Example  $f_n = 2^n + n!$

- $a_n = 2^n$        $a_{n+1} - 2a_n = 0$       (order 1)

Example  $f_n = 2^n + n!$

- $a_n = 2^n$        $a_{n+1} - 2a_n = 0$       (order 1)
- $b_n = n!$        $b_{n+1} - (n+1)b_n = 0$       (order 1)

**Example**  $f_n = 2^n + n!$

- $a_n = 2^n$        $a_{n+1} - 2a_n = 0$       (order 1)
- $b_n = n!$        $b_{n+1} - (n+1)b_n = 0$       (order 1)

We expect a recurrence of order two



**Example**  $f_n = 2^n + n!$

- $a_n = 2^n$        $a_{n+1} - 2a_n = 0$       (order 1)
- $b_n = n!$        $b_{n+1} - (n+1)b_n = 0$       (order 1)

We expect a recurrence of order two

$$p_2(n)f_{n+2} + p_1(n)f_{n+1} + p_0(n)f_n = 0$$

**Example**  $f_n = 2^n + n!$

- $a_n = 2^n$        $a_{n+1} - 2a_n = 0$       (order 1)

- $b_n = n!$        $b_{n+1} - (n+1)b_n = 0$       (order 1)

We expect a recurrence of order two

$$p_2(n)(a_{n+2} + b_{n+2}) + p_1(n)(a_{n+1} + b_{n+1}) + p_0(n)(a_n + b_n) = 0$$

**Example**  $f_n = 2^n + n!$

- $a_n = 2^n$        $a_{n+1} - 2a_n = 0$       (order 1)

- $b_n = n!$        $b_{n+1} - (n+1)b_n = 0$       (order 1)

We expect a recurrence of order two

$$\begin{aligned} & p_2(n)(4a_n + (n+2)(n+1)b_n) \\ & + p_1(n)(2a_n + (n+1)b_n) \\ & + p_0(n)(a_n + b_n) = 0 \end{aligned}$$

**Example**  $f_n = 2^n + n!$

- $a_n = 2^n$        $a_{n+1} - 2a_n = 0$       (order 1)

- $b_n = n!$        $b_{n+1} - (n+1)b_n = 0$       (order 1)

We expect a recurrence of order two

$$\begin{aligned} & (p_2(n) \cdot 4 + p_1(n) \cdot 2 + p_0(n))a_n \\ & + (p_2(n) \cdot (n+2)(n+1) + p_1(n) \cdot (n+1) + p_0(n))b_n = 0 \end{aligned}$$

**Example**  $f_n = 2^n + n!$

- $a_n = 2^n$        $a_{n+1} - 2a_n = 0$       (order 1)
- $b_n = n!$        $b_{n+1} - (n+1)b_n = 0$       (order 1)

We expect a recurrence of order two

$$\begin{pmatrix} 4 & 2 & 1 \\ (n+2)(n+1) & n+1 & 1 \end{pmatrix} \begin{pmatrix} p_2(n) \\ p_1(n) \\ p_0(n) \end{pmatrix} = 0$$

**Example**  $f_n = 2^n + n!$

- $a_n = 2^n$        $a_{n+1} - 2a_n = 0$       (order 1)
- $b_n = n!$        $b_{n+1} - (n+1)b_n = 0$       (order 1)

We expect a recurrence of order two

$$\begin{pmatrix} 4 & 2 & 1 \\ (n+2)(n+1) & n+1 & 1 \end{pmatrix} \begin{pmatrix} p_2(n) \\ p_1(n) \\ p_0(n) \end{pmatrix} = 0$$

More columns than rows  $\Rightarrow$  must have nontrivial solution.

**Example**  $f_n = 2^n + n!$

- $a_n = 2^n$        $a_{n+1} - 2a_n = 0$       (order 1)
- $b_n = n!$        $b_{n+1} - (n+1)b_n = 0$       (order 1)

We expect a recurrence of order two

$$\begin{pmatrix} 4 & 2 & 1 \\ (n+2)(n+1) & n+1 & 1 \end{pmatrix} \begin{pmatrix} p_2(n) \\ p_1(n) \\ p_0(n) \end{pmatrix} = 0$$

More columns than rows  $\Rightarrow$  must have nontrivial solution.

$$(p_2(n), p_1(n), p_0(n)) = (n-1, -n^2-3n+2, 2n(n+1)).$$

**Example**  $f_n = 2^n + n!$

- $a_n = 2^n$        $a_{n+1} - 2a_n = 0$       (order 1)
- $b_n = n!$        $b_{n+1} - (n+1)b_n = 0$       (order 1)

We expect a recurrence of order two

$$\begin{pmatrix} 4 & 2 & 1 \\ (n+2)(n+1) & n+1 & 1 \end{pmatrix} \begin{pmatrix} p_2(n) \\ p_1(n) \\ p_0(n) \end{pmatrix} = 0$$

More columns than rows  $\Rightarrow$  must have nontrivial solution.

$$(n-1)f_{n+2} - (n^2 + 3n - 2)f_{n+1} + 2n(n+1)f_n = 0.$$



- D-finite

- Solutions of linear recurrences / differential equations with polynomial coefficients.

- Closure properties

- If  $f$  and  $g$  are D-finite, then also  $f + g$  and  $fg$  are D-finite (and some others, too).

- Bounds

- Want to know the sizes of the resulting recurrences / differential equations in advance.

- D-finite

- Solutions of linear recurrences / differential equations with polynomial coefficients.

- Closure properties

- If  $f$  and  $g$  are D-finite, then also  $f + g$  and  $fg$  are D-finite (and some others, too).

- Bounds

- Want to know the sizes of the resulting recurrences / differential equations in advance.

A typical element of  $\mathbb{Z}[x][\partial]$  looks like this:

$$\begin{aligned} & (20x^6 + 36x^5 + 14x^4 + 8x^3 - 7x^2 - 43x - 46)\partial^4 \\ & - (24x^6 + 38x^5 - 29x^4 - 18x^3 + 14x^2 + 19x - 43)\partial^3 \\ & + (5x^6 + 36x^5 + 50x^4 + 16x^3 - 36x^2 - 41x + 43)\partial^2 \\ & - (19x^6 - x^5 - 40x^4 - 11x^3 + 9x^2 + 25x + 27)\partial \\ & - (28x^6 + 18x^5 - 40x^4 + 11x^3 + 48x^2 + 27x + 38) \end{aligned}$$

A typical element of  $\mathbb{Z}[x][\partial]$  looks like this:

$$\begin{aligned} & (20x^6 + 36x^5 + 14x^4 + 8x^3 - 7x^2 - 43x - 46)\partial^4 \\ & - (24x^6 + 38x^5 - 29x^4 - 18x^3 + 14x^2 + 19x - 43)\partial^3 \\ & + (5x^6 + 36x^5 + 50x^4 + 16x^3 - 36x^2 - 41x + 43)\partial^2 \\ & - (19x^6 - x^5 - 40x^4 - 11x^3 + 9x^2 + 25x + 27)\partial \\ & - (28x^6 + 18x^5 - 40x^4 + 11x^3 + 48x^2 + 27x + 38) \end{aligned}$$

A typical element of  $\mathbb{Z}[x][\partial]$  looks like this:

$$\begin{aligned} & (20x^6 + 36x^5 + 14x^4 + 8x^3 - 7x^2 - 43x - 46)\partial^4 \\ & - (24x^6 + 38x^5 - 29x^4 - 18x^3 + 14x^2 + 19x - 43)\partial^3 \\ & + (5x^6 + 36x^5 + 50x^4 + 16x^3 - 36x^2 - 41x + 43)\partial^2 \\ & - (19x^6 - x^5 - 40x^4 - 11x^3 + 9x^2 + 25x + 27)\partial \\ & - (28x^6 + 18x^5 - 40x^4 + 11x^3 + 48x^2 + 27x + 38) \end{aligned}$$

order

degree

A typical element of  $\mathbb{Z}[x][\partial]$  looks like this:

$$\begin{aligned}
 & (20x^6 + 36x^5 + 14x^4 + 8x^3 - 7x^2 - 43x - 46)\partial^4 \\
 & - (24x^6 + 38x^5 - 29x^4 - 18x^3 + 14x^2 + 19x - 43)\partial^3 \\
 & + (5x^6 + 36x^5 + 50x^4 + 16x^3 - 36x^2 - 41x + 43)\partial^2 \\
 & - (19x^6 - x^5 - 40x^4 - 11x^3 + 9x^2 + 25x + 27)\partial \\
 & - (28x^6 + 18x^5 - 40x^4 + 11x^3 + 48x^2 + 27x + 38)
 \end{aligned}$$

Diagram annotations:

- exp(height)**: Points to the  $x^4$  term in the third polynomial.
- order**: Points to the  $\partial^4$  term in the first polynomial.
- degree**: Points to the  $x^6$  term in the first polynomial.

A typical element of  $\mathbb{Z}[x][\partial]$  looks like this:

$$\begin{aligned}
 & (20x^6 + 36x^5 + 14x^4 + 8x^3 - 7x^2 - 43x - 46)\partial^4 \\
 & - (24x^6 + 38x^5 - 29x^4 - 18x^3 + 14x^2 + 19x - 43)\partial^3 \\
 & + (5x^6 + 36x^5 + 50x^4 + 16x^3 - 36x^2 - 41x + 43)\partial^2 \\
 & - (19x^6 - x^5 - 40x^4 - 11x^3 + 9x^2 + 25x + 27)\partial \\
 & - (28x^6 + 18x^5 - 40x^4 + 11x^3 + 48x^2 + 27x + 38)
 \end{aligned}$$

Diagram annotations:

- exp(height)**: Points to the  $x^4$  term in the third polynomial.
- order**: Points to the  $\partial^4$  term in the first polynomial.
- degree**: Points to the  $x^6$  term in the fifth polynomial.

Want: bounds on the order, degree, and height of operators obtained from executing closure properties.





**Theorem A.** If  $L_1, \dots, L_n$  are operators of degree  $\leq d$  and height  $\leq h$  for  $f_1, \dots, f_n$ , then there is an operator  $L$  for  $f_1 + f_2 + \dots + f_n$  with

$$\text{ord}(L) \leq r := \sum_{k=1}^n \text{ord}(L_k)$$

$$\text{deg}(L) \leq (n(r+1) - r)d$$

$$\begin{aligned} \text{height}(L) \leq & \text{height}(r) + \text{height}((n(r+1) - r - 1)!) \\ & + (n(r+1) - r)(\text{height}(d) + c^{(r)}(d, h)) \end{aligned}$$

**Theorem A.** If  $L_1, \dots, L_n$  are operators of degree  $\leq d$  and height  $\leq h$  for  $f_1, \dots, f_n$ , then there is an operator  $L$  for  $f_1 + f_2 + \dots + f_n$  with

$$\text{ord}(L) \leq r := \sum_{k=1}^n \text{ord}(L_k) \quad \text{— classic}$$

$$\text{deg}(L) \leq (n(r+1) - r)d$$

$$\begin{aligned} \text{height}(L) \leq & \text{height}(r) + \text{height}((n(r+1) - r - 1)!) \\ & + (n(r+1) - r)(\text{height}(d) + c^{(r)}(d, h)) \end{aligned}$$

**Theorem A.** If  $L_1, \dots, L_n$  are operators of degree  $\leq d$  and height  $\leq h$  for  $f_1, \dots, f_n$ , then there is an operator  $L$  for  $f_1 + f_2 + \dots + f_n$  with

$$\text{ord}(L) \leq r := \sum_{k=1}^n \text{ord}(L_k) \quad \text{— classic}$$

$$\text{deg}(L) \leq (n(r+1) - r)d \quad \text{— Bostan et al. ISSAC'12}$$

$$\begin{aligned} \text{height}(L) \leq & \text{height}(r) + \text{height}((n(r+1) - r - 1)!) \\ & + (n(r+1) - r)(\text{height}(d) + c^{(r)}(d, h)) \end{aligned}$$

**Theorem A.** If  $L_1, \dots, L_n$  are operators of degree  $\leq d$  and height  $\leq h$  for  $f_1, \dots, f_n$ , then there is an operator  $L$  for  $f_1 + f_2 + \dots + f_n$  with

$$\text{ord}(L) \leq r := \sum_{k=1}^n \text{ord}(L_k) \quad \text{— classic}$$

$$\text{deg}(L) \leq (n(r+1) - r)d \quad \text{— Bostan et al. ISSAC'12}$$

$$\left. \begin{aligned} \text{height}(L) &\leq \text{height}(r) + \text{height}((n(r+1) - r - 1)!) \\ &\quad + (n(r+1) - r)(\text{height}(d) + c^{(r)}(d, h)) \end{aligned} \right\} \text{new}$$

**Theorem A.** If  $L_1, \dots, L_n$  are operators of degree  $\leq d$  and height  $\leq h$  for  $f_1, \dots, f_n$ , then there is an operator  $L$  for  $f_1 + f_2 + \dots + f_n$  with

$$\text{ord}(L) \leq r := \sum_{k=1}^n \text{ord}(L_k) \quad \text{— classic}$$

$$\text{deg}(L) \leq (n(r+1) - r)d \quad \text{— Bostan et al. ISSAC'12}$$

$$\left. \begin{aligned} \text{height}(L) \leq & \text{height}(r) + \text{height}((n(r+1) - r - 1)!) \\ & + (n(r+1) - r)(\text{height}(d) + c^{(r)}(d, h)) \end{aligned} \right\} \text{new}$$

$\approx$  bound on  $\text{height}(\sigma^r(p))$  and  $\text{height}(\delta^r(p))$  for  
all  $p \in C[x]$  with  $\text{deg}(p) \leq d$  and  $\text{height}(p) \leq h$

**Theorem A.** If  $L_1, \dots, L_n$  are operators of degree  $\leq d$  and height  $\leq h$  for  $f_1, \dots, f_n$ , then there is an operator  $L$  for  $f_1 + f_2 + \dots + f_n$  with

from “more columns than rows”

$$\text{ord}(L) \leq r := \sum_{k=1}^n \text{ord}(L_k) \quad \text{— classic}$$

$$\deg(L) \leq (n(r+1) - r)d \quad \text{— Bostan et al. ISSAC'12}$$

$$\left. \begin{aligned} \text{height}(L) &\leq \text{height}(r) + \text{height}((n(r+1) - r - 1)!) \\ &\quad + (n(r+1) - r)(\text{height}(d) + c^{(r)}(d, h)) \end{aligned} \right\} \text{new}$$

**Theorem A.** If  $L_1, \dots, L_n$  are operators of degree  $\leq d$  and height  $\leq h$  for  $f_1, \dots, f_n$ , then there is an operator  $L$  for  $f_1 + f_2 + \dots + f_n$  with

from “more columns than rows”

$$\text{ord}(L) \leq r := \sum_{k=1}^n \text{ord}(L_k) \quad \text{— classic}$$

$$\text{deg}(L) \leq (n(r+1) - r)d \quad \text{— Bostan et al. ISSAC'12}$$

$$\left. \begin{aligned} \text{height}(L) \leq & \text{height}(r) + \text{height}((n(r+1) - r - 1)!) \\ & + (n(r+1) - r)(\text{height}(d) + c^{(r)}(d, h)) \end{aligned} \right\} \text{new}$$

from degrees and heights of the matrix entries

**Theorem B.** If  $L_1, \dots, L_n$  are operators for  $f_1, \dots, f_n$ , then there is an operator  $L$  for  $f_1 + f_2 + \dots + f_n$  with

$$\text{ord}(L) = r \quad \text{and} \quad \deg(L) = d$$

for every  $r, d$  with

$$r \geq \sum_{k=1}^n \text{ord}(L_k) \quad \text{and} \quad d \geq \frac{(r+1) \sum_{k=1}^n \deg(L_k) - \sum_{k=1}^n \text{ord}(L_k) \deg(L_k)}{r+1 - \sum_{k=1}^n \text{ord}(L_k)}.$$



**Theorem B.** If  $L_1, \dots, L_n$  are operators for  $f_1, \dots, f_n$ , then there is an operator  $L$  for  $f_1 + f_2 + \dots + f_n$  with

$$\text{ord}(L) = r \quad \text{and} \quad \deg(L) = d$$

for every  $r, d$  with

$$r \geq \sum_{k=1}^n \text{ord}(L_k) \quad \text{and} \quad d \geq \frac{(r+1) \sum_{k=1}^n \deg(L_k) - \sum_{k=1}^n \text{ord}(L_k) \deg(L_k)}{r+1 - \sum_{k=1}^n \text{ord}(L_k)}.$$



Let  $L_1, \dots, L_n$  be operators for  $f_1, \dots, f_n$ .

Let  $L_1, \dots, L_n$  be operators for  $f_1, \dots, f_n$ .

We now consider operators  $L$  for functions  $f$  of the form

$$(3x+8)f_1(x)^3f_2'(x) + (1+x)f_1(x)f_1''(x)f_2(x) + (2x+5)f_1'(x)^2f_2''(x).$$

Let  $L_1, \dots, L_n$  be operators for  $f_1, \dots, f_n$ .

We now consider operators  $L$  for functions  $f$  of the form

$$(3x+8)f_1(x)^3f_2'(x) + (1+x)f_1(x)f_1''(x)f_2(x) + (2x+5)f_1'(x)^2f_2''(x).$$

- polynomial expressions in  $\partial^j f_i$  with coefficients in  $C[x]$ .

Let  $L_1, \dots, L_n$  be operators for  $f_1, \dots, f_n$ .

We now consider operators  $L$  for functions  $f$  of the form

$$(3x + 8)f_1(x)^3 f_2'(x) + (1 + x)f_1(x)f_1''(x)f_2(x) + (2x + 5)f_1'(x)^2 f_2''(x).$$

- polynomial expressions in  $\partial^j f_i$  with coefficients in  $C[x]$ .
- for every  $i$ , the expressions are homogeneous in  $f_i, \partial f_i, \dots$

Let  $L_1, \dots, L_n$  be operators for  $f_1, \dots, f_n$ .

We now consider operators  $L$  for functions  $f$  of the form

$$(3x + 8)f_1(x)^3 f_2'(x) + (1 + x)f_1(x)f_1''(x)f_2(x) + (2x + 5)f_1'(x)^2 f_2''(x).$$

- polynomial expressions in  $\partial^j f_i$  with coefficients in  $C[x]$ .
- for every  $i$ , the expressions are homogeneous in  $f_i, \partial f_i, \dots$
- only  $\partial^j f_i$  with  $j < \text{ord}(L_i)$  appear (w.l.o.g.).

Let  $L_1, \dots, L_n$  be operators for  $f_1, \dots, f_n$ .

We now consider operators  $L$  for functions  $f$  of the form

$$(3x + 8)f_1(x)^3 f_2'(x) + (1 + x)f_1(x)f_1''(x)f_2(x) + (2x + 5)f_1'(x)^2 f_2''(x).$$

- polynomial expressions in  $\partial^j f_i$  with coefficients in  $C[x]$ .
- for every  $i$ , the expressions are homogeneous in  $f_i, \partial f_i, \dots$
- only  $\partial^j f_i$  with  $j < \text{ord}(L_i)$  appear (w.l.o.g.).

Let  $f = P(f_1, \partial f_1, \dots, f_2, \dots)$  be such an expression, and let  $D_i$  be the total degree of  $P$  with respect to the variables corresponding to  $f_i, \partial f_i, \dots, \partial^{\text{ord}(L_i)-1} f_i$ .



**Theorem C.** There exists an operator  $L$  for  $f$  with

$$\text{ord}(L) \leq m := \prod_{i=1}^n \binom{D_i + \text{ord}(L_i) - 1}{D_i}$$

$$\deg(L) \leq m \deg(P) + m^2 \sum_{i=1}^n D_i \deg(L_i)$$

$$\text{height}(L) \leq \text{height}(m!) + m c^{(m)}(\deg(P), \text{height}(P))$$

$$+ (m - 1) \text{height}\left(\deg(P) + m \sum_{i=1}^n D_i \deg(L_i)\right)$$

$$+ m^2 \sum_{i=1}^n \left( \text{height}(4) D_i + \text{height}(D_i + 1) + D_i \text{height}(\text{ord}(L_i) + m) \right. \\ \left. + \text{height}(\deg(L_i)) + c^{(m)}(\deg(L_i), \text{height}(L_i)) \right).$$

**Theorem D.** Furthermore, there exist operators  $L$  for  $f$  with

$$\text{ord}(L) = r \quad \text{and} \quad \deg(L) = d$$

for every  $r, d$  with

$$r \geq m := \prod_{i=1}^n \binom{D_i + \text{ord}(L_i) - 1}{D_i} \quad \text{and}$$
$$d \geq \frac{r m \sum_{i=1}^n D_i \deg(L_i) + m \deg(P)}{r + 1 - m}$$

**Theorem D.** Furthermore, there exist operators  $L$  for  $f$  with

$$\text{ord}(L) = r \quad \text{and} \quad \deg(L) = d$$

for every  $r, d$  with

$$r \geq m := \prod_{i=1}^n \binom{D_i + \text{ord}(L_i) - 1}{D_i} \quad \text{and}$$

$$d \geq \frac{r m \sum_{i=1}^n D_i \deg(L_i) + m \deg(P)}{r + 1 - m}$$

- D-finite

- Solutions of linear recurrences / differential equations with polynomial coefficients.

- Closure properties

- If  $f$  and  $g$  are D-finite, then also  $f + g$  and  $fg$  are D-finite (and some others, too).

- Bounds

- Want to know the sizes of the resulting recurrences / differential equations in advance.

- D-finite

- Solutions of linear recurrences / differential equations with polynomial coefficients.

- Closure properties

- If  $f$  and  $g$  are D-finite, then also  $f + g$  and  $fg$  are D-finite (and some others, too).

- Bounds

- Want to know the sizes of the resulting recurrences / differential equations in advance.