



Complex Variables, Theory and Application: An International Journal: An International Journal

Publication details, including instructions for authors and subscription information:
<http://www.tandfonline.com/loi/gcov19>

Differentially transcendental formal power series

Dmitry Gokhman ^a

^a Division of Mathematics and Statistics , University of Texas at San Antonio , San Antonio, TX, 78249-0664, USA
Published online: 29 May 2007.

To cite this article: Dmitry Gokhman (1996) Differentially transcendental formal power series, Complex Variables, Theory and Application: An International Journal: An International Journal, 29:1, 41-44

To link to this article: <http://dx.doi.org/10.1080/17476939608814872>

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the “Content”) contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden. Terms & Conditions of access and use can be found at <http://www.tandfonline.com/page/terms-and-conditions>

Differentially Transcendental Formal Power Series

DMITRY GOKHMAN

*Division of Mathematics and Statistics, University of Texas at San Antonio,
 San Antonio, TX 78249-0664, USA*

We prove that a formal power series in $1/x$, whose coefficients are in a field extension of \mathbf{Q} and are algebraically independent over \mathbf{Q} , is differentially transcendental (i.e. not differentially algebraic) over this field extension. This is stated without proof in [2]. This result provides a source of functions analytic at ∞ that are not differentially algebraic over \mathbf{R} . Such functions are of particular interest, because their germs belong to Hardy fields, but not to the class E of [1]—the intersection of all maximal Hardy fields.

AMS No. 34E05, 41A60, 12H05, 12J25, 26A12

Communicated: R. P. Gilbert

(Received January 3, 1994; in final form February 4, 1994)

Suppose F is a field extension of the field of rational numbers \mathbf{Q} . Let x be an indeterminate and let $u = 1/x$. Let $F[[u]] = F[[1/x]]$ denote the ring of formal power series in u with coefficients in F . Then its field of quotients $F((u))$ is a differential field with at least two possible derivations: formal differentiation with respect to x and u denoted by D_x and D_u .

DEFINITION 1 (see e.g. §VI.1 [7]) Suppose F is an extension field of K and $S \subseteq F$.

- (i) S is algebraically dependent over K if a nonzero polynomial in finitely many variables with coefficients in K is annulled by elements of S .
- (ii) S is a transcendence basis over K means that S is algebraically independent (i.e. not algebraically dependent) over K and is maximal with respect to this property (i.e. F is algebraic over $K(S)$).
- (iii) A transcendence basis has unique cardinality (Theorem VI.1.9 [7]) which is called the transcendence degree of F over K and is denoted by $\text{tr.deg}_K F$.

DEFINITION 2 (see e.g. §I.6 [8]) Suppose F is a field, K is a differential field, $F \subseteq K$, and $f \in K$. To say that f is differentially algebraic over F means that $\{f, f', f'', \dots\}$ is algebraically dependent over K , i.e. f is a root of a nonzero differential polynomial with coefficients in F .

PROPOSITION 1 (Proposition 7.4, [2]) If the set $\{a_i \in F, i = 0, 1, \dots\}$ is algebraically independent over \mathbf{Q} then

$$f = \sum_{i=0}^{\infty} a_i x^{-i} \in F\left(\left(\frac{1}{x}\right)\right)$$

is not differentially algebraic over F with respect to D_x .

Proof Suppose f is a root of a non-zero differential polynomial p over F with respect to D_x . Let $b_j \in F$ ($j = 0, 1, \dots, \bar{j}$) be the coefficients of p . Then f is differentially algebraic with respect to D_x over K , where $K = Q(\{b_j\})$, so

$$\text{tr.deg}_K K(f, D_x f, D_x^2 f, \dots) < \infty.$$

Since $\text{tr.deg}_Q K < \infty$, we have

$$\text{tr.deg}_Q K(f, D_x f, D_x^2 f, \dots) = \text{tr.deg}_Q K + \text{tr.deg}_K K(f, D_x f, D_x^2 f, \dots) < \infty.$$

Therefore

$$\begin{aligned} \text{tr.deg}_Q Q(f, D_x f, D_x^2 f, \dots) &= \text{tr.deg}_Q K(f, D_x f, D_x^2 f, \dots) \\ &\quad - \text{tr.deg}_{Q(f, D_x f, D_x^2 f, \dots)} K(f, D - x f, D_x^2 f, \dots) < \infty. \end{aligned}$$

Note that

$$D_x f = D_u f \left(-\frac{1}{x^2} \right) = D_u f(-u^2),$$

$$D_x^2 f = (D_u^2 f(-u^2) + D_u f(-2u))(-u^2)$$

and so on, so $Q(f, D_x f, D_x^2 f, \dots) = Q(u, f, D_u f, D_u^2 f, \dots)$.

Now since $\text{tr.deg}_Q Q(u) = 1$ and $\text{tr.deg}_Q Q(f, D_x f, D_x^2 f, \dots) < \infty$, we have

$$\text{tr.deg}_Q Q(u, f, D_u f, D_u^2 f, \dots) < \infty \quad \text{and} \quad \text{tr.deg}_Q Q(f, D_u f, D_u^2 f, \dots) < \infty.$$

Therefore, f satisfies some non-zero differential polynomial q over Q with respect to D_u . Since the field Q is infinite, there exist $t_k \in Q$ ($k = 0, 1, \dots, k$) such that $q(t_0, t_1, \dots, t_{\bar{k}}) \neq 0$. Let

$$g(u) = \sum_{k=0}^{\infty} \frac{t_k}{k!} u^k.$$

Then $q(g, D_u g, \dots) \neq 0$, because it is non-zero when evaluated at $u = 0$ (it equals $q(\{t_k\})$).

If h is a formal sum $\sum_{i=0}^{\infty} y_i u^i$ with y_i indeterminate, then $q(h, D_u h, \dots)$ is a power series in u , whose coefficients are polynomials in y_i over \mathbf{Z} . Since evaluating y_i at $(t_0, t_1, \dots, t_{\bar{k}}, 0, 0, \dots)$ produces a non-zero answer, one of the above coefficients is a non-zero polynomial in y_i over Q . Let r denote this polynomial. Evaluating y_i at a_i gives $q(f, D_u f, \dots) = 0$, so, in particular, $r(a_i) = 0$, which contradicts the assumption that a_i are algebraically independent over Q . ■

The value of this result lies in providing a class of functions analytic at ∞ that are differentially transcendental over \mathbf{R} . The germs of such functions necessarily belong to Hardy fields (see the definition below), but not to Boshernitzan's class $E[1]$ —the intersection of all maximal Hardy fields.¹

¹ E is an extension of Hardy's class L of logarithmico-exponential functions [5] and is the maximal scale for functions in Hardy fields (functions of *regular growth*).

DEFINITION 3 (see [4]) A differential field of continuous germs² of real functions at $+\infty$, where the derivation is ordinary differentiation, is called a Hardy field.

COROLLARY 1 Suppose f is a real function that is analytic at ∞ and the set of coefficients of its Taylor series at ∞ is algebraically independent over \mathbf{Q} . Then

- (i) f is differentially transcendental over \mathbf{R} ,
- (ii) $\mathbf{R}(f, f' \dots)$ is a Hardy field,
- (iii) $f \notin E$, where E is the intersection of all maximal Hardy fields.

Proof (i) is a special case of Proposition 1 with $F = \mathbf{R}$ and the series actually convergent. (ii) is a consequence of the fact that the zeros of an analytic function are necessarily isolated and, thus, cannot have ∞ as an accumulation point. This means that every nonzero element of $\mathbf{R}[f, f' \dots]$, i.e. a nonzero differential polynomial of f , is invertible in a (punctured) neighborhood of ∞ (see Theorem 7.1[1]). (iii) is a consequence of the fact that E is a differentially algebraic extension of \mathbf{R} (Theorem 14.4 [2]). ■

The few other known examples of classes of functions satisfying the conditions of Corollary 1 include

- (i) Euler's Γ -function, which is not differentially algebraic over \mathbf{R} by Hölder's theorem [6] and generates a Hardy field [9];
- (ii) Functions represented by a Dirichlet series

$$\sum_{n=1}^{\infty} \frac{a_n}{n^x},$$

where $a_n \in \mathbf{R}$ are subpolynomial in n and the set of all prime divisors of n in the support of a_n ³ is infinite, e.g. the Riemann ζ -function on a positive half-line [9];

- (iii) The function

$$\sum_{n=1}^{\infty} \frac{1}{e_n(x)},$$

where $e_1(x) = e^x$ and $e_n(x) = e^{e_{n-1}(x)}$ for $n > 1$ [9];

- (iv) Certain fractional iterates of e^x [3];
- (v) Certain ultimately⁴ \mathcal{C}^∞ transexponential solutions of two difference equations: $f(x+1) = e^{f(x)}$ and $f(x+1) = e^{f(x)} - 1$ [3].

References

- [1] M. Boshernitzan, An extension of Hardy's class L of "Orders of Infinity", *J. d'Analyse Math.* **39** (1981), 235–255.
- [2] M. Boshernitzan, New "Orders of Infinity", *J. d'Analyse Math.* **41** (1982), 130–167.

²A *germ* is an equivalence class of functions, where two functions are equivalent, exactly when they agree in a neighborhood of the point of interest, in our case $+\infty$.

³The *support* of a_n is the set of all n such that $a_n \neq 0$.

⁴A property is said to hold *ultimately*, exactly when it holds in a neighborhood of $+\infty$.

- [3] M. Boshernitzan, Hardy fields and existence of transexponential functions, *Aequationes Mathematicae* **30**, 2–3 (1986), 258–280.
- [4] N. Bourbaki, *Fonctions d'une variable réelle*, Hermann, Paris, second edition, 1961. (Ch. V, Étude Locale des Fonctions).
- [5] G. H. Hardy, *Orders of infinity, the 'Infinitärcalcul' of Paul du Bois-Reymond*, Cambridge University Press, 1910.
- [6] O. Hölder, Über die Eigenschaft der Γ -Funktion, keiner algebraischen Differentialgleichung zu genügen, *Math. Ann.* **28** (1887), 1–13.
- [7] T. Hungerford, *Algebra*, Springer-Verlag, 1974.
- [8] E. R. Kolchin, *Differential algebra and algebraic groups*, Academic Press, 1973.
- [9] M. Rosenlicht, The rank of a Hardy field, *Trans. AMS* **280** (1983), 659–671.