

# Resolving finite indeterminacy

## A definitive constructive universal prime ideal theorem

Peter Schuster

Daniel Wessel

Dipartimento di Informatica  
Università degli Studi di Verona  
Verona, Italy  
peter.schuster@univr.it  
daniel.wessel@univr.it

### Abstract

Dynamical methods were designed to eliminate the ideal objects abstract algebra abounds with. Typically granted by an incarnation of Zorn's Lemma, those ideal objects often serve for proving the semantic conservation of additional non-deterministic sequents, that is, with finite but not necessarily singleton succedents. Eliminating ideal objects dynamically was possible also because (finitary) coherent or geometric logic predominates in that area: the use of a non-deterministic axiom can be captured by a finite branching of the proof tree.

Incidentally, a paradigmatic case has widely been neglected in dynamical algebra: Krull's Lemma for prime ideals. Digging deeper just about that case, which we have dealt with only recently (with Yengui), has now brought us to unearth the general phenomenon underlying dynamical algebra: Given a claim of computational nature that usually is proved by the semantic conservation of certain additional non-deterministic axioms, there is a finite labelled tree belonging to a suitable inductively generated class which tree encodes the desired computation. Our characterisation works in the fairly universal setting of a consequence relation enriched with non-deterministic axioms; uniformises many of the known instances of the dynamical method; generalises the proof-theoretic conservation criterion we have offered before (with Rinaldi); and last but not least links the syntactical with the semantic approach: every ideal object used for the customary proof of a concrete claim can be approximated by one of the corresponding tree's branches.

**CCS Concepts:** • Theory of computation → Constructive mathematics; Proof theory.

**Keywords:** dynamical proof, non-deterministic axiom, geometric logic, proof-theoretic conservation, finite tree, computational content, inductive generation, Krull's Lemma, prime ideals

### ACM Reference Format:

Peter Schuster and Daniel Wessel. 2020. Resolving finite indeterminacy: A definitive constructive universal prime ideal theorem. In *Proceedings of the 35th Annual ACM/IEEE Symposium on Logic in Computer Science (LICS '20)*, July 8–11, 2020, Saarbrücken, Germany. ACM, New York, NY, USA, 11 pages. <https://doi.org/10.1145/3373718.3394777>

## 1 Introduction

Abstract algebra abounds with ideal objects and the invocations of transfinite methods, typically Zorn's Lemma [114], that grant those object's existence. Put under logical scrutiny, ideal objects often serve for proving the semantic conservation of additional non-deterministic sequents, that is, with finite but not necessarily singleton succedents.

By design, dynamical methods in algebra [34, 65, 113] allow to eliminate the ideal objects upon shifting focus from semantic model extension principles to their corresponding syntactic conservation theorems. This move in line with Hilbert's programme has shaped modern constructive algebra and has seen tremendous success, not least because (finitary) coherent or geometric logic [25, 58, 59, 68, 71, 72, 82, 112] predominates in that area: the use of a non-deterministic axiom can be captured by a finite branching of the proof tree [34]. Coherent theories, on the other hand, lend themselves to automated theorem proving [10, 11, 35, 42, 50, 105].

A paradigmatic case, which to a certain extent has been neglected in dynamical algebra proper, is Krull's Lemma for prime ideals. A particular form of this asserts that a multiplicative subset of a commutative ring contains the zero element if and only if the set at hand meets every prime ideal. Prompted by certain aspects in the novel treatment of valuative dimension [60], Krull's Lemma has seen a constructive treatment only recently [99]. The latter, however, has brought us to unearth the underlying general phenomenon

---

Permission to make digital or hard copies of all or part of this work for personal or classroom use is granted without fee provided that copies are not made or distributed for profit or commercial advantage and that copies bear this notice and the full citation on the first page. Copyrights for components of this work owned by others than ACM must be honored. Abstracting with credit is permitted. To copy otherwise, to republish, to post on servers or to redistribute to lists, requires prior specific permission and/or a fee. Request permissions from [permissions@acm.org](mailto:permissions@acm.org).

LICS '20, July 8–11, 2020, Saarbrücken, Germany

© 2020 Association for Computing Machinery.

ACM ISBN 978-1-4503-7104-9/20/07...\$15.00

<https://doi.org/10.1145/3373718.3394777>

in the present paper: Given a claim of computational nature that usually is proved by the semantic conservation of certain additional non-deterministic axioms, there is a finite labelled tree belonging to a suitable inductively generated class which tree encodes the desired computation.

Our characterisation works in the fairly universal setting of consequence relations, a cornerstone of universal logic and algebra with a long and rich history that can be traced back to Hertz [46–48, 62] and Tarski [12, 107], and, in the guise of covering relations, has played an important role in the development of formal topology [20, 21, 69, 70, 91–93].

Consequence relations serve here to capture the basic structures—ideals of commutative rings, propositional theories, and partial orders—on top of which we consider certain non-deterministic axioms that describe “ideal” refinements of those structures: prime ideals, complete theories, and linear order extensions.

A decisive aspect of our approach is the notion of a regular set for certain non-deterministic axioms over a fixed consequence relation. Abstracted from the multiplicative subsets occurring in Krull’s Lemma, regular sets turn out to be the fundamental ingredient of our Universal Prime Ideal Theorem (UPIT, Proposition 3.2). They allow to calibrate precisely UPIT’s gearing and account for its constructive version (Proposition 4.6).

In this manner we uniformise many of the known instances of the dynamical method [34, 65, 113]. We further generalise (Proposition 6.2) the proof-theoretic conservation criterion we have offered before [87, 88], using Scott-style entailment relations [100–102], to unify the many phenomena present in the literature, e.g. [19, 31, 64, 67, 75].

Last but not least, we thus link the syntactical with the semantic approach: every ideal object used for the traditional proof of a concrete claim can be approximated by one of the corresponding tree’s branches (Proposition 4.2).

It is worth emphasizing at this point that, as compared to (propositional) dynamical algebra [27], the prime ideal theorem we offer is not only constructive but also definitive and universal. We succeed in unearthing the one common pattern of how the related trees are to be grown, by passing to the logical setting of consequence. Our approach, moreover, is ready for use in customary mathematical practice without any need to adapt first the axioms, which is not untypical for dynamical algebra. Last but not least, we identify regularity as both sufficient and necessary for the prime ideal theorem under consideration.

## Structure of this paper

In Section 2 we discuss consequence relations, non-deterministic axioms, as well as regular subsets, which relate to both the former concepts. In Section 3 we present UPIT, a straightforward consequence of Zorn’s Lemma but equivalent, as will be seen later on, to the Prime Ideal Theorem (for commutative rings, say). In Section 4 we introduce certain

inductively defined classes of finite trees which then help us to provide the constructive counterpart (CUPIT) of the Universal Prime Ideal Theorem. Three quite different applications will be discussed in Section 5, putting emphasis on the universality of our approach: a constructive version of Krull’s Lemma for commutative rings, Glivenko’s theorem for propositional logic, as well as an order extension principle will all follow immediately from CUPIT. Connections with existing work on multi-conclusion entailment relations will be discussed in the final Section 6.

## On method and foundations

Unless specified otherwise, we work in a suitable fragment of Aczel’s *Constructive Zermelo–Fraenkel Set Theory* (**CZF**) [2–6] based on intuitionistic first-order predicate logic. When we occasionally need to invoke a fragment of the principle of Excluded Middle or even a form of the Axiom of Choice (AC), and thus go beyond **CZF**, we simply switch to **ZF** and **ZFC**, respectively, and indicate this accordingly.

By a *finite* set we understand a set that can be written as  $\{a_1, \dots, a_n\}$  for some  $n \geq 0$ . Given any set  $S$ , let  $\text{Pow}(S)$  (respectively,  $\text{Fin}(S)$ ) consist of the (finite) subsets of  $S$ . We refer to [87, 88] for further provisos to carry over to this note.<sup>1</sup>

From formal topology [92] we borrow the *overlap* symbol: the notation  $U \nabla V$  is to say that the sets  $U$  and  $V$  have an element in common.

## 2 Key notions

### 2.1 Consequence relations

By a *consequence relation* or a *single-conclusion entailment relation* we understand a relation

$$\triangleright \subseteq \text{Fin}(S) \times S$$

which is *reflexive*, *monotone* and *transitive* in the following sense:

$$\frac{U \ni a}{U \triangleright a} \text{ (R)} \quad \frac{U \triangleright a}{U, V \triangleright a} \text{ (M)} \quad \frac{U \triangleright b \quad U, b \triangleright a}{U \triangleright a} \text{ (T)}$$

where as usual  $U, V \equiv U \cup V$  and  $U, b \equiv U \cup \{b\}$ . We also sometimes write  $a_1, \dots, a_n$  in place of  $\{a_1, \dots, a_n\}$  even if  $n = 0$ .

It is of course well-known that every consequence relation  $\triangleright$  gives way to an algebraic closure operator

$$\langle - \rangle : \text{Pow}(S) \rightarrow \text{Pow}(S)$$

defined by

$$a \in \langle T \rangle \equiv (\exists U \in \text{Fin}(T)) U \triangleright a.$$

<sup>1</sup> For example, we deviate from the terminology prevalent in constructive mathematics and set theory [5, 6, 13, 14, 65, 66]: to reserve the term ‘finite’ to sets which are in *bijection* with  $\{1, \dots, n\}$  for a necessarily unique  $n \geq 0$ . Those exactly are the sets which are finite in our sense and are *discrete* too, i.e. have decidable equality [66].

Conversely, given  $\langle - \rangle$ , by stipulating

$$U \triangleright a \equiv a \in \langle U \rangle$$

we gain back a consequence relation from an algebraic closure operator.

The *ideals* of a consequence relation  $\triangleright$  are the subsets  $\alpha$  of  $S$  which are closed with respect to the corresponding closure operator, which is to say that  $\alpha = \langle \alpha \rangle$ . These are precisely the subsets  $\alpha$  of  $S$  such that if  $\alpha \supseteq U$  and  $U \triangleright a$ , then  $a \in \alpha$ . We say that  $\alpha$  is *finitely generated* if  $\alpha = \langle U \rangle$  for some  $U \in \text{Fin}(S)$ .

## 2.2 Non-deterministic axioms

By a *non-deterministic axiom*<sup>2</sup> on  $S$  we understand a pair  $(A, B) \in \text{Fin}(S) \times \text{Fin}(S)$ , which we often put in turnstile notation:

$$A \vdash B.$$

A subset  $\mathfrak{p}$  of  $S$  is *closed* for  $(A, B)$  if  $A \subseteq \mathfrak{p}$  implies  $\mathfrak{p} \not\vdash B$ .

Let  $\mathcal{E}$  be a set of non-deterministic axioms. An ideal of  $\triangleright$  that is closed for every axiom of  $\mathcal{E}$  will be called a *prime ideal*.<sup>3</sup> We denote with

$$\text{Spec}(\mathcal{E})$$

the class of prime ideals of  $\mathcal{E}$ . Given an ideal  $\alpha$  of  $\triangleright$ , let

$$\text{Spec}(\mathcal{E})/\alpha = \{ \mathfrak{p} \in \text{Spec}(\mathcal{E}) \mid \mathfrak{p} \supseteq \alpha \}.$$

## 2.3 Regular subsets

**Convention.** From now on, and throughout the following Sections 3 and 4, let  $S$  be a set with consequence relation  $\triangleright$ , and let  $\mathcal{E}$  be a set of non-deterministic axioms on  $S$ .

We say that a subset  $R$  of  $S$  is *regular* if, for all  $U \in \text{Fin}(S)$  and  $(A, B) \in \mathcal{E}$ ,

$$\frac{(\forall b \in B) \langle U, b \rangle \not\vdash R}{\langle U, A \rangle \not\vdash R}$$

An element  $r$  of  $S$  is said to be *regular* if so is  $\{r\}$ . Hence  $r$  is regular precisely when, for all  $U \in \text{Fin}(S)$  and  $(A, B) \in \mathcal{E}$ ,

$$\frac{(\forall b \in B) U, b \triangleright r}{U, A \triangleright r}$$

Regularity of an element  $r$  of  $S$  thus means to require *disjunction elimination* [87, 88]

$$\frac{U, b_1 \triangleright r \quad \dots \quad U, b_\ell \triangleright r}{U, a_1, \dots, a_k \triangleright r}$$

for the succedent of every  $a_1, \dots, a_k \vdash b_1, \dots, b_\ell$  in  $\mathcal{E}$ .

<sup>2</sup>Our terminology borrows from van den Berg's principle of *non-deterministic inductive definitions* [108], variants of which have recently come to play a role in constructive reverse mathematics [49, 56].

<sup>3</sup>We say “prime ideal” to stress that variants of the prime ideal theorem (e.g., for commutative rings, distributive lattices, Boolean algebras) form the ground for our abstract version (Proposition 3.2).

## 3 A universal prime ideal theorem

The following is an abstraction of the usual proof of Krull's Lemma [61] and related prime ideal principles [86].

**Lemma 3.1 (ZFC).** *Let  $R \subseteq S$  be regular and let  $\alpha$  be an ideal. If  $R \cap \alpha = \emptyset$ , then there is a prime ideal  $\mathfrak{p} \supseteq \alpha$  such that  $R \cap \mathfrak{p} = \emptyset$ .*

*Proof.* Zorn's Lemma yields an ideal  $\mathfrak{p}$  over  $\alpha$  which is maximal with respect to the property that  $R \cap \mathfrak{p} = \emptyset$ . Every such  $\mathfrak{p}$  is a prime ideal! To see this, let  $(A, B) \in \mathcal{E}$  be such that  $A \subseteq \mathfrak{p}$  yet  $\mathfrak{p} \cap B = \emptyset$ . Maximality implies that  $\langle \mathfrak{p}, b \rangle \not\vdash R$  for every  $b \in B$ . Since  $R$  is regular, it follows that  $\mathfrak{p} = \langle \mathfrak{p}, A \rangle \not\vdash R$ , a contradiction.  $\square$

Notice that the proof of Lemma 3.1 establishes that if  $R$  is regular, then every ideal which is maximal among those avoiding  $R$  is prime. It is necessary for this that  $R$  be regular.

Here is our semantic classical *Universal Prime Ideal Theorem* (UPIT):

**Proposition 3.2 (ZFC).** *Let  $R \subseteq S$ . The following are equivalent.*

1.  $R$  is regular.
2. For every (finitely generated) ideal  $\alpha$ , the following are equivalent:
  - a.  $R \not\vdash \alpha$ .
  - b.  $(\forall \mathfrak{p} \in \text{Spec}(\mathcal{E})/\alpha) R \not\vdash \mathfrak{p}$ .

*Proof.* Suppose that  $R$  is regular. For each ideal  $\alpha$ , every element witnessing item 2.a witnesses item 2.b just as well. The reverse implication is the contrapositive of Lemma 3.1.

Conversely, suppose that for every finitely generated ideal the equivalence of item 2 holds, let  $U \in \text{Fin}(S)$  and  $(A, B) \in \mathcal{E}$  such that  $\langle U, b \rangle \not\vdash R$  for every  $b \in B$ . To show that  $\langle U, A \rangle \not\vdash R$  it now suffices to check that every prime ideal over  $\langle U, A \rangle$  meets  $R$ . In fact, if  $\mathfrak{p}$  is prime and  $\mathfrak{p} \supseteq \langle U, A \rangle \supseteq A$ , then there is  $b \in \mathfrak{p} \cap B$ , for which  $\langle U, b \rangle \not\vdash R$  and thus  $\mathfrak{p} \not\vdash R$ .  $\square$

UPIT is a weak form of the Axiom of Choice, equivalent to the prime ideal theorem for distributive lattices (cf. Proposition 6.8). There is no way around excluded middle to prove Proposition 3.2, see Proposition 6.6 below.

**Corollary 3.3 (ZFC).** *The following are equivalent.*

1. Every element of  $S$  is regular.
2. For every (finitely generated) ideal  $\alpha$ ,

$$\alpha = \bigcap \text{Spec}(\mathcal{E})/\alpha.$$

## 4 Trees for prime ideals

Given an ideal  $\alpha$ , we consider next a certain collection  $T_\alpha$  of finite labelled trees, generated in such a manner that the root of every  $t \in T_\alpha$  be labelled with a finite subset  $U$  of  $\alpha$ , and the nodes be labelled with elements of  $S$ . The latter will be determined successively by consequences of  $U$  along the additional axioms of  $\mathcal{E}$ .

Given a path  $\pi$  of such a tree  $t \in T_a$ , we write

$$\pi \triangleright r$$

to say that  $r$  is a consequence of the set labelling the root of  $t$  together with the labels occurring at the nodes of  $\pi$ . Set

$$\langle \pi \rangle = \{ r \in S \mid \pi \triangleright r \}.$$


Note that we understand paths to lead from the root of a tree to one of its leaves.

**Definition 4.1.** Let  $a$  be an ideal. We generate  $T_a$  inductively according to the following rules:

1. For every  $U \in \text{Fin}(a)$ , the trivial tree (i.e., the root-only tree) labelled with  $U$  belongs to  $T_a$ .
2. If  $(A, B) \in \mathcal{E}$  and if  $t \in T_a$  has a path  $\pi$  such that  $\langle \pi \rangle \supseteq A$ , i.e.,  $\pi \triangleright a$  for every  $a \in A$ , then add, for every  $b \in B$ , a child labelled with  $b$  at the leaf of  $\pi$ .

This is a so-called *generalised inductive definition* [1, 81, 103].

For instance, if  $(A, \{b_1, \dots, b_\ell\}) \in \mathcal{E}$  and  $U \in \text{Fin}(S)$ , then the following tree belongs to  $T_{(U,A)}$ :



By a slight abuse of notation, we say that a tree  $t \in T_a$  is *trivial* if it results from an application of the base rule in Definition 4.1 only. The trivial trees in  $T_a$  thus correspond with the elements of  $\text{Fin}(a)$ .

It is instructive to think of the given ideal  $a$  as a set of initial data, of which just a finite amount  $U$  be used for computation; with this we label the root. The paths of a tree  $t \in T_a$  then represent the possible courses of a computation as if the ideal  $a$  were prime.

**Proposition 4.2.** Let  $a$  be an ideal, and  $t \in T_a$  a tree. For every prime ideal  $p \supseteq a$  there is a path  $\pi$  through  $t$  such that  $p \supseteq \langle \pi \rangle$ .

*Proof.* The path can be constructed by induction as follows. To begin with,  $p$  contains the finite subset of  $a$  that labels the root of  $t$ . Now suppose that the path has been constructed up to a node  $v$  at which  $t$  branches with respect to  $(A, B) \in \mathcal{E}$ . By induction,  $p \supseteq A$  and thus  $p \ni b$  for some  $b \in B$ . We then add to the path the child of  $v$  labelled by this  $b$ .  $\square$

The paths of every  $t \in T_a$  may thus be considered finite approximations of the prime ideals that contain  $a$ .

**Definition 4.3.** Let  $a$  be an ideal. We say that a tree  $t \in T_a$  *terminates* in a subset  $R \subseteq S$ , in short

$$t \downarrow R,$$

if for every path  $\pi$  of  $t$  there is  $r \in R$  such that  $\pi \triangleright r$ , that is,  $\langle \pi \rangle \cap R \neq \emptyset$ . We say that a tree  $t \in T_a$  *terminates* in an element  $r$  of  $S$  if  $t$  terminates in the singleton set  $\{r\}$ .

**Example 4.4.**  $R \not\Downarrow a$  if and only if there is a trivial tree in  $T_a$  terminating in  $R$ .

The main result of this paper, Proposition 4.6 below, boils down to the following key observation.

**Lemma 4.5.** Let  $R$  be a regular subset of  $S$  and let  $a$  be an ideal. If some  $t \in T_a$  terminates in  $R$ , then  $R \not\Downarrow a$ .

*Proof.* By induction on the construction of  $t \in T_a$ . The base case is trivial. Consider next the case in which a tree in  $T_a$  has been extended at the leaf of one of its paths  $\pi$  with children labelled with  $b \in B$ , where  $(A, B) \in \mathcal{E}$  and  $\langle \pi \rangle \supseteq A$ . Suppose then that for every  $b \in B$  there is  $r \in R$  such that  $\pi, b \triangleright r$ . Regularity implies that there is  $r_0 \in R$  such that  $\pi, A \triangleright r_0$ . Since  $\langle \pi \rangle \supseteq A$ , it follows that  $\pi \triangleright r_0$ , whence the induction hypothesis applies.  $\square$

Here is our *Constructive Universal Prime Ideal Theorem* (CUPIT), the constructive counterpart of Proposition 3.2.

**Proposition 4.6.** Let  $R \subseteq S$ . The following are equivalent.

1.  $R$  is regular.
2. For every (finitely generated) ideal  $a$ , the following are equivalent:
  - a.  $R \not\Downarrow a$ .
  - b. There is a tree  $t \in T_a$  which terminates in  $R$ .

*Proof.* Suppose that  $R$  is regular. If  $a \in R \cap a$ , then the trivial tree, labelled with  $a$ , terminates in  $R$ ; conversely, if  $t \in T_a$  terminates in  $R$ , then  $R \not\Downarrow a$  by Lemma 4.5.

As regards the converse, to show that  $R$  is regular let  $U \in \text{Fin}(S)$  and  $(A, B) \in \mathcal{E}$  such that  $\langle U, b \rangle \not\Downarrow R$  for every  $b \in B$ . To see that  $\langle U, A \rangle \not\Downarrow R$ , by item 2 it is enough to observe that the tree displayed in (1) terminates in  $R$ .  $\square$

CUPIT is fairly versatile, as will be emphasised by means of three applications in Section 5.

Here is the constructive counterpart of Corollary 3.3.

**Corollary 4.7.** The following are equivalent.

1. Every element of  $S$  is regular.
2. For every (finitely generated) ideal  $a$ ,
$$a = \{ r \in S \mid (\exists t \in T_a) t \downarrow r \}.$$

Corollary 4.7 corresponds to a certain conservation result of [87, 88] as will briefly be outlined in Section 6.

## 5 Applications

### 5.1 Krull's Lemma

Our first case study concerns prime ideals of commutative rings. These have already been considered from a similar angle [99], into which we have recently been led by certain aspects of the novel treatment of valuative dimension [60]. Our concept of regular subset now allows us to go beyond. For an algorithmic approach via proof mining to the existence of ideal objects in commutative algebra we refer to [80].



Let  $\mathbf{A}$  be a commutative ring with 1. On  $S = \mathbf{A}$  we consider the entailment relation of *radical ideal*:

$$a_1, \dots, a_k \triangleright a \equiv (\exists n \in \mathbb{N}) a^n \in \sum_{i=1}^k \mathbf{A}a_i.$$

Note that an ideal for  $\triangleright$  is nothing but a radical ideal of  $\mathbf{A}$ . On top of  $\triangleright$  we consider the non-deterministic axiom of *prime ideal*, i.e., we let  $\mathcal{E}$  consist of all the instances, with  $a, b \in \mathbf{A}$ , of

$$ab \vdash a, b.$$

We say that a subset  $M$  of  $\mathbf{A}$  is *weakly multiplicative* if for all  $a, b \in M$  there is  $x \in M$  such that  $ab \in \text{Fil}(x)$ , where  $\text{Fil}(x)$  denotes the principal filter generated by  $x$ . In other words,  $M$  is weakly multiplicative if for every pair of elements  $a, b \in M$  there is  $x \in M$  along with  $n \in \mathbb{N}$  and  $c \in \mathbf{A}$  such that  $x^n = abc$ . In particular, every multiplicative subset is weakly multiplicative.

**Lemma 5.1.** *A subset  $R$  of  $\mathbf{A}$  is regular if and only if it is weakly multiplicative.*

*Proof.* Suppose that  $R$  is regular and let  $a, b \in R$ . By regularity, there is  $r \in R$  such that  $ab \triangleright r$  which translates as claimed. Conversely, if  $R$  is weakly multiplicative, suppose that  $U, a \triangleright x$  and  $U, b \triangleright y$  for certain  $x, y \in R$ . This is to say that there are  $k, \ell \in \mathbb{N}$  and  $r, s \in \mathbf{A}$  as well as  $u, v \in \langle U \rangle$  such that

$$x^k = ra + u \quad \text{and} \quad y^\ell = sb + v.$$

It follows that

$$(xy)^{k+\ell} = tab + w$$

for certain  $t \in \mathbf{A}$  and  $w \in \langle U \rangle$ . Since  $R$  is weakly multiplicative there is  $z \in R$  along with  $n \in \mathbb{N}$  and  $c \in \mathbf{A}$  such that  $z^n = xyc$ . Thus,

$$z^{n(k+\ell)} = c^{k+\ell}(tab + w),$$

which witnesses  $U, ab \triangleright z \in R$ , whence  $R$  is regular.  $\square$

The following is a constructive version of Krull's Lemma that every radical ideal is the intersection of all containing prime ideals [61]. It is a direct consequence of CUPIT and the preceding Lemma 5.1.

**Proposition 5.2.** *Let  $M \subseteq \mathbf{A}$ . The following are equivalent.*

1.  $M$  is weakly multiplicative.
2. For every radical ideal  $\mathfrak{a}$  of  $\mathbf{A}$ , the following are equivalent:
  - a.  $M \not\subseteq \mathfrak{a}$ .
  - b. There is a tree  $t \in T_{\mathfrak{a}}$  which terminates in  $M$ .

**Corollary 5.3.** *For every  $a \in \mathbf{A}$ , the following are equivalent.*

1.  $a$  is nilpotent, i.e., there is  $n \in \mathbb{N}$  such that  $a^n = 0$ .
2. There is a tree  $t \in T_0$  terminating in  $\{a^n \mid n \in \mathbb{N}\}$ .

As an application of Corollary 5.3 we consider the well-known theorem that every non-constant coefficient of an invertible polynomial is nilpotent. This result has an elegant proof by reduction to the integral case, and has already seen

many a constructive treatment—see, e.g., [8, 27, 65, 79, 83, 96, 97]. Thus, suppose that


$$f = \sum_{i=0}^m a_i X^i \quad \text{and} \quad g = \sum_{j=0}^n b_j X^j$$

are such that

$$1 = fg = \sum_{k=0}^{m+n} c_k X^k, \quad \text{where } c_k = \sum_{i+j=k} a_i b_j. \quad (2)$$

It suffices to check by induction on  $m > 1$  that  $a_m$  is nilpotent. This is enough because in every commutative ring the sum of an invertible and a nilpotent element is in turn invertible.

From (2) we infer that  $a_m b_n = 0$ ,  $a_0 b_0 = 1$  as well as that  $a_m b_j \in \langle b_{j+1}, \dots, b_n \rangle$  for  $0 \leq j < n$ . With this information we construct a tree which terminates in  $a_m$ , thus witnessing the latter's nilpotency:



In fact, the rightmost path entails  $a_m$  since  $b_0$  is invertible; all other paths trivially entail  $a_m$ .

## 5.2 Glivenko's Theorem

Let  $\vdash_i$  and  $\vdash_c$  stand for intuitionistic and classical logic in a propositional language  $S$ . It is known [55, 74] that

$$\Gamma \vdash_c \varphi \quad \text{if and only if} \quad \Gamma, \Delta \vdash_i \varphi$$

for a suitable finite set  $\Delta$  of formulas  $\psi \vee \neg\psi$ , where  $\psi$  is a propositional variable occurring in  $\Gamma$  or  $\varphi$ .


Let  $\triangleright = \vdash_i$  on  $S$  and consider on top of  $\triangleright$  the non-deterministic axiom of excluded middle, i.e., let  $\mathcal{E}$  consist of all the instances, with  $\varphi \in S$ , of

$$\vdash \varphi, \neg\varphi$$

Let us consider an example. If, say,

$$\Gamma, \psi_1 \vee \neg\psi_1, \psi_2 \vee \neg\psi_2 \vdash_i \varphi,$$

then the tree



belongs to  $T_{\langle \Gamma \rangle}$  and terminates in  $\varphi$ . Proposition 4.6 implies that if  $\varphi$  is regular, then  $\Gamma \triangleright \varphi$ , which is to say that  $\Gamma \vdash_i \varphi$ .

**Lemma 5.4.** *A formula  $\varphi$  is regular if and only if it is stable, i.e., such that  $\neg\neg\varphi \triangleright \varphi$ .*

*Proof.* Clearly, if  $\varphi$  is regular, then  $\neg\neg\varphi \triangleright \varphi$  follows from  $\neg\neg\varphi, \varphi \triangleright \varphi$  and  $\neg\neg\varphi, \neg\varphi \triangleright \varphi$ . For the converse use the intuitionistic properties of (double) negation, due to Brouwer [16, 17], that  $\triangleright\neg\neg(\psi \vee \neg\psi)$ , and that  $\Gamma, \psi \triangleright \neg\chi$  implies  $\Gamma, \neg\psi \triangleright \neg\chi$ .  $\square$

Since every negated formula  $\neg\chi$  is stable, we regain the following version of Glivenko's theorem [44] from Proposition 4.6 with  $\alpha = \langle \Gamma \rangle$  and the above observations for  $\neg\chi$  as  $\varphi$ .

**Proposition 5.5 (Glivenko).** *If  $\Gamma \vdash_c \neg\chi$ , then  $\Gamma \vdash_i \neg\chi$ .*

We hasten to say that proofs of Glivenko's theorem usually go along similar lines. Recent literature about Glivenko's result includes [37, 38, 45, 57, 63, 76, 78].<sup>4</sup>

But what has Glivenko's Theorem to do with transfinite methods? In fact Proposition 5.5 is the syntactical underpinning of the following special case of Proposition 3.2, which in turn is a variant [38] of Lindenbaum's Lemma [107].

**Proposition 5.6 (ZFC).** *The intersection of all complete theories over  $\Gamma$  equals  $\{ \varphi \in S \mid \Gamma \vdash_i \neg\neg\varphi \}$ .*

As usual, by a *complete theory* we mean a deductively closed subset  $\Gamma$  of  $S$  such that for every  $\varphi \in S$  either  $\varphi \in \Gamma$  or  $\neg\varphi \in \Gamma$ . By the latter condition it is irrelevant whether deductive closure is understood for  $\vdash_i$  or  $\vdash_c$ , but of course the theories of the former are exactly the ideals of  $\triangleright = \vdash_i$ .

### 5.3 Order extension

Here is another application, this one in the context of order relations. Let  $E$  be a set. We say that a binary relation  $R$  on  $S$  is *order-regular* if, for all  $a, b, c, d \in E$ ,

$$(a, b) \in R \wedge (c, d) \in R \implies (a, d) \in R \vee (c, b) \in R. \quad (3)$$

An *interval order* is an order-regular relation which, in addition, is irreflexive.<sup>5</sup> Hence every interval order is transitive.

<sup>4</sup>This list of references is by no means meant exhaustive.

<sup>5</sup>The concept of interval order can be traced back to early work of Wiener's on the theory of measurement [40]. Fishburn coined the term "interval

By a *quasiorder* we understand a reflexive transitive relation. A *linear order*  $L$  is a transitive relation such that, for all  $a, b \in E$  either  $(a, b) \in L$  or  $(b, a) \in L$ , by way of which it is reflexive. A quasiorder is order-regular precisely when it is linear.

If  $P$  and  $P'$  are quasiorders such that  $P \subseteq P'$ , then we say that the latter *extends* the former. With this terminology, the problem of finding a *linear extension* of a quasiorder becomes trivial: the cartesian product  $E \times E$  will do. Thus a more restrictive concept of extension is at work, e.g., in social choice theory [15, 18], yet to achieve linearity often requires transfinite methods [98, 106]. The following results force non-triviality by demanding that a linear order be in the complement of a given order-regular relation.

Let  $S = E \times E$  and let  $\triangleright$  correspond to *transitive closure*, i.e., put

$$U \triangleright (a, b) \equiv (a, b) \in \bigcup_{i \geq 1} U^i$$

where  $U^1 = U$  and  $U^{i+1} = U^i \circ U$ , where

$$U \circ V = \{ (a, c) \mid (\exists b \in E)((a, b) \in U \wedge (b, c) \in V) \}.$$

On top of this we now consider the non-deterministic axiom of order-regularity, i.e. we let  $\mathcal{E}$  consist of all the instances, with  $a, b, c, d \in E$ , of

$$(a, b), (c, d) \vdash (a, d), (c, b).$$

**Lemma 5.7.** *Every order-regular relation is regular.*

*Proof.* Suppose that  $R$  is order-regular. We need to show that

$$\frac{\langle U, (a, b) \rangle \not\vdash R \quad \langle U, (c, d) \rangle \not\vdash R}{\langle U, (a, d), (c, b) \rangle \not\vdash R}$$

where  $U \in \text{Fin}(S)$ . It suffices to settle the case in which the two assumptions are witnessed by chains through  $(a, b)$  and  $(c, d)$ , respectively. Invoking order-regularity immediately yields the result. At one glance:

$$\begin{array}{ccccccc} x & \text{----} & a & \text{-----} & b & \text{----} & y \\ & & & \searrow & & \nearrow & \\ & & & c & \text{-----} & d & \text{----} & w \\ & & z & \text{----} & & & & \end{array}$$

Here  $(x, y), (z, w) \in R$ , and dashed lines indicate chains through  $U$ . Depending on whether  $(x, w) \in R$  or  $(z, y) \in R$ , glue along  $(a, d)$  or  $(c, b)$ , accordingly.  $\square$

In **ZFC**, UPIT implies that if  $R$  is an order-regular relation, and  $P$  is a quasiorder on  $E$ , then  $R$  and  $P$  are disjoint if and only if there is a linear order  $L$  that extends  $P$  yet keeps off  $R$ . The particular case where  $P$  is the diagonal relation yields that an order-regular relation  $R$  is an interval order if and only if its complement contains a linear order. In fact, a prime quasiorder is nothing but a linear one.

order" for what Wiener had called "relation of complete sequence" [39–41]. Yet another occurrence of this concept goes under the name of an irreflexive "Ferrers relation" due to Riguet [84].

We further note the following consequence of CUPIT, which is the constructive counterpart of the aforesaid.

**Proposition 5.8.** *Let  $R$  be an order-regular relation and let  $P$  be a quasiorder on  $E$ . The following are equivalent.*

1.  $R \not\sqsubseteq P$ .
2. *There is a tree  $t \in T_P$  which terminates in  $R$ .*

## 6 Multi-conclusion entailment

In this final section we shed some light on certain aspects of multi-conclusion entailment relations as extending their single-conclusion counterparts. The relevance of the notion of entailment relation to constructive algebra and point-free topology has been pointed out in [19], and has been used very widely, e.g. in [22–24, 26, 28, 31, 33, 75, 85, 89, 95, 109, 110]. Lorenzen's precedence is currently under scrutiny [29, 30]. Consequence and entailment have further caught interest from various angles [7, 33, 36, 43, 52–54, 77, 90, 94, 104, 111].

We begin with a brief summary, closely referring to [87, 88] for a thorough account which builds on ideas and results from proof theory [75] and dynamical algebra [31, 34, 65].

Let  $S$  be a set. Recall that a *multi-conclusion entailment relation* [100–102] is a relation

$$\vdash \subseteq \text{Fin}(S) \times \text{Fin}(S)$$

between finite subsets  $U$  and  $V$  of  $S$  which is *reflexive* and *monotone*:

$$\frac{U \not\sqsubseteq V}{U \vdash V} \text{ (R)} \quad \frac{U \vdash V}{U, U' \vdash V, V'} \text{ (M)}$$

as well as *transitive*:

$$\frac{U \vdash V, a \quad U, a \vdash V}{U \vdash V} \text{ (T)}$$

where we make use of the usual shorthand notations.

Given  $\vdash$ , its *trace*  $\triangleright_\vdash$  is defined by

$$U \triangleright_\vdash a \equiv U \vdash a$$

and in fact is a consequence relation.

A *model* of  $\vdash$  is a subset  $p$  of  $S$  such that if  $p \supseteq U$  and  $U \vdash V$  then  $p \not\sqsubseteq V$ . The class of all models of  $\vdash$  will be denoted by

$$\text{Mod}(\vdash).$$

It is a consequence of the prime ideal theorem for distributive lattices (PIT) that every multi-conclusion entailment relation is determined by its models [19], which is to say that

$$(\forall p \in \text{Mod}(\vdash))(U \subseteq p \implies p \not\sqsubseteq V) \implies U \vdash V. \quad (4)$$

Every set  $\mathcal{E}$  of non-deterministic axioms gives rise to a multi-conclusion entailment relation  $\vdash_{\mathcal{E}}$  which is least among those  $\vdash$  for which  $A \vdash B$  for all  $(A, B) \in \mathcal{E}$ . Following [90], this  $\vdash_{\mathcal{E}}$  can be generated inductively by the rules for reflexivity and transitivity on axioms:

$$\frac{U \not\sqsubseteq V}{U \vdash_{\mathcal{E}} V} \text{ (R)} \quad \frac{(A, B) \in \mathcal{E} \quad (\forall b \in B) U, b \vdash_{\mathcal{E}} V}{U, A \vdash_{\mathcal{E}} V} \text{ (Ax)} \quad (5)$$

much akin to [32, 73]. Every model of  $\vdash_{\mathcal{E}}$  is apparently closed for every member of  $\mathcal{E}$ , keeping in mind that the elements of the latter are turned into entailments. Conversely, by induction it is easy to see that if a subset of  $S$  is closed for every member of  $\mathcal{E}$ , then it is in fact a model of  $\vdash_{\mathcal{E}}$ .

### 6.1 Regularity as conservation

A multi-conclusion entailment relation  $\vdash$  *extends* a single-conclusion entailment relation  $\triangleright$  on  $S$  if, for every  $U \in \text{Fin}(S)$  and  $a \in S$ ,  $U \triangleright a$  implies  $U \vdash a$ , which is to say that  $\triangleright \subseteq \triangleright_\vdash$ . In case the converse holds as well, i.e. altogether  $\triangleright = \triangleright_\vdash$ , any such extension is said to be *conservative*. An extension  $\vdash$  of  $\triangleright$  is conservative if and only if [87, 88], for all entailments  $a_1, \dots, a_k \vdash b_1, \dots, b_\ell$  and  $U \in \text{Fin}(S)$ ,

$$\frac{U, b_1 \triangleright c \quad \dots \quad U, b_\ell \triangleright c}{U, a_1, \dots, a_k \triangleright c} \quad (6)$$

It suffices to consider only initial entailments in place of  $a_1, \dots, a_k \vdash b_1, \dots, b_\ell$  whenever  $\vdash$  is inductively generated.

Consider again our default set  $\mathcal{E}$  of non-deterministic axioms on top of a single-conclusion entailment relation  $\triangleright$ . Passing to the union of  $\mathcal{E}$  and  $\triangleright$ , we may assume that  $(U, \{a\}) \in \mathcal{E}$  whenever  $U \triangleright a$ . Next let  $\vdash_{\mathcal{E}}$  denote the multi-conclusion entailment relation which is inductively generated by  $\mathcal{E}$  according to the rules (5) laid out before. In other words, this  $\vdash_{\mathcal{E}}$  is the least multi-conclusion entailment relation which extends  $\triangleright$  such that  $A \vdash_{\mathcal{E}} B$  for all  $(A, B) \in \mathcal{E}$ . By the above remarks on the models  $\vdash_{\mathcal{E}}$ , we immediately know about the semantics of this entailment relation:

**Proposition 6.1.**  $\text{Spec}(\mathcal{E}) = \text{Mod}(\vdash_{\mathcal{E}})$ .

The following is a mere rephrasing of the conservation criterion (6) recalled before.

**Proposition 6.2.** *The following are equivalent.*

1. *Every element of  $S$  is regular.*
2.  *$\vdash_{\mathcal{E}}$  is conservative over  $\triangleright$ .*

### 6.2 Intersecting prime ideals

Next we aim at identifying those elements which are common to every prime ideal over a given one. If  $t \in T_{\mathfrak{a}}$  for a certain ideal  $\mathfrak{a}$ , let  $\text{paths}(t)$  denote the set of all paths of  $t$ . Next we introduce an auxiliary relation

$$U \triangleright_T a \equiv (\exists t \in T_{(U)}) (\forall \pi \in \text{paths}(t)) \pi \triangleright_0 a,$$

where  $\triangleright_0$  is the least consequence relation that extends  $\triangleright$  with additional axioms

$$A \triangleright_0 a \quad \text{for } (A, \emptyset) \in \mathcal{E},$$

and where we understand  $\pi \triangleright_0 a$  according to the conventions laid out in the first paragraphs of Section 4. It will turn out that  $\triangleright_T$  is a consequence relation. Notice that

$$U \triangleright_0 a \quad \text{implies} \quad U \vdash_{\mathcal{E}} a, \quad (7)$$

simply because  $\vdash_{\mathcal{E}}$  contains the generating axioms of  $\triangleright_0$ .

**Lemma 6.3.** For all  $U \in \text{Fin}(S)$  and  $c \in S$  and  $(A, B) \in \mathcal{E}$ ,

$$\frac{(\forall b \in B) U, b \triangleright_T c}{U, A \triangleright_T c}$$

*Proof.* Let  $B = \{b_1, \dots, b_n\}$ . If  $n = 0$ , then the trivial tree labelled with  $U, A$  witnesses  $U, A \triangleright_T c$ . Suppose next that  $n \geq 1$ , and that trees  $t_i \in T_{\langle U, b_i \rangle}$  witness  $U, b_i \triangleright_T c$  for  $1 \leq i \leq n$ . These trees can be grafted, correspondingly, at the leaves of the tree displayed in (1) so as to obtain a witness for  $U, A \triangleright_T c$ .  $\square$

**Proposition 6.4.** For all  $U \in \text{Fin}(S)$  and  $a \in S$ ,

$$U \vdash_{\mathcal{E}} a \quad \text{if and only if} \quad U \triangleright_T a,$$

i.e.,  $\triangleright_T$  is the trace of  $\vdash_{\mathcal{E}}$ . In particular,  $\triangleright_T$  is a consequence relation.

*Proof.* For the left-to-right implication we argue by induction on the generation of  $U \vdash_{\mathcal{E}} a$ . Reflexivity (R) is clear, and the case of transitivity (Ax) is taken care of by Lemma 6.3.

As regards the converse, we argue by induction on the tree  $t \in T_{\langle U \rangle}$  witnessing  $U \triangleright_T a$ . The base case boils down to (7). Consider next the case in which  $t$  has been extended at the leaf of one of its paths  $\pi$  with children labelled with the  $b \in B$ , where  $(A, B) \in \mathcal{E}$  and  $\langle \pi \rangle \supseteq A$ . For every  $b \in B$ , relabelling the root of  $t$  with  $U \cup \{b\}$  yields a witness for  $U, b \triangleright_T a$  for which  $U, b \vdash_{\mathcal{E}} a$  by induction, whence  $U, A \vdash_{\mathcal{E}} a$  by (Ax). It remains to observe that the assumption that  $\langle \pi \rangle \supseteq A$  lifts along the generation of  $t$  with successive cuts (T), so that we may assume  $\langle U \rangle \supseteq A$  and conclude  $U \vdash_{\mathcal{E}} a$ .  $\square$

By Proposition 6.4,  $\vdash_{\mathcal{E}}$  is conservative at least over  $\triangleright_T$ . This means (Proposition 6.2) that every element of  $S$  is regular for  $\vdash_{\mathcal{E}}$  over  $\triangleright_T$  (Lemma 6.3). In particular, the trace of  $\vdash_{\mathcal{E}}$  equals  $\triangleright_T$  and thus can be computed in terms of the inductively defined tree class.

**Corollary 6.5 (ZFC).** For every ideal  $\mathfrak{a}$ ,

$$\bigcap \text{Spec}(\mathcal{E})/\mathfrak{a} = \{a \in S \mid (\exists U \in \text{Fin}(\mathfrak{a})) U \triangleright_T a\}. \quad (8)$$

*Proof.* Combine completeness (4) with Proposition 6.1 and Proposition 6.4.  $\square$

If every element of  $S$  is regular for  $\mathcal{E}$  over  $\triangleright$ , then already  $\triangleright$  is the trace of  $\vdash_{\mathcal{E}}$ , and thus equals  $\triangleright_T$  by Proposition 6.4. So (8) boils down to Corollary 3.3.2.

### 6.3 Some reverse mathematics

The *Restricted Law of Excluded Middle* (REM) is not part of **CZF**. This REM means  $\varphi \vee \neg\varphi$  for every set-theoretic formula  $\varphi$  that is *bounded* in the sense that only set-bounded quantifiers  $\forall x \in y$  and  $\exists x \in y$  occur in  $\varphi$ .

The following essentially rests on an argument of Bell's [9] and has already been used to show that completeness (4) implies REM [88].

**Proposition 6.6.** UPIT implies REM.

*Proof.* Given a Boolean algebra  $B$ , consider on  $S = B$  the entailment relation  $\triangleright$  of filter:

$$a_1, \dots, a_k \triangleright a \equiv a_1 \wedge \dots \wedge a_k \leq a.$$

On top of  $\triangleright$  we consider the non-deterministic axioms of *proper prime filter*, viz.

$$0 \vdash$$

$$a \vee b \vdash a, b$$

for all  $a, b \in B$ . Distributivity ensures that every element is regular. Corollary 3.3 (which is a direct consequence of UPIT) thus implies that *in each Boolean algebra the intersection of the family of all its prime filters is  $\{1\}$* . Bell has shown that over intuitionistic set theory **IZ** this statement implies excluded middle [9, p. 161ff.]. The argument goes through over **CZF** if restricted to deal with bounded formulas only.  $\square$

**Lemma 6.7.** Suppose that  $R \subseteq S$  is regular. If  $U \vdash_{\mathcal{E}} V$  and  $V \subseteq R$ , then  $\langle U \rangle \not\triangleright R$ .

*Proof.* By induction on  $U \vdash_{\mathcal{E}} V$ . The case of transitivity (Ax) boils down to regularity.  $\square$

In particular, we see once again that if  $a$  is regular, then  $U \triangleright a$  if and only if  $U \vdash_{\mathcal{E}} a$ , and so  $\vdash_{\mathcal{E}}$  is conservative over  $\triangleright$  if (and only if) every element of  $S$  is regular.

As indicated in Section 5.1, Krull's Lemma, which is equivalent to the prime ideal theorem for distributive lattices (PIT) [51], is a consequence of UPIT. The reader has perhaps anticipated the following result which calibrates UPIT.

**Proposition 6.8.** Over **ZF**, UPIT is equivalent to PIT.

*Proof.* While it is evident that UPIT implies PIT, to show the converse, let  $\mathfrak{a}$  be an ideal and let  $R$  be regular. Put

$$U \vdash' V \equiv (\exists U_0 \in \text{Fin}(\mathfrak{a}))(\exists V_0 \in \text{Fin}(R)) U, U_0 \vdash_{\mathcal{E}} V, V_0.$$

This  $\vdash'$  is an entailment relation [19] the models of which are the prime ideals that contain  $\mathfrak{a}$  but avoid  $R$ . Now, if  $R$  and  $\mathfrak{a}$  are disjoint, then  $\vdash'$  is consistent according to Lemma 6.7: that is,  $\emptyset \not\vdash' \emptyset$ . Hence  $\vdash'$  has a model by (4), which is a consequence of PIT.  $\square$

## Acknowledgments

The present study was carried out within the projects “A New Dawn of Intuitionism: Mathematical and Philosophical Advances” (ID 60842) funded by the John Templeton Foundation, and “Reducing complexity in algebra, logic, combinatorics - REDCOM” belonging to the programme “Ricerca Scientifica di Eccellenza 2018” of the Fondazione Cariverona. The authors are members of the Gruppo Nazionale per le Strutture Algebriche, Geometriche e le loro Applicazioni (GNSAGA) within the Italian Istituto Nazionale di Alta Matematica (INdAM).<sup>6</sup> Last but not least, the authors wish to express their

<sup>6</sup>The opinions expressed in this paper are those of the authors and do not necessarily reflect the views of those foundations.



gratitude to Ulrich Berger, Stefan Neuwirth and Iosif Petrakis for interesting discussions, as well as to the anonymous referees for expertly and insightful remarks that helped to improve the paper.

## References

- [1] Peter Aczel. 1977. An introduction to inductive definitions. In *Handbook of Mathematical Logic*, Jon Barwise (Ed.). Studies in Logic and the Foundations of Mathematics, Vol. 90. Elsevier Science B.V., Amsterdam, 739–782.
- [2] Peter Aczel. 1978. The type theoretic interpretation of constructive set theory. In *Logic Colloquium '77 (Proc. Conf., Wrocław, 1977)*. Stud. Logic Foundations Math., Vol. 96. North-Holland, Amsterdam, 55–66.
- [3] Peter Aczel. 1982. The type theoretic interpretation of constructive set theory: choice principles. In *The L. E. J. Brouwer Centenary Symposium (Noordwijkerhout, 1981)*. Stud. Logic Found. Math., Vol. 110. North-Holland, Amsterdam, 1–40.
- [4] Peter Aczel. 1986. The type theoretic interpretation of constructive set theory: inductive definitions. In *Logic, methodology and philosophy of science, VII (Salzburg, 1983)*. Stud. Logic Found. Math., Vol. 114. North-Holland, Amsterdam, 17–49.
- [5] Peter Aczel and Michael Rathjen. 2000. *Notes on Constructive Set Theory*. Technical Report. Institut Mittag-Leffler. Report No. 40.
- [6] Peter Aczel and Michael Rathjen. 2010. Constructive set theory. (2010). <https://www1.maths.leeds.ac.uk/~rathjen/book.pdf> Book draft.
- [7] Arnon Avron. 1991. Simple consequence relations. *Inform. and Comput.* 92 (1991), 105–139.
- [8] B. Banaschewski and J. J. C. Vermeulen. 1996. Polynomials and radical ideals. *J. Pure Appl. Algebra* 113, 3 (1996), 219–227.
- [9] John L. Bell. 2005. *Set Theory. Boolean-Valued Models and Independence Proofs* (third ed.). Oxford University Press, Oxford.
- [10] Marc Bezem and Thierry Coquand. 2005. Automating coherent logic. In *International Conference on Logic for Programming Artificial Intelligence and Reasoning*. Springer, 246–260.
- [11] Marc Bezem and Dimitri Hendriks. 2008. On the mechanization of the proof of Hessenberg’s theorem in coherent logic. *J. Automat. Reason.* 40, 1 (2008), 61–85.
- [12] Jean-Yves Béziau. 2006. Les axiomes de Tarski. In *La philosophie en Pologne 1919-1939*, Roger Pouivet and Manuel Resbuschi (Eds.). Librairie Philosophique J. Vrin, Paris.
- [13] Errett Bishop. 1967. *Foundations of Constructive Analysis*. McGraw-Hill, New York.
- [14] Errett Bishop and Douglas Bridges. 1985. *Constructive Analysis*. Springer.
- [15] Walter Bossert and Kotaro Suzumura. 2010. *Consistency, Choice, and Rationality*. Harvard University Press, Cambridge.
- [16] L.E.J. Brouwer. 1908. De onbetrouwbaarheid der logische principes. *Tijdschrift voor Wijsbegeerte* 2 (1908), 152–158.
- [17] L.E.J. Brouwer. 1925. Intuitionistische Zerlegung mathematischer Grundbegriffe. *Jahresbericht der Deutschen Mathematiker-Vereinigung* 33 (1925), 251–256.
- [18] Susumu Cato. 2012. Szpilrajn, Arrow and Suzumura: Concise proofs of extension theorems and an extension. *Metroeconomica* 63 (2012), 235–249.
- [19] Jan Cederquist and Thierry Coquand. 2000. Entailment relations and distributive lattices. In *Logic Colloquium '98. Proceedings of the Annual European Summer Meeting of the Association for Symbolic Logic, Prague, Czech Republic, August 9–15, 1998*, Samuel R. Buss, Petr Hájek, and Pavel Pudlák (Eds.). Lect. Notes Logic, Vol. 13. A. K. Peters, Natick, MA, 127–139.
- [20] Francesco Ciraulo, Maria Emilia Maietti, and Giovanni Sambin. 2013. Convergence in formal topology: a unifying notion. *J. Log. Anal.* 5 (2013).
- [21] Francesco Ciraulo and Giovanni Sambin. 2008. Finitary formal topologies and Stone’s representation theorem. *Theoret. Comput. Sci.* 405, 1–2 (2008), 11–23.
- [22] Thierry Coquand. 2000. A direct proof of the localic Hahn-Banach theorem. <http://www.cse.chalmers.se/~coquand/formal.html>
- [23] Thierry Coquand. 2000. Lewis Carroll, Gentzen and entailment relations. <http://www.cse.chalmers.se/~coquand/formal.html>
- [24] Thierry Coquand. 2005. About Stone’s notion of spectrum. *J. Pure Appl. Algebra* 197, 1–3 (2005), 141–158.
- [25] Thierry Coquand. 2005. A completeness proof for geometrical logic. In *Logic, Methodology and Philosophy of Science. Proceedings of the Twelfth International Congress*, P. Hájek, L. Valdés-Villanueva, and D. Westerståhl (Eds.). King’s College Publications, London, 79–90.
- [26] Thierry Coquand. 2009. Space of valuations. *Ann. Pure Appl. Logic* 157 (2009), 97–109.
- [27] Thierry Coquand, Lionel Ducos, Henri Lombardi, and Claude Quitté. 2003. L’idéal des coefficients du produit de deux polynômes. *Revue des Mathématiques de l’Enseignement Supérieur* 113, 3 (2003), 25–39.
- [28] Thierry Coquand and Henri Lombardi. 2002. Hidden constructions in abstract algebra: Krull dimension of distributive lattices and commutative rings. In *Commutative Ring Theory and Applications (Lect. Notes Pure Appl. Mathematics)*, M. Fontana, S.-E. Kabbaj, and S. Wiegand (Eds.), Vol. 231. Addison-Wesley, Reading, MA, 477–499.
- [29] Thierry Coquand, Henri Lombardi, and Stefan Neuwirth. 2019. Lattice-ordered groups generated by an ordered group and regular systems of ideals. *Rocky Mountain J. Math.* 49, 5 (2019), 1449–1489.
- [30] Thierry Coquand and Stefan Neuwirth. 2017. An introduction to Lorenzen’s “Algebraic and logistic investigations on free lattices” (1951). (2017). <https://arxiv.org/abs/1711.06139> Preprint.
- [31] Thierry Coquand and Henrik Persson. 2001. Valuations and Dedekind’s Prague theorem. *J. Pure Appl. Algebra* 155, 2–3 (2001), 121–129.
- [32] Thierry Coquand, Giovanni Sambin, Jan Smith, and Silvio Valentini. 2003. Inductively generated formal topologies. *Ann. Pure Appl. Logic* 124 (2003), 71–106.
- [33] Thierry Coquand and Guo-Qiang Zhang. 2000. Sequents, frames, and completeness. In *Computer Science Logic (Fischbachau, 2000)*, Peter G. Clote and Helmut Schwichtenberg (Eds.). Lecture Notes in Comput. Sci., Vol. 1862. Springer, Berlin, 277–291.
- [34] Michel Coste, Henri Lombardi, and Marie-Françoise Roy. 2001. Dynamical method in algebra: Effective Nullstellensätze. *Ann. Pure Appl. Logic* 111, 3 (2001), 203–256.
- [35] Hans De Nivelle and Jia Meng. 2006. Geometric resolution: A proof procedure based on finite model search. In *International Joint Conference on Automated Reasoning*. Springer, 303–317.
- [36] Kosta Došen. 1999. On passing from singular to plural consequences. In *Logic at Work: Essays Dedicated to the Memory of Helena Rasiowa*, Ewa Orłowska (Ed.). Stud. Fuzziness Soft Comput., Vol. 24. Physica, Heidelberg, 533–547.
- [37] Christian Espindola. 2013. A short proof of Glivenko theorems for intermediate predicate logics. *Arch. Math. Logic* 52, 7–8 (2013), 823–826.
- [38] Giulio Fellin, Peter Schuster, and Daniel Wessel. 2019. The Jacobson Radical of a Propositional Theory. In *Proof-Theoretic Semantics: Assessment and Future Perspectives. Proceedings of the Third Tübingen Conference on Proof-Theoretic Semantics, 27–30 March 2019*, Thomas Piecha and Peter Schroeder-Heister (Eds.). University of Tübingen, 287–299. <http://dx.doi.org/10.15496/publikation-35319>
- [39] Peter C. Fishburn. 1970. *Utility Theory for Decision Making*. Wiley, New York.
- [40] Peter Fishburn and Bernard Monjardet. 1992. Norbert Wiener on the Theory of Measurement (1914, 1915, 1921). *J. Math. Psych.* 36 (1992), 165–184.

- [41] Peter C. Fishburn. 1970. Intransitive indifference with unequal indifference intervals. *J. Math. Psych.* 7 (1970), 144–149.
- [42] John Fisher and Marc Bezem. 2009. Skolem machines. *Fund. Inform.* 91, 1 (2009), 79–103.
- [43] Dov M. Gabbay. 1981. *Semantical investigations in Heyting's intuitionistic logic*. Synthese Library, Vol. 148. D. Reidel Publishing Co., Dordrecht-Boston, Mass.
- [44] Valery Glivenko. 1929. Sur quelques points de la Logique de M. Brouwer. *Acad. Roy. Belg. Bull. Cl. Sci. (5)* 15 (1929), 183–188.
- [45] Giulio Guerrieri and Alberto Naibo. 2019. Postponement of raa and Glivenko's theorem, revisited. *Studia Logica* 107, 1 (2019), 109–144.
- [46] Paul Hertz. 1922. Über Axiomensysteme für beliebige Satzsysteme. I. Teil. Sätze ersten Grades. *Math. Ann.* 87, 3 (1922), 246–269.
- [47] Paul Hertz. 1923. Über Axiomensysteme für beliebige Satzsysteme. II. Teil. Sätze höheren Grades. *Math. Ann.* 89, 1 (1923), 76–102.
- [48] Paul Hertz. 1929. Über Axiomensysteme für beliebige Satzsysteme. *Math. Ann.* 101, 1 (1929), 457–514.
- [49] Ayana Hirata, Hajime Ishihara, Tatsuji Kawai, and Takako Nemoto. 2019. Equivalents of the finitary non-deterministic inductive definitions. *Ann. Pure Appl. Logic* 170, 10 (2019), 1256–1272.
- [50] Bjarne Holen, Dag Hovland, and Martin Giese. 2013. Efficient rule-matching for hyper-tableaux. In *IWIL 2012 (EPiC Series, vol. 22)*, K. Korovin, S. Schulz, and E. Ternovska (Eds.), 4–17.
- [51] Paul Howard and Jean Rubin. 1998. *Consequences of the Axiom of Choice*. American Mathematical Society, Providence, RI.
- [52] Lloyd Humberstone. 2011. On a conservative extension argument of Dana Scott. *Log. J. IGPL* 19 (2011), 241–288.
- [53] Lloyd Humberstone. 2012. Dana Scott's work with generalized consequence relations. In *Universal Logic: An Anthology. From Paul Hertz to Dov Gabbay*, Jean-Yves Béziau (Ed.), Birkhäuser, Basel, 263–279.
- [54] Rosalie Iemhoff. 2016. Consequence Relations and Admissible Rules. *J. Philosophical Logic* 45, 3 (2016), 327–348.
- [55] Hajime Ishihara. 2014. Classical propositional logic and decidability of variables in intuitionistic propositional logic. *Log. Methods Comput. Sci.* 10, 3 (2014), 3:1, 7. [https://doi.org/10.2168/LMCS-10\(3:1\)2014](https://doi.org/10.2168/LMCS-10(3:1)2014)
- [56] Hajime Ishihara and Takako Nemoto. 2016. Non-Deterministic Inductive Definitions and Fullness. In *Concepts of Proof in Mathematics, Philosophy, and Computer Science*, D. Probst and P. Schuster (Eds.), Ontos Mathematical Logic, Vol. 6. Walter de Gruyter, Berlin, 163–170.
- [57] Hajime Ishihara and Helmut Schwichtenberg. 2016. Embedding classical in minimal implicational logic. *MLQ Math. Log. Q.* 62, 1-2 (2016), 94–101.
- [58] P. T. Johnstone. 1977. *Topos Theory*. Academic Press [Harcourt Brace Jovanovich Publishers], London. London Mathematical Society Monographs, Vol. 10.
- [59] Peter T. Johnstone. 2002. *Sketches of an Elephant: A Topos Theory Compendium. Vol. 2*. Oxford Logic Guides, Vol. 44. The Clarendon Press Oxford University Press, Oxford.
- [60] Gregor Kemper and Ihnen Yengui. 2019. Valiative dimension and monomial orders. (2019). <https://arxiv.org/abs/1906.12067>
- [61] Wolfgang Krull. 1929. Idealtheorie in Ringen ohne Endlichkeitsbedingung. *Math. Ann.* 101 (1929), 729–744.
- [62] Javier Legris. [n.d.]. Paul Hertz and the origins of structural reasoning. In *Universal Logic: An Anthology. From Paul Hertz to Dov Gabbay*, Jean-Yves Béziau (Ed.), Birkhäuser, Basel, 3–10.
- [63] Tadeusz Litak, Miriam Polzer, and Ulrich Rabenstein. 2017. Negative translations and normal modality. In *2nd International Conference on Formal Structures for Computation and Deduction*. LIPICs. Leibniz Int. Proc. Inform., Vol. 84. Schloss Dagstuhl. Leibniz-Zent. Inform., Wadern, Art. No. 27, 18.
- [64] Henri Lombardi. 2002. Hidden constructions in abstract algebra. I. Integral dependance. *J. Pure Appl. Algebra* 167 (2002), 259–267.
- [65] Henri Lombardi and Claude Quitté. 2015. *Commutative Algebra: Constructive Methods. Finite Projective Modules*. Algebra and Applications, Vol. 20. Springer Netherlands, Dordrecht.
- [66] Ray Mines, Fred Richman, and Wim Ruitenburg. 1988. *A Course in Constructive Algebra*. Springer, New York. Universitext.
- [67] Christopher J. Mulvey and Joan Wick-Pelletier. 1991. A globalization of the Hahn–Banach theorem. *Adv. Math.* 89 (1991), 1–59.
- [68] Sara Negri. [n.d.]. Geometric rules in infinitary logic. In *Arnon Avron on Semantics and Proof Theory of Non-Classical Logics*. Springer. Forthcoming.
- [69] Sara Negri. 1996. Stone bases alias the constructive content of Stone representation. In *Logic and algebra. Proceedings of the international conference dedicated to the memory of Roberto Magari, April 26–30, 1994, Pontignano, Italy*, Aldo Ursini and Paolo Aglianò (Eds.), Lecture Notes in Pure and Applied Mathematics, Vol. 180. Marcel Dekker, New York, 617–636.
- [70] Sara Negri. 2002. Continuous domains as formal spaces. *Math. Structures Comput. Sci.* 12, 1 (2002), 19–52.
- [71] Sara Negri. 2003. Contraction-free sequent calculi for geometric theories with an application to Barr's theorem. *Arch. Math. Logic* 42, 4 (2003), 389–401.
- [72] Sara Negri. 2014. Proof analysis beyond geometric theories: from rule systems to systems of rules. *J. Logic Comput.* 26, 2 (2014), 513–537.
- [73] Sara Negri and Jan von Plato. 1998. Cut Elimination in the Presence of Axioms. *Bull. Symb. Log.* 4, 4 (1998), 418–435.
- [74] Sara Negri and Jan von Plato. 2001. *Structural Proof Theory*. Cambridge University Press, Cambridge.
- [75] Sara Negri, Jan von Plato, and Thierry Coquand. 2004. Proof-theoretical analysis of order relations. *Arch. Math. Logic* 43 (2004), 297–309.
- [76] Hiroakira Ono. 2009. Glivenko theorems revisited. *Ann. Pure Appl. Logic* 161, 2 (2009), 246–250.
- [77] Gillman Payette and Peter K. Schotch. 2014. Remarks on the Scott–Lindenbaum Theorem. *Studia Logica* 102, 5 (2014), 1003–1020.
- [78] Luiz Carlos Pereira and Edward Hermann Haeusler. 2015. On constructive fragments of classical logic. In *Dag Prawitz on proofs and meaning*. Outst. Contrib. Log., Vol. 7. Springer, Cham, 281–292.
- [79] Henrik Persson. 1999. An application of the constructive spectrum of a ring. In *Type Theory and the Integrated Logic of Programs*. Chalmers University and University of Göteborg. PhD thesis.
- [80] Thomas Powell, Peter Schuster, and Franziskus Wiesnet. 2019. An algorithmic approach to the existence of ideal objects in commutative algebra. In *26th Workshop on Logic, Language, Information and Computation (WoLLIC 2019), Utrecht, Netherlands, 2–5 July 2019, Proceedings (Lect. Notes Comput. Sci.)*, R. Iemhoff and M. Moortgat (Eds.), Vol. 11541. Springer, Berlin, 533–549.
- [81] Michael Rathjen. 2005. Generalized inductive definitions in Constructive Set Theory. In *From Sets and Types to Topology and Analysis: Towards Practicable Foundations for Constructive Mathematics*, Laura Crosilla and Peter Schuster (Eds.), Oxford Logic Guides, Vol. 48. Clarendon Press, Oxford, Chapter 16.
- [82] Michael Rathjen. 2016. Remarks on Barr's theorem. Proofs in Geometric Theories. In *Concepts of Proof in Mathematics, Philosophy, and Computer Science*, D. Probst and P. Schuster (Eds.), Ontos Mathematical Logic, Vol. 6. Walter de Gruyter, Berlin, 347–374.
- [83] Fred Richman. 1988. Nontrivial uses of trivial rings. *Proc. Amer. Math. Soc.* 103, 4 (1988), 1012–1014.
- [84] Jacques Riguet. 1951. Les relations de Ferrers. *C. R. Acad. Sci.* 232 (1951), 1729–1730.
- [85] Davide Rinaldi. 2014. *Formal Methods in the Theories of Rings and Domains*. Doctoral dissertation. Universität München.
- [86] Davide Rinaldi and Peter Schuster. 2016. A universal Krull–Lindenbaum theorem. *J. Pure Appl. Algebra* 220 (2016), 3207–3232.
- [87] Davide Rinaldi, Peter Schuster, and Daniel Wessel. 2017. Eliminating disjunctions by disjunction elimination. *Bull. Symb. Logic* 23, 2 (2017), 181–200.

- [88] Davide Rinaldi, Peter Schuster, and Daniel Wessel. 2018. Eliminating disjunctions by disjunction elimination. *Indag. Math. (N.S.)* 29, 1 (2018), 226–259.
- [89] Davide Rinaldi and Daniel Wessel. 2018. Extension by conservation. Sikorski’s theorem. *Log. Methods Comput. Sci.* 14, 4:8 (2018), 1–17.
- [90] Davide Rinaldi and Daniel Wessel. 2019. Cut elimination for entailment relations. *Arch. Math. Logic* 58, 5–6 (2019), 605–625.
- [91] Giovanni Sambin. 1987. Intuitionistic formal spaces—a first communication. In *Mathematical Logic and its Applications, Proc. Adv. Internat. Summer School Conf., Druzhba, Bulgaria, 1986*, D. Skordev (Ed.). Plenum, New York, 187–204.
- [92] Giovanni Sambin. 2003. Some points in formal topology. *Theoret. Comput. Sci.* 305, 1–3 (2003), 347–408.
- [93] Giovanni Sambin. forthcoming. *The Basic Picture. Structures for Constructive Topology*. Clarendon Press, Oxford.
- [94] Tor Sandqvist. 2018. Preservation of structural properties in intuitionistic extensions of an inference relation. *Bull. Symb. Log.* 24, 3 (2018), 291–305.
- [95] Konstantin Schlagbauer, Peter Schuster, and Daniel Wessel. 2019. Der Satz von Hahn–Banach per Disjunktionselimination. *Confluentes Math.* 11, 1 (2019), 79–93.
- [96] Peter Schuster. 2012. Induction in algebra: a first case study. In *2012 27th Annual ACM/IEEE Symposium on Logic in Computer Science*. IEEE Computer Society Publications, 581–585. Proceedings, LICS 2012, Dubrovnik, Croatia.
- [97] Peter Schuster. 2013. Induction in algebra: a first case study. *Log. Methods Comput. Sci.* 9, 3 (2013), 20.
- [98] Peter Schuster and Daniel Wessel. 2018. Suzumura consistency, an alternative approach. *J. Appl. Logics – IfCoLog* 5, 1 (2018), 263–286.
- [99] Peter Schuster, Daniel Wessel, and Ihnen Yengui. 2019. Dynamic evaluation of integrity and the computational content of Krull’s lemma. (2019). Preprint.
- [100] Dana Scott. 1971. On engendering an illusion of understanding. *J. Philos.* 68 (1971), 787–807.
- [101] Dana Scott. 1974. Completeness and axiomatizability in many-valued logic. In *Proceedings of the Tarski Symposium (Proc. Sympos. Pure Math., Vol. XXV, Univ. California, Berkeley, Calif., 1971)*, Leon Henkin, John Addison, C.C. Chang, William Craig, Dana Scott, and Robert Vaught (Eds.). Amer. Math. Soc., Providence, RI, 411–435.
- [102] Dana S. Scott. 1973. Background to formalization. In *Truth, syntax and modality (Proc. Conf. Alternative Semantics, Temple Univ., Philadelphia, Pa., 1970)*, Hugues Leblanc (Ed.). North-Holland, Amsterdam, 244–273. Studies in Logic and the Foundations of Math., Vol. 68.
- [103] Joseph R. Shoenfield. 2001. *Mathematical Logic*. Association for Symbolic Logic, Urbana, IL. Reprint of the 1973 second printing.
- [104] D. J. Shoesmith and T. J. Smiley. 1978. *Multiple-Conclusion Logic*. Cambridge University Press, Cambridge.
- [105] Sana Stojanović, Vesna Pavlović, and Predrag Janičić. 2010. A coherent logic based geometry theorem prover capable of producing formal and readable proofs. In *International Workshop on Automated Deduction in Geometry*. Springer, 201–220.
- [106] Edward Szpilrajn. 1930. Sur l’extension de l’ordre partiel. *Fund. Math.* 16 (1930), 368–389.
- [107] Alfred Tarski. 1930. Fundamentale Begriffe der Methodologie der deduktiven Wissenschaften. I. *Monatsh. Math. Phys.* 37 (1930), 361–404.
- [108] Benno van den Berg. 2013. Non-deterministic inductive definitions. *Arch. Math. Logic* 52, 1–2 (2013), 113–135.
- [109] Daniel Wessel. 2019. Ordering groups constructively. *Comm. Algebra* 47, 12 (2019), 4853–4873.
- [110] Daniel Wessel. 2019. Point-free spectra of linear spreads. In *Mathesis Universalis, Computability and Proof*, S. Centrone, S. Negri, D. Sarikaya, and P. Schuster (Eds.). Springer, 353–374.
- [111] Ryszard Wójcicki. 1988. *Theory of logical calculi. Basic theory of consequence operations*. Synthese Library, Vol. 199. Kluwer Academic Publishers Group, Dordrecht. xviii+473 pages.
- [112] Gavin C. Wraith. 1980. Intuitionistic algebra: some recent developments in topos theory. In *Proceedings of the International Congress of Mathematicians (Helsinki, 1978)*. Acad. Sci. Fennica, Helsinki, 331–337.
- [113] Ihnen Yengui. 2015. *Constructive Commutative Algebra. Projective Modules over Polynomial Rings and Dynamical Gröbner Bases*. Lecture Notes in Mathematics, Vol. 2138. Springer, Cham.
- [114] Max Zorn. 1935. A remark on method in transfinite algebra. *Bull. Amer. Math. Soc.* 41 (1935), 667–670.