

Size-Change Abstraction and Max-Plus Automata

Laure Daviaud (Liafa)

joint work with Thomas Colcombet (Liafa)
and Florian Zuleger (TU Vienna)





- Max-plus automata: definition and example
- Asymptotic behaviour of a max-plus automaton
- Application to the computational time complexity of terminating size-change abstraction instances
- Ideas of the main proof

Max-Plus Automata

Max-Plus Automata

Max-plus automaton: Non deterministic finite automaton for which each transition is also **labelled by a non-negative integer** called the weight of the transition.


$$(\mathbb{A}, Q, I, T, E) \text{ with } E \subseteq (Q \times \mathbb{A} \times \mathbb{N} \times Q))$$




Max-Plus Automata

Max-plus automaton: Non deterministic finite automaton for which each transition is also **labelled by a non-negative integer** called the weight of the transition.

$$(\mathbb{A}, Q, I, T, E) \text{ with } E \subseteq (Q \times \mathbb{A} \times \mathbb{N} \times Q)$$




Weight of a run:
sum of the weights of the transitions



Max-Plus Automata

Max-plus automaton: Non deterministic finite automaton for which each transition is also **labelled by a non-negative integer** called the weight of the transition.

$$(\mathbb{A}, Q, I, T, E) \text{ with } E \subseteq (Q \times \mathbb{A} \times \mathbb{N} \times Q)$$




Weight of a run:
sum of the weights of the transitions

Computed function:


$$\mathbb{A}^* \rightarrow \mathbb{N} \cup \{-\infty\}$$

$w \mapsto$ maximum of the weights of the runs labelled by w going from an initial state to a final state
($-\infty$ if no such run)




Max-Plus Automata

Example: $a^m b a^n b a^p$




Max-Plus Automata



Example: $a^m b a^n b a^p$

weight of run (1): 2

Max-Plus Automata




Example: $a^m b a^n b a^p$

weight of run (1): 2

weight of run (2): n

Max-Plus Automata




Example: $a^m b a^n b a^p$

weight of run (1): 2

weight of run (2): n

weight of run (3): p

Max-Plus Automata




Example: $a^m b a^n b a^p$

weight of run (1): 2


weight of run (2): n

weight of run (3): p

weight of run (4): m



Max-Plus Automata



Example: $a^m b a^n b a^p$

weight of run (1): 2


weight of run (2): n

weight of run (3): p

weight of run (4): m

$$a^m b a^n b a^p \mapsto \max(2, m, n, p)$$

Max-Plus Automata



Example: $a^m b a^n b a^p$

weight of run (1): 2


weight of run (2): n

weight of run (3): p

weight of run (4): m

$$a^m b a^n b a^p \mapsto \max(2, m, n, p)$$

Max-Plus Automata



Example: $a^m b a^n b a^p$

weight of run (1): 2


weight of run (2): n

weight of run (3): p

weight of run (4): m

$$a^m b a^n b a^p \mapsto \max(2, m, n, p)$$

Max-Plus Automata



$$a^{n_0} b a^{n_1} b \dots b a^{n_k} \mapsto \max(n_0, n_1, \dots, n_k, k)$$

Asymptotic behaviour of a max-plus automaton

Theorem [Krob, 92 (equivalent to min-plus automata)]

The following problems are undecidable:

Given f and g computed by max-plus automata,

- is $f \leq g$?
- is $f = g$?

\rightsquigarrow Find other ways to look at the behaviour of functions computed by max-plus automata.

Asymptotic behaviour of a max-plus automaton

$$f : a^{n_0} b a^{n_1} b \cdots b a^{n_k} \mapsto \max(n_0, n_1, \dots, n_k, k)$$

Asymptotic behaviour of a max-plus automaton

$$f : a^{n_0} b a^{n_1} b \cdots b a^{n_k} \mapsto \max(n_0, n_1, \dots, n_k, k)$$

- $w_n = a^n \rightsquigarrow f(w_n) = n \rightsquigarrow f(w_n) = |w_n|$

Asymptotic behaviour of a max-plus automaton

$$f : a^{n_0} b a^{n_1} b \cdots b a^{n_k} \mapsto \max(n_0, n_1, \dots, n_k, k)$$

- $w_n = a^n \rightsquigarrow f(w_n) = n \rightsquigarrow f(w_n) = |w_n|$
- $w_n = a^n b a^n \rightsquigarrow f(w_n) = n \rightsquigarrow f(w_n) \sim \frac{|w_n|}{2}$

Asymptotic behaviour of a max-plus automaton


$$f : a^{n_0} b a^{n_1} b \cdots b a^{n_k} \mapsto \max(n_0, n_1, \dots, n_k, k)$$

- $w_n = a^n \rightsquigarrow f(w_n) = n \rightsquigarrow f(w_n) = |w_n|$
- $w_n = a^n b a^n \rightsquigarrow f(w_n) = n \rightsquigarrow f(w_n) \sim \frac{|w_n|}{2}$
- $w_n = (a^n b)^n a^n \rightsquigarrow f(w_n) = n \rightsquigarrow f(w_n) \sim |w_n|^{1/2}$

Asymptotic behaviour of a max-plus automaton

$$f : a^{n_0} b a^{n_1} b \dots b a^{n_k} \mapsto \max(n_0, n_1, \dots, n_k, k)$$

- $w_n = a^n \rightsquigarrow f(w_n) = n \rightsquigarrow f(w_n) = |w_n|$
- $w_n = a^n b a^n \rightsquigarrow f(w_n) = n \rightsquigarrow f(w_n) \sim \frac{|w_n|}{2}$
- $w_n = (a^n b)^n a^n \rightsquigarrow f(w_n) = n \rightsquigarrow f(w_n) \sim |w_n|^{1/2}$



Asymptotic behaviour of a max-plus automaton

$f : \mathbb{A}^* \mapsto \mathbb{N} \cup \{-\infty\}$ computed by a max-plus automaton.

$$\begin{array}{ll} \overline{f} : \mathbb{N} & \mapsto \mathbb{N} \cup \{-\infty\} \\ n & \rightarrow \min\{f(w) \mid |w| = n\} \end{array}$$

Asymptotic behaviour of a max-plus automaton

$f : \mathbb{A}^* \mapsto \mathbb{N} \cup \{-\infty\}$ computed by a max-plus automaton.

$$\begin{aligned} \bar{f} : \mathbb{N} &\mapsto \mathbb{N} \cup \{-\infty\} \\ n &\rightarrow \min \{ f(w) \mid |w| = n \} \end{aligned}$$

Theorem

There exists effectively $\alpha \in (\mathbb{Q} \cap [0, 1]) \cup \{-\infty\}$ such that $\bar{f}(n) = \Theta(n^\alpha)$.

- $\alpha = -\infty$: there is an infinite number of words of weight $-\infty$
- $\alpha = 0$: there is an infinite sequence of words that is bounded
- $\alpha = 1$: all infinite sequences of words are equivalent to the length

Asymptotic behaviour of a max-plus automaton

$f : \mathbb{A}^* \mapsto \mathbb{N} \cup \{-\infty\}$ computed by a max-plus automaton.

$$\begin{aligned} \bar{f} : \mathbb{N} &\mapsto \mathbb{N} \cup \{-\infty\} \\ n &\rightarrow \min \{ f(w) \mid |w| = n \} \end{aligned}$$

Theorem

There exists effectively $\alpha \in (\mathbb{Q} \cap [0, 1]) \cup \{-\infty\}$ such that $\bar{f}(n) = \Theta(n^\alpha)$.

- $\alpha = -\infty$: there is an infinite number of words of weight $-\infty$
- $\alpha = 0$: there is an infinite sequence of words that is bounded
- $\alpha = 1$: all infinite sequences of words are equivalent to the length

\rightsquigarrow Length of the longest word having value at most n : $\Theta(n^{1/\alpha})$.

*Application to the computational time
complexity of terminating size-change
abstraction instances*

Size-Change Abstraction

```
Input x,y:
  while (x>=0){
    y--;
    if (x=0){
      x--;
      y=read_input();
    }
  }
```

Size-Change Abstraction


Input x, y :

```
while (x>=0){  
  y--;  
  if (x=0){  
    x--;  
    y=read_input();  
  }  
}
```

Variables : x, y

Primed versions : x', y'

$t_1: x \geq x', y > y'$



$t_2: x > x'$

Size-Change Abstraction

Input x, y :

```
while (x >= 0) {  
    y--;  
    if (x = 0) {  
        x--;  
        y = read_input();  
    }  
}
```

Variables : x, y
Primed versions : x', y'

$t_1: x \geq x', y > y'$



$t_2: x > x'$

$(5, 5) \xrightarrow{t_1} (5, 4) \xrightarrow{t_2} (4, 8) \xrightarrow{t_1} (3, 6) \xrightarrow{t_1} (3, 2) \xrightarrow{t_2} (2, 10) \dots$

Size-Change Abstraction

Input x, y :

```
while (x>=0){  
    y--;  
    if (x=0){  
        x--;  
        y=read_input();  
    }  
}
```

Variables : x, y
Primed versions : x', y'

$t_1: x \geq x', y > y'$



$t_2: x > x'$

$(5, 5) \xrightarrow{t_1} (5, 4) \xrightarrow{t_2} (4, 8) \xrightarrow{t_1} (3, 6) \xrightarrow{t_1} (3, 2) \xrightarrow{t_2} (2, 10) \dots$

- A finite number of variables that can take values in \mathbb{N} .
- Transitions: conjunction of a finite number of predicates of the form $x_i > x'_j$ or $x_i \geq x'_j$.
- A trace: sequence of transitions and valuations compatible with the transitions.

Size-Change Abstraction

Terminating SCA instance: no infinite trace.

Theorem [Lee, Jones, Ben-Amram]

It is decidable whether a given SCA instance is terminating.


Size-Change Abstraction

Terminating SCA instance: no infinite trace.

Theorem [Lee, Jones, Ben-Amram]

It is decidable whether a given SCA instance is terminating.

$$t_1: x \geq x', y > y'$$



$$t_2: x > x'$$


Size-Change Abstraction

Terminating SCA instance: no infinite trace.

Theorem [Lee, Jones, Ben-Amram]

It is decidable whether a given SCA instance is terminating.

$$t_1: x \geq x', y > y'$$



$$t_2: x > x'$$

Restriction to $[0, n]$: What is the length of the longest trace ?

Theorem

Given a terminating SCA instance, the **longest trace** is of order $\Theta(n^\alpha)$, for some rational number α no smaller than 1.


Moreover, **α is computable**.

Theorem

Given a terminating SCA instance, the **longest trace** is of order $\Theta(n^\alpha)$, for some rational number α no smaller than 1.

Moreover, α is **computable**.

$$t_1: x \geq x', y > y'$$




$$t_2: x > x'$$


$$\begin{aligned} (n, n) &\xrightarrow{t_1} (n, n-1) \xrightarrow{t_1} \dots \xrightarrow{t_1} (n, 0) \\ &\xrightarrow{t_2} (n-1, n) \xrightarrow{t_1} \dots \xrightarrow{t_1} (n-1, 0) \\ &\quad \xrightarrow{t_2} (n-2, n) \xrightarrow{t_1} \dots \\ &\quad \quad \quad \xrightarrow{t_1} (0, 0) \end{aligned}$$

Size-Change Abstraction

$t_1: x \geq x', y > y'$



$t_2: x > x'$





Ideas of the main proof


Asymptotic behaviour of max-plus automata

Given f computed by a max-plus automaton, there exists effectively $\alpha \in (\mathbb{Q} \cap [0, 1]) \cup \{-\infty\}$ such that $\bar{f}(n) = \Theta(n^\alpha)$.


Max-Plus Automata: an algebraic view




Max-Plus Automata: an algebraic view




$$\mu(a) = \begin{pmatrix} 0 & -\infty & -\infty & -\infty \\ -\infty & 0 & -\infty & -\infty \\ -\infty & -\infty & 1 & -\infty \\ -\infty & -\infty & -\infty & 0 \end{pmatrix}$$



Max-Plus Automata: an algebraic view





$$\mu(a) = \begin{pmatrix} 0 & -\infty & -\infty & -\infty \\ -\infty & 0 & -\infty & -\infty \\ -\infty & -\infty & 1 & -\infty \\ -\infty & -\infty & -\infty & 0 \end{pmatrix}$$



$$\mu(b) = \begin{pmatrix} 1 & -\infty & -\infty & -\infty \\ -\infty & 0 & 0 & -\infty \\ -\infty & -\infty & -\infty & 0 \\ -\infty & -\infty & -\infty & 0 \end{pmatrix}$$

Max-Plus Automata: an algebraic view




$$\mu(a) = \begin{pmatrix} 0 & -\infty & -\infty & -\infty \\ -\infty & 0 & -\infty & -\infty \\ -\infty & -\infty & 1 & -\infty \\ -\infty & -\infty & -\infty & 0 \end{pmatrix}$$


$$\mu(b) = \begin{pmatrix} 1 & -\infty & -\infty & -\infty \\ -\infty & 0 & 0 & -\infty \\ -\infty & -\infty & -\infty & 0 \\ -\infty & -\infty & -\infty & 0 \end{pmatrix}$$

Semi-ring $(\mathbb{N} \cup \{-\infty\}, \max, +)$
 $(M \otimes N)_{i,j} = \max_{1 \leq k \leq n} (M_{i,k} + N_{k,j})$

Max-Plus Automata: an algebraic view



$$\mu(a) = \begin{pmatrix} 0 & -\infty & -\infty & -\infty \\ -\infty & 0 & -\infty & -\infty \\ -\infty & -\infty & 1 & -\infty \\ -\infty & -\infty & -\infty & 0 \end{pmatrix}$$




$$\mu(b) = \begin{pmatrix} 1 & -\infty & -\infty & -\infty \\ -\infty & 0 & 0 & -\infty \\ -\infty & -\infty & -\infty & 0 \\ -\infty & -\infty & -\infty & 0 \end{pmatrix}$$


Semi-ring $(\mathbb{N} \cup \{-\infty\}, \max, +)$
 $(M \otimes N)_{i,j} = \max_{1 \leq k \leq n} (M_{i,k} + N_{k,j})$

$$\mu(a_1 a_2 \cdots a_k) = \mu(a_1) \otimes \mu(a_2) \otimes \cdots \otimes \mu(a_k)$$

Max-Plus Automata: an algebraic view



$$\mu(a) = \begin{pmatrix} 0 & -\infty & -\infty & -\infty \\ -\infty & 0 & -\infty & -\infty \\ -\infty & -\infty & 1 & -\infty \\ -\infty & -\infty & -\infty & 0 \end{pmatrix}$$




$$\mu(b) = \begin{pmatrix} 1 & -\infty & -\infty & -\infty \\ -\infty & 0 & 0 & -\infty \\ -\infty & -\infty & -\infty & 0 \\ -\infty & -\infty & -\infty & 0 \end{pmatrix}$$

Semi-ring $(\mathbb{N} \cup \{-\infty\}, \max, +)$
 $(M \otimes N)_{i,j} = \max_{1 \leq k \leq n} (M_{i,k} + N_{k,j})$

$$\mu(a_1 a_2 \cdots a_k) = \mu(a_1) \otimes \mu(a_2) \otimes \cdots \otimes \mu(a_k)$$


$\mu(w)_{i,j}$ is the maximal weight of runs going from i to j labelled by w .

Max-Plus Automata: an algebraic view




$$\mu(a) = \begin{pmatrix} 0 & -\infty & -\infty & -\infty \\ -\infty & 0 & -\infty & -\infty \\ -\infty & -\infty & 1 & -\infty \\ -\infty & -\infty & -\infty & 0 \end{pmatrix}$$

$$\mu(b) = \begin{pmatrix} 1 & -\infty & -\infty & -\infty \\ -\infty & 0 & 0 & -\infty \\ -\infty & -\infty & -\infty & 0 \\ -\infty & -\infty & -\infty & 0 \end{pmatrix}$$




$$I = (0, 0, 0, -\infty) \quad F = \begin{pmatrix} 0 \\ -\infty \\ 0 \\ 0 \end{pmatrix}$$

Max-Plus Automata: an algebraic view



$$\mu(a) = \begin{pmatrix} 0 & -\infty & -\infty & -\infty \\ -\infty & 0 & -\infty & -\infty \\ -\infty & -\infty & 1 & -\infty \\ -\infty & -\infty & -\infty & 0 \end{pmatrix}$$

$$\mu(b) = \begin{pmatrix} 1 & -\infty & -\infty & -\infty \\ -\infty & 0 & 0 & -\infty \\ -\infty & -\infty & -\infty & 0 \\ -\infty & -\infty & -\infty & 0 \end{pmatrix}$$



$$I = (0, 0, 0, -\infty) \quad F = \begin{pmatrix} 0 \\ -\infty \\ 0 \\ 0 \end{pmatrix}$$


$$f(w) = I \otimes \mu(w) \otimes F$$

Ideas for the proof

$f : \mathbb{A}^* \mapsto \mathbb{N} \cup \{-\infty\}$ computed by a max-plus automaton.

$$\begin{aligned} \bar{f} : \mathbb{N} &\mapsto \mathbb{N} \cup \{-\infty\} \\ n &\rightarrow \min \{ f(w) \mid |w| = n \} \end{aligned}$$

\leadsto describe the asymptotic behaviour of infinite sequences of words.



Ideas for the proof

$f : \mathbb{A}^* \mapsto \mathbb{N} \cup \{-\infty\}$ computed by a max-plus automaton.

$$\begin{array}{rcl} \overline{f} : \mathbb{N} & \mapsto & \mathbb{N} \cup \{-\infty\} \\ n & \rightarrow & \min\{f(w) \mid |w| = n\} \end{array}$$

Describe the set $\{(\mu(w), |w|) \mid w \in \mathbb{A}^*\}$:

Ideas for the proof

$f : \mathbb{A}^* \mapsto \mathbb{N} \cup \{-\infty\}$ computed by a max-plus automaton.

$$\begin{array}{rcl} \overline{f} : \mathbb{N} & \mapsto & \mathbb{N} \cup \{-\infty\} \\ n & \rightarrow & \min\{f(w) \mid |w| = n\} \end{array}$$

Describe the set $\{(\mu(w), |w|) \mid w \in \mathbb{A}^*\}$:

- Represent the asymptotic behaviour of infinite sequences of words:
presentable sets.

Ideas for the proof

$f : \mathbb{A}^* \mapsto \mathbb{N} \cup \{-\infty\}$ computed by a max-plus automaton.

$$\begin{array}{rcl} \overline{f} : \mathbb{N} & \mapsto & \mathbb{N} \cup \{-\infty\} \\ n & \rightarrow & \min\{f(w) \mid |w| = n\} \end{array}$$

Describe the set $\{(\mu(w), |w|) \mid w \in \mathbb{A}^*\}$:

- Represent the asymptotic behaviour of infinite sequences of words:
presentable sets.
- **Approximate** the smallest pairs.



Ideas of the proof: presentable sets

Represent the asymptotic behaviour of an infinite sequence of words

$$\left\{ \left(\begin{pmatrix} 0 & -\infty & n^{1/2} & n^{1/3} \\ 0 & 0 & n & 1 \\ n^{3/5} & -\infty & 1 & -\infty \\ -\infty & n^{2/3} & -\infty & 0 \end{pmatrix}, n \right) \mid n \in \mathbb{N} \right\}$$

Ideas of the proof: presentable sets

Represent the asymptotic behaviour of an infinite sequence of words



$$(a^n b)^n a^n$$

$$\begin{pmatrix} n^{1/2} & -\infty & -\infty & -\infty \\ -\infty & 0 & n^{1/2} & n^{1/2} \\ -\infty & -\infty & -\infty & n^{1/2} \\ -\infty & -\infty & -\infty & 0 \end{pmatrix}$$



Ideas of the proof: presentable sets

Represent a convex polyhedra of exponents

$$\left\{ \left(\begin{pmatrix} 0 & -\infty & n^\lambda & n^\mu \\ 0 & 0 & n & 1 \\ n^\nu & -\infty & 1 & -\infty \\ -\infty & n^\eta & -\infty & 0 \end{pmatrix}, n \right) \mid \begin{cases} n \in \mathbb{N} \\ \lambda, \eta, \mu, \nu \in [0, 1] \\ \mu + \eta \geq 1 \\ 5\lambda + 10\nu \geq 8 \end{cases} \right\}$$

Ideas of the proof: presentable sets

Represent a convex polyhedra of exponents




$$(a^*b)^*a^*$$

$$\begin{pmatrix} n^{1-\lambda} & -\infty & -\infty & -\infty \\ -\infty & 0 & 1 & n^\lambda \\ -\infty & -\infty & -\infty & n^\lambda \\ -\infty & -\infty & -\infty & 0 \end{pmatrix}$$


for all $\lambda \in [0, 1]$

$$\begin{pmatrix} 1 & -\infty & -\infty & -\infty \\ -\infty & 0 & n & 1 \\ -\infty & -\infty & -\infty & 1 \\ -\infty & -\infty & -\infty & 0 \end{pmatrix}$$

Ideas of the proof: approximation




Ideas of the proof: approximation



$$(M, \ell) \preceq_a (N, k) \quad \text{if} \quad \begin{aligned} M &\leq aN \\ k &\leq a\ell \\ \tilde{M} &= \tilde{N} \end{aligned}$$


Ideas of the proof: approximation



$$(M, \ell) \preccurlyeq_a (N, k) \quad \text{if} \quad \begin{aligned} M &\leq aN \\ k &\leq a\ell \\ \tilde{M} &= \tilde{N} \end{aligned}$$

$X \preccurlyeq_a Y$ if for all $(N, k) \in Y$, there is $(M, \ell) \in X$ such that $(M, \ell) \preccurlyeq_a (N, k)$

Ideas of the proof: approximation




$$(M, \ell) \preccurlyeq_a (N, k) \quad \text{if} \quad \begin{aligned} M &\leq aN \\ k &\leq a\ell \\ \tilde{M} &= \tilde{N} \end{aligned}$$

$X \preccurlyeq_a Y$ if for all $(N, k) \in Y$, there is $(M, \ell) \in X$ such that $(M, \ell) \preccurlyeq_a (N, k)$

$X \approx_a Y$ if $X \preccurlyeq_a Y$ and $Y \preccurlyeq_a X$

Ideas of the proof: approximation



$$(M, \ell) \preccurlyeq_a (N, k) \quad \text{if} \quad \begin{aligned} M &\leq aN \\ k &\leq a\ell \\ \tilde{M} &= \tilde{N} \end{aligned}$$

$X \preccurlyeq_a Y$ if for all $(N, k) \in Y$, there is $(M, \ell) \in X$ such that $(M, \ell) \preccurlyeq_a (N, k)$

$X \approx_a Y$ if $X \preccurlyeq_a Y$ and $Y \preccurlyeq_a X$

$X \approx Y$ if there is a such that $X \approx_a Y$

Structure of the proof: forest factorization theorem

Approximate by presentable sets:



- the "product" of two presentable sets
- the "closure under product" of "idempotent" presentable sets

↪ use of the forest factorization theorem of Simon:

- start with matrices corresponding to letters
- apply the two previous operations
- after a finite number of steps, we get a presentable set that approximates the set $\{(\mu(w), |w|) \mid w \in \mathbb{A}^*\}$



Conclusion and further questions

- What about min-plus automata?



Conclusion and further questions

- What about min-plus automata?



- Compute the multiplicative coefficient.
(done for min-plus and $|\cdot|$ up to an ε -approximation)