

# Simple strategies for Banach-Mazur games and fairly correct systems

Thomas Brihaye and Quentin Menet

Université de Mons – Belgium

HIGHLIGHTS of Logic, Games and Automata – Paris

# Outline of the talk


- 1 Motivations
- 2 Banach-Mazur games and large sets
- 3 Simple strategies for Banach-Mazur games
- 4 The concept of  $\alpha$ -strategy

# Motivations

## Classical Model-Checking

Given a model  $M$  and a property  $\varphi$ , decide whether :

$M \models \varphi$ , i.e.  $\{\rho \text{ execution of } M \mid \rho \models \varphi\}$  is **empty**.



$M_{\text{coin}} \not\models \mathbf{F} \text{ head}$  ;  $M_{\text{coin}} \not\models \mathbf{GF} \text{ tails}$

# Motivations

## Fair Model-Checking

Given a model  $M$  and a property  $\varphi$ , decide whether :

$M \models \varphi$ , i.e.  $\{\rho \text{ execution of } M \mid \rho \models \varphi\}$  is “**very small**”.

# Motivations

## Fair Model-Checking

Given a model  $M$  and a property  $\varphi$ , decide whether :

$M \models \varphi$ , i.e.  $\{\rho \text{ execution of } M \mid \rho \models \varphi\}$  is “**very small**”.

How to formalise the **fair model-checking**?

- Via probability

$$\begin{aligned} M \models_{\mathbb{P}} \varphi & \text{ iff } \mathbb{P}(\{\rho \text{ of } M \mid \rho \models \varphi\}) = 1 \\ & \text{ iff } \mathbb{P}(\{\rho \text{ of } M \mid \rho \not\models \varphi\}) = 0 \end{aligned}$$

- Via topology

$$\begin{aligned} M \models_T \varphi & \text{ iff } \{\rho \text{ of } M \mid \rho \models \varphi\} \text{ is large} \\ & \text{ iff } \{\rho \text{ of } M \mid \rho \not\models \varphi\} \text{ is meagre} \end{aligned}$$

# Motivations

## A natural question

Given a model  $M$  and property  $\varphi$ , do we have that

$$M \models_{\mathbb{P}} \varphi \quad \Leftrightarrow \quad M \models_T \varphi \quad ???$$

# Motivations

## A natural question

Given a model  $M$  and property  $\varphi$ , do we have that

$$M \models_{\mathbb{P}} \varphi \quad \Leftrightarrow \quad M \models_T \varphi \quad ???$$

In general, the answer is **NO** (e.g. *fat Cantor set*.)

# Motivations

## A natural question

Given a model  $M$  and property  $\varphi$ , do we have that

$$M \models_{\mathbb{P}} \varphi \quad \Leftrightarrow \quad M \models_T \varphi \quad ???$$

In general, the answer is **NO** (e.g. *fat Cantor set*.)

## Theorem [VV06]

Given a finite system  $M$  and an  $\omega$ -regular property  $\varphi$ , we have that

$$M \models_{\mathbb{P}} \varphi \quad \Leftrightarrow \quad M \models_T \varphi,$$

for bounded Borel measures.



# Motivations

## A natural question

Given a model  $M$  and property  $\varphi$ , do we have that

$$M \models_{\mathbb{P}} \varphi \quad \Leftrightarrow \quad M \models_T \varphi \quad ???$$

In general, the answer is **NO** (e.g. *fat Cantor set*.)

## Theorem [VV06]

Given a finite system  $M$  and an  $\omega$ -regular property  $\varphi$ , we have that

$$M \models_{\mathbb{P}} \varphi \quad \Leftrightarrow \quad M \models_T \varphi,$$

for bounded Borel measures.

Can we go further?

# Outline of the talk

- 1 Motivations
- 2 Banach-Mazur games and large sets
- 3 Simple strategies for Banach-Mazur games
- 4 The concept of  $\alpha$ -strategy

# Banach-Mazur games

## Definition

A *Banach-Mazur game*  $\mathcal{G}$  on a finite graph is a triplet  $(G, v_0, W)$  where

- $G = (V, E)$  is a finite directed graph with no deadlock,
- $v_0 \in V$  is the initial state,
- $W \subset V^\omega$ .

Given  $(G, v_0, W)$ , **Pl. 0** and **Pl. 1** play as follows :

# Banach-Mazur games

## Definition

A *Banach-Mazur game*  $\mathcal{G}$  on a finite graph is a triplet  $(G, v_0, W)$  where

- $G = (V, E)$  is a finite directed graph with no deadlock,
- $v_0 \in V$  is the initial state,
- $W \subset V^\omega$ .

Given  $(G, v_0, W)$ , **Pl. 0** and **Pl. 1** play as follows :

- **Pl. 1** begins with choosing a finite path  $\rho_1$  starting in  $v_0$  ;

# Banach-Mazur games

## Definition

A *Banach-Mazur game*  $\mathcal{G}$  on a finite graph is a triplet  $(G, v_0, W)$  where

- $G = (V, E)$  is a finite directed graph with no deadlock,
- $v_0 \in V$  is the initial state,
- $W \subset V^\omega$ .

Given  $(G, v_0, W)$ , **Pl. 0** and **Pl. 1** play as follows :

- **Pl. 1** begins with choosing a finite path  $\rho_1$  starting in  $v_0$  ;
- **Pl. 0** prolongs  $\rho_1$  by choosing another finite path  $\rho_2$  ;

# Banach-Mazur games

## Definition

A *Banach-Mazur game*  $\mathcal{G}$  on a finite graph is a triplet  $(G, v_0, W)$  where

- $G = (V, E)$  is a finite directed graph with no deadlock,
- $v_0 \in V$  is the initial state,
- $W \subset V^\omega$ .

Given  $(G, v_0, W)$ , **Pl. 0** and **Pl. 1** play as follows :

- **Pl. 1** begins with choosing a finite path  $\rho_1$  starting in  $v_0$  ;
- **Pl. 0** prolongs  $\rho_1$  by choosing another finite path  $\rho_2$  ;
- **Pl. 1** prolongs  $\rho_1\rho_2$  by choosing another finite path  $\rho_3$  ;

# Banach-Mazur games

## Definition

A *Banach-Mazur game*  $\mathcal{G}$  on a finite graph is a triplet  $(G, v_0, W)$  where

- $G = (V, E)$  is a finite directed graph with no deadlock,
- $v_0 \in V$  is the initial state,
- $W \subset V^\omega$ .

Given  $(G, v_0, W)$ , **Pl. 0** and **Pl. 1** play as follows :

- **Pl. 1** begins with choosing a finite path  $\rho_1$  starting in  $v_0$ ;
- **Pl. 0** prolongs  $\rho_1$  by choosing another finite path  $\rho_2$ ;
- **Pl. 1** prolongs  $\rho_1\rho_2$  by choosing another finite path  $\rho_3$ ;
- ...

# Banach-Mazur games

## Definition

A *Banach-Mazur game*  $\mathcal{G}$  on a finite graph is a triplet  $(G, v_0, W)$  where

- $G = (V, E)$  is a finite directed graph with no deadlock,
- $v_0 \in V$  is the initial state,
- $W \subset V^\omega$ .

Given  $(G, v_0, W)$ , **Pl. 0** and **Pl. 1** play as follows :

- **Pl. 1** begins with choosing a finite path  $\rho_1$  starting in  $v_0$ ;
- **Pl. 0** prolongs  $\rho_1$  by choosing another finite path  $\rho_2$ ;
- **Pl. 1** prolongs  $\rho_1\rho_2$  by choosing another finite path  $\rho_3$ ;
- ...

A play  $\rho = \rho_1\rho_2\rho_3 \cdots$  is won by **Pl. 0** wins iff  $\rho \in W$ .



# Strategies for Banach-Mazur games

Let  $G = (V, E)$  be a graph and  $\mathcal{G} = (G, v_0, W)$  be a Banach-Mazur game.


## Definition

A strategy for **Pl. 0** is a function  $f : V^* \rightarrow V^*$  such that whenever

$$f(\rho) = \rho'$$

we have that  $\rho'$  prolongs  $\rho$  in  $G$ .

$\rho_1$



# Strategies for Banach-Mazur games

Let  $G = (V, E)$  be a graph and  $\mathcal{G} = (G, v_0, W)$  be a Banach-Mazur game.

## Definition


A strategy for [Pl. 0](#) is a function  $f : V^* \rightarrow V^*$  such that whenever

$$f(\rho) = \rho'$$

we have that  $\rho'$  prolongs  $\rho$  in  $G$ .

$\rho_1 \rho_2$

$$f(\rho_1) = \rho_2$$



# Strategies for Banach-Mazur games

Let  $G = (V, E)$  be a graph and  $\mathcal{G} = (G, v_0, W)$  be a Banach-Mazur game.

## Definition


A strategy for [Pl. 0](#) is a function  $f : V^* \rightarrow V^*$  such that whenever

$$f(\rho) = \rho'$$

we have that  $\rho'$  prolongs  $\rho$  in  $G$ .

$\rho_1 \rho_2 \rho_3$

$$f(\rho_1) = \rho_2$$



# Strategies for Banach-Mazur games

Let  $G = (V, E)$  be a graph and  $\mathcal{G} = (G, v_0, W)$  be a Banach-Mazur game.

## Definition

A strategy for **Pl. 0** is a function  $f : V^* \rightarrow V^*$  such that whenever


$$f(\rho) = \rho'$$

we have that  $\rho'$  prolongs  $\rho$  in  $G$ .

$\rho_1 \rho_2 \rho_3 \rho_4$

$$f(\rho_1) = \rho_2$$

$$f(\rho_1 \rho_2 \rho_3) = \rho_4$$



# Strategies for Banach-Mazur games

Let  $G = (V, E)$  be a graph and  $\mathcal{G} = (G, v_0, W)$  be a Banach-Mazur game.

## Definition

A strategy for **Pl. 0** is a function  $f : V^* \rightarrow V^*$  such that whenever


$$f(\rho) = \rho'$$

we have that  $\rho'$  prolongs  $\rho$  in  $G$ .

$\rho_1 \rho_2 \rho_3 \rho_4 \rho_5$

$$f(\rho_1) = \rho_2$$

$$f(\rho_1 \rho_2 \rho_3) = \rho_4$$



# Strategies for Banach-Mazur games

Let  $G = (V, E)$  be a graph and  $\mathcal{G} = (G, v_0, W)$  be a Banach-Mazur game.

## Definition

A strategy for **Pl. 0** is a function  $f : V^* \rightarrow V^*$  such that whenever

$$f(\rho) = \rho'$$


we have that  $\rho'$  prolongs  $\rho$  in  $G$ .

$\rho_1 \rho_2 \rho_3 \rho_4 \rho_5 \rho_6$

$$f(\rho_1) = \rho_2$$

$$f(\rho_1 \rho_2 \rho_3) = \rho_4$$

$$f(\rho_1 \rho_2 \rho_3 \rho_4 \rho_5) = \rho_6$$



# Strategies for Banach-Mazur games

Let  $G = (V, E)$  be a graph and  $\mathcal{G} = (G, v_0, W)$  be a Banach-Mazur game.

## Definition

A strategy for **Pl. 0** is a function  $f : V^* \rightarrow V^*$  such that whenever

$$f(\rho) = \rho'$$


we have that  $\rho'$  prolongs  $\rho$  in  $G$ .

$\rho_1 \rho_2 \rho_3 \rho_4 \rho_5 \rho_6 \rho_7$

$$f(\rho_1) = \rho_2$$

$$f(\rho_1 \rho_2 \rho_3) = \rho_4$$

$$f(\rho_1 \rho_2 \rho_3 \rho_4 \rho_5) = \rho_6$$







# Banach-Mazur games and large sets

Let  $(V, E)$  be a graph. We consider  $V^\omega$  equipped with the Cantor topology.

## Definitions

A set  $S \subseteq V^\omega$  is said

- *nowhere dense* if the closure of  $S$  has empty interior.
- *meagre* if it can be seen as a countable union of nowhere dense sets.
- *large* if  $S^c$  is meagre.

## Theorem [Oxtoby57]

Let  $\mathcal{G} = (G, v_0, W)$  be a Banach-Mazur game on a finite graph.

Pl. 0 has a winning strategy for  $\mathcal{G}$  if and only if  $W$  is large.

# Outline of the talk

- 1 Motivations
- 2 Banach-Mazur games and large sets
- 3 Simple strategies for Banach-Mazur games
- 4 The concept of  $\alpha$ -strategy

## Back to our motivation

### Theorem [VV06]

Given a finite system  $M$  and an  $\omega$ -regular property  $\varphi$ , we have that

$$M \models_{\mathbb{P}} \varphi \iff M \models_T \varphi,$$

for bounded Borel measures.

The key ingredient to prove the above result is the following result :

### Theorem [BGK03]

Given  $\mathcal{G} = (G, v_0, W)$  where  $W$  is an  $\omega$ -regular property, we have that

Pl. 0 has a winning strategy for  $\mathcal{G}$

iff

Pl. 0 has a **positional** winning strategies for  $\mathcal{G}$ .

# Simple strategies for Banach-Mazur game

Given  $\mathcal{G} = (G, v_0, W)$ , let  $f : V^* \rightarrow V^*$  be a strategy for Pl. 0.

$$f(\underbrace{\rho_1 \rho_2 \cdots \rho_{2n+1}}_{\text{What is observed}}) = \underbrace{\rho_{2n+2}}_{\text{What is played}}$$

We say that  $f$  is

# Simple strategies for Banach-Mazur game

Given  $\mathcal{G} = (G, v_0, W)$ , let  $f : V^* \rightarrow V^*$  be a strategy for Pl. 0.

$$\underbrace{f(\rho_1 \rho_2 \cdots \rho_{2n+1})}_{\text{What is observed}} = \underbrace{\rho_{2n+2}}_{\text{What is played}}$$

We say that  $f$  is

- *positional* if it only depends on  $\text{Last}(\rho_{2n+1})$ .

# Simple strategies for Banach-Mazur game

Given  $\mathcal{G} = (G, v_0, W)$ , let  $f : V^* \rightarrow V^*$  be a strategy for Pl. 0.

$$\underbrace{f(\rho_1 \rho_2 \cdots \rho_{2n+1})}_{\text{What is observed}} = \underbrace{\rho_{2n+2}}_{\text{What is played}}$$

We say that  $f$  is

- *positional* if it only depends on  $\text{Last}(\rho_{2n+1})$ .
- *finite memory* if it only depends on  $\text{Last}(\rho_{2n+1})$  and a finite memory.

## Simple strategies for Banach-Mazur game

Given  $\mathcal{G} = (G, v_0, W)$ , let  $f : V^* \rightarrow V^*$  be a strategy for Pl. 0.

$$\underbrace{f(\rho_1 \rho_2 \cdots \rho_{2n+1})}_{\text{What is observed}} = \underbrace{\rho_{2n+2}}_{\text{What is played}}$$

We say that  $f$  is

- *positional* if it only depends on  $\text{Last}(\rho_{2n+1})$ .
- *finite memory* if it only depends on  $\text{Last}(\rho_{2n+1})$  and a finite memory.
- *b-bounded* if  $|\rho_{2n+2}| \leq b$ .

## Simple strategies for Banach-Mazur game

Given  $\mathcal{G} = (G, v_0, W)$ , let  $f : V^* \rightarrow V^*$  be a strategy for Pl. 0.

$$\underbrace{f(\rho_1 \rho_2 \cdots \rho_{2n+1})}_{\text{What is observed}} = \underbrace{\rho_{2n+2}}_{\text{What is played}}$$

We say that  $f$  is

- *positional* if it only depends on  $\text{Last}(\rho_{2n+1})$ .
- *finite memory* if it only depends on  $\text{Last}(\rho_{2n+1})$  and a finite memory.
- *b-bounded* if  $|\rho_{2n+2}| \leq b$ .
- *bounded* if there is  $b \geq 1$  such that  $f$  is b-bounded.



## Simple strategies for Banach-Mazur game

Given  $\mathcal{G} = (G, v_0, W)$ , let  $f : V^* \rightarrow V^*$  be a strategy for Pl. 0.

$$\underbrace{f(\rho_1 \rho_2 \cdots \rho_{2n+1})}_{\text{What is observed}} = \underbrace{\rho_{2n+2}}_{\text{What is played}}$$

We say that  $f$  is

- *positional* if it only depends on  $\text{Last}(\rho_{2n+1})$ .
- *finite memory* if it only depends on  $\text{Last}(\rho_{2n+1})$  and a finite memory.
- *b-bounded* if  $|\rho_{2n+2}| \leq b$ .
- *bounded* if there is  $b \geq 1$  such that  $f$  is  $b$ -bounded.
- *move-counting* if it only depends on  $\text{Last}(\rho_{2n+1})$  and the number of moves already played.

# Simple strategies for Banach-Mazur game


Given  $\mathcal{G} = (G, v_0, W)$ , let  $f : V^* \rightarrow V^*$  be a strategy for Pl. 0.

$$\underbrace{f(\rho_1 \rho_2 \cdots \rho_{2n+1})}_{\text{What is observed}} = \underbrace{\rho_{2n+2}}_{\text{What is played}}$$

We say that  $f$  is

- *positional* if it only depends on  $\text{Last}(\rho_{2n+1})$ .
- *finite memory* if it only depends on  $\text{Last}(\rho_{2n+1})$  and a finite memory.
- *b-bounded* if  $|\rho_{2n+2}| \leq b$ .
- *bounded* if there is  $b \geq 1$  such that  $f$  is  $b$ -bounded.
- *move-counting* if it only depends on  $\text{Last}(\rho_{2n+1})$  and the number of moves already played.
- *length-counting* if it only depends on the  $\text{Last}(\rho_{2n+1})$  and the length of the prefix already played.

# Simple strategies for Pl. 0 on finite graphs



Combining simple observations and results from [BGK03], [VV06], [GL12], [BM13]

# Relations with the sets of probability one

## Proposition

Let  $\mathcal{G} = (G, v_0, W)$  be a Banach-Mazur game on a finite graph and  $\mathbb{P}$  a reasonable probability measure.

If Pl. 0 has  $\begin{cases} \text{a move-counting} \\ \text{a bounded} \end{cases}$  winning strategy for  $\mathcal{G}$ , then  $\mathbb{P}(W) = 1$ .

## Relations with the sets of probability one

### Proposition

Let  $\mathcal{G} = (G, v_0, W)$  be a Banach-Mazur game on a finite graph and  $\mathbb{P}$  a reasonable probability measure.

If Pl. 0 has  $\begin{cases} \text{a move-counting} \\ \text{a bounded} \end{cases}$  winning strategy for  $\mathcal{G}$ , then  $\mathbb{P}(W) = 1$ .

There exist large **open** set of probability 1 without a positional/ bounded/ move-counting winning strategy.


We look for a new concept of “simple strategy”

# Outline of the talk

- 1 Motivations
- 2 Banach-Mazur games and large sets
- 3 Simple strategies for Banach-Mazur games
- 4 The concept of  $\alpha$ -strategy


# Banach-Mazur game

A play consists in concatenating **finite paths**,



# Banach-Mazur game


A play consists in concatenating **finite paths**,  
or equivalently in building a decreasing sequence of **open sets**.






# Banach-Mazur game

A play consists in concatenating **finite paths**,  
or equivalently in building a decreasing sequence of **open sets**.




## Banach-Mazur game

A play consists in concatenating **finite paths**,  
or equivalently in building a decreasing sequence of **open sets**.




## Banach-Mazur game

A play consists in concatenating **finite paths**,  
or equivalently in building a decreasing sequence of **open sets**.




## Banach-Mazur game

A play consists in concatenating **finite paths**,  
or equivalently in building a decreasing sequence of **open sets**.




## Banach-Mazur game

A play consists in concatenating **finite paths**,  
or equivalently in building a decreasing sequence of **open sets**.




# Banach-Mazur game

A play consists in concatenating **finite paths**,  
or equivalently in building a decreasing sequence of **open sets**.



# Banach-Mazur game

A play consists in concatenating **finite paths**,  
or equivalently in building a decreasing sequence of **open sets**.



## Another simple strategy for Banach-Mazur game

Given  $\mathcal{G} = (G, v_0, W)$ , a strategy for Pl. 0 can be seen as  $f : \mathcal{O} \rightarrow \mathcal{O}$ .

$$f(\underbrace{O_1 O_2 \cdots O_{2n+1}}_{\text{What is observed}}) = \underbrace{O_{2n+2}}_{\text{What is played}},$$

where  $O_1 \supseteq O_2 \supseteq \cdots \supseteq O_{2n+1} \supseteq O_{2n+2}$  are open sets.



## Another simple strategy for Banach-Mazur game

Given  $\mathcal{G} = (G, v_0, W)$ , a strategy for Pl. 0 can be seen as  $f : \mathcal{O} \rightarrow \mathcal{O}$ .

$$\underbrace{f(\underbrace{O_1 O_2 \cdots O_{2n+1}}_{\text{What is observed}})}_{\text{What is played}} = \underbrace{O_{2n+2}}_{\text{What is played}},$$

where  $O_1 \supseteq O_2 \supseteq \cdots \supseteq O_{2n+1} \supseteq O_{2n+2}$  are open sets.

Assuming that  $G$  is equipped with a probability distribution on edges.

### The notion of $\alpha$ -strategy

Given  $0 < \alpha < 1$ , we say that  $f$  is an  $\alpha$ -strategy if and only if

$$\mathbb{P}(O_{2n+2} | O_{2n+1}) \geq \alpha.$$

# Our results

## Theorem

Let  $\mathcal{G} = (G, v_0, W)$  be a Banach-Mazur game on a finite graph and  $\mathbb{P}$  a reasonable probability measure.

If Pl. 0 has a winning  $\alpha$ -strategy for some  $\alpha > 0$ , then  $\mathbb{P}(W) = 1$ .

# Our results

## Theorem

Let  $\mathcal{G} = (G, v_0, W)$  be a Banach-Mazur game on a finite graph and  $\mathbb{P}$  a reasonable probability measure.

If Pl. 0 has a winning  $\alpha$ -strategy for some  $\alpha > 0$ , then  $\mathbb{P}(W) = 1$ .

## Theorem

When  $W$  is a **countable intersection of open sets**, the following assertions are equivalent :

- 1  $P(W) = 1$ ,
- 2 Pl. 0 has a winning  $\alpha$ -strategy for some  $\alpha > 0$ ,
- 3 Pl. 0 has a winning  $\alpha$ -strategy for all  $0 < \alpha < 1$ .

# Summary



Thank you !!!