

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/265463977>

# Linear recurrent sequences and powers of a square matrix

Article · January 2006

CITATIONS

41

READS

419

2 authors:



[Hacène Belbachir](#)

University of Science and Technology Houari Boumediene

157 PUBLICATIONS 888 CITATIONS

[SEE PROFILE](#)



[Farid Bencherif](#)

University of Science and Technology Houari Boumediene

29 PUBLICATIONS 124 CITATIONS

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:



on a constant related to prime counting function [View project](#)



Matrix functions and linear recurrent sequences [View project](#)

# LINEAR RECURRENT SEQUENCES AND POWERS OF A SQUARE MATRIX

Hacène Belbachir<sup>1</sup>

USTHB, Department of Mathematics, P.O.Box 32 El Alia, 16111, Bab Ezzouar, Algiers, Algeria  
hbelbachir@usthb.dz and hacenebelbachir@gmail.com

Farid Bencherif<sup>2</sup>

USTHB, Department of Mathematics, P.O.Box 32 El Alia, 16111, Bab Ezzouar, Algiers, Algeria  
fbencherif@usthb.dz and fbencherif@yahoo.fr

Received: 8/10/05, Revised: 3/20/06, Accepted: 4/4/06, Published: 4/24/06

## Abstract

In this paper, we establish a formula expressing explicitly the general term of a linear recurrent sequence, allowing us to generalize the original result of J. McLaughlin [7] concerning powers of a matrix of size 2, to the case of a square matrix of size  $m \geq 2$ . Identities concerning Fibonacci and Stirling numbers and various combinatorial relations are derived.

## 1. Introduction

The main theorem of J. McLaughlin [7] states the following:

**Theorem 1.** Let  $A = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$  be a square matrix of order two, let  $T = a + d$  be its trace, and let  $D = ad - bc$  be its determinant. Let

$$y_n = \sum_{i=0}^{\lfloor n/2 \rfloor} \binom{n-i}{i} T^{n-2i} (-D)^i. \quad (1)$$

Then, for  $n \geq 1$ ,

$$A^n = \begin{pmatrix} y_n - dy_{n-1} & by_{n-1} \\ cy_{n-1} & y_n - ay_{n-1} \end{pmatrix}. \quad (2)$$

---

<sup>1</sup>Partially supported by the laboratory LAID3.

<sup>2</sup>Partially supported by the laboratory LATN.

We remark that in this theorem,  $(y_n)_{n \geq -1}$  is a linear recurrent sequence that satisfies

$$\begin{cases} y_{-1} = 0, \\ y_0 = 1, \\ y_n = Ty_{n-1} - Dy_{n-2} \text{ for all integer } n \geq 1. \end{cases} \quad (3)$$

By setting  $A_0 = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$  and  $A_1 = \begin{pmatrix} -d & b \\ c & -a \end{pmatrix}$ , relation (2) of Theorem 1 may be written as follows:

$$A^n = y_n A_0 + y_{n-1} A_1 \text{ for all integers } n \geq 0, \quad (4)$$

with  $A_0 = I_2$  and  $A_1 = A - TI_2$  (where  $I_2$  is the unit matrix).

In Section 3, we extend this result (relation (4)) to any matrix  $A \in \mathcal{M}_m(\mathcal{A})$  of order  $m \geq 2$ ,  $\mathcal{A}$  being a unitary commutative ring.

We prove the following result:

Let  $A \in M_m(\mathcal{A})$  and let  $P(X) = X^m - a_1 X^{m-1} - \dots - a_{m-1} X - a_m$  be the characteristic polynomial of  $A$ . Let  $A_0, A_1, \dots, A_{m-1}$  be matrices of  $M_m(\mathcal{A})$  defined by

$$A_k = - \sum_{i=0}^k a_i A^{k-i}, \text{ for } 0 \leq k \leq m-1, \text{ with } a_0 = -1,$$

and let  $(y_n)_{n > -m}$  be the sequence of elements of  $\mathcal{A}$  satisfying

$$y_n = \sum_{k_1+2k_2+\dots+mk_m=n} \binom{k_1+k_2+\dots+k_m}{k_1, k_2, \dots, k_m} a_1^{k_1} a_2^{k_2} \dots a_m^{k_m}, \text{ for } n > -m.$$

Then, for all integers  $n \geq 0$ ,  $A^n = y_n A_0 + y_{n-1} A_1 + \dots + y_{n-m+1} A_{m-1}$ .

The proof of this result is based on Theorem 2 given in Section 2.

In this section, we generalize Theorem 2, which permits us to express the general term  $u_n$  of a recurrent linear sequence satisfying the relation

$$u_n = a_1 u_{n-1} + a_2 u_{n-2} + \dots + a_m u_{n-m} \text{ for all } n \geq 1,$$

in terms of the coefficients  $a_1, a_2, \dots, a_m$  and  $u_0, u_{-1}, u_{-2}, \dots, u_{-(m-1)}$ . Applications to Fibonacci, generalized Fibonacci and “multibonacci” sequences are also given.

Finally, in Section 4, further combinatorial identities are derived, including identities concerning the Stirling numbers of the first and second kind.

As an illustration, we give a nice duality between the two following relations (Corollaries 5 and 7):

$$\sum_{k_1+2k_2+\dots+mk_m=n} \binom{k_1+\dots+k_m}{k_1, \dots, k_m} (-1)^{n-(\sum_{i=1}^m k_i)} \binom{m}{m-1}^{k_1} \dots \binom{m}{0}^{k_m} = \binom{n+m-1}{m-1},$$

$$\sum_{k_1+2k_2+\dots+mk_m=n} \binom{k_1+\dots+k_m}{k_1,\dots,k_m} (-1)^{n-(\sum_{i=1}^m k_i)} \begin{bmatrix} m \\ m-1 \end{bmatrix}^{k_1} \cdots \begin{bmatrix} m \\ 0 \end{bmatrix}^{k_m} = \begin{Bmatrix} n+m-1 \\ m-1 \end{Bmatrix},$$
 where  $\binom{k_1+\dots+k_m}{k_1,\dots,k_m}$  is the multinomial coefficient (Section 2), and  $\begin{bmatrix} n \\ k \end{bmatrix}$  and  $\begin{Bmatrix} n \\ k \end{Bmatrix}$  are, respectively, the Stirling numbers of the first and second kind as defined in [5].

## 2. Explicit Expression of the General Term of a Recurrent Linear Sequence

In this section, we let  $m \geq 2$  be an integer,  $\mathcal{A}$  a unitary commutative ring,  $a_1, a_2, \dots, a_m, \alpha_0, \alpha_1, \dots, \alpha_{m-1}$  elements of  $\mathcal{A}$ , and  $(u_n)_{n \geq -m}$  a sequence of elements of  $\mathcal{A}$  defined by

$$\begin{cases} u_{-j} = \alpha_j & \text{for } 0 \leq j \leq m-1, \\ u_n = a_1 u_{n-1} + a_2 u_{n-2} + \dots + a_m u_{n-m} & \text{for } n \geq 1. \end{cases} \quad (5)$$

The aim of this section is to give an explicit expression of  $u_n$  in terms of  $n, a_1, a_2, \dots, a_m, \alpha_0, \alpha_1, \dots, \alpha_{m-1}$  (Theorem 3).

Let us define the sequence  $(y_n)_{n \in \mathbb{Z}}$  of elements of  $\mathcal{A}$ , with the convention that an empty sum is zero, by

$$y_n = \sum_{k_1+2k_2+\dots+mk_m=n} \binom{k_1+k_2+\dots+k_m}{k_1,k_2,\dots,k_m} a_1^{k_1} a_2^{k_2} \dots a_m^{k_m}, \quad \text{for } n \in \mathbb{Z}, \quad (6)$$

the summation being taken over all  $m$ -tuples  $(k_1, k_2, \dots, k_m)$  of integers  $k_j \geq 0$  satisfying  $k_1 + 2k_2 + \dots + mk_m = n$ . With the previous convention, we have  $y_n = 0$  for  $n < 0$ . The multinomial coefficient that appears in the summation is defined for integers  $k_1, k_2, \dots, k_m \geq 0$ , by

$$\binom{k_1+k_2+\dots+k_m}{k_1,k_2,\dots,k_m} = \frac{(k_1+k_2+\dots+k_m)!}{k_1!k_2!\dots k_m!},$$

and can always be written as a product of binomial coefficients

$$\binom{k_1+k_2+\dots+k_m}{k_1,k_2,\dots,k_m} = \binom{k_1+k_2+\dots+k_m}{k_1+k_2+\dots+k_{m-1}} \cdots \binom{k_1+k_2+k_3}{k_1+k_2} \binom{k_1+k_2}{k_1}.$$

Let us adopt the following convention. For  $k_1 + k_2 + \dots + k_m \geq 1$ , we put

$$\binom{k_1+k_2+\dots+(k_j-1)+\dots+k_m}{k_1,k_2,\dots,k_j-1,\dots,k_m} = 0 \quad \text{when } k_j = 0,$$

for any  $j \in \{1, 2, \dots, m\}$ . We can now state the following lemma [p. 80 (Vol. 1), 4].

**Lemma 1.** *Let  $k_j \geq 0$  be integers for  $j \in \{1, 2, \dots, m\}$ , such that  $k_1 + \dots + k_m \geq 1$ . Then*

$$\binom{k_1+k_2+\dots+k_m}{k_1,k_2,\dots,k_m} = \sum_{j=1}^m \binom{k_1+\dots+(k_j-1)+\dots+k_m}{k_1,\dots,k_j-1,\dots,k_m}.$$

This lemma permits us to easily prove the following result.

**Lemma 2.** *The sequence  $(y_n)_{n \in \mathbb{Z}}$ , defined by relation (6), satisfies the recurrence relation*

$$y_n = a_1 y_{n-1} + a_2 y_{n-2} + \cdots + a_m y_{n-m} \quad \text{for } n \geq 1$$

with  $y_0 = 1$  and  $y_n = 0$  for  $n < 0$ .

*Proof.* First notice that, for  $n \geq 1$ , for all  $j \in \{1, 2, \dots, m\}$  we have

$$a_j y_{n-j} = \sum_{k_1 + 2k_2 + \cdots + mk_m = n} \binom{k_1 + \cdots + (k_j - 1) + \cdots + k_m}{k_1, \dots, k_j - 1, \dots, k_m} a_1^{k_1} \cdots a_j^{k_j} \cdots a_m^{k_m}.$$

Applying Lemma 1, we obtain  $\sum_{j=1}^m a_j y_{n-j} = y_n$ . The relations  $y_0 = 1$  and  $y_n = 0$  for  $n < 0$  follow immediately.  $\square$

We can now state the following result.

**Theorem 2.** *Let  $(u_n)_{n > -m}$  the sequence of elements of  $\mathcal{A}$  defined by*

$$\begin{cases} u_{-j} = 0 & \text{for } 1 \leq j \leq m-1, \\ u_0 = 1, \\ u_n = a_1 u_{n-1} + a_2 u_{n-2} + \cdots + a_m u_{n-m} & \text{for } n \geq 1. \end{cases}$$

Then for all integers  $n > -m$ ,

$$u_n = \sum_{k_1 + 2k_2 + \cdots + mk_m = n} \binom{k_1 + k_2 + \cdots + k_m}{k_1, k_2, \dots, k_m} a_1^{k_1} a_2^{k_2} \cdots a_m^{k_m}.$$

**Corollary 1.** *Let  $q \geq 1$  be an integer,  $a, b \in \mathcal{A}$ , and let  $(v_n)_{n \geq -q}$  be the sequence of elements of  $\mathcal{A}$  defined by*

$$\begin{cases} v_{-j} = 0 & \text{for } 1 \leq j \leq q, \\ v_0 = 1, \\ v_{n+1} = av_n + bv_{n-q} & \text{for } n \geq 0. \end{cases} \quad (7)$$

Then, for all  $n \geq -q$ ,

$$v_n = \sum_{k=0}^{\lfloor \frac{n}{q+1} \rfloor} \binom{n - kq}{k} a^{n-k(q+1)} b^k, \quad (8)$$

and, for all  $n \geq 0$ ,

$$v_{n+1} + bv_{n-q} = 2v_{n+1} - av_n = \sum_{k=0}^{\lfloor \frac{n+1}{q+1} \rfloor} \frac{n+1-k(q-1)}{n+1-kq} \binom{n+1-kq}{k} a^{n+1-k(q+1)} b^k. \quad (9)$$

*Proof.* We deduce relation (8) directly from Theorem 2, with  $m = q + 1$ ,  $a_1 = a$ ,  $a_m = b$  and  $a_k = 0$  for  $1 < k < m$ . From (8), we deduce (9) as follows:

$$\begin{aligned} v_{n+1} + bv_{n-q} &= \sum_{k=0}^{\lfloor \frac{n+1}{q+1} \rfloor} \binom{n+1-kq}{k} a^{n+1-k(q+1)} b^k + \sum_{k=0}^{\lfloor \frac{n-q}{q+1} \rfloor} \binom{n-(k+1)q}{k} a^{n-q-(q+1)k} b^{k+1} \\ &= \sum_{k=0}^{\lfloor \frac{n+1}{q+1} \rfloor} \binom{n+1-kq}{k} a^{n+1-k(q+1)} b^k + \sum_{k=1}^{\lfloor \frac{n+1}{q+1} \rfloor} \binom{n-kq}{k-1} a^{n+1-k(q+1)} b^k \\ &= \sum_{k=0}^{\lfloor \frac{n+1}{q+1} \rfloor} \binom{n+1-kq}{k} a^{n+1-k(q+1)} b^k + \sum_{k=0}^{\lfloor \frac{n+1}{q+1} \rfloor} \frac{k}{n+1-kq} \binom{n+1-kq}{k} a^{n+1-k(q+1)} b^k \\ &= \sum_{k=0}^{\lfloor \frac{n+1}{q+1} \rfloor} \frac{n+1-k(q-1)}{n+1-kq} \binom{n+1-kq}{k} a^{n+1-k(q+1)} b^k. \end{aligned}$$

□

We now give some applications of the above corollary.

**Application 1.** Let  $(F_n)_{n \geq 0}$  be the Fibonacci sequence

$$\begin{cases} F_0 = 0, \\ F_1 = 1, \\ F_{n+1} = F_n + F_{n-1} \quad \text{for } n \geq 1. \end{cases}$$

Then, by setting  $\varphi_n = F_{n+1}$  for  $n \geq -1$ , we see that  $(\varphi_n)_{n \geq -1}$  is also defined by

$$\begin{cases} \varphi_{-1} = 0, \\ \varphi_0 = 1, \\ \varphi_n = \varphi_{n-1} + \varphi_{n-2} \quad \text{for } n \geq 1. \end{cases}$$

The application of Corollary 1 gives us that

$$\varphi_n = F_{n+1} = \sum_{k=0}^{\lfloor n/2 \rfloor} \binom{n-k}{k}, \quad \text{for } n \geq -1,$$

the relation given in [pp. 18-20, 12], and announced in [9]. Also

$$F_n + F_{n+2} = \sum_{k=0}^{\lfloor \frac{n+1}{2} \rfloor} \frac{n+1}{n+1-k} \binom{n+1-k}{k}, \quad \text{for } n \geq 0,$$

and we find the relations given in Problem 6.98 of [10], which state that

$$F_{2n-1} + F_{2n+1} = \sum_{k=0}^n \frac{2n}{2n-k} \binom{2n-k}{k}$$

and

$$F_{2n} + F_{2n+2} = \sum_{k=0}^n \frac{2n+1}{2n+1-k} \binom{2n+1-k}{k}.$$

**Application 2.** For  $q \geq 1$ , let  $(G_n^{(q)})_{n \geq 0}$  be the generalized Fibonacci sequence as cited in [8], and let  $(H_n^{(q)})_{n \geq 0}$  be a sequence of numbers defined as follows:

$$\begin{cases} G_0^{(q)} = G_1^{(q)} = \dots = G_q^{(q)} = 1, \\ G_{n+1}^{(q)} = G_n^{(q)} + G_{n-q}^{(q)} \quad \text{for } n \geq q, \end{cases} \quad \text{and} \quad \begin{cases} H_0^{(q)} = H_1^{(q)} = \dots = H_q^{(q)} = 1, \\ H_{n+1}^{(q)} = H_n^{(q)} - H_{n-q}^{(q)} \quad \text{for } n \geq q. \end{cases}$$

We can extend easily the above sequences to  $(G_n)_{n \geq -q}$  and  $(H_n)_{n \geq -q}$  by

$$\begin{cases} G_{-j}^{(q)} = 0 \quad \text{for } 1 \leq j \leq q, \\ G_0^{(q)} = 1, \\ G_{n+1}^{(q)} = G_n^{(q)} + G_{n-q}^{(q)} \quad \text{for } n \geq 0, \end{cases} \quad \text{and} \quad \begin{cases} H_{-j}^{(q)} = 0 \quad \text{for } 1 \leq j \leq q, \\ H_0^{(q)} = 1, \\ H_{n+1}^{(q)} = H_n^{(q)} - H_{n-q}^{(q)} \quad \text{for } n \geq 0. \end{cases}$$

The application of Corollary 1 gives us, for  $n \geq -q$ , the relations

$$G_n^{(q)} = \sum_{k=0}^{\lfloor \frac{n}{q+1} \rfloor} \binom{n-kq}{k}, \quad \text{and} \quad H_n^{(q)} = \sum_{k=0}^{\lfloor \frac{n}{q+1} \rfloor} (-1)^k \binom{n-kq}{k}.$$

Notice that  $G_n^{(1)} = F_{n+1} = \varphi_n$ , and  $H_n^{(2)}$  is the integer function studied by L. Bernstein [1], who showed that the only zeros of  $H_n^{(2)}$  are at  $n = 3$  and  $n = 12$ . This result was treated also by L. Carlitz [2, 3] and recently by J. McLaughlin and B. Sury [8].

The following theorem gives us an explicit formulation for  $u_n$  in terms of  $n, a_1, a_2, \dots, a_m, \alpha_0, \alpha_1, \dots, \alpha_{m-1}$ , and thus generalizes Theorem 2.

**Theorem 3.** Let  $(u_n)_{n > -m}$  be a sequence of elements of  $\mathcal{A}$  defined by

$$\begin{cases} u_{-j} = \alpha_j \quad \text{for } 0 \leq j \leq m-1, \\ u_n = a_1 u_{n-1} + a_2 u_{n-2} + \dots + a_m u_{n-m} \quad \text{for } n \geq 1. \end{cases} \quad (10)$$

Let  $(\lambda_j)_{0 \leq j \leq m-1}$  and  $(y_n)_{n > -m}$  be the sequences of elements of  $\mathcal{A}$  defined by

$$\lambda_j = - \sum_{k=j}^{m-1} a_{k-j} \alpha_k, \quad \text{for } 0 \leq j \leq m-1, \quad \text{with } a_0 = -1, \quad (11)$$

and

$$y_n = \sum_{k_1+2k_2+\dots+mk_m=n} \binom{k_1+k_2+\dots+k_m}{k_1, k_2, \dots, k_m} a_1^{k_1} a_2^{k_2} \dots a_m^{k_m}, \quad \text{for } n > -m.$$

Then for all integer  $n > -m$ , we have  $u_n = \lambda_0 y_n + \lambda_1 y_{n+1} + \cdots + \lambda_{m-1} y_{n+m-1}$ .

*Remark.* Note that (11) is equivalent to

$$\begin{pmatrix} \lambda_0 \\ \lambda_1 \\ \lambda_2 \\ \vdots \\ \lambda_{m-2} \\ \lambda_{m-1} \end{pmatrix} = \begin{pmatrix} 1 & -a_1 & -a_2 & \cdots & -a_{m-2} & -a_{m-1} \\ 0 & 1 & -a_1 & -a_2 & \cdots & -a_{m-2} \\ 0 & 0 & 1 & -a_1 & \cdots & -a_{m-3} \\ \vdots & \vdots & \ddots & \ddots & \ddots & \vdots \\ 0 & 0 & \cdots & 0 & 1 & -a_1 \\ 0 & 0 & 0 & \cdots & 0 & 1 \end{pmatrix} \begin{pmatrix} \alpha_0 \\ \alpha_1 \\ \alpha_2 \\ \vdots \\ \alpha_{m-2} \\ \alpha_{m-1} \end{pmatrix}.$$

or

$$[\lambda_0, \lambda_1, \dots, \lambda_{m-1}]^t = C [\alpha_0, \alpha_1, \dots, \alpha_{m-1}]^t, \quad (12)$$

where

$$C = (c_{ij})_{1 \leq i, j \leq m}, \quad \text{with } c_{ij} = \begin{cases} 0 & \text{if } i > j, \\ 1 & \text{if } i = j, \\ -a_{j-i} & \text{if } i < j. \end{cases}$$

We deduce also from relations (10) and (12) the matrix equality

$$\begin{pmatrix} u_n \\ u_{n+1} \\ \vdots \\ u_{n+m-2} \\ u_{n+m-1} \end{pmatrix} = \begin{pmatrix} y_n & y_{n+1} & \cdots & y_{n+m-2} & y_{n+m-1} \\ y_{n+1} & y_{n+2} & \cdots & y_{n+m-1} & y_{n+m} \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ y_{n+m-2} & y_{n+m-1} & \cdots & y_{n+2m-4} & y_{n+2m-3} \\ y_{n+m-1} & y_{n+m} & \cdots & y_{n+2m-3} & y_{n+2m-2} \end{pmatrix} \times$$

$$\times \begin{pmatrix} 1 & -a_1 & -a_2 & \cdots & -a_{m-1} \\ 0 & 1 & -a_1 & \ddots & \vdots \\ \vdots & \ddots & \ddots & \ddots & -a_2 \\ 0 & & 0 & 1 & -a_1 \\ 0 & 0 & \cdots & 0 & 1 \end{pmatrix} \begin{pmatrix} \alpha_0 \\ \alpha_1 \\ \vdots \\ \alpha_{m-2} \\ \alpha_{m-1} \end{pmatrix}$$

*Proof.* Let  $\mathcal{S}$  be the  $\mathcal{A}$ -module of sequences  $(v_n)_{n > -m}$  satisfying the recurrence relation

$$v_n - a_1 v_{n-1} - a_2 v_{n-2} - \cdots - a_m v_{n-m} = 0 \quad \text{for all } n \geq 1.$$

Let us consider the family of sequences  $(v_n^{(k)})_{n > -m}$ , for  $0 \leq k \leq m-1$ , defined by

$$\begin{cases} v_n^{(0)} &= y_n, \\ v_n^{(1)} &= y_{n+1} - a_1 y_n, \\ v_n^{(2)} &= y_{n+2} - a_1 y_{n+1} - a_2 y_n, \\ &\vdots \\ v_n^{(m-1)} &= y_{n+m-1} - a_1 y_{n+m-2} - a_2 y_{n+m-3} - \cdots - a_{m-1} y_n, \end{cases}$$



i.e.,

$$\begin{aligned} v_n^{(k)} &= y_{n+k} - (a_1 y_{n+k-1} + a_2 y_{n+k-2} + \cdots + a_k y_n) \\ &= \sum_{i=0}^k -a_i y_{n+k-i}, \quad \text{with } a_0 = -1. \end{aligned} \quad (13)$$

By Theorem 2, we have  $(y_n)_{n > -m} \in \mathcal{S}$ . Consequently,  $(y_{n+q})_{n > -m} \in \mathcal{S}$  for  $q \geq 0$ , and finally we deduce that

$$(v_n^{(k)})_{n > -m} \in \mathcal{S}, \quad \text{for } 0 \leq k \leq m-1. \quad (14)$$

It is easy to observe that we also have, for  $j, k \in \{0, 1, 2, \dots, m-1\}$ ,

$$v_{-j}^{(k)} = \delta_{kj} = \begin{cases} 1 & \text{if } k = j \\ 0 & \text{if } k \neq j \end{cases} \quad (15)$$

In fact, with  $a_0 = -1$ , we can write, for  $j, k \in \{0, 1, \dots, m-1\}$ ,  $v_{-j}^{(k)} = -\sum_{i=0}^k a_i y_{-j+k-i}$ .

If  $k < j$ , then  $-j + k - i < 0$  for  $1 \leq i \leq k$  and  $v_{-j}^{(k)} = 0$  (because  $y_q = 0$  for  $q < 0$ ). If  $k = j$ , then  $v_{-j}^{(k)} = v_{-j}^{(j)} = -\sum_{i=0}^j a_i y_{-i} = -a_0 y_0 = 1$ . If  $k > j$ , then for  $r = k - j < 0$ , we have  $r \geq 1$  and

$$\begin{aligned} v_{-j}^{(k)} &= y_r - (a_1 y_{r-1} + a_2 y_{r-2} + \cdots + a_k y_{r-k}), \quad \text{and by using } r - k = -j \leq 0, \\ &= y_r - (a_1 y_{r-1} + a_2 y_{r-2} + \cdots + a_k y_{r-k} + \cdots + a_m y_{r-m}) \\ &= 0 \quad \text{because } (y_n)_{n > -m} \in \mathcal{S} \text{ and } r \geq 1. \end{aligned}$$

Relations (14) and (15) give easily, that for all  $n > -m$ ,  $u_n = \sum_{k=0}^{m-1} \alpha_k v_n^{(k)}$  and, with (13),

$$\begin{aligned} u_n &= \sum_{k=0}^{m-1} \alpha_k \left( \sum_{i=0}^k -a_i y_{n+k-i} \right) \\ &= \sum_{k=0}^{m-1} \sum_{j=0}^k -a_{k-j} \alpha_k y_{n+j} \\ &= \sum_{j=0}^{m-1} \left( \sum_{k=j}^{m-1} -a_{k-j} \alpha_k \right) y_{n+j} \\ &= \sum_{j=0}^{m-1} \lambda_j y_{n+j}, \end{aligned}$$

where  $\lambda_j$  is as defined in (11). □

Theorem 3 allows us to find various formulae for the Fibonacci numbers.

**Corollary 2.** *For all integers  $n \geq 1$ ,*

$$F_n = \frac{1}{2^{n+2}} \sum_{k=1}^n \frac{n+1-k}{k} \binom{k}{n+1-2k} 8^k.$$

*Remark.* In this summation, we may, in fact, restrict the sum to those integers  $i$  ranging between  $\frac{n+1}{3}$  and  $\frac{n+1}{2}$ , the binomial coefficient of the formula being zero for the other integers.

*Proof.* Note that  $F_n = 2F_{n-2} + F_{n-3}$  for  $n \geq 3$ . Denoting by  $(y_n)_{n \geq -2}$  the sequence defined by

$$\begin{cases} y_{-2} = y_{-1} = 0, \\ y_0 = 1, \\ y_n = 2y_{n-2} + y_{n-3} \quad \text{for } n \geq 1, \end{cases}$$

we see that, for  $n \geq 1$ ,  $F_n = y_{n-1} + y_{n-2}$ . Theorem 3 allows us to state that, for  $n \geq -2$ ,

$$y_n = \sum_{2k+3l=n} \binom{k+l}{l} 2^k = \sum_{0 \leq t \leq n} \binom{t}{n-2t} 2^{3t-n},$$

and Corollary 2 follows by simple calculations. □

The following result can also be easily deduced from Theorem 3.

**Corollary 3.** *For all integers  $m \geq 1$ ,  $F_{2m-1} = \sum_{k=0}^{\lfloor m/3 \rfloor} \frac{m+k}{m-k} \binom{m-k}{2k} 2^{m-1-3k}$ .*

Now, let us consider for  $q > 1$ , the “multibonacci” sequence  $(\phi_n^{(q)})_{n > -q}$  defined by

$$\begin{cases} \phi_{-(q-1)}^{(q)} = \cdots = \phi_{-2}^{(q)} = \phi_{-1}^{(q)} = 0, \\ \phi_0^{(q)} = 1, \\ \phi_n^{(q)} = \phi_{n-1}^{(q)} + \phi_{n-2}^{(q)} + \cdots + \phi_{n-q}^{(q)} \quad \text{for } n \geq 1, \end{cases} \quad (16)$$

where  $\phi_n^{(2)} = \varphi_n = F_{n+1} = G_n^{(1)}$ . Theorem 3 also implies that, for all  $n \geq 0$ ,

$$\phi_n^{(q)} = \sum_{k_1+2k_2+\cdots+qk_q=n} \binom{k_1+k_2+\cdots+k_q}{k_1, k_2, \dots, k_q}.$$

Thus, for  $q = 3$ , we obtain

$$\phi_n^{(3)} = \sum_{i+2j+3k=n} \binom{i+j+k}{i, j, k} = \sum_{2i+3j \leq n} \binom{n-i-2j}{i+j} \binom{i+j}{i}.$$

We deduce from (16) that  $\phi_{n+1}^{(q)} = 2\phi_n^{(q)} - \phi_{n-q}^{(q)}$ , for  $n \geq 1$ . Let us consider  $(\psi_n)_{n \geq -q}$ , the sequence defined by  $\psi_n := \phi_n^{(q)}$ , for  $n > -q$  and  $\psi_{-q} = 1$ , which satisfies the recurrence relation  $\psi_{n+1} = 2\psi_n - \psi_{n-q}$ , for  $n \geq 0$ . After applying Theorem 3, we find that, for  $n \geq 0$ ,

$$\begin{aligned} \psi_n (= \phi_n^{(q)}) &= \lambda_0 z_n + \lambda_1 z_{n+1} + \cdots + \lambda_q z_{n+q} \\ &= z_n - 2z_{n+q-1} + z_{n+q}, \end{aligned}$$

with  $z_n = \sum_{k=0}^{\lfloor \frac{n}{q+1} \rfloor} \binom{n-kq}{k} 2^{n-k(q+1)} (-1)^k$ , for  $n \geq -q$ . We know, via Theorem 3 and Corollary 1, that the sequence  $(z_n)_{n \geq -q}$  satisfies the recurrence relation  $z_{n+1} - 2z_n + z_{n-q} = 0$ , for  $n \geq 0$ . This gives, for  $n \geq 0$ ,

$$\begin{aligned} \phi_n^{(q)} &= z_n - z_{n-1} + (z_{n+q} - 2z_{n+q-1} + z_{n-1}) \\ &= z_n - z_{n-1}. \end{aligned}$$

Applying relation (9) in Corollary 1, we can write, for  $n \geq 1$ ,

$$\phi_n^{(q)} = \sum_{k=0}^{\lfloor \frac{n}{q+1} \rfloor} \frac{n-k(q-1)}{n-kq} \binom{n-kq}{k} 2^{n-1-k(q+1)} (-1)^k.$$

Thus,

$$\sum_{k_1+2k_2+\cdots+qk_q=n} \binom{k_1+\cdots+k_q}{k_1, \dots, k_q} = \sum_{k=0}^{\lfloor \frac{n}{q+1} \rfloor} \frac{n-k(q-1)}{n-kq} \binom{n-kq}{k} 2^{n-1-k(q+1)} (-1)^k. \quad (17)$$

For  $q = 3$ , we obtain

$$\begin{aligned} \sum_{i+2j+3k=n} \binom{i+j+k}{i, j, k} &= \sum_{2i+3j \leq n} \binom{n-i-2j}{i+j} \binom{i+j}{i} \\ &= \sum_{k=0}^{\lfloor \frac{n}{4} \rfloor} \frac{n-2k}{n-3k} (-1)^k \binom{n-3k}{k} 2^{n-1-4k}. \end{aligned}$$

### 3. Powers of a Square Matrix of Order $m$

We start this section with the main result of this paper.

**Theorem 4.** Let  $A \in \mathcal{M}_m(\mathcal{A})$  and let  $P(X) = X^m - a_1 X^{m-1} - \cdots - a_{m-1} X - a_m$  be the characteristic polynomial of  $A$ . Let  $A_0, A_1, \dots, A_{m-1}$  be matrices of  $\mathcal{M}_m(\mathcal{A})$  defined by

$$A_k = - \sum_{i=0}^k a_i A^{k-i}, \quad \text{for } 0 \leq k \leq m-1, \text{ with } a_0 = -1, \quad (18)$$

and let  $(y_n)_{n > -m}$  be the sequence of elements of  $A$  satisfying

$$y_n = \sum_{k_1+2k_2+\dots+mk_m=n} \binom{k_1+k_2+\dots+k_m}{k_1, k_2, \dots, k_m} a_1^{k_1} a_2^{k_2} \dots a_m^{k_m}, \quad \text{for } n > -m.$$

Then, for all integers  $n \geq 0$ ,

$$A^n = y_n A_0 + y_{n-1} A_1 + \dots + y_{n-m+1} A_{m-1} \quad (19)$$

*Proof.* Define, for  $0 \leq k \leq m$ ,

$$P_k(X) = - \sum_{i=0}^k a_i X^{k-i}, \quad \text{with } a_0 = -1. \quad (20)$$

For  $0 \leq k \leq m-1$ , we have

$$X P_k(X) = P_{k+1}(X) + a_{k+1}. \quad (21)$$

Relation (20) shows that the degree of  $P_k$  is  $k$  for  $0 \leq k \leq m$ . It implies that  $(P_0, P_1, \dots, P_{m-1})$  is a basis of the free  $\mathcal{A}$ -module  $\mathcal{A}_{m-1}[X]$  consisting of polynomials of  $\mathcal{A}[X]$  of degree  $\leq m-1$ .

For all  $n \geq 0$ , the remainder  $R_n$  of the euclidean division of  $X^n$  by  $P_m$  is written as a linear combination of polynomials  $P_0, P_1, \dots, P_{m-1}$ . Then, for all  $n \geq 0$ , there exists a unique family  $(\alpha_{n,k})_{0 \leq k \leq m-1}$  such that

$$R_n = \sum_{k=0}^{m-1} \alpha_{n,k} P_k. \quad (22)$$

For  $0 \leq n \leq m-1$ , we have  $R_n(X) = X^n$ , where  $X^n$  is a linear combination of  $P_0, \dots, P_{m-1}$ , and

$$\begin{cases} \alpha_{0,0} = 1 \\ \alpha_{n,k} = 0 \quad \text{for } n < k \leq m-1. \end{cases} \quad (23)$$

Relations (21) and (22) imply that

$$\begin{aligned} X R_n(X) &= \sum_{k=0}^{m-1} \alpha_{n,k} (P_{k+1}(X) + a_{k+1}) \\ &= \sum_{k=0}^{m-1} a_{k+1} \alpha_{n,k} + \sum_{k=1}^{m-1} \alpha_{n,k-1} P_k(X) + \alpha_{n,m-1} P_m(X). \end{aligned}$$

As a consequence, the polynomial  $R_{n+1}(X) - X R_n(X) - \alpha_{n,m-1} P_m(X)$ , of degree  $\leq m-1$ , is divisible by  $P_m(X)$ , which is of degree  $m$ . This polynomial is thus the zero polynomial as

well as its components in the basis  $(P_0, P_1, \dots, P_{m-1})$ . This provides us with the following relations:

$$\alpha_{n+1,0} = \sum_{k=0}^{m-1} a_{k+1} \alpha_{n,k}, \quad \text{for } n \geq 0, \quad (24)$$

$$\alpha_{n+1,k} = \alpha_{n,k-1} \quad \text{for } 1 \leq k \leq m-1. \quad (25)$$

Let us set, for all integers  $n \in \mathbb{Z}$ ,

$$z_n = \begin{cases} \alpha_{n,0} & \text{for } n \geq 0, \\ 0 & \text{for } n < 0. \end{cases} \quad (26)$$

One checks easily that for all integers  $n \geq 0$  and for  $0 \leq k \leq m-1$ ,

$$\alpha_{n,k} = z_{n-k}. \quad (27)$$

Indeed, if  $n \geq k$  this relation follows from (25) and (26), and if  $0 \leq n < k \leq m-1$  it follows from (23) and (26). From (25), (26) and (27), we find that

$$\begin{cases} z_n = 0 & \text{for } n < 0, \\ z_0 = 1, \\ z_n = a_1 z_{n-1} + a_2 z_{n-2} + \dots + a_{m-1} z_{n-m+1} + a_m z_{n-m} & \text{for } n \geq 1. \end{cases}$$

Theorem 2 implies that  $z_n = y_n$ , for all  $n \in \mathbb{Z}$ , where

$$y_n = \sum_{k_1+2k_2+\dots+mk_m=n} \binom{k_1+k_2+\dots+k_m}{k_1, k_2, \dots, k_m} a_1^{k_1} a_2^{k_2} \dots a_m^{k_m}.$$

This last fact, together with the Cayley-Hamilton Theorem, (22), (20), (18) and (27) now give that

$$A^n = R_n(A) = \sum_{i=0}^{m-1} \alpha_{n,i} P_i(A) = \sum_{i=0}^{m-1} z_{n-i} A_i = \sum_{i=0}^{m-1} y_{n-i} A_i.$$

which completes the proof of (19). □

#### 4. Further Combinatorial Identities

Some nice combinatorial identities can be derived from Theorem 4 by considering various particular matrices with simple forms.

**Corollary 4.** *Let  $n$  be a positive integer. Then*

$$\sum_{k_1+2k_2+\dots+mk_m=n} \binom{k_1+\dots+k_m}{k_1, \dots, k_m} (-1)^{n-(k_1+\dots+k_m)} \binom{m}{m-1}^{k_1} \dots \binom{m}{0}^{k_m} = \binom{n+m-1}{m-1},$$

where  $\binom{n+m-1}{m-1}$  is the number of  $m$ -combinations with repetition of finite set  $\{1, \dots, n\}$ , and the summation is taken over all  $m$ -tuples  $(k_1, k_2, \dots, k_m)$  of integers  $k_j \geq 0$  satisfying the relation  $k_1 + 2k_2 + \dots + mk_m = n$ .

*Proof.* Let  $J_m$  be the  $m \times m$  Jordan matrix,

$$J_m = \begin{pmatrix} 1 & 1 & 0 & \cdots & 0 \\ 0 & 1 & 1 & \ddots & \vdots \\ 0 & 0 & \ddots & \ddots & 0 \\ \vdots & \vdots & \ddots & 1 & 1 \\ 0 & 0 & \cdots & 0 & 1 \end{pmatrix}.$$

The characteristic polynomial of  $J_m$  is  $(X - 1)^m$ . We also have  $J_m^n = \left( \binom{n}{j-i} \right)_{1 \leq i, j \leq m}$ .

Applying Theorem 4 with  $A = J_m$ , and considering the  $(1, m)$  entries of both sides of (19), we obtain the relation  $\binom{n}{m-1} = y_{n-(m-1)}$ , which leads to the result.  $\square$

From Corollary 4, we obtain the following combinatorial identities, for  $m = 2, 3, 4$ .

$$\begin{aligned} \sum_{2i \leq n} (-1)^i \binom{n-i}{i} 2^{n-2i} &= n+1, \\ \sum_{2i+3j \leq n} (-1)^i \binom{n-i-2j}{i+j} \binom{i+j}{i} 3^{n-i-3j} &= \binom{n+1}{2} \\ \sum_{2i+3j+4k \leq n} (-1)^{i+k} \binom{n-i-2j-3k}{i+j+k} \binom{i+j+k}{i+j} \binom{i+j}{i} 2^{2n-3i-4j-8k} 3^i &= \binom{n+3}{3} \end{aligned}$$

Like J. Mc Laughlin and B. Sury [8, Corollary 6], we can also derive Corollary 4 from the following result.

**Corollary 5.** ([8, Theorem 1]) *Let  $x_1, x_2, \dots, x_m$  be elements of the unitary commutative ring  $\mathcal{A}$  with  $s_k = \sum_{1 \leq i_1 < i_2 < \dots < i_k \leq m} x_{i_1} x_{i_2} \cdots x_{i_k}$ , for  $1 \leq k \leq m$ . Then, for each positive integer  $n$ ,*

$$\sum_{k_1+k_2+\dots+k_m=n} x_1^{k_1} \cdots x_m^{k_m} = \sum_{k_1+2k_2+\dots+mk_m=n} \binom{k_1+\dots+k_m}{k_1, \dots, k_m} (-1)^{n-k_1-\dots-k_m} s_1^{k_1} \cdots s_m^{k_m},$$

with the summations being taken over all  $m$ -tuples  $(k_1, k_2, \dots, k_m)$  of integers  $k_j \geq 0$  satisfying the relations  $k_1 + k_2 + \dots + k_m = n$  for the left-hand side and  $k_1 + 2k_2 + \dots + mk_m = n$  for the right-hand side.

*Proof.* Let us give a proof of this result by using Theorem 2. For  $n > -m$  and  $1 \leq l \leq m$ , consider  $q_n^{(l)} := \sum_{k_1+k_2+\dots+k_l=n} x_1^{k_1} x_2^{k_2} \dots x_l^{k_l}$ , and  $q_n := q_n^{(m)}$ . Let  $E : \beta_n \mapsto \beta_{n+1}$  be the shift operator which acts on any sequence  $(\beta_n)_n$ , and, for  $1 \leq l \leq m$ , let  $Q_l$  be the operator given by  $Q_l := (E - x_1)(E - x_2) \dots (E - x_l)$ . Notice that  $Q_m = E^m - s_1 E^{m-1} + \dots + (-1)^m s_m$ . Then, for  $n \geq 1$  and  $2 \leq l \leq m$ , we have  $(E - x_l) \cdot q_{n-l}^{(l)} = q_{n-(l-1)}^{(l-1)}$ . Therefore,

$$Q_m \left( q_{n-m}^{(m)} \right) = Q_{m-1} \left( q_{n-(m-1)}^{(m-1)} \right) = \dots = Q_1 \left( q_{n-1}^{(1)} \right) = 0.$$

Thus,

$$q_n = s_1 q_{n-1} - s_2 q_{n-2} + \dots + (-1)^{m-1} s_m q_{n-m}, \quad \text{for } n \geq 1. \quad (28)$$

By the definition of  $q_n$ , we also have

$$q_n = 0 \quad \text{for } n < 0 \quad \text{and} \quad q_0 = 1. \quad (29)$$

Applying Theorem 2 to the sequence  $(q_n)_{n > -m}$ , we obtain the result.  $\square$

Corollary 4 follows immediately by setting  $x_i = 1$  for all  $i$  in the previous corollary.

The following theorem is an extension of a result of J. Mc Laughlin and B. Sury [8, Theorem 3].

**Theorem 5.** *Let  $K$  be a field of characteristic zero, and let  $x_1, x_2, \dots, x_m$  be independent variables. Then, in  $K(x_1, x_2, \dots, x_m)$ ,*

$$\sum_{k_1+\dots+k_m=n} x_1^{k_1} x_2^{k_2} \dots x_m^{k_m} = \sum_{i=1}^m \frac{x_i^{n+m-1}}{\prod_{j \neq i} (x_i - x_j)}.$$

*Proof.* For  $\gamma_1, \gamma_2, \dots, \gamma_m \in K(x_1, x_2, \dots, x_m)$ , let  $V(\gamma_1, \gamma_2, \dots, \gamma_m)$  denote the Vandermonde determinant defined by  $V(\gamma_1, \gamma_2, \dots, \gamma_m) = \det(\gamma_i^{j-1})_{1 \leq i, j \leq m}$ . It is well known that  $V(\gamma_1, \gamma_2, \dots, \gamma_m) = \prod_{1 \leq i < j \leq m} (\gamma_j - \gamma_i)$ . By (28), the sequence  $(q_n)_{n > -m}$  defined by

$$q_n = \sum_{k_1+k_2+\dots+k_m=n} x_1^{k_1} x_2^{k_2} \dots x_m^{k_m}$$

is a recurrent sequence with characteristic polynomial

$$X^m - s_1 X^{m-1} + \dots + (-1)^m s_m = (X - x_1)(X - x_2) \dots (X - x_m).$$

This polynomial has  $m$  distinct roots  $x_1, x_2, \dots, x_m$ . We deduce that there exist  $m$  elements  $A_i = A_i(x_1, x_2, \dots, x_m) \in K(x_1, x_2, \dots, x_m)$ ,  $1 \leq i \leq m$ , such that

$$q_n = \sum_{i=1}^m A_i x_i^n \quad \text{for } n \geq -m.$$

The initial conditions given by (28) lead to the Cramer system  $\sum_{i=1}^m \frac{A_i}{x_i^j} = \delta_{j,0}$  for  $0 \leq j \leq m-1$ . The resolution of this system gives

$$A_i = \frac{V\left(\frac{1}{x_1}, \dots, \frac{1}{x_{i-1}}, 0, \frac{1}{x_{i+1}}, \dots, \frac{1}{x_m}\right)}{V\left(\frac{1}{x_1}, \dots, \frac{1}{x_{i-1}}, \frac{1}{x_i}, \frac{1}{x_{i+1}}, \dots, \frac{1}{x_m}\right)} = \frac{x_i^{m-1}}{\prod_{j \neq i} (x_i - x_j)}, \text{ for } 1 \leq i \leq m.$$

This completes the proof.  $\square$

Let us now give an application to Stirling numbers. For  $n \geq 0$ , with the notations of [5], the Stirling numbers of the first kind  $\left[ \begin{smallmatrix} n \\ k \end{smallmatrix} \right]$ , and the Stirling numbers of the second kind  $\left\{ \begin{smallmatrix} n \\ k \end{smallmatrix} \right\}$ , can be defined by the equations

$$X^{\overline{n}} = \sum_{k=0}^n \left[ \begin{smallmatrix} n \\ k \end{smallmatrix} \right] X^k, \quad (30)$$

with  $X^{\overline{n}} := \begin{cases} 1 & \text{if } n = 0, \\ X(X+1)(X+2) \cdots (X+n-1) & \text{if } n \geq 1, \end{cases}$  and

$$X^n = \sum_{k=0}^n \left\{ \begin{smallmatrix} n \\ k \end{smallmatrix} \right\} X^{\underline{k}}, \quad (31)$$

with  $X^{\underline{k}} := \begin{cases} 1 & \text{if } k = 0, \\ X(X-1)(X-2) \cdots (X-k+1) & \text{if } k \geq 1. \end{cases}$

It is well known [p. 38 (Vol. 2), 4] that  $\left\{ \begin{smallmatrix} n \\ k \end{smallmatrix} \right\} = \frac{1}{k!} \sum_{i=0}^k (-1)^{k-i} \binom{k}{i} i^n$ . From Theorem 5 and Corollary 5, we deduce

$$\sum_{k_1+2k_2+\dots+mk_m=n} \binom{k_1+\dots+k_m}{k_1, \dots, k_m} (-1)^{n-k_1-\dots-k_m} s_1^{k_1} \dots s_m^{k_m} = \sum_{i=1}^m \frac{x_i^{n+m-1}}{\prod_{j \neq i} (x_i - x_j)}. \quad (32)$$

If we take  $x_k = -(k-1)$  for  $1 \leq k \leq m$ , we have  $s_k = (-1)^k \left[ \begin{smallmatrix} m \\ m-k \end{smallmatrix} \right]$ , for  $1 \leq k \leq m$ , and relation (32) gives the following result

**Corollary 6.** *For all positive integers  $m$  and  $n$ ,*

$$\sum_{k_1+2k_2+\dots+mk_m=n} \binom{k_1+\dots+k_m}{k_1, \dots, k_m} (-1)^{n-(k_1+\dots+k_m)} \left[ \begin{smallmatrix} m \\ m-1 \end{smallmatrix} \right]^{k_1} \cdots \left[ \begin{smallmatrix} m \\ 0 \end{smallmatrix} \right]^{k_m} = \left\{ \begin{smallmatrix} n+m-1 \\ m-1 \end{smallmatrix} \right\}.$$



Note that Theorem 5 gives the following relation, which is stated in [p. 42, (Vol. 2), 4].

For all positive integers  $m$  and  $n$ ,  $\left\{ \begin{matrix} n+m-1 \\ m-1 \end{matrix} \right\} = \sum_{k_1+\dots+k_{m-1}=n} 1^{k_1} 2^{k_2} \dots (m-1)^{k_{m-1}}.$

For  $m = 3, 4$  and  $5$ , we obtain

$$\sum_{2i \leq n} (-1)^i \binom{n-i}{i} 2^i 3^{n-2i} = 2^{n+1} - 1,$$

$$\sum_{2i+3j \leq n} (-1)^i \binom{n-i-2j}{i+j} \binom{i+j}{i} 6^{n-2i-2j} 11^i = \frac{3^{n+2} - 2^{n+3} + 1}{2},$$

$$\begin{aligned} \sum_{2i+3j+4k \leq n} (-1)^{i+k} \binom{n-i-2j-3k}{i+j+k} \binom{i+j+k}{i+j} \binom{i+j}{i} 2^{n-2i-2j-k} 3^k 5^{n-i-j-4k} 7^i \\ = \frac{4^{n+3} - 3^{n+4} + 3 \cdot 2^{n+3} - 1}{6}. \end{aligned}$$

**Corollary 7.** *Let  $n$  be a positive integer and let  $x$  and  $y$  be indeterminates. Then*

$$\begin{aligned} \sum_{2i+3j+4k \leq n} (-1)^{i+k} \binom{n-i-2j-3k}{i+j+k} \binom{i+j+k}{i+j} \binom{i+j}{i} (2x+2y)^{n-2i-2j-4k} \times \\ \times (xy)^{j+2k} (x^2+4xy+y^2)^i = \frac{(n+1)(x^{n+3}-y^{n+3}) - (n+3)xy(x^{n+1}-y^{n+1})}{(x-y)^3}. \end{aligned}$$

**Corollary 8.** *Let  $n$  a positive integer and  $x, y$  be indeterminates. Then*

$$\begin{aligned} \sum_{2i+3j+4k \leq n} (-1)^{i+k} \binom{n-i-2j-3k}{i+j+k} \binom{i+j+k}{i+j} \binom{i+j}{i} \times \\ \times (3x+y)^{n-2i-3j-4k} x^{i+2j+3k} y^k (3x+3y)^i (x+3y)^j \\ = \frac{(n+1)(n+2)(x^{n+3}-y^{n+3}) - (n+3)y[n(x^{n+2}-y^{n+2}) + (n+2)x^{n+1}(x-y)]}{2(x-y)^3}. \end{aligned}$$

*Proof of Corollaries 7 and 8.* It suffices to put, in Corollary 4,  $x_1 = x_2 = x$  and  $x_3 = x_4 = y$  to obtain Corollary 7;  $x_1 = x_2 = x_3 = x$  and  $x_4 = y$  to obtain Corollary 8.  $\square$

**Acknowledgments.** The authors are grateful to the referee and would like to thank him/her for comments which improved the quality of this paper.

## References

- [1] L. BERNSTEIN: Zeros of the function  $f(n) = \sum_i (-1)^i \binom{n-2i}{i}$ . J. Number Theory, **6** (1974), 264-270.
- [2] L. CARLITZ: Some combinatorial identities of Bernstein. SIAM J. Math. Anal., **9** (1978), no. 1, 65–75.
- [3] L. CARLITZ: Recurrences of the third order and related combinatorial identities. Fibonacci Quarterly, **16** (1978), no. 1, 11–18.
- [4] L. COMTET: Analyse combinatoire. Puf, Coll. Sup. Paris, (1970), Vol. 1 & Vol. 2.
- [5] R. L. GRAHAM, D. E. KNUTH, O. PATASHNIK: Concrete Mathematics. Addison Wesley Publishing Company, Inc., (1994).
- [6] R. C. JOHNSON: Matrix methods for Fibonacci and related sequences. [www.dur.ac.uk/bob.johnson/fibonacci/](http://www.dur.ac.uk/bob.johnson/fibonacci/), (2004).
- [7] J. MC LAUGHLIN: Combinatorial identities deriving from the n-th power of  $2 \times 2$  matrix. Electronic Journal of Combinatorial Number Theory, A19, **4** (2004).
- [8] J. MC LAUGHLIN, B. SURY: Powers of matrix and combinatorial identities. Electronic Journal of Combinatorial Number Theory, A13, **5** (2005).
- [9] RAJESH RAM: Fibonacci number formulae. <http://users.tellurian.net/hsejar/maths/fibonacci/> (2003).
- [10] Z. F. STARC: Solution of the Problem 6.98. Univ. Beograd. Publikac. Elektrotehn. Fak. Ser. Mat. **13** (2002), 103-105.
- [11] S. TANNY, M. ZUKER: Analytic methods applied to a sequence of binomial coefficients. Discrete Math., **24** (1978), 299-310.
- [12] N. N. VOROB'EV: Fibonacci numbers. Translated from the russian by Halina Moss, Translation editor Ian N. Sneddon Blaisdell Publishing Co., New York-London, (1961).