

The Diagonal of a D -Finite Power Series Is D -Finite

L. LIPSHITZ

Department of Mathematics, Purdue University, West Lafayette, Indiana 47907

Communicated by N. Jacobson

Received May 15, 1985

ⓑ ⇒ ⓐ: easy
ⓐ ⇒ ⓑ?

Does it work for K a (division) ring of characteristic 0?

Let K be a field of characteristic zero, $x = x_1, \dots, x_n$ several variables, and $K[[x]]$ the ring of formal power series in x_1, \dots, x_n over K . We call $f \in K[[x]]$ D -finite (or differentially finite) if the set of all derivatives $(\partial/\partial x_1)^{i_1} \cdots (\partial/\partial x_n)^{i_n} f$ ($i_j \in \mathbb{N}$) lie in a finite-dimensional vector space over $K(x)$, the field of rational functions in x_1, \dots, x_n . This is equivalent to saying that f satisfies a system of linear partial differential equations of the form

$$\textcircled{B} \quad \left\{ a_{in_i}(x) \left(\frac{\partial}{\partial x_i} \right)^{n_i} + a_{in_i-1}(x) \left(\frac{\partial}{\partial x_i} \right)^{n_i-1} + \cdots + a_{i0}(x) \right\} f = 0, \quad i = 1, \dots, n, \quad (1)$$

where the $a_{ij}(x) \in K[x]$. We shall also write these equations as $A_i(x_1, \dots, x_n; \partial/\partial x_i) f = 0$, $i = 1, \dots, n$. The theory of D -finite power series in one variable is worked out in [9]. We call $f \in K[[x]]$ rational if $f \in K(x)$ and algebraic if it is algebraic over $K(x)$. If $f = \sum a_{i_1 \dots i_n} x_1^{i_1} \cdots x_n^{i_n}$ we define the primitive diagonal $I_{12}(f) = \sum a_{i_1 i_1 i_3 \dots i_n} x_1^{i_1} x_3^{i_3} \cdots x_n^{i_n}$. The other primitive diagonals I_{ij} (for $i < j$) are defined similarly. By a diagonal we mean any composition of the I_{ij} , and by the complete diagonal (or just the diagonal) of f we mean $I_{12} I_{23} \cdots I_{n-1n}(f) = \sum a_{ii \dots i} x^i$.

In this paper we will show (Theorem 1) that any diagonal of a D -finite power series is again D -finite. In [6] it is shown that the diagonal of a rational power series in two variables is algebraic and that in the case that K has characteristic $p \neq 0$ any diagonal of a rational power series in any number of variables is algebraic. (In characteristic 0 the diagonal of a rational power series in three variables need not be algebraic.) In [2, 3] it is shown, in the case that K has characteristic $p \neq 0$, that the diagonal of an algebraic power series in any number of variables is algebraic and that if $f \in \mathbb{Z}_p[[x]]$ is algebraic (\mathbb{Z}_p the p -adic integers) then any diagonal of f is algebraic mod p^s (for all s). In [7, 10] it is claimed that the diagonal of a rational function in any number of variables is D -finite, but the proofs con-

climination?

Diagonal of $f \leadsto s^{-2}$ section of F

tain gaps which do not seem easy to fill. Doron Zeilberger has informed me that he is able to prove that the diagonal of a rational function is D -finite using Bernstein theory. We shall use a clever counting argument introduced in [7, 10] in Lemma 3 below. In [1] it is shown that the complete diagonals of a restricted class of rational power series are D -finite. The restriction can be avoided by the use of Dwork's paper [4]—see also [5]. Deligne has also pointed out (see the footnote on p. 5 of [1]) that the D -finiteness of the diagonals of rational power series can be deduced via resolution of singularities from the finiteness of cohomology for the complement of a hypersurface. While our proof below is elementary and more general, these methods give more information about the differential equations satisfied by the diagonals of rational power series.

Let $f \in K[[x_1, \dots, x_n]]$ satisfy Eq. (1), and let s be a new variable. Define

$$F(s, x_1, x_3, \dots, x_n) = \frac{1}{s} f\left(s, \frac{x_1}{s}, x_3, \dots, x_n\right).$$

F is not a formal power series in s, x_1, x_3, \dots, x_n , but is an element of the $K[s, x_1, x_3, \dots, x_n]$ -module M of all

$$\begin{cases} x_1 \mapsto s \\ x_2 \mapsto x_1/s \end{cases}$$

$$G = \sum_{\substack{j \in \mathbb{Z} \\ i_2, \dots, i_n \in \mathbb{N} \\ j + i_2 \geq -k}} a_{ji_2 \dots i_n} s^j x_1^{i_2} x_3^{i_3} \dots x_n^{i_n} \Rightarrow j \in \mathbb{Z} \cap [i_2 - k, \infty)$$

for some $k \in \mathbb{N}$, depending on G . Let \mathcal{D} be the ring of all linear partial differential operators in $\partial/\partial s, \partial/\partial x_1, \partial/\partial x_3, \dots, \partial/\partial x_n$ with coefficients from $K[s, x_1, x_3, \dots, x_n]$. Then M is a \mathcal{D} -module in the natural way. Notice that the coefficient of $1/s$ in F is just $I_{12}(f)$. Later we shall need

$$I_{12}(f) = [s^{-2}]F$$

LEMMA 1. If $0 \neq p \in K[s, x_1, x_3, \dots, x_n]$ and $G \in M$ satisfy $pG = 0$ then $G = 0$.

Proof. For suitable k , $s^k G \in K[[s, x_1/s, x_3, \dots, x_n]]$. Make the substitution $x_1 = su$, u a new variable, to get $0 = p(s, su, x_3, \dots, x_n) s^k G(s, u, x_3, \dots, x_n) \in K[[s, u, x_3, \dots, x_n]]$. The conclusion now follows from the observations that multiplication by s , and the substitution $x_1 = su$, are both one-to-one.

LEMMA 2. F is D -finite (in the variables s, x_1, x_3, \dots, x_n).

Proof. This is immediate from the fact that f is D -finite, by the chain rule. \square

Hence there are nonzero linear partial differential operators, with polynomial coefficients,

$$S : \quad A \left(s, x_1, \dots, x_n; \frac{\partial}{\partial s} \right) = L(s, x_1, x_3, \dots, x_n) \left(\frac{\partial}{\partial s} \right)^m \\ + \text{lower-order terms in } \frac{\partial}{\partial s}$$

and

$$X_{\mu} : \quad B_i \left(s, x_1, \dots, x_n; \frac{\partial}{\partial x_i} \right) = L_i(s, x_1, x_3, \dots, x_n) \left(\frac{\partial}{\partial x_i} \right)^{m_i} \\ + \text{lower-order terms in } \frac{\partial}{\partial x_i},$$

for $i = 1, 3, \dots, n$ such that

$$\begin{cases} AF = 0 \\ B_i F = 0 \end{cases} \quad \text{for } i = 1, 3, \dots, n. \quad (2)$$

LEMMA 3. *There are nonzero linear partial differential operators $P_i(x_1, x_3, \dots, x_n; \partial/\partial s, \partial/\partial x_i)$, for $i = 1, 3, \dots, n$, with coefficients from $K[x_1, x_3, \dots, x_n]$, P_i containing only derivatives of the form $(\partial/\partial s)^\beta (\partial/\partial x_i)^\gamma$ such that*

$$P_i \left(x_1, x_3, \dots, x_n; \frac{\partial}{\partial s}, \frac{\partial}{\partial x_i} \right) F = 0 \quad \text{for } i = 1, 3, \dots, n.$$

Proof. Without loss of generality we may assume that A and the B_i in (2) above all have the same leading coefficient, i.e., that $L_i = L$ for $i = 1, 3, \dots, n$. Let all the coefficients in A and the B_i have total degrees $\leq d$. Let $D = (\partial/\partial s)^\beta (\partial/\partial x_1)^\gamma$. If $\beta \geq m$ we have

$$LDF = \sum P_\delta D_\delta F,$$

where the sum on the right-hand side is over $D_\delta = (\partial/\partial s)^{\delta_1} (\partial/\partial x_1)^{\delta_2}$ with $\delta_1 < \beta$ and $\delta_2 \leq \gamma$ and the P_δ are polynomials in s, x_1, x_3, \dots, x_n of total degree $\leq d$. We obtain this by applying $(\partial/\partial s)^{\beta-m} (\partial/\partial x_1)^\gamma$ to $AF = 0$. The similar statement holds if $\gamma \geq m_1$, but then we must use $B_1 F = 0$ and the sum is over $\delta_1 \leq \beta$ and $\delta_2 < \gamma$.

Iterating the above we see that if $\beta + \gamma \leq N$ then

$$L^N \left(\frac{\partial}{\partial s} \right)^\beta \left(\frac{\partial}{\partial x_1} \right)^\gamma F = \sum P_\delta D_\delta F,$$

where now the sum is over all $D_\delta = (\partial/\partial s)^{\delta_1} (\partial/\partial x_1)^{\delta_2}$ with $\delta_1 < m$ and $\delta_2 < m_1$ and the polynomials P_δ have total degrees $\leq Nd$.

Now let

$$D = x_1^{\alpha_1} x_3^{\alpha_3} \dots x_n^{\alpha_n} \left(\frac{\partial}{\partial s} \right)^\beta \left(\frac{\partial}{\partial x_1} \right)^\gamma, \quad (3)$$

where $\sum \alpha_i + \beta + \gamma \leq N$. Then

$$L^N D F = \sum \bar{P}_\delta D_\delta F, \quad (4)$$

where the sum is over all $D_\delta = (\partial/\partial s)^{\delta_1} (\partial/\partial x_1)^{\delta_2}$ with $\delta_1 < m$, $\delta_2 < m_1$ and the total degrees of the \bar{P}_δ are all $\leq N(d+1)$. The number of monomials in s, x_1, x_3, \dots, x_n of degree $\leq N(d+1)$ is $\binom{N(d+1)+n+1}{n+1}$. Hence the vector space of all such $\bar{P}_\delta D_\delta$ has dimension $mm_1 \binom{N(d+1)+n+1}{n+1} \leq c_1 N^{n+1}$ for some fixed c_1 and all $N \geq 1$. On the other hand, the number of D 's of the type (3) above is $\binom{N+n+2}{n+1}$, which is $> c_2 N^{n+2}$ for some $c_2 > 0$. Hence for N large enough there are $a_{\alpha_1 \alpha_3 \dots \alpha_n \beta \gamma} \in K$, not all zero, such that

$$L^N \sum_{\alpha_1 + \dots + \alpha_n + \beta + \gamma \leq N} a_{\alpha_1 \dots \alpha_n \beta \gamma} x_1^{\alpha_1} x_3^{\alpha_3} \dots x_n^{\alpha_n} \left(\frac{\partial}{\partial s} \right)^\beta \left(\frac{\partial}{\partial x_1} \right)^\gamma F = 0.$$

Let

$$P_1 = \sum a_{\alpha_1 \dots \alpha_n \beta \gamma} x_1^{\alpha_1} x_3^{\alpha_3} \dots x_n^{\alpha_n} \left(\frac{\partial}{\partial s} \right)^\beta \left(\frac{\partial}{\partial x_1} \right)^\gamma.$$

Then we have $L^N P_1 F = 0$ and hence, by Lemma 1, that $P_1 F = 0$. The P_3, \dots, P_n are found in a similar way, using $\partial/\partial x_i$ and $B_i = 0$ in place of $\partial/\partial x_1$ and $B_1 = 0$. This completes the proof of Lemma 3. \square

Now let the P_i be as in Lemma 3 and let $P_i = \sum_{j=\alpha_i}^{\beta_i} P_{ij}(x_1, x_3, \dots, x_n; \partial/\partial x_i)(\partial/\partial s)^j$ with $P_{i\alpha_i} \neq 0$. Notice that the coefficient of $1/s^{\alpha_i+1}$ in $P_i F$ is $(-1)^{\alpha_i} \alpha_i! P_{i\alpha_i} I_{12}(f)$. Hence $I_{12}(f)$ satisfies the equations

$$P_{i\alpha_i} \left(x_1, x_3, \dots, x_n; \frac{\partial}{\partial x_i} \right) I_{12}(f) = 0 \quad \text{for } i = 1, 3, \dots, n,$$

i.e., $I_{12}(f)$ is D -finite.

Iterating we have

$$[s^{-(\alpha_i+1)}] P_i F = (-1)^{\alpha_i} \alpha_i! P_{i\alpha_i} I_{12}(f)$$

THEOREM 1. If $f \in K[[x_1, \dots, x_n]]$ is D -finite, and I is any diagonal, then $I(f)$ is D -finite.

Remarks. (1) In the case that f is convergent for $|x_i| < a$ for all i , F is analytic for $0 < |s| < a$, $|x_1| < |s| a$, and $|x_i| < a$ for $i = 3, \dots, n$, and we can avoid the use of module M and Lemma 1 is trivial.

Closure
under
Hadamard
product

(2) If $f = \sum a_v x^v$, $g = \sum b_v x^v$, v a multi-index, then the Hadamard product $f * g = \sum a_v b_v x^v$. Since $f * g = I_{1n+1} I_{2n+2} \cdots I_{nn} f(x_1, \dots, x_n) g(x_{n+1}, \dots, x_{2n})$, it follows from Theorem 1 that if f and g are D -finite then so is $f * g$. (If f is just differentially algebraic and g is D -finite it doesn't follow that $f * g$ is differentially algebraic—see Proposition 6.3 of [8].)

(3) Instead of iterating the argument given for Theorem 1 one can do several steps at once. For example, if $f(x_1, x_2, x_3)$ is D -finite and one wants to show that the complete diagonal $I(f)$ is D -finite, one can consider $F(s, t, x_1) = (1/st) f(s, t/s, x_1/t)$ and use the argument in Lemma 3 to show that F satisfies an equation of the form

$$P\left(x_1; \frac{\partial}{\partial s}, \frac{\partial}{\partial t}, \frac{\partial}{\partial x_1}\right) F = 0, \quad P \neq 0.$$

If $P = P_{\alpha\beta}(x_1; \partial/\partial x_1)(\partial/\partial s)^\alpha (\partial/\partial t)^\beta + \text{higher-order terms in } \partial/\partial s \text{ or } \partial/\partial t$ with $P_{\alpha\beta} \neq 0$ then by considering the coefficient of $(1/s^{\alpha+1})(1/t^{\beta+1})$ we get that $P_{\alpha\beta}(x_1; \partial/\partial x_1) I(f) = 0$. This will give us smaller bounds for the order and degree of the equation satisfied by $I(f)$ than those obtained by iterating.

(4) If $f(x) = \sum_{n_1, \dots, n_k \geq 0} f(n_1, \dots, n_k) x_1^{n_1} \cdots x_k^{n_k}$ is D -finite and $C \subseteq \mathbf{N}^k$ is defined by a finite set of inequalities of the form $\sum a_i n_i + b \geq 0$, where the $a_i, b \in \mathbf{Z}$, then

$$h(x) = \sum_{\substack{n_1, \dots, n_k \geq 0 \\ \tilde{m} \in C}} f(n_1, \dots, n_k) x_1^{n_1} \cdots x_k^{n_k}$$

is also D -finite. To see this consider the case that C is defined by just one inequality $\sum_{i=1}^l \alpha_i n_i + \alpha_0 \geq \sum_{i=l+1}^k \beta_i n_i$, where the $\alpha_i, \beta_i \in \mathbf{N}$. Let

$$g(x, s, t) = s^{\alpha_0} \prod_{i=1}^l \frac{1}{1 - x_i s^{\alpha_i}} \prod_{i=l+1}^k \frac{1}{1 - x_i t^{\beta_i}}$$

$$a(s, t) = \frac{1}{1-s} \frac{1}{1-st} = \sum_{i \geq j} s^i t^j$$

$$b(x, s, t) = a(s, t) \prod_{i=1}^n \frac{1}{1 - x_i}$$

Then

$$\tilde{g}(x) = (g * b)(x, 1, 1) = \sum_{\substack{n_i \geq 0 \\ \tilde{m} \in C}} x_1^{n_1} \cdots x_k^{n_k}$$

and $h(x) = f(x) * \tilde{g}(x)$. Iterating we get the result for general C .

(5) If $f(x)$, C are as above and the $m_i(n_1, \dots, n_k) = \sum_{j=1}^k a_{ij}n_j + b_i$, where the a_{ij} , b_i are nonnegative elements of \mathbb{Q} , then

$$l(x) = \sum_{\substack{n_1, \dots, n_k \geq 0 \\ c}} f(n_1, \dots, n_k) x_1^{m_1(n_1, \dots, n_k)} \dots x_k^{m_k(n_1, \dots, n_k)}$$

is also D -finite. Let $y_i(x) = x_1^{a_{i1}} \dots x_k^{a_{ik}}$ and notice that $l(x) = x_1^{b_1} \dots x_k^{b_k} h(y_1, \dots, y_k)$, where h is as above.

(6) Remark (5) gives a positive answer to question 4(e) of [9]. For example, if the sequences $f_i(n)$, $i = 1, 2, 3$, are p -recursive (i.e., the $\sum f_i(n)x^n$ are D -finite) then $f(x, y, z) = \sum_{i, j, k \geq 0} f_1(i) f_2(j) f_3(k) x^i y^j z^k$ is D -finite. Let C be defined by $i + 2j = k$ (i.e., $i + 2j \geq k$ and $k \geq i + 2j$). Then

$$F(x) = \sum_{\substack{i, j, k \geq 0 \\ i + 2j = k}} f_1(i) f_2(j) f_3(k) x^{(i+k)/2}$$

is D -finite (taking $m_1(i, j, k) = (i+k)/2$ and $m_2 = m_3 = 0$). But $F(x) = \sum_{n \geq 0} \sum_{k=0}^n f_1(n-k) f_2(k) f_3(n+k) x^n$.

ACKNOWLEDGMENTS

I thank J. Denef and B. Dwork for their comments on an earlier draft of this paper. This work was supported in part by a grant from the NSF.

Note added in proof. H. Sharif (University of Kent, Canterbury, England) has informed me (April 1987), that he has independently proved the theorem of this paper, by somewhat different methods.

REFERENCES

1. G. CHRISTOL, "Diagonals de fractions rationnelles et équations différentielles," Groupe d'étude d'Analyse ultramétrique, No. 18, 1982/1983.
2. P. DELIGNE, Intégration sur un cycle évanescant, *Invent. Math.* **76** (1984), 129–143.
3. J. DENEFF AND L. LIPSHITZ, Algebraic power series and diagonals, *J. Number Theory* **26** (1987), 46–67.
4. B. DWORK, On the zeta function of a hypersurface, IV, *Ann. of Math.* **90** (1969), 335–352.
5. B. DWORK, Differential systems associated with families of singular hypersurfaces, preprint.
6. H. FURSTENBERG, Algebraic functions over finite fields, *J. Algebra* **7** (1967), 271–277.
7. I. GESSEL, Two theorems on rational power series, *Utilitas Math.* **19** (1981), 247–254.
8. L. LIPSHITZ AND L. RUBEL, A gap theorem for power series solutions of algebraic differential equations, *Amer. J. Math.* **108** (1986), 1193–1214.
9. R. P. STANLEY, Differentiably finite power series, *Eur. J. Combin.* **1** (1980), 175–188.
10. D. ZEILBERGER, Sister Celine's technique and its generalizations, *J. Math. Anal. Appl.* **85** (1982), 114–145.