

## Quickly Excluding a Planar Graph

NEIL ROBERTSON\*

*Department of Mathematics, Ohio State University,  
231 West 18th Avenue, Columbus, Ohio 43210*

PAUL SEYMOUR

*Bellcore, 445 South Street,  
Morristown, New Jersey 07960*

AND

ROBIN THOMAS†

*DIMACS Center, Hill Center, Busch Campus, Rutgers University,  
New Brunswick, New Jersey 08903*

Received October 18, 1989; revised March 14, 1994

In an earlier paper, the first two authors proved that for any planar graph  $H$ , every graph with no minor isomorphic to  $H$  has bounded tree width; but the bound given there was enormous. Here we prove a much better bound. We also improve the best known bound on the tree-width of planar graphs with no minor isomorphic to a  $g \times g$  grid. © 1994 Academic Press, Inc.

### 1. INTRODUCTION

A graph  $H$  is a *minor* of a graph  $G$  if  $H$  can be obtained from a subgraph of  $G$  by contracting edges. (In this paper, all graphs are finite and may have loops or multiple edges.) A *tree-decomposition* of a graph  $G$  is a pair  $(T, W)$ , where  $T$  is a tree and  $W = (W_t : t \in V(T))$  is a family of subsets of  $V(G)$ , such that

(i)  $\bigcup_{t \in V(T)} W_t = V(G)$ , and for every edge  $e \in E(G)$  there exists  $t \in V(T)$  such that  $W_t$  contains both ends of  $e$

(ii) if  $t, t', t'' \in V(T)$  and  $t'$  lies on the path of  $T$  between  $t$  and  $t''$  then  $W_t \cap W_{t''} \subseteq W_{t'}$ .

\* This research was partially supported by NSF Grant DMS8903132 and was performed under a consulting agreement with Bellcore.

† Current address: School of Mathematics, Georgia Institute of Technology, Atlanta, Georgia 30332.

( $V(G)$  and  $E(G)$  denote the vertex- and edge-sets of  $G$ .) The *width* of  $(T, W)$  is  $\max_{t \in V(T)}(|W_t| - 1)$ , and the *tree-width* of  $G$  is the minimum width of a tree-decomposition of  $G$ . The following was proved in [6]:

(1.1). *For any planar graph  $H$  there is a number  $N$  such that every graph with no minor isomorphic to  $H$  has tree-width  $\leq N$ .*

An explicit formula for a number  $N$  with this property was given (in terms of  $|V(H)|$ ,  $|E(H)|$ ) but it was enormous; it involved iterated exponentiation where the number of iterations also involved iterated exponentiation (and so on, twice more). Our main objective here is to give a much better bound on  $N$  in the following:

(1.2). *Let  $H$  be a planar graph and let  $n = 2|V(H)| + 4|E(H)|$ . Then every graph with no minor isomorphic to  $H$  has tree-width  $\leq 20^{2n^5}$ .*

To prove this, we first reduce the problem to the case when  $H$  is a  $g \times g$  grid, that is, the simple graph with  $g^2$  vertices  $\{(i, j) : 1 \leq i, j \leq g\}$ , where  $(i, j)$  and  $(i', j')$  are adjacent if  $|i' - i| + |j' - j| = 1$ . This reduction will occupy the remainder of this section, and Sections 2–5 are concerned with the grid case. Section 6 gives a similar improvement of a theorem of [5], concerning the tree-width of planar graphs with no minor isomorphic to a  $g \times g$  grid.

The main part of the proof of (1.2), in Sections 2–5, relies heavily on the concept of a “tangle” in a graph. This is defined in Section 2, but, roughly, a tangle in  $G$  is a part of  $G$  which is highly coherent, so that for every small-order separation of  $G$  the tangle clearly resides on one side of it or the other. It is shown in [8] that a graph has large tree-width if and only if it has a large tangle, and so our problem is to show that a graph with a large tangle has a large grid minor. The way we construct the large grid minor is rather like the way one knits a scarf, stitch by stitch and row by row. The main lemmas assert that if part of the minor has been knitted and we still have a large tangle in its complement, then we can add another stitch to our minor without destroying too much of the remaining tangle.

Now let us reduce (1.2) to the case when  $H$  is a grid. We begin with the following:

(1.3). *If  $H$  is a simple planar Hamiltonian graph with  $g$  vertices then  $H$  is isomorphic to a minor of a  $g \times g$  grid.*

*Proof.* We may assume that  $H$  is a triangulation, that is, every region of  $H$  is bordered by a three-edge circuit. Let  $V(H) = \{v_1, \dots, v_g\}$ , where  $v_i$  and  $v_{i+1}$  are adjacent ( $1 \leq i \leq g$ ), where  $v_{g+1}$  means  $v_1$ . Let  $C$  be the

corresponding Hamiltonian circuit and take a drawing of  $H$  in a sphere. Let  $A, B$  be the subgraphs of  $H$  drawn in the two closed discs bounded by  $C$  (thus,  $A$  and  $B$  are both "outerplanar" graphs and  $C$  is a Hamiltonian circuit of both of them). Let  $G$  be the  $g \times g$  grid with vertex set  $\{(i, j) : 1 \leq i, j \leq g\}$  as before. For  $2 \leq k \leq g$ , choose  $i(k), j(k)$  with  $1 \leq i(k), j(k) \leq k-1$ , minimum such that  $v_{i(k)}, v_k$  are adjacent in  $A$ , and  $v_{j(k)}, v_k$  are adjacent in  $B$ . For  $1 \leq k \leq g-1$ , choose  $i'(k), j'(k)$  with  $k+1 \leq i'(k), j'(k) \leq g$ , maximum such that  $v_k, v_{i'(k)}$  are adjacent in  $A$  and  $v_k, v_{j'(k)}$  are adjacent in  $B$ . For  $2 \leq k \leq g-1$ , let  $X_k$  be the subgraph of  $G$  induced by

$$\{(i, k) : i(k) < i < i'(k)\} \cup \{(k, j) : j(k) < j < j'(k)\},$$

and let  $X_1, X_g$  be the subgraphs induced respectively by

$$\begin{aligned} &\{(i, 1) : 1 \leq i < i'(1)\} \cup \{(1, j) : 1 \leq j < j'(1)\} \\ &\{(i, g) : i(g) \leq i \leq g\} \cup \{(g, j) : j(g) \leq j \leq g\}. \end{aligned}$$

Then each  $X_k$  is connected, and  $X_1, \dots, X_g$  are mutually disjoint, as is easily seen. Moreover, if  $k < k'$  and  $j'(k) = k'$  and  $i(k') = k$  then  $k = 1$  and  $k' = g$ , since  $H$  is a triangulation; and the same holds if  $i'(k) = k'$  and  $j(k') = k$ . It follows that if  $v_k, v_{k'}$  are adjacent in  $H$ , where  $k' > k$ , then there is an edge of  $G$  between  $V(X_k)$  and  $V(X_{k'})$ . (For instance, if  $v_k, v_{k'}$  are adjacent in  $A$  then  $(k+1, k') \in V(X_{k'})$ ,  $(k, k'-1) \in V(X_k)$ , and  $(k, k')$  belongs to one of  $V(X_k), V(X_{k'})$ .) Thus  $G$  has a minor isomorphic to  $H$ , as required. ■

(1.4). If  $n \geq 2$  and  $H$  is a simple planar graph with  $\leq n$  vertices, then there is a simple planar Hamiltonian graph  $G$  with  $\leq 2n$  vertices such that  $G$  has a minor isomorphic to  $H$ .

*Proof.* We may assume that  $|V(H)| \geq 4$  and  $H$  is a triangulation. It follows easily by induction that a triangulation  $T$  with  $n \geq 4$  vertices has at most  $n-4$  "separating triangles," that is, circuits of length 3 whose deletion disconnects  $T$ . Choose an edge  $e$  from a separating triangle of  $H$  if there is one, and let  $r_1, r_2$  be the two regions bordered by  $e$ . Let the four vertices incident with  $r_1$  or  $r_2$  be  $v_1, v_2, v_3, v_4$ . Delete  $e$  and add a new vertex adjacent to  $v_1, v_2, v_3, v_4$ . This procedure reduces the number of separating triangles by at least one, and so by applying this procedure at most  $n-4$  times, we obtain a triangulation  $G$  with no separating triangle which has a minor isomorphic to  $H$ . By Whitney's theorem [9]  $G$  is Hamiltonian, as required. ■

(1.5). If  $H$  is a planar graph with  $|V(H)| + 2|E(H)| \leq n$ , then  $H$  is isomorphic to a minor of the  $2n \times 2n$  grid.

*Proof.* We may assume that  $n \geq 2$ . By subdividing each edge of  $H$  twice, we see that  $H$  is isomorphic to a minor of a simple planar graph  $H_1$  with  $|V(H_1)| \leq n$ . By (1.4)  $H_1$  is isomorphic to a minor of a simple planar Hamiltonian graph with  $\leq 2n$  vertices. By (1.3) the result follows. ■

Thus, to prove (1.2) it suffices to prove the following.

(1.6) *For  $g \geq 1$ , every graph with no minor isomorphic to the  $g \times g$  grid has tree-width  $\leq 20^{2g^5}$ .*

Although (1.6) gives a much better bound than was previously known, there is still no reason to think that it is close to the right answer. By a probabilistic argument based on girth considerations (a slight variant on a construction of Erdős [2]), we have proved the existence of graphs which have no  $g \times g$  grid minor and have tree-width at least proportional to  $g^2 \log g$ . We suspect that  $O(g^2 \log g)$  is closer to the right answer than the bound of (1.6); indeed, it may even be the right answer.

## 2. TANGLES

The proof of (1.6) is based on the concept of a “tangle,” developed in [8]. A *separation* of a graph  $G$  is a pair  $(A, B)$  of subgraphs with  $A \cup B = G$  and  $E(A \cap B) = \emptyset$ ; its *order* is  $|V(A \cap B)|$ . If  $\theta \geq 1$  is an integer, a *tangle of order  $\theta$*  in  $G$  is a set  $\mathcal{T}$  of separations of order  $< \theta$ , such that

- (i) if  $(A, B)$  is a separation of  $G$  of order  $< \theta$ , then  $\mathcal{T}$  contains one of  $(A, B)$ ,  $(B, A)$
- (ii) if  $(A_1, B_1), (A_2, B_2), (A_3, B_3) \in \mathcal{T}$  then  $A_1 \cup A_2 \cup A_3 \neq G$
- (iii) if  $(A, B) \in \mathcal{T}$  then  $V(A) \neq V(G)$ .

(We refer to these as the *first*, *second*, and *third axioms*.) We define  $\text{ord}(\mathcal{T}) = \theta$ .

In this section we shall present some preliminary lemmas about tangles. It is proved in [8, Theorem (5.2)] that

(2.1) *If  $G$  has a tangle of order  $\geq \theta$ , then the tree-width of  $G$  is at least  $\theta - 1$ . Conversely, if  $G$  has no tangle of order  $\geq \theta$ , then its tree-width is at most  $3\theta/2$ .*

This will enable us to use the presence of tangles rather than the absence of tree-decompositions to prove the existence of a large grid minor. We shall prove the following, from which (1.6) follows via (2.1).

(2.2) *For  $g \geq 1$ , every graph with a tangle of order  $\geq 20^{g^4(2g-1)}$  has a minor isomorphic to the  $g \times g$  grid.*

Let  $H$  be a  $g \times g$  grid minor of  $G$ , where  $g \geq 2$ , with vertex set  $\{(i, j) : 1 \leq i, j \leq g\}$ , labelled in the natural way. For  $1 \leq i \leq g$  the set of edges of  $H$  joining two vertices of  $H$  with first coordinate  $i$  is called a *row* of  $H$ , and the *columns* of  $H$  are defined similarly. Since  $E(H) \subseteq E(G)$ , the rows and columns of  $H$  are subsets of  $E(G)$ . Let  $\mathcal{T}_H$  be the set of all separations  $(A, B)$  of  $G$  of order  $< g$  such that  $E(B)$  includes a row of  $H$  (or equivalently, by [8, Theorem (7.1)], includes a column of  $H$ ). It follows easily from [8, Theorems (7.3) and (6.1)] that  $\mathcal{T}_H$  is a tangle in  $G$  of order  $g$ . A tangle  $\mathcal{T}$  in  $G$  *dominates*  $H$  if  $\mathcal{T}_H \subseteq \mathcal{T}$ .

For applications in future papers of the Graph Minors series, it is convenient to prove a strengthening of (2.2), the following:

(2.3). *Let  $g \geq 2$ , and let  $\mathcal{T}$  be a tangle in  $G$  of order  $\geq 20g^{4(2g-1)}$ . Then  $\mathcal{T}$  dominates a  $g \times g$  grid minor.*

A (numerically weaker) form of (2.3) was included in an early version of [8], but it turns out to be easier to prove it here instead. We see that (2.3) implies (2.2), because (2.2) is obvious when  $g = 1$ . We shall prove (2.3) in Section 5.

Let  $\mathcal{T}$  be a tangle in  $G$ . For  $X \subseteq V(G)$ , we define  $\text{rk}_{\mathcal{T}}(X)$  to be the minimum order of a separation  $(A, B) \in \mathcal{T}$  with  $X \subseteq V(A)$ , if there is such an  $(A, B)$ , and otherwise  $\text{rk}_{\mathcal{T}}(X) = \text{ord}(\mathcal{T})$ . It follows from the third axiom that  $\text{rk}_{\mathcal{T}}(V(G)) = \text{ord}(\mathcal{T})$ . Let us say that  $X \subseteq V(G)$  is *free* (relative to  $\mathcal{T}$ ) if  $|X| \leq \text{ord}(\mathcal{T})$  and there is no  $(A, B) \in \mathcal{T}$  of order  $< |X|$  with  $X \subseteq V(A)$ . From [8, Theorems (12.1) and (12.2)] and the fact that  $\text{rk}_{\mathcal{T}}(V(G)) = \text{ord}(\mathcal{T})$ , it follows that

(2.4).  *$\text{rk}_{\mathcal{T}}$  is the rank function of a matroid on  $V(G)$  with rank  $\text{ord}(\mathcal{T})$ , and its independent sets are the free sets.*

Let  $H$  be a subgraph of  $G$  and let  $\mathcal{T}'$  be a tangle in  $H$ . Let  $\mathcal{T}$  be the set of all separations  $(A, B)$  of  $G$  of order  $< \text{ord}(\mathcal{T}')$  such that  $(A \cap H, B \cap H) \in \mathcal{T}'$ . It is easy to see that

(2.5).  *$\mathcal{T}$  thus defined is a tangle in  $G$  of order  $\text{ord}(\mathcal{T}')$ .*

We call  $\mathcal{T}$  the tangle in  $G$  induced by  $\mathcal{T}'$ . If  $\mathcal{T}''$  is any tangle in  $G$  with  $\mathcal{T} \subseteq \mathcal{T}''$ , we say that  $\mathcal{T}'$  is *conformal* with  $\mathcal{T}''$ . Certainly  $\mathcal{T}'$  is conformal with  $\mathcal{T}$  itself.

Since we shall have many occasions to verify that some tangle is conformal with another, let us point out the following short-cut.

(2.6). *Let  $H$  be a subgraph of  $G$ , let  $\mathcal{T}$  be a tangle in  $G$ , and let  $\mathcal{T}'$  be a tangle in  $H$ . Then  $\mathcal{T}'$  is conformal with  $\mathcal{T}$  if and only if  $\text{ord}(\mathcal{T}') \leq \text{ord}(\mathcal{T})$  and  $(A \cap H, B \cap H) \in \mathcal{T}'$  for every  $(A, B) \in \mathcal{T}$  of order  $< \text{ord}(\mathcal{T}')$ .*

*Proof.* We assume  $\text{ord}(\mathcal{T}') \leq \text{ord}(\mathcal{T})$ , for otherwise  $\mathcal{T}'$  is not conformal with  $\mathcal{T}$ . Now  $\mathcal{T}'$  is conformal with  $\mathcal{T}$  if and only if for every separation  $(A, B)$  of  $G$  of order  $< \text{ord}(\mathcal{T}')$ , if  $(A \cap H, B \cap H) \in \mathcal{T}'$  then  $(A, B) \in \mathcal{T}$ ; or equivalently (since exactly one of  $(A \cap H, B \cap H)$ ,  $(B \cap H, A \cap H) \in \mathcal{T}'$ , and exactly one of  $(A, B)$ ,  $(B, A) \in \mathcal{T}$  because  $\text{ord}(\mathcal{T}') \leq \text{ord}(\mathcal{T})$ ) if  $(B, A) \in \mathcal{T}$  then  $(B \cap H, A \cap H) \in \mathcal{T}'$ . The result follows. ■

We shall also need the following.

(2.7). *Let  $\mathcal{T}$  be a tangle in  $G$ , let  $H$  be a subgraph of  $G$ , and let  $\mathcal{T}'$  be a tangle in  $H$  conformal with  $\mathcal{T}$ . For  $X \subseteq V(H)$ ,  $\text{rk}_{\mathcal{T}'}(X) \leq \text{rk}_{\mathcal{T}}(X)$ .*

*Proof.* Since  $\text{rk}_{\mathcal{T}'}(X) \leq \text{ord}(\mathcal{T}')$ , we may assume that  $\text{rk}_{\mathcal{T}}(X) < \text{ord}(\mathcal{T}')$ , and since  $\text{ord}(\mathcal{T}') \leq \text{ord}(\mathcal{T})$ , there exists  $(A, B) \in \mathcal{T}$  of order  $\text{rk}_{\mathcal{T}}(X)$  with  $X \subseteq V(A)$ . Now  $(A, B)$  has order  $\text{rk}_{\mathcal{T}}(X) < \text{ord}(\mathcal{T}')$ , and by (2.6),  $(A \cap H, B \cap H) \in \mathcal{T}'$ . Since  $X \subseteq V(A \cap H)$  it follows that

$$\text{rk}_{\mathcal{T}'}(X) \leq |V((A \cap H) \cap (B \cap H))| \leq |V(A \cap B)| = \text{rk}_{\mathcal{T}}(X)$$

as required. ■

The next result is proved in [8, Theorem (2.9)].

(2.8). *Let  $\mathcal{T}$  be a tangle in a graph  $G$ , let  $(A_1, B_1) \in \mathcal{T}$ , and let  $(A_2, B_2)$  be a separation of  $G$  of order  $< \text{ord}(\mathcal{T})$ . If either  $V(B_1) \subseteq V(B_2)$  or  $V(A_2) \subseteq V(A_1)$  then  $(A_2, B_2) \in \mathcal{T}$ .*

We denote by  $G \setminus Z$  the graph obtained from  $G$  by deleting  $Z$  (here  $Z$  may be a vertex or edge, or a set of vertices or edges). Let  $\mathcal{T}$  be a tangle in  $G$ , and let  $Z \subseteq V(G)$  with  $|Z| < \text{ord}(\mathcal{T})$ . We denote by  $\mathcal{T} \setminus Z$  the set of all separations  $(A', B')$  of  $G \setminus Z$  such that there exists  $(A, B) \in \mathcal{T}$  with  $Z \subseteq V(A \cap B)$  and  $A \setminus Z = A'$ ,  $B \setminus Z = B'$ .

(2.9).  *$\mathcal{T} \setminus Z$  is a tangle in  $G \setminus Z$  of order  $\text{ord}(\mathcal{T}) - |Z|$ , conformal with  $\mathcal{T}$ .*

*Proof.* The first assertion is proved in [8, Theorem (8.5)]. For the second, we use (2.6). Let  $(A, B) \in \mathcal{T}$  have order  $< \text{ord}(\mathcal{T}) - |Z|$ . Choose a separation  $(A^+, B^+)$  of  $G$  with  $Z \subseteq V(A^+ \cap B^+)$ ,  $A \subseteq A^+$ ,  $B \subseteq B^+$  and  $A^+ \setminus Z = A \cap (G \setminus Z)$ ,  $B^+ \setminus Z = B \cap (G \setminus Z)$ . Then  $(A^+, B^+)$  has order  $< \text{ord}(\mathcal{T})$ . Now  $V(B) \subseteq V(B^+)$ , and so by (2.8),  $(A^+, B^+) \in \mathcal{T}$ . Hence  $(A^+ \setminus Z, B^+ \setminus Z) \in \mathcal{T} \setminus Z$  by definition of  $\mathcal{T} \setminus Z$ , and so  $(A \cap (G \setminus Z), B \cap (G \setminus Z)) \in \mathcal{T} \setminus Z$ . By (2.6),  $\mathcal{T} \setminus Z$  is conformal with  $\mathcal{T}$ . ■

(2.10). Let  $\mathcal{T}$  be a tangle in a graph  $G$ , and let  $z_1, \dots, z_k \in V(G)$  be distinct, such that  $\{z_1, \dots, z_k\}$  is free relative to  $\mathcal{T}$  and  $\text{ord}(\mathcal{T}) \geq 2k$ . Then there exist  $y_1, \dots, y_k \in V(G \setminus \{z_1, \dots, z_k\})$ , distinct, such that

- (i) for  $1 \leq i \leq k$ ,  $y_i$  is adjacent in  $G$  to  $z_i$ , and
- (ii)  $\{y_1, \dots, y_k\}$  is free relative to  $\mathcal{T} \setminus \{z_1, \dots, z_k\}$ .

*Proof.* Let  $Z = \{z_1, \dots, z_k\}$ , let  $\mathcal{T}' = \mathcal{T} \setminus Z$ , let  $G' = G \setminus Z$  and for  $1 \leq i \leq k$ , let  $Y_i$  be the set of vertices of  $G'$  adjacent to  $z_i$  in  $G$ . First we show the following:

- (1) For all  $I \subseteq \{1, \dots, k\}$ ,  $\text{rk}_{\mathcal{T}'}(\bigcup (Y_i : i \in I)) \geq |I|$ .

*Subproof.* Suppose not. Since  $\text{ord}(\mathcal{T}') \geq k \geq |I|$ , there exists  $(A', B') \in \mathcal{T}'$  of order  $< |I|$  such that  $\bigcup (Y_i : i \in I) \subseteq V(A')$ . From the definition of  $\mathcal{T} \setminus Z$  there exists  $(A, B) \in \mathcal{T}$  with  $Z \subseteq V(A \cap B)$  and  $A \setminus Z = A'$ ,  $B \setminus Z = B'$ . By choosing  $A$  maximal, it follows from (2.8) that for each  $i \in I$ , every edge incident with  $z_i$  belongs to  $E(A)$ . Hence  $(A, B'')$  is a separation of  $G$ , where  $B'' = B \setminus \{z_i : i \in I\}$ . Its order is

$$|V(A \cap B)| - |I| = |V(A' \cap B')| + |Z| - |I| < |Z|;$$

and  $(B'', A) \notin \mathcal{T}$ , since  $(A, B) \in \mathcal{T}$  and  $A \cup B'' = G$ . Thus  $(A, B'') \in \mathcal{T}$ . But that contradicts the hypothesis that  $Z$  is free relative to  $\mathcal{T}$ . Thus there is no such  $(A', B')$ . This proves (1). ■

From a theorem of Rado [4], (2.4), and (1), it follows that there exist  $y_i \in Y_i$  ( $1 \leq i \leq k$ ) such that  $\text{rk}_{\mathcal{T}'}\{y_1, \dots, y_k\} = k$ , and so, by (2.4),  $\{y_1, \dots, y_k\}$  is free relative to  $\mathcal{T}'$ . ■

- (2.11). Let  $\mathcal{T}$  be a tangle in  $G$  and let  $(C, D) \in \mathcal{T}$  with order 0. Let

$$\mathcal{T}' = \{(A \cap D, B \cap D) : (A, B) \in \mathcal{T}\}.$$

Then  $\mathcal{T}'$  is a tangle in  $D$  with the same order as  $\mathcal{T}$ , and it is conformal with  $\mathcal{T}$ .

*Proof.* We must verify the three axioms. Let  $(A', B')$  be a separation of  $D$  of order  $< \text{ord}(\mathcal{T})$ . Then  $(A' \cup C, B')$  is a separation of  $G$  of the same order, and so one of  $(A' \cup C, B')$ ,  $(B', A' \cup C) \in \mathcal{T}$ . Hence one of  $(A', B')$ ,  $(B', A') \in \mathcal{T}'$ . This verifies the first axiom.

Let  $(A'_i, B'_i) \in \mathcal{T}'$  ( $i = 1, 2, 3$ ) and suppose that  $A'_1 \cup A'_2 \cup A'_3 = D$ . There exists  $(A_i, B_i) \in \mathcal{T}$  with  $A'_i = A_i \cap D$ ,  $B'_i = B_i \cap D$  ( $i = 1, 2, 3$ ). Now  $(A_i \cup C, B_i \cap D)$  has order at most that of  $(A_i, B_i)$ , and  $(B_i \cap D, A_i \cup C) \notin \mathcal{T}$  since

$(C, D) \in \mathcal{T}$  and  $A_i \cup C \cup (B_i \cap D) = G$ . Thus  $(A_i \cup C, B_i \cap D) \in \mathcal{T}$  ( $i = 1, 2, 3$ ). But  $(A_1 \cup C) \cup (A_2 \cup C) \cup (A_3 \cup C) = G$ , a contradiction. This verifies the second axiom.

Let  $(A', B') \in \mathcal{T}'$  and choose  $(A, B) \in \mathcal{T}$  with  $A \cap D = A'$ ,  $B \cap D = B'$ . Then  $(A \cup C, B \cap D) \in \mathcal{T}$  as before, and so  $V(A \cup C) \neq V(G)$ , that is,  $V(A) \neq V(D)$ . This verifies the third axiom.

Thus  $\mathcal{T}'$  is a tangle in  $D$ ; and by (2.6) it is conformal with  $\mathcal{T}$ , as required. ■

(2.12). *Let  $\mathcal{T}$  be a tangle in  $G$ , and let  $Z \subseteq V(G)$  be free relative to  $\mathcal{T}$ , with  $|Z| < \text{ord}(\mathcal{T})$ . Then there is a component  $C$  of  $G \setminus Z$  such that each vertex in  $Z$  has a neighbour in  $V(C)$ .*

*Proof.* Choose  $(A, B) \in \mathcal{T}$  with  $V(A \cap B) = Z$  and with  $B$  minimal. (This is possible, for setting  $B = G$  and  $V(A) = Z$ ,  $E(A) = \emptyset$  satisfies  $(A, B) \in \mathcal{T}$ , by the first and third axioms, since  $|Z| < \text{ord}(\mathcal{T})$ .)

(1) *There do not exist subgraphs  $B_1, B_2$  of  $B$  with  $B_1 \cup B_2 = B$ ,  $E(B_1 \cap B_2) = \emptyset$ ,  $V(B_1 \cap B_2) = Z$ , and  $V(B_1), V(B_2) \neq Z$ .*

*Subproof.* If  $B_1, B_2$  have these properties, then  $(A \cup B_1, B_2)$  is a separation of  $G$ , and  $V((A \cup B_1) \cap B_2) = Z$ . From the minimality of  $B$ ,  $(A \cup B_1, B_2) \notin \mathcal{T}$ , and so  $(B_2, A \cup B_1) \in \mathcal{T}$ . Similarly  $(B_1, A \cup B_2) \in \mathcal{T}$ ; but  $(A, B) \in \mathcal{T}$  and  $A \cup B_1 \cup B_2 = G$ , contrary to the second axiom.

Let  $C = B \setminus Z$ . Then  $C$  is non-null, from the third axiom. From (1), (2.8) and the minimality of  $B$ , we deduce that  $C$  is connected and  $V(C)$  meets every edge of  $B$ . Now let  $v \in Z$ ; to complete the proof, we must show that  $v$  has a neighbour in  $V(C)$ . Suppose not. Since  $V(C)$  meets every edge of  $B$ ,  $v$  is an isolated vertex of  $B$ , and so  $(A, B \setminus v)$  is a separation of  $G$ . By (2.8),  $(A, B \setminus v) \in \mathcal{T}$ . But  $Z \subseteq V(A)$ , and  $Z$  is free relative to  $\mathcal{T}$ , a contradiction, as required. ■

### 3. LINKING TWO VERTICES

The object of this section is to prove that if  $G$  has a large order tangle and  $s, t \in V(G)$ , and  $\{s, t\}$  is free, then there is a path  $P$  between  $s$  and  $t$  such that  $G \setminus V(P)$  still has a large order tangle (not as large as the original, but of order at least  $\frac{1}{18}$  the original). If  $H$  is a subgraph of  $G$  we write  $H \subseteq G$ . We begin with the following lemma.

(3.1). *Let  $k \geq 1$  be an integer, let  $G$  be a graph and let  $Z \subseteq V(G)$  with  $|Z| = 18k$ . Suppose that for all  $X, Y \subseteq Z$  with  $|X| = |Y|$  there are  $|X|$  mutually vertex-disjoint paths between  $X$  and  $Y$ . Let  $(S, T)$  be a partition of  $Z$  with  $|S| = |T| = 9k$ . Then there is a path  $P \subseteq G$  with one end in  $S$  and the*



other in  $T$ , and with no internal vertex in  $Z$ , such that there is no separation  $(A, B)$  of  $G \setminus V(P)$  of order  $< k$  with  $|V(A) \cap Z|, |V(B) \cap Z| \geq 6k$ .

*Proof.* Let  $\mathcal{T}$  be the tangle of order  $6k$  consisting of all separations  $(A, B)$  of order  $< 6k$  with  $|V(A) \cap Z| \leq |V(A \cap B)|$ . (We observe that this is a tangle, for if  $(A, B)$  is a separation then one of  $|V(A) \cap Z|, |V(B) \cap Z| \leq |V(A \cap B)|$  from our hypothesis.)

(1) For any path  $P$  from  $S$  to  $T$  with no internal vertex in  $Z$ , if  $\text{rk}_{\mathcal{T}}(V(P)) \leq 5k$  then  $P$  satisfies the theorem.

*Subproof.* Let  $(C, D) \in \mathcal{T}$  with  $V(P) \subseteq V(C)$  and  $|V(C \cap D)| \leq 5k$ . Suppose that  $P$  does not satisfy the theorem; then there is a separation  $(A, B)$  of  $G$  with  $V(P) \subseteq V(A \cap B)$  and  $|V(A \cap B) - V(P)| < k$  such that  $|V(A) \cap Z|, |V(B) \cap Z| \geq 6k + 2$ . Now since  $(C, D) \in \mathcal{T}$  it follows that  $|V(C) \cap Z| \leq |V(C \cap D)| \leq 5k$ , and so  $|V(D) \cap Z| \geq 13k$ . From the symmetry between  $A$  and  $B$ , we may assume that

$$|V(A) \cap V(D) \cap Z| \geq 13k/2 > 6k.$$

Now  $(C', D') = (A \cap D, B \cup C)$  is a separation of  $G$  of order  $< 6k$ , because

$$\begin{aligned} V((A \cap D) \cap (B \cup C)) &\subseteq V(C \cap D) \cup (V(A \cap B) - V(C)) \\ &\subseteq V(C \cap D) \cup (V(A \cap B) - V(P)). \end{aligned}$$

But  $|V(C') \cap Z| > 6k$ , and  $|V(D') \cap Z| \geq |V(B) \cap Z| > 6k$ . Thus  $(C', D')$ ,  $(D', C') \notin \mathcal{T}$ , a contradiction. This proves (1).

Let  $\mathcal{P}$  be a set of  $9k$  mutually disjoint paths between  $S$  and  $T$ . Then no member of  $\mathcal{P}$  has an internal vertex in  $Z$ , and we shall prove that one of them satisfies the theorem. If  $P_1, P_2, P_3 \in \mathcal{P}$  are distinct, let us say that  $P_2$  splits  $P_1$  and  $P_3$  if there do not exist  $k$  mutually disjoint paths of  $G$  from  $V(P_1)$  to  $V(P_3)$  all disjoint from  $V(P_2)$ . We claim that we may assume that

(2) If  $P_1, P_2, P_3 \in \mathcal{P}$  are distinct and  $P_2$  splits  $P_1$  and  $P_3$ , then  $P_3$  does not split  $P_1$  and  $P_2$ .

*Subproof.* By (1) we may assume that  $\text{rk}_{\mathcal{T}}(V(P_1)), \text{rk}_{\mathcal{T}}(V(P_2)) > 5k$ , and so there are  $> 5k$  mutually disjoint paths of  $G$  from  $V(P_1)$  to  $V(P_2)$ . Thus, either there are  $> 5k/2$  disjoint paths between  $V(P_1)$  and  $V(P_2)$  avoiding  $V(P_3)$ , or  $> 5k/2$  disjoint paths between  $V(P_1)$  and  $V(P_3)$  avoiding  $V(P_2)$ . The result follows.

By (2), there are at most  $\binom{9k}{3}$  pairs  $(P_2, \{P_1, P_3\})$ , where  $P_1, P_2, P_3 \in \mathcal{P}$  are distinct and  $P_2$  splits  $P_1$  and  $P_3$ . Hence the average number of unordered pairs  $\{P_1, P_3\}$  split by  $P_2 \in \mathcal{P}$  (averaged over all  $P_2 \in \mathcal{P}$ ) is at most

$\binom{9k}{3}/9k < 55k^2/4$ . Consequently, there exists  $P \in \mathcal{P}$  which splits fewer than  $55k^2/4$  unordered pairs  $P_1, P_2$ . We claim that  $P$  satisfies the theorem. For suppose not, and let  $(A, B)$  be a separation of  $G$  with  $V(P) \subseteq V(A \cap B)$  and  $|V(A \cap B)| - |V(P)| < k$ , such that  $|V(A) \cap Z|, |V(B) \cap Z| \geq 6k + 2$ . Let  $a$  be the number of members  $P' \in \mathcal{P}$  with  $V(P') \subseteq V(A)$  and  $P' \neq P$ , and define  $b$  similarly. Then  $a + b + k \geq 9k$  since at most  $k$  members of  $\mathcal{P}$  intersect  $V(A \cap B)$ . Moreover, since some path of  $\mathcal{P}$  begins at each vertex of  $V(A) \cap Z$ , and at most  $k - 1$  of them leave  $V(A)$ , and all the rest (except for  $P$ ) contribute to  $a$ , it follows that

$$|V(A) \cap Z| \leq 2(a + 1) + k - 1.$$

Since  $|V(A) \cap Z| \geq 6k + 2$ , we deduce that  $2a \geq 5k$ , and similarly  $2b \geq 5k$ . We have shown then that  $a + b \geq 8k$ ,  $a \geq 5k/2$ ,  $b \geq 5k/2$ , and  $ab < 55k^2/4$  from the choice of  $P$ . But these four inequalities are contradictory. Thus  $P$  satisfies the theorem as required. ■

The main result of this section is the following.

(3.2). *Let  $k \geq 1$  be an integer, let  $s, t$  be distinct vertices of a graph  $G$ , and let  $\mathcal{T}$  be a tangle in  $G$  of order  $\geq 18k + 3$ , such that  $\{s, t\}$  is free relative to  $\mathcal{T}$ . Then there is a path  $P$  between  $s$  and  $t$ , and a tangle in  $G \setminus V(P)$  of order  $k$ , conformal with  $\mathcal{T}$ .*

*Proof.* Choose  $(A, B) \in \mathcal{T}$  with  $A$  maximal such that there is a partition  $(S, T)$  of  $V(A \cap B)$ , and two disjoint connected subgraphs  $J, K$  of  $A$ , satisfying the following conditions (1)–(4):

(1) *There is no  $(A', B') \in \mathcal{T}$  of order  $< |V(A \cap B)|$  such that  $A \subseteq A'$  and  $B' \subseteq B$ .*

(2)  $|S|, |T| \leq 9k + 1$ .

(3)  $s \in V(J)$ ,  $t \in V(K)$ ,  $V(J) \cap S \neq \emptyset \neq V(K) \cap T$ , and  $J, K$  both contain exactly one vertex of  $V(A \cap B)$ .

(4) *For each  $v \in V(A \cap B) - V(J \cup K)$ , if  $v \in S$  then  $v$  is adjacent in  $A$  to a vertex of  $J$ , and if  $v \in T$  then  $v$  is adjacent in  $A$  to a vertex of  $K$ .*

Such a choice of  $(A, B)$  is possible, because setting  $B = G$  and  $V(A) = \{s, t\}$ ,  $E(A) = \emptyset$ ,  $V(J) = \{s\}$ ,  $V(K) = \{t\}$  satisfies (1)–(4). Let  $Z = V(A \cap B)$ .

(5) *There is no  $(A', B') \in \mathcal{T}$  of order  $|Z|$  such that  $A' \neq A$ ,  $A \subseteq A'$  and  $B' \subseteq B$ .*

*Subproof.* Suppose that such a choice of  $(A', B')$  exists. Let  $|Z| = n$ . Suppose first that there is a separation  $(C, D)$  of  $B \cap A'$  of order  $< n$  with

$Z \subseteq V(C)$  and  $V(A' \cap B') \subseteq V(D)$ . Let  $A'' = A \cup C$ ,  $B'' = B' \cup D$ ; then  $(A'', B'')$  is a separation of  $G$  of order  $< n$  with  $A \subseteq A'' \subseteq A'$  and  $B' \subseteq B'' \subseteq B$ . But  $(A', B') \in \mathcal{T}$  and so  $(A'', B'') \in \mathcal{T}$  by (2.8), contrary to (1). Thus there is no such  $(C, D)$ ; and so by Menger's theorem, there are mutually disjoint paths  $P_1, \dots, P_n$  of  $B \cap A'$  from  $Z$  to  $V(A' \cap B')$ . Let  $P_i$  have first vertex  $v_i$  and last vertex  $v'_i$  ( $1 \leq i \leq n$ ). Let  $J'$  be the union of  $J$ , the unique path  $P_i$  such that  $v_i \in V(J)$ , and for each  $j$  such that  $v_j \in S - V(J)$  and  $v_j \neq v'_j$ , the path  $P_j \setminus v'_j$  and an edge of  $A$  joining  $v_j$  to a vertex of  $J$  (which exists by (4)). Define  $K'$  similarly. Let  $S' = \{v'_i : 1 \leq i \leq n, v_i \in S\}$ , and define  $T'$  similarly. Then  $(A', B')$ ,  $S'$ ,  $T'$ ,  $J'$ ,  $K'$  satisfy (1)–(4), contrary to the maximality of  $A$ . This proves (5).

(6)  $|S| = |T| = 9k + 1$ , and hence  $|Z| = 18k + 2$ .

*Subproof.* Suppose that  $|S| < 9k + 1$ , say. Let  $V(J) \cap Z = \{v\}$ . If  $A \cup (B \setminus v) = G$ , then  $(A, B \setminus v)$  is a separation of order smaller than  $|Z|$ , and by (2.8),  $(A, B \setminus v) \in \mathcal{T}$ , contrary to (1). Hence  $A \cup (B \setminus v) \neq G$ , and so there is an edge  $e$  of  $B$  incident with  $v$ . Some end,  $u$ , say, of  $e$  is not in  $V(A)$ , by (2.8) and the maximality of  $A$ . Let  $V(A') = V(A) \cup \{u\}$ ,  $E(A') = E(A) \cup \{e\}$ ,  $B' = B \setminus e$ ; then  $(A', B')$  has order  $1 + |Z| \leq 18k + 2$  and  $(A', B') \in \mathcal{T}$  by (2.8). Let  $S' = S \cup \{u\}$ ; then  $A'$ ,  $B'$ ,  $S'$ ,  $T$ ,  $J$ ,  $K$  satisfy (1)–(4), because of (5), contrary to the maximality of  $A$ . This proves (6).

(7) If  $X, Y \subseteq Z$  with  $|X| = |Y|$ , there are  $|X|$  mutually disjoint paths of  $B$  between  $X$  and  $Y$ , each with no internal vertex in  $Z$ .

*Subproof.* Suppose not. Then by Menger's theorem, there is a separation  $(C, D)$  of  $B$  with  $Z - (X \cup Y) \subseteq V(C \cap D)$  and

$$|V(C \cap D) - (Z - (X \cup Y))| < |X|$$

such that  $X \subseteq V(C)$  and  $Y \subseteq V(D)$ . Now  $(A \cup C, D)$  has order  $< |Z|$ , for  $V(A \cap D) - V(C) \subseteq Y - X$ , and so

$$\begin{aligned} |V((A \cup C) \cap D)| &= |V(C \cap D) - (Z - (X \cup Y))| + |Z - (X \cup Y)| \\ &\quad + |V(A \cap D) - V(C)| \\ &< |X| + |Z - (X \cup Y)| + |Y - X| = |Z|. \end{aligned}$$

From (1), it follows that  $(A \cup C, D) \notin \mathcal{T}$ , and so  $(D, A \cup C) \in \mathcal{T}$ . Similarly,  $(C, A \cup D) \in \mathcal{T}$ ; but  $(A, B) \in \mathcal{T}$ , and  $A \cup C \cup D = G$ , contrary to the second axiom. This proves (7).

Let  $V(J) \cap Z = \{v\}$  and  $V(K) \cap Z = \{w\}$ , and let  $G' = B \setminus \{v, w\}$ ,  $Z' = Z - \{v, w\}$ ,  $S' = S - \{v\}$ ,  $T' = T - \{w\}$ . By (6), (7), and (3.1) applied to  $G'$ ,  $Z'$ ,  $S'$ ,  $T'$  we deduce

(8) *There is a path  $P'$  of  $G'$  from  $S'$  to  $T'$  with no internal vertex in  $Z'$ , such that there is no separation  $(A', B')$  of  $G' \setminus V(P')$  of order  $< k$  with  $|V(A') \cap Z'|, |V(B') \cap Z'| \geq 6k$ .*

Now let  $P$  be a path of  $G$  from  $s$  to  $t$  with  $P \cap G' = P'$  (this exists and can be chosen with  $V(P) \subseteq V(J \cup K \cup P')$  because of (4)). Let  $\mathcal{T}'$  be the set of all separations  $(A', B')$  of  $G' \setminus V(P)$  of order  $< k$  such that  $|V(A') \cap Z'| < 6k$ . We claim that  $\mathcal{T}'$  is a tangle in  $G' \setminus V(P)$  of order  $k$ . For the second and third axioms are clearly satisfied, since  $|Z' \cap V(G' \setminus V(P))| = 18k - 2 > 3(6k - 1)$ . To see the first axiom, let  $(A', B')$  be a separation of  $G' \setminus V(P)$  of order  $< k$ ; then  $(A' \cap G', B' \cap G')$  is a separation of  $G' \setminus V(P')$  of order  $< k$ , and so by (8) one of  $|V(A' \cap G') \cap Z'|, |V(B' \cap G') \cap Z'| < k$ ; that is, one of  $|V(A') \cap Z'|, |V(B') \cap Z'| < k$ . It follows that one of  $(A', B')$ ,  $(B', A')$  belongs to  $\mathcal{T}'$ , and this verifies the first axiom. Hence  $\mathcal{T}'$  is a tangle in  $G' \setminus V(P)$  of order  $k$ .

(9)  *$\mathcal{T}'$  is conformal with  $\mathcal{T}$ .*

*Subproof.* Let  $(A_0, B_0)$  be a separation of  $G$  of order  $< k$ , such that  $(A', B') \in \mathcal{T}'$ , where  $A' = A_0 \cap (G' \setminus V(P))$ ,  $B' = B_0 \cap (G' \setminus V(P))$ . We must show that  $(A_0, B_0) \in \mathcal{T}$ . Now

$$V(P) \cap Z' = V(P \cap G') \cap Z' = V(P') \cap Z',$$

and so  $|V(P) \cap Z'| \leq 2$  by (8). But  $V(P) \cap Z \subseteq (V(P) \cap Z') \cup \{v, w\}$ , and so  $|V(P) \cap Z| \leq 4$ . Since  $|V(A') \cap Z'| \leq 6k - 1$  from the definition of  $\mathcal{T}'$  and

$$Z - V(B_0) \subseteq V(A_0) \cap Z \subseteq (V(A') \cap Z) \cup (V(P) \cap Z),$$

it follows that  $|Z - V(B_0)| \leq 6k + 3$ . But

$$V((A \cup B_0) \cap B \cap A_0) \subseteq V(A_0 \cap B_0) \cup (Z - V(B_0))$$

and  $|V(A_0 \cap B_0)| \leq k - 1$ , and so  $(A \cup B_0, B \cap A_0)$  has order  $\leq (6k + 3) + (k - 1) = 7k + 2$ . By (1) and (6),  $(A \cup B_0, B \cap A_0) \notin T$ , and so  $(B \cap A_0, A \cup B_0) \in \mathcal{T}$  by the first tangle axiom. It follows that  $(B_0, A_0) \notin \mathcal{T}$ , by the second axiom, because  $(A, B)$ ,  $(B \cap A_0, A \cup B_0) \in \mathcal{T}$  and  $A \cup (B \cap A_0) \cup B_0 = G$ . Hence  $(A_0, B_0) \in \mathcal{T}$ , as required. ■

From (9), the result follows. ■

#### 4. PRESERVING FREEDOM

As we said earlier, the purpose of (3.2) is to enable us to “knit” a grid minor, pulling out one stitch at a time from a tangle in the remainder of the graph, in such a way that after each stitch there still remains a large

order tangle in the remainder, so that we can continue. However, it is important that the part of the grid that we have already knitted (or at least the most recently knitted row of it) remain well connected to the tangle in the complement, for otherwise we shall not be able to add subsequent stitches where we want them. This is a serious problem (it must be, for otherwise one could construct clique minors just as well as grid minors), and to resolve it we shall modify our method of specifying exactly where the next stitch should attach. To this end, we notice that if the next stitch cannot be placed where we want it, we can infer a "non-planarity" in the graph, and enough non-planarities will give us a large clique minor. So either the stitches mostly come along where we want them, and we knit the grid we want, or we produce a large clique minor (which, of course, contains a large grid anyway).

Let  $G$  be a graph,  $\mathcal{T}$  a tangle in  $G$ , and  $Z \subseteq V(G)$ , free relative to  $\mathcal{T}$ . We call the triple  $G, \mathcal{T}, Z$  a *nodule* of order  $k$ , where  $k = \text{ord}(\mathcal{T})$ . Let  $G, \mathcal{T}, Z$  and  $G', \mathcal{T}', Z'$  be nodules. We say that the second is a *residual* nodule of the first if  $G'$  is a subgraph of  $G \setminus Z$ ,  $\mathcal{T}'$  is conformal with  $\mathcal{T}$ ,  $|Z'| = |Z|$ , and there are  $|Z|$  mutually vertex-disjoint paths of  $G$  between  $Z$  and  $Z'$ , each with no vertex in  $G'$  except its last vertex. We call such a set of paths a *linkage* (for  $G', \mathcal{T}', Z'$  from  $G, \mathcal{T}, Z$ ). We observe that, by (2.6), if  $G', \mathcal{T}', Z'$  is a residual nodule of  $G, \mathcal{T}, Z$  and  $G'', \mathcal{T}'', Z''$  is a residual nodule of  $G', \mathcal{T}', Z'$  then  $G'', \mathcal{T}'', Z''$  is a residual nodule of  $G, \mathcal{T}, Z$ .

If  $e \in E(G)$ ,  $G/e$  denotes the graph obtained by contracting  $e$ .

(4.1). *Let  $G$  be a graph, let  $Z \subseteq V(G)$ , and let  $C$  be a connected subgraph of  $G$  with  $V(C) \cap Z \neq \emptyset$ . Let  $\mathcal{T}$  be a tangle in  $G \setminus V(C)$  of order  $\geq 2|Z|$ , inducing a tangle  $\mathcal{T}_0$  in  $G$  such that  $G, \mathcal{T}_0, Z$  is a nodule. Then there is a residual nodule  $G', \mathcal{T}', Z'$  of order  $\text{ord}(\mathcal{T}) - |Z|$  such that  $C \cap G'$  is null.*

*Proof.* We proceed by induction on  $|V(C) - Z|$ . Let  $|Z| = h$ , and choose  $(A, B) \in \mathcal{T}_0$  of order  $h$  with  $Z \subseteq V(A)$  and  $A$  maximal. (This is possible, for  $V(A) = Z$ ,  $E(A) = \emptyset$ ,  $B = G$  is one such choice.) There are two cases.

*Case 1.*  $E(C) \not\subseteq E(A)$ . Choose  $e \in E(C) - E(A)$ . Let  $\mathcal{T}'_0$  be the tangle in  $G/e$  induced by  $\mathcal{T}$  (we see that  $G \setminus V(C)$  is a subgraph of  $G/e$  since  $e \in E(C)$ ). Now not both ends of  $e$  belong to  $Z$ , by the maximality of  $A$  and (2.8), and so no two members of  $Z$  are identified in forming  $G/e$ . Thus we may regard  $Z$  as a subset of  $V(G/e)$ . We claim that  $Z$  is free relative to  $\mathcal{T}'_0$ .

For if not, there is a separation  $(A'_0, B'_0) \in \mathcal{T}'_0$  of order  $< h$  with  $Z \subseteq V(A'_0)$ . Let  $w$  be the vertex of  $G/e$  formed by identifying the ends of  $e$ . If  $w \in V(A'_0) - V(B'_0)$  let  $B' = B'_0$ , and let  $A'$  be the subgraph of  $G$  with  $e \in E(A')$  and  $A'/e = A'_0$ . If  $w \in V(B'_0) - V(A'_0)$  let  $A' = A'_0$  and let  $B'$  be the

subgraph of  $G$  with  $e \in E(B')$  and  $B'/e = B'_0$ . If  $w \in V(A'_0 \cap B'_0)$  let  $A'$ ,  $B'$  be the subgraphs of  $G$  with  $e \in E(A')$ ,  $A'/e = A'_0$ ,  $e \notin E(B')$ ,  $(B' + e)/e = B'_0$ , with the natural definition of  $B' + e$ . In each case  $(A', B')$  is a separation of  $G$  and

$$(A' \cap (G \setminus V(C)), B' \cap (G \setminus V(C))) = (A'_0 \cap (G \setminus V(C)), B'_0 \cap (G \setminus V(C))) \in \mathcal{T}.$$

Thus  $(A', B') \in \mathcal{T}_0$ , because the order of  $(A', B')$  is at most one more than the order  $(A'_0, B'_0)$  and hence at most  $h + \text{ord}(\mathcal{T}_0)$ . Since  $Z$  is free with respect to  $\mathcal{T}_0$  and  $Z \subseteq V(A')$  it follows that  $|V(A' \cap B')| \geq h$ ; hence,  $|V(A' \cap B')| = h$  and  $e \in E(A')$ . Yet  $(A, B) \in \mathcal{T}_0$ , and  $(A \cup A', B \cap B')$  has order  $> h$ ; for otherwise it belongs to  $\mathcal{T}_0$ , contrary to the maximality of  $A$ , since  $e \in E(A') - E(A)$ . Since the sum of the orders of  $(A \cup A', B \cap B')$  and  $(A \cap A', B \cup B')$  equals the sum of the orders of  $(A, B)$  and  $(A', B')$  and, hence, is  $2h$ , it follows that  $(A \cap A', B \cup B')$  has order  $< h$ . Now  $(A, B) \in \mathcal{T}_0$  and so by (2.8),  $(A \cap A', B \cup B') \in \mathcal{T}_0$ , contrary to the freedom of  $Z$  relative to  $\mathcal{T}_0$ . This proves the freedom of  $Z$  relative to  $\mathcal{T}'_0$ .

From the inductive assumption applied to  $G/e$ ,  $C/e$  and  $\mathcal{T}$ , there exists a residual nodule  $G'$ ,  $\mathcal{T}'$ ,  $Z'$  of  $G/e$ ,  $\mathcal{T}'_0$ ,  $Z$  of order  $\text{ord}(\mathcal{T}) - |Z|$ , with  $G' \cap (C/e)$  null. We claim that  $\mathcal{T}'$  is conformal with  $\mathcal{T}_0$ . For let  $(A', B')$  be a separation of  $G$  of order  $< \text{ord}(\mathcal{T}') = \text{ord}(\mathcal{T}) - |Z|$ , such that  $(A' \cap G', B' \cap G') \in \mathcal{T}'$ . We must show that  $(A', B') \in \mathcal{T}_0$ . Let  $(A^*, B^*)$  be the separation of  $G/e$ , where  $A^* = A'/e$  if  $e \in E(A')$ ,  $A^* = A'$  if some end of  $e$  is not in  $V(A')$ , and  $A^*$  is obtained from  $A'$  by identifying the ends of  $e$  if  $e \in E(B')$  and both ends of  $e$  are in  $V(A')$  (and  $B^*$  is defined similarly). Now  $A' \cap G' = A^* \cap G'$  and  $B' \cap G' = B^* \cap G'$ , and  $(A^*, B^*)$  has order  $< \text{ord}(\mathcal{T}')$ ; and so  $(A^*, B^*) \in \mathcal{T}'_0$  since  $(A' \cap G', B' \cap G') \in \mathcal{T}'$  and  $\mathcal{T}'$  is conformal with  $\mathcal{T}'_0$ . By the definition of  $\mathcal{T}'_0$ ,

$$(A' \cap (G \setminus V(C)), B' \cap (G \setminus V(C))) = (A^* \cap (G \setminus V(C)), B^* \cap (G \setminus V(C))) \in \mathcal{T},$$

and so  $(A', B') \in \mathcal{T}_0$ . Thus  $\mathcal{T}'$  is conformal with  $\mathcal{T}_0$ . Hence  $G'$ ,  $\mathcal{T}'$ ,  $Z'$  is a residual nodule of  $G$ ,  $\mathcal{T}_0$ ,  $Z$  satisfying the theorem.

*Case 2.*  $E(C) \subseteq E(A)$ . Let  $V(A \cap B) = X = \{x_1, \dots, x_h\}$ . We claim that  $X$  is free relative to  $\mathcal{T}_0$ . For suppose not and let  $(A_0, B_0) \in \mathcal{T}_0$  with order  $< h$ , such that  $X \subseteq V(A_0)$ . Now

$$V((A \cup A_0) \cap (B \cap B_0)) \subseteq V(A_0 \cap B_0) \cup (X \cap V(B_0)) = V(A_0 \cap B_0)$$

and so  $(A \cup A_0, B \cap B_0)$  has order  $< h$ . Since  $Z \subseteq V(A \cup A_0)$  we deduce from the freedom of  $Z$  relative to  $\mathcal{T}_0$  that  $(A \cup A_0, B \cap B_0) \notin \mathcal{T}_0$ . Thus  $(B \cap B_0, A \cup A_0) \in \mathcal{T}_0$ ; but  $(A, B)$ ,  $(A_0, B_0) \in \mathcal{T}_0$  and  $A \cup A_0 \cup (B \cap B_0) = G$ , contrary to the second axiom. Thus  $X$  is free relative to  $\mathcal{T}_0$ .

By (2.10) there exist distinct  $y_1, \dots, y_h \in V(G) - X$  such that  $\{y_1, \dots, y_h\}$  is free relative to  $\mathcal{T}_0 \setminus X$  and  $y_i$  is adjacent to  $x_i$  ( $1 \leq i \leq h$ ). Since  $(A \setminus X, B \setminus X) \in \mathcal{T}_0 \setminus X$  and, hence,

$$\text{rk}_{\mathcal{T}_0 \setminus X}(V(A \setminus X)) = 0,$$

it follows by (2.4) that  $y_1, \dots, y_h \in V(B)$ . Let  $G' = B \setminus X$ . Since  $C$  is connected,  $V(C) \cap Z \neq \emptyset$  and  $E(C) \subseteq E(A)$ , it follows that  $V(C \cap G') = \emptyset$ . By (2.9) and (2.11) there is a tangle  $\mathcal{T}'$  in  $G'$  of order  $\text{ord}(\mathcal{T}) - h$ , conformal with  $\mathcal{T}_0 \setminus X$  and hence with  $\mathcal{T}_0$  (since "conformality" is transitive and  $\mathcal{T}_0 \setminus X$  is conformal with  $\mathcal{T}_0$ ).

We claim that  $\{y_1, \dots, y_h\}$  is free relative to  $\mathcal{T}'$ . For suppose not; then since  $\text{ord}(\mathcal{T}') = \text{ord}(\mathcal{T}) - h \geq h$ , there exists  $(A', B') \in \mathcal{T}'$  of order  $< h$  with  $\{y_1, \dots, y_h\} \subseteq V(A')$ . Since  $\mathcal{T}'$  is conformal with  $\mathcal{T}_0 \setminus X$  it follows that  $(A' \cup (A \setminus X), B') \in \mathcal{T}_0 \setminus X$ , and yet this separation has the same order as  $(A', B')$ , contrary to the freedom of  $\{y_1, \dots, y_h\}$  relative to  $\mathcal{T}_0 \setminus X$ . Thus  $\{y_1, \dots, y_h\}$  is free relative to  $\mathcal{T}'$ , and so  $G', \mathcal{T}', \{y_1, \dots, y_h\}$  is a nodule.

We claim that there are  $h$  disjoint paths of  $A$  between  $Z$  and  $X$ . For suppose not; then by Menger's theorem there is a separation  $(A_1, B_1)$  of  $A$  of order  $< h$  with  $Z \subseteq V(A_1)$  and  $X \subseteq V(B_1)$ . Then  $(A_1, B_1 \cup B)$  is a separation of  $G$  of order  $< h$ , and since  $(A, B) \in \mathcal{T}_0$  it follows from (2.8) that  $(A_1, B_1 \cup B) \in \mathcal{T}_0$ , contrary to the freedom of  $Z$  relative to  $\mathcal{T}_0$ . This proves our claim that there are  $h$  disjoint paths  $P_1, \dots, P_h$  of  $A$  between  $Z$  and  $X$ . Let the vertex of  $P_i$  in  $X$  be  $x_i$  ( $1 \leq i \leq h$ ); then by extending  $P_i$  by an edge joining  $x_i$  and  $y_i$ , for each  $i$ , we deduce that there is a linkage for  $G', \mathcal{T}', \{y_1, \dots, y_h\}$ . The result follows. ■

From (4.1) and (3.2) we deduce the following.

(4.2. *Let  $G, \mathcal{T}, Z$  be a nodule with  $|Z| = h$ , of order  $\geq 18(h+k)+3$ , where  $k \geq h$ , and let  $s, t \in Z$  be distinct. Then there is a residual nodule  $G', \mathcal{T}', Z'$  of order  $k$ , and a linkage  $\mathcal{P}$  for it from  $G, \mathcal{T}, Z$ , and a path  $Q$  of  $G$  from  $s$  to  $t$  disjoint from  $G'$ , such that for each  $P \in \mathcal{P}$ ,  $P \cap Q$  is either null or a path.*

*Proof.* By (3.2) there is a path  $C$  between  $s$  and  $t$  and a tangle  $\mathcal{T}_1$  in  $G \setminus V(C)$  of order  $h+k$ , conformal with  $\mathcal{T}$ . Let  $\mathcal{T}_0$  be the tangle in  $G$  induced by  $\mathcal{T}_1$ ; then  $\mathcal{T}_0 \subseteq \mathcal{T}$ . Since  $|Z| \leq \text{ord}(\mathcal{T}_0)$  and  $\mathcal{T}_0 \subseteq \mathcal{T}$  and  $G, \mathcal{T}, Z$  is a nodule, it follows that  $G, \mathcal{T}_0, Z$  is a nodule. Since  $\text{ord}(\mathcal{T}_1) \geq 2|Z|$ , we may apply (4.1). We deduce that there is a residual nodule  $G', \mathcal{T}', Z'$  of  $G, \mathcal{T}_0, Z$  and hence of  $G, \mathcal{T}, Z$ , of order  $\text{ord}(\mathcal{T}_1) - h = k$ , such that  $C \cap G'$  is null. Let  $\mathcal{P}$  be a linkage for it from  $G, \mathcal{T}, Z$  and choose a path  $Q$  of  $G$  from  $s$  to  $t$ , disjoint from  $G'$ , with  $Q \cup \bigcup \{P : P \in \mathcal{P}\}$  minimal. (Such

a choice is possible because  $Q = C$  is one such choice.) It follows that  $P \cap Q$  is either null or a path for each  $P \in \mathcal{P}$ , as required. ■

(4.3). Let  $G, \mathcal{T}, Z$  be a nodule of order  $\geq k20^n$ , where  $|Z| = n \geq 2$  and  $k \geq n/2$ . Then there is a residual nodule  $G', \mathcal{T}', Z'$  of order  $\geq k$  such that  $Z$  is a subset of the vertex set of one component of  $G \setminus V(G')$ .

*Proof.* Choose  $r$  with  $1 \leq r \leq n-1$ , minimum such that there is a residual nodule  $G', \mathcal{T}', Z'$  of order  $\geq k20^r$  and  $Z$  is a subset of the union of the vertex sets of at most  $r$  components of  $G \setminus V(G')$ . Such a choice of  $r$  is possible, because by (4.2) such a residual nodule exists when  $r = n-1$ . Suppose that  $r \geq 2$ , and let  $s, t \in Z$  belong to different components of  $G \setminus V(G')$ . Choose a linkage for  $G', \mathcal{T}', Z'$ , and let  $s', t' \in Z'$  be the last vertices of the paths in this linkage with first vertices  $s, t$ . Let  $h' = n, k' = k20^{r-1}$ ; then by applying (4.2) to  $G', \mathcal{T}', Z', s', t', h', k'$ , we obtain a contradiction to the minimality of  $r$ . Hence  $r = 1$ , and the result follows. ■

(4.4). Let  $2 \leq h \leq n \leq 2k$ , and let  $G, \mathcal{T}, Z$  be a nodule of order  $\geq k20^{nh}$ , with  $|Z| = h$ . Then there is a nodule  $G', \mathcal{T}', Y$  of order  $\geq k$ , with  $G'$  a subgraph of  $G \setminus Z$  and with  $|Y| = n$ , such that  $G', \mathcal{T}', Z'$  is a residual nodule of  $G, \mathcal{T}, Z$  for every  $Z' \subseteq Y$  with  $|Z'| = h$ .

*Proof.* By (2.4), since  $\text{ord}(\mathcal{T}) \geq n$  there is a set  $Z_0 \subseteq V(G)$  with  $Z \subseteq Z_0$  and  $|Z_0| = n$ , free relative to  $\mathcal{T}$ . By  $h$  applications of (4.3), we deduce that there are nodules  $G_r, \mathcal{T}_r, Z_r$  ( $0 \leq r \leq h$ ), where  $G_0 = G$  and  $\mathcal{T}_0 = \mathcal{T}$ , such that for  $1 \leq r \leq h$

- (i)  $G_r, \mathcal{T}_r, Z_r$  is a residual nodule of  $G_{r-1}, \mathcal{T}_{r-1}, Z_{r-1}$
- (ii)  $\mathcal{T}_r$  has order  $\geq k20^{n(h-r)}$ , and
- (iii)  $Z_{r-1}$  is a subset of the vertex set of one component  $C_r$  of  $G_{r-1} \setminus V(G_r)$ .

We show that setting  $G' = G_h, \mathcal{T}' = \mathcal{T}_h, Y = Z_h$  satisfies the theorem. For let  $Z' \subseteq Z_h$  with  $|Z'| = h$ , and let  $J$  be the graph  $G \setminus (V(G') - Z')$ . Certainly  $\mathcal{T}'$  is conformal with  $\mathcal{T}$ , so it suffices to show that there is a linkage for  $G', \mathcal{T}', Z'$  from  $G, \mathcal{T}, Z$ ; that is, there are  $h$  disjoint  $ZZ'$ -paths in  $J$ . To see this, let  $X \subseteq V(J)$  with  $|X| < h$ . For  $r = 1, \dots, h$ , by (i)  $G_r, \mathcal{T}_r, Z_r$  is a residual nodule of  $G_{r-1}, \mathcal{T}_{r-1}, Z_{r-1}$ ; let  $\mathcal{P}_r$  be a corresponding linkage. Then  $\mathcal{P}_r$  is a set of  $n$  disjoint  $Z_{r-1}Z_r$ -paths in  $G_{r-1}$ , each with just one vertex in  $V(G_r)$ , namely its last vertex. Since  $|Z_0| = |Z_1| = \dots = |Z_h| = n$ , there are  $n$  disjoint  $Z_0Z_h$ -paths  $P_1, \dots, P_n$  in  $G$  such that each  $P_i$  is of the form  $P_i^1 \cup P_i^2 \cup \dots \cup P_i^h$ , where  $P_i^r \in \mathcal{P}_r$  ( $r = 1, \dots, h$ ). Since  $Z \subseteq Z_0, Z' \subseteq Z_h$



and  $|Z| = |Z'| = h > |X|$ , there exist  $i, j \in \{1, \dots, h\}$  such that  $P_i$  is a  $ZZ_h$ -path,  $P_j$  is a  $Z_0Z'$ -path, and neither  $P_i$  nor  $P_j$  meets  $X$ . Since  $C_1, \dots, C_h$  are disjoint and  $|X| < h$  there exists  $r$  with  $1 \leq r \leq h$  such that  $X \cap V(C_r) = \emptyset$ . Let the last vertex of  $P_i$  be  $v$ . Then  $P_i \setminus v$ ,  $P_j$  and  $C_r$  are all subgraphs of  $J$ , they are all connected, and both  $P_i \setminus v$  and  $P_j$  meet  $V(C_r)$ ; and so  $(P_i \setminus v) \cup P_j \cup C_r$  is a connected subgraph of  $J$  meeting  $Z$  and  $Z'$  and not meeting  $X$ . Since this holds for all choices of  $X$ , it follows by Menger's theorem that there are  $h$  disjoint  $ZZ'$ -paths in  $J$ ; that is,  $G', \mathcal{T}', Z'$  is a residual nodule of  $G, \mathcal{T}, Z$ , as required. ■

## 5. PRODUCING A GRID

Now we use (4.2) to deduce properties of tangles not dominating a large grid minor. We begin with the following lemma.

(5.1). *Let  $\mathcal{T}$  be a tangle in  $G$  of order  $\geq g \geq 2$ , and for  $1 \leq i, j \leq g$  let  $W_{ij} \subseteq V(G)$  be such that*

(i) *the sets  $W_{ij}$  are mutually disjoint, and each is the vertex set of a connected subgraph of  $G$*

(ii) *for all  $i, j, i', j'$  with  $|i' - i| + |j' - j| = 1$  there is an edge of  $G$  joining  $W_{ij}$  and  $W_{i'j'}$ .*

(iii) *for  $1 \leq j \leq g$ ,  $\text{rk}_{\mathcal{T}}(W_{1j} \cup \dots \cup W_{gj}) \geq g$ .*

*Let  $H$  be the corresponding  $g \times g$  grid minor; then  $\mathcal{T}$  dominates  $H$ .*

*Proof.* Let  $(A, B) \in \mathcal{T}_H$  (we recall that  $\mathcal{T}_H$  was defined before (2.3)); we must show that  $(A, B) \in \mathcal{T}$ . For  $1 \leq j \leq g$  let  $W_j = W_{1j} \cup \dots \cup W_{gj}$ . Now  $E(B)$  includes a row of  $H$ , say the  $i$ th row. Thus for  $1 \leq j \leq g$ ,  $W_{ij} \cap V(B) \neq \emptyset$ , and so  $W_j \cap V(B) \neq \emptyset$ . Since  $|V(A \cap B)| < g$  and the sets  $W_1, \dots, W_g$  are mutually disjoint, there exists  $j$  such that  $W_j \cap V(A \cap B) = \emptyset$ . Since the restriction of  $G$  to  $W_j$  is connected and  $W_j \cap V(B) \neq \emptyset$ , it follows that  $W_j \subseteq V(B) - V(A)$ . Since  $\text{rk}_{\mathcal{T}}(W_j) \geq g$  and  $|V(A \cap B)| < g$ , we deduce that  $(B, A) \notin \mathcal{T}$ , and so  $(A, B) \in \mathcal{T}$ , as required. ■

(5.2). *If  $\mathcal{T}$  is a tangle in  $G$  dominating no  $g \times g$  grid minor, and  $\mathcal{T}'$  is a tangle in a subgraph  $G' \subseteq G$  which is conformal with  $\mathcal{T}$ , then  $\mathcal{T}'$  dominates no  $g \times g$  grid minor of  $G'$ .*

*Proof.* Let  $H$  be a  $g \times g$  grid minor of  $G'$  and suppose that  $\mathcal{T}'$  dominates  $H$ . Then  $H$  is a  $g \times g$  grid minor of  $G$ , and we claim that  $\mathcal{T}$

dominates  $H$ . For let  $(A, B)$  be a separation of  $G$  of order  $< g$  such that  $E(B)$  includes a row of  $H$ . Then  $(A \cap G', B \cap G')$  is a separation of  $G'$  of order  $< g$ , and  $E(B \cap G')$  includes a row of  $H$ . Since  $\mathcal{T}'$  dominates  $H$ , it follows that  $(A \cap G', B \cap G') \in \mathcal{T}'$ , and hence  $(A, B) \in \mathcal{T}$  since  $\mathcal{T}'$  is conformal with  $\mathcal{T}$ . Thus  $\mathcal{T}$  dominates  $H$ , a contradiction. ■

(5.3). Let  $g, k, n$  be integers with  $2k \geq n \geq 2g - 1 \geq 3$ . Let  $G, \mathcal{T}, Z$  be a nodule of order  $\geq k20^{2g}$ , such that  $\mathcal{T}$  dominates no  $g \times g$  grid minor and  $|Z| = n$ . Let  $Z = \{z_1, \dots, z_n\}$ . Then there is a residual nodule  $G', \mathcal{T}', Z'$  of order  $\geq k$  and a linkage  $\{P_1, \dots, P_n\}$ , where  $P_i$  has first vertex  $z_i$  ( $1 \leq i \leq n$ ), and a path  $Q$  of  $G \setminus V(G')$ , and there exist  $i, j$  with  $2 \leq i \leq 2g - 2$ ,  $1 \leq j \leq n$ , and  $j \neq i - 1$ ,  $i, i + 1$ , such that  $Q$  has first vertex in  $P_i$ , last vertex in  $P_j$ , and no other vertex in  $P_1 \cup \dots \cup P_n$ .

*Proof.* Let  $G_0 = G$ ,  $\mathcal{T}_0 = \mathcal{T}$ ,  $Z_0 = Z$ , and  $z_i^0 = z_i$  ( $1 \leq i \leq n$ ). By  $2g - 1$  applications of (4.2), we may define inductively for all  $r$  with  $1 \leq r \leq 2g - 1$ :

(i) a nodule  $G_r, \mathcal{T}_r, Z_r$  of order  $\geq k20^{2g-r}$  which is a residual nodule of  $G_{r-1}, \mathcal{T}_{r-1}, Z_{r-1}$  (and hence of  $G, \mathcal{T}, Z$ , by transitivity), and a linkage  $\{P'_1, \dots, P'_n\}$  for it from  $G_{r-1}, \mathcal{T}_{r-1}, Z_{r-1}$

(ii) an ordering  $\{z'_1, \dots, z'_n\}$  of  $Z_r$  such that  $P'_i$  has ends  $z_i^{r-1}, z'_i$  for  $1 \leq i \leq n$ , and

(iii) a path  $Q_r$  of  $G_{r-1} \setminus V(G_r)$  from  $z_g^{r-1}$  to  $z_1^{r-1}$ , such that for  $1 \leq i \leq n$ ,  $P'_i \cap Q_r$  is null or a path.

For  $1 \leq i \leq n$ , let  $P_i = P'_i \cup \dots \cup P_i^{2g-1}$ .

*Case 1.* For at least  $g$  values of  $r$  with  $1 \leq r \leq 2g - 1$ ,  $Q_r$  meets  $P'_g, P'_{g-1}, \dots, P'_1$  in order before meeting any other  $P'_i$ .

Let  $1 \leq r_1 < r_2 < \dots < r_g \leq 2g - 1$  be such that for  $r = r_1, \dots, r_g$ ,  $Q_r$  meets  $P'_g, P'_{g-1}, \dots, P'_1$  in order before meeting any other  $P'_i$ . Let  $r_0 = 0$ , and for  $1 \leq i, j \leq g$  let  $W_{ij}$  be the union of the following two sets:

(a)  $V(P_i) \cap (V(G_{r_{j-1}}) - V(G_{r_j}))$

(b) all vertices of  $Q_{r_j}$  which lie in  $Q_{r_j}$  strictly between the intersections with  $P_{i-1}$  and  $P_i$  (or  $\emptyset$ , if  $i = 1$ ).

We see that  $W_{ij}$  satisfy the hypothesis of (5.1), for if  $1 \leq j \leq g$  then since  $\mathcal{T}_r$  is conformal with  $\mathcal{T}$  by (i), it follows from (2.7) (writing  $r$  for  $r_{j-1}$ ) that

$$\text{rk}_{\mathcal{T}}(W_{1j} \cup \dots \cup W_{gj}) \geq \text{rk}_{\mathcal{T}}\{z'_1, \dots, z'_g\} \geq \text{rk}_{\mathcal{T}_r}\{z'_1, \dots, z'_g\} = g.$$

Thus, by (5.1),  $\mathcal{T}$  dominates a  $g \times g$  grid minor, a contradiction.

*Case 2.* For at least  $g$  values of  $r$  with  $1 \leq r \leq 2g-1$ ,  $Q_r$  meets  $P_g^r$ ,  $P_{g+1}^r, \dots, P_{2g-1}^r$  in order, before meeting any other  $P_i^r$ .

Again,  $\mathcal{T}$  dominates a  $g \times g$  grid minor; a contradiction.

*Case 3.* Neither Case 1 nor Case 2 applies.

Then there exists  $r$  with  $1 \leq r \leq 2g-1$  such that for some  $i$  with  $2 \leq i \leq 2g-2$ ,  $Q_r$  meets  $P_i^r$  and next meets  $P_j^r$  where  $j \neq i-1, i, i+1$ . Let  $Q$  be a minimal subpath of  $Q_r$  between  $P_i^r$  and  $P_j^r$ ; then the theorem is satisfied. ■

A leaf of a tree is a vertex of valency 1. Two disjoint subgraphs  $A, B$  of  $G$  are said to *touch* if there is a vertex of  $A$  adjacent in  $G$  to some vertex of  $B$ . Let  $G', \mathcal{T}', Z'$  be a residual nodule of  $G, \mathcal{T}, Z$ , where  $|Z| = m$ . A graph  $S$  with  $V(S) = Z$  is a *shadow* of  $G', \mathcal{T}', Z'$  if there are  $m$  disjoint connected subgraphs  $C_1, \dots, C_m$  of  $G \setminus (V(G') - Z')$  such that

- (i) for  $1 \leq i \leq m$ ,  $|V(C_i) \cap Z| = 1 = |V(C_i) \cap Z'|$ ; let  $V(C_i) \cap Z = \{z_i\}$
- (ii) for  $1 \leq i < j \leq m$ , if  $z_i$  and  $z_j$  are adjacent in  $S$  then  $C_i \setminus Z'$  and  $C_j \setminus Z'$  touch.

(5.4). Let  $g, m, k$  be integers with  $g \geq 2$ ,  $m \geq 3$ , and  $k \geq n/2$ , where  $n = (2g+1)(2m-5) + 2$ . Let  $G, \mathcal{T}, Z$  be a nodule of order  $\geq k20^{2g(m-2)+n}$ , such that  $\mathcal{T}$  dominates no  $g \times g$  grid minor and  $|Z| = n$ . Then there is a residual nodule  $G', \mathcal{T}', Z'$  of order  $\geq k$  with a shadow which is a tree with  $\geq m$  leaves.

*Proof.* First we show the following.

(1) There is a residual nodule of  $G, \mathcal{T}, Z$  of order  $\geq k20^{2g(m-2)}$  with a shadow which is a tree with  $\geq 2$  leaves.

*Subproof.* By (4.3) there is a residual nodule  $G', \mathcal{T}', Z'$  of order  $\geq k20^{2g(m-2)}$  such that  $Z$  is a subset of one component  $C$  of  $G \setminus V(G')$ . Let  $\{P_1, \dots, P_n\}$  be a linkage for  $G', \mathcal{T}', Z'$  from  $G, \mathcal{T}, Z$ , where  $P_i$  has first vertex  $z_i$  ( $1 \leq i \leq n$ ). Let  $J = C \cup P_1 \cup \dots \cup P_n$ , and for  $1 \leq i \leq n$  let  $C_i$  be a connected subgraph of  $J$  with  $P_i \subseteq C_i$ , such that  $C_1, \dots, C_n$  are mutually vertex-disjoint and their union is maximal. Since  $J$  is connected, it follows that  $V(C_1 \cup \dots \cup C_n) = V(J)$ . Let  $S_0$  be the graph with  $V(S_0) = Z$  in which  $z_i$  and  $z_j$  are adjacent if  $i \neq j$  and  $C_i \setminus Z', C_j \setminus Z'$  touch. Since  $C$  is connected it follows that  $S_0$  is connected and, hence, has a spanning tree  $S$ , which has  $\geq 2$  leaves since  $n \geq 2$ . But  $S$  is a shadow, and so (1) holds.

We shall prove the following statement (2) for all  $r$  with  $2 \leq r \leq m$ , by induction on  $r$ .

(2) There is a residual nodule of  $G, \mathcal{T}$ , of  $Z$  of order  $\geq k20^{2g(m-r)}$  with a shadow which is a tree with  $\geq r$  leaves.

*Subproof.* (2) holds when  $r = 2$ ; suppose then that (2) holds for some  $r$  with  $2 \leq r < m$ ; we shall prove that it holds for  $r + 1$ . Let  $G'$ ,  $\mathcal{T}'$ ,  $Z'$  be the residual nodule in (2), and let  $S$  be the shadow and  $C_1, \dots, C_n$  the corresponding disjoint connected subgraphs of  $G \setminus (V(G') - Z')$ . Let  $C_i \cap Z = \{z_i\}$ ,  $C_i \cap Z' = \{z'_i\}$  ( $1 \leq i \leq n$ ). Now we may assume that  $S$  has exactly  $r$  leaves, for otherwise the nodule  $G'$ ,  $\mathcal{T}'$ ,  $Z'$  satisfies (2) with  $r$  replaced by  $r + 1$ ; and so  $S$  is a subdivision of a tree with  $\leq 2r - 3$  edges. Since

$$|E(S)| = n - 1 > (2g + 1)(2m - 5),$$

and  $2r - 3 \leq 2m - 5$ , it follows that one of these edges is subdivided at least  $2g + 1$  times. Thus we may choose the numbering of  $C_1, \dots, C_n$  (and hence of  $z_1, \dots, z_n$ ), such that  $z_n, z_1, z_2, \dots, z_{2g}$  all have valency 2 in  $S$  and such that for  $0 \leq i \leq 2g - 1$ ,  $z_i$  is adjacent in  $S$  to  $z_{i+1}$  (where  $z_0$  means  $z_n$ ). By (5.2),  $\mathcal{T}'$  dominates no  $g \times g$  grid minor of  $G'$ , since  $\mathcal{T}'$  is conformal with  $\mathcal{T}$ . By (5.3) applied to  $G'$ ,  $\mathcal{T}'$ ,  $Z'$ , there is a residual nodule  $G''$ ,  $\mathcal{T}''$ ,  $Z''$  of  $G'$ ,  $\mathcal{T}'$ ,  $Z'$  of order  $\geq k20^{2g(m-(r+1))}$  and a linkage  $\{P'_1, \dots, P'_n\}$  for it from  $G'$ ,  $\mathcal{T}'$ ,  $Z'$ , where  $P'_i$  has first vertex  $z'_i$  ( $1 \leq i \leq n$ ), and a path  $Q$  of  $G' \setminus V(G'')$ , such that  $Q$  has first vertex in some  $P'_i$  with  $2 \leq i \leq 2g - 2$ , last vertex in some  $P'_j$  where  $j \neq i - 1, i, i + 1$ , and it has no other vertex in  $P'_1 \cup \dots \cup P'_n$ .

Let  $S'$  be obtained from  $S$  by adding the edge between  $z_i$  and  $z_j$ . Then  $S'$  is a shadow of  $G''$ ,  $\mathcal{T}''$ ,  $Z''$  (regarded as a residual nodule of  $G$ ,  $\mathcal{T}$ ,  $Z$ ). But  $S'$  has a spanning tree with  $\geq r + 1$  leaves (delete either the edge between  $z_{i-2}$  and  $z_{i-1}$ , or that between  $z_{i+1}$  and  $z_{i+2}$ , depending which gives a tree). Thus (2) holds for  $r + 1$ . In particular, (2) holds with  $r = m$ , as required. ■

(5.5). Let  $g, m, k$  be integers with  $g \geq 2$ ,  $m \geq 3$ , and  $k \geq n/2$ , where  $n = (2g + 1)(2m - 5) + 2$ . Let  $t = 2g(m - 2) + n(m + 1)$ , and let  $G, \mathcal{T}, Z$  be a nodule of order  $\geq k20^t$ , such that  $\mathcal{T}$  dominates no  $g \times g$  grid minor and  $|Z| = m$ . Then there is a residual nodule  $G', \mathcal{T}', Z'$  of order  $\geq k$  with a linkage  $\mathcal{P}$ , and a connected subgraph  $C$  of  $G \setminus V(G')$  which touches but does not intersect  $P \setminus Z'$  for each  $P \in \mathcal{P}$ .

*Proof.* By (4.4), there is a nodule  $G'', \mathcal{T}'', Y$  of order  $\geq k20^{2g(m-2)+n}$ , such that  $G''$  is a subgraph of  $G \setminus Z$ , and  $|Y| = n$ , and  $G'', \mathcal{T}'', Z''$  is a residual nodule of  $G, \mathcal{T}, Z$  for every  $Z'' \subseteq Y$  with  $|Z''| = m$ . By (5.2),  $\mathcal{T}''$  dominates no  $g \times g$  grid minor of  $G''$ . By (5.4) applied to  $G'', \mathcal{T}'', Y$  we deduce that it has a residual nodule  $G', \mathcal{T}', Y'$  of order  $\geq k$  with a linkage  $\mathcal{P}'$ , and there is a connected subgraph  $C$  of  $G'' \setminus V(G')$  which touches but does not meet  $P \setminus Y'$  for at least  $m$  members  $P$  of  $\mathcal{P}'$ . Choose  $Z' \subseteq Y'$  with  $|Z'| = m$  such that  $C$  touches but does not meet  $P \setminus Y'$  for each  $P \in \mathcal{P}'$  with

last vertex in  $Z'$ . Let  $Z''$  be the set of first vertices of the members of  $\mathcal{P}'$  with last vertex in  $Z'$ . Since  $Z'' \subseteq Y$  and  $|Z''| = m$ , it follows that  $G'', \mathcal{T}'', Z''$  is a residual nodule of  $G, \mathcal{T}, Z$ ; let  $\mathcal{P}''$  be a corresponding linkage. Each member of  $\mathcal{P}''$  has its last vertex in  $Z''$  and no other vertex in  $V(G'')$ , while each member of  $\mathcal{P}'$  is a path of  $G''$ . Let  $\mathcal{P}$  be the set of paths formed by taking, for each  $P \in \mathcal{P}''$ , the union of  $P$  and the path of  $\mathcal{P}'$  which meets  $P$ ; then  $\mathcal{P}$  is a linkage for  $G', \mathcal{T}', Z'$  satisfying the theorem. ■

Let  $m, n \geq 1$  be integers. An  $m \times n$  array in a graph  $G$  is a family of subsets  $W_{ij}$  ( $0 \leq i \leq m, 1 \leq j \leq n$ ) of  $V(G)$  such that

(i) the sets  $W_{ij}$  are mutually disjoint, and each is the vertex set of a connected subgraph of  $G$

(ii) for  $1 \leq i \leq m$  and  $1 \leq j \leq n$  there is an edge of  $G$  between  $W_{0j}$  and  $W_{ij}$ ; and for  $1 \leq i \leq m$  and  $1 \leq j \leq n-1$  there is an edge of  $G$  between  $W_{ij}$  and  $W_{i,j+1}$ .

If  $\mathcal{T}$  is a tangle in  $G$ , we say  $\mathcal{T}$  dominates this array if for  $0 \leq i \leq m$ ,

$$\text{rk}_{\mathcal{T}}(W_{i1} \cup \dots \cup W_{in}) \geq \min(m, n).$$

(5.6). Let  $g \geq 2, m \geq 3, n \geq 1$  be integers, and let

$$t = 2g(m-2) + ((2g+1)(2m-5)+2)(m+1).$$

If  $\mathcal{T}$  is a tangle of order  $\geq 20^m$  in a graph  $G$  and  $\mathcal{T}$  dominates no  $g \times g$  grid minor, then  $\mathcal{T}$  dominates an  $m \times n$  array in  $G$ .

*Proof.* Let  $\mathcal{T}$  be a tangle of order  $\geq 20^m$ . By (2.4) we can choose  $Z \subseteq V(G)$  with  $|Z| = m$ , free relative to  $\mathcal{T}$ . Thus  $G, \mathcal{T}, Z$  is a nodule, and  $\mathcal{T}$  dominates no  $g \times g$  grid minor. By  $n-1$  applications of (5.2) and (5.5) we deduce that there are nodules  $G_r, \mathcal{T}_r, Z_r$  ( $0 \leq r \leq n-1$ ), where  $G_0 = G, \mathcal{T}_0 = \mathcal{T}, Z_0 = Z$ , such that for  $1 \leq r \leq n-1$ ,

(i)  $G_r, \mathcal{T}_r, Z_r$  is a residual nodule of  $G_{r-1}, \mathcal{T}_{r-1}, Z_{r-1}$

(ii)  $\mathcal{T}_r$  has order  $\geq 20^{(n-r)t}$  and dominates no  $g \times g$  grid minor of  $G_r$

(iii) there is a linkage  $\mathcal{P}_r$  for  $G_r, \mathcal{T}_r, Z_r$  from  $G_{r-1}, \mathcal{T}_{r-1}, Z_{r-1}$  and a connected subgraph  $C_r$  of  $G_{r-1} \setminus V(G_r)$ , such that for each  $P \in \mathcal{P}_r$ ,  $C_r \cap P$  is null and some vertex of  $C_r$  is adjacent to some vertex of  $P \setminus Z_r$ .

Now  $Z_{n-1}$  is free relative to  $\mathcal{T}_{n-1}$ , and since  $|Z_{n-1}| = m < \text{ord}(\mathcal{T}_{n-1})$  there is a component  $C_n$  of  $G_{n-1} \setminus Z_{n-1}$  such that each vertex in  $Z_{n-1}$  has a neighbour in  $V(C_n)$ , by (2.12). If  $n=1$ , let  $P_1, \dots, P_m$  be the  $m$  one-vertex

paths with a vertex in  $Z_0$ . If  $n \geq 2$ , let  $P_1, \dots, P_m$  be the components of the union of all the members of  $\mathcal{P}_1 \cup \dots \cup \mathcal{P}_{n-1}$ . For  $1 \leq i \leq m$  and  $1 \leq j \leq n-1$  let

$$W_{ij} = V(P_i) \cap (V(G_{j-1}) - V(G_j));$$

for  $1 \leq i \leq m$  let  $W_{in} = V(P_i) \cap V(G_{n-1})$ ; and for  $1 \leq j \leq n$  let  $W_{0j} = V(C_j)$ . Then the sets  $W_{ij}$  form an  $m \times n$  array, and we claim that  $\mathcal{T}$  dominates it. For let  $0 \leq i \leq m$  and suppose that there exists  $(A, B) \in \mathcal{T}$  of order  $< \min(m, n)$  with

$$W_{i1} \cup \dots \cup W_{in} \subseteq V(A).$$

Since  $|V(A \cap B)| < n$ , there exists  $j$  with  $1 \leq j \leq n$  such that  $X \cap V(A \cap B) = \emptyset$ , where

$$X = W_{0j} \cup W_{1j} \cup \dots \cup W_{mj}.$$

Since the restriction of  $G$  to  $X$  is connected and since  $X \cap V(A) \neq \emptyset$  because  $\emptyset \neq W_{ij} \subseteq X$ , it follows that  $X \subseteq V(A)$ . But  $Z_{j-1} \subseteq X$ , and so  $Z_{j-1} \subseteq V(A)$ , and hence  $\text{rk}_{\mathcal{T}}(Z_{j-1}) \leq |V(A \cap B)| < m$ , contrary to (2.7). Hence there is no such  $(A, B)$ , as required. ■

Now we can prove our main result (2.3), which we restate.

(5.7). *Let  $g \geq 2$ , and let  $\mathcal{T}$  be a tangle in  $G$  of order  $\geq 20g^{4(2g-1)}$ . Then  $\mathcal{T}$  dominates a  $g \times g$  grid minor.*

*Proof.* Any tangle of order  $\geq 3$  dominates a  $2 \times 2$  grid, as can be shown for instance as follows. By [8, Theorem (8.3)] we may assume that  $G$  is two-connected. But  $|V(G)| \geq 4$  since  $\mathcal{T}$  has order  $\geq 3$ , and so there is a circuit of length  $\geq 4$ . Hence  $G$  has a  $2 \times 2$  grid minor, and  $\mathcal{T}$  dominates it since  $G$  is two-connected.

We may therefore assume that  $g \geq 3$ . Let  $t = \lfloor \frac{1}{2}g \rfloor$ ; then, since

$$g^4(2g-1) \geq 2g(g-2)gt + ((2g+1)(2g-5)+2)(g+1)gt,$$

we may assume from (5.6) (taking  $m = g \geq 3$  and  $n = gt$ ) that  $\mathcal{T}$  dominates a  $g \times gt$  array  $W_{ij}$  ( $0 \leq i \leq g$ ,  $1 \leq j \leq gt$ ). For  $1 \leq i \leq g$  and  $1 \leq j \leq g$ , let

$$U_{ij} = \bigcup (W_{ik} : (j-1)t < k \leq jt)$$

if  $i$  is odd, and let

$$U_{ij} = \bigcup (W_{ik} : (j-1)t < k \leq jt) \cup W_{0, (j-1)t+i/2}$$

if  $i$  is even. Then for all  $i, j, i', j'$  with  $|i - i'| + |j - j'| = 1$  there is an edge between  $U_{ij}$  and  $U_{i'j'}$ , and each  $U_{ij}$  is the vertex set of a connected subgraph of  $G$ . Moreover, for  $1 \leq i \leq g$ ,

$$U_{i1} \cup \dots \cup U_{ig} \supseteq W_{i1} \cup \dots \cup W_{i,gt}$$

and so  $\text{rk}_{\mathcal{T}}(U_{i1} \cup \dots \cup U_{ig}) \geq \min(g, gt) = g$ , since  $\mathcal{T}$  dominates the array. From (5.1) (with rows and columns exchanged)  $\mathcal{T}$  dominates a  $g \times g$  grid minor, as required. ■

In some circumstances, the maximum size of a complete minor of  $G$  is much smaller than the maximum size of a grid minor, and if so we can obtain an improved bound on tree-width, as follows.

(5.8). *If  $G$  has no  $g \times g$  grid minor and no  $K_h$  minor, then the tree-width of  $G$  is at most  $20^{5gh^3}$ .*

*Proof.* We assume that  $g \geq 2$  and  $h \geq 3$ , for otherwise the result is trivial. Suppose that  $W_{ij}$  ( $0 \leq i \leq h$ ,  $1 \leq j \leq h$ ) is an  $h \times h$  array in  $G$ . For  $1 \leq i \leq h$ , let

$$V_i = W_{0i} \cup (W_{i1} \cup \dots \cup W_{ih}).$$

Then  $V_1, \dots, V_h$  are disjoint, each is the vertex set of a connected subgraph of  $G$ , and for  $1 \leq i < j \leq h$  there is an edge between  $V_i$  and  $V_j$ . Hence  $G$  has a  $K_h$  minor, a contradiction. It follows that  $G$  has no  $h \times h$  array, and so from (5.6), no tangle of order  $\geq 20^k$  where

$$k = 2gh(h-2) + ((2g+1)(2h-5) + 2)h(h+1) \leq 5gh^3 - 1.$$

The result follows from (2.1). ■

## 6. GRIDS IN PLANAR GRAPHS

So far, we have been using tangles to obtain a numerically improved version of the theorem of [6], concerning the tree-width of general graphs with no  $g \times g$  grid minor. One can also use tangles to improve the theorem of [5], concerning the tree-width of planar graphs with no  $g \times g$  grid minor. Since the " $r \times r$  cylinder" of [5] has the  $r \times r$  grid as a subgraph, it follows from [5, Theorem (5.2)] that

(6.1). *Let  $g \geq 1$  be an integer. Every planar graph with no  $g \times g$  grid minor has tree-width  $< \frac{3}{2}(g^2 + 2g) - 2$ .*

We shall prove the following improvement.

(6.2). *Let  $g \geq 1$  be an integer. Every planar graph with no  $g \times g$  grid minor has tree-width  $\leq 6g - 5$ .*

(6.2) follows from (2.1) and the following.

(6.3). *Let  $g \geq 1$  be an integer. Every planar graph with a tangle of order  $4g - 3$  has a  $g \times g$  grid minor.*

*Proof.* We may assume that  $g \geq 2$  and by (2.11) it suffices to prove the result for connected graphs. Thus, let  $G$  be a connected planar graph, drawn in a sphere  $\Sigma$ , and let  $\mathcal{T}$  be a tangle in  $G$  of order  $4g - 3$ . Choose closed discs  $\Delta_1, \Delta_2 \subseteq \Sigma$  with  $\Delta_1 \cup \Delta_2 = \Sigma$  and  $\Delta_1 \cap \Delta_2 = bd(\Delta_1) = bd(\Delta_2)$ , such that

- (i)  $\Delta_1 \cap \Delta_2$  meets the drawing only in vertices,
- (ii)  $(G \cap \Delta_1, G \cap \Delta_2) \in \mathcal{T}$ , where  $G \cap \Delta_i$  denotes the subgraph of  $G$  drawn in  $\Delta_i$
- (iii) subject to (i) and (ii),  $|V(G \cap \Delta_1)| + |E(G \cap \Delta_1)| - |V(G \cap \Delta_2)|$  is maximum.

(Certainly this is possible, for if we choose  $\Delta_1, \Delta_2$  with  $G \cap \Delta_1$  null then (i) and (ii) are satisfied.) We refer to (iii) as the *optimality* of  $\Delta_1$ . A *line* in  $\Sigma$  is a subset homeomorphic to  $[0, 1]$  and its *ends* are defined in the obvious way.

(1) *Let  $a, b \in \Delta_1 \cap \Delta_2$  be distinct, and let  $F_1, F_2$  be the two lines in  $\Delta_1 \cap \Delta_2$  with ends  $a, b$ . Let  $F \subseteq \Delta_2$  be a line with ends  $a, b$ , meeting the drawing only in vertices, and with  $F \cap \Delta_1 = \{a, b\}$ . Then*

$$|F \cap V(G)| \geq \min(|F_1 \cap V(G)|, |F_2 \cap V(G)|).$$

*Subproof.* Let  $D_i$  be the closed disc in  $\Delta_2$  bounded by  $F \cup F_i$  ( $i = 1, 2$ ). Then  $D_1 \cup \Delta_1, D_2 \cup \Delta_1$  are both closed discs. From the second axiom,  $\mathcal{T}$  does not contain both

$$(G \cap D_1, G \cap (D_2 \cup \Delta_1)), \quad (G \cap D_2, G \cap (D_1 \cup \Delta_1)),$$

because  $(G \cap \Delta_1) \cup (G \cap D_1) \cup (G \cap D_2) = G$ . Consequently we may assume that  $(G \cap D_1, G \cap (D_2 \cup \Delta_1)) \notin \mathcal{T}$ . If  $(G \cap (D_2 \cup \Delta_1), G \cap D_1) \in \mathcal{T}$ , then from the optimality of  $\Delta_1$  we deduce that  $V(G \cap \Delta_2) = V(G \cap D_1)$ , and in



particular  $F_2 \cap V(G) \subseteq F \cap V(G)$  and the result holds. If  $(G \cap (D_2 \cup \Delta_1), G \cap \Delta_1) \notin \mathcal{T}$ , then from the first axiom it has order  $\geq 4g - 3$  and, hence, greater than the order of  $(G \cap \Delta_1, G \cap \Delta_2)$ ; and so

$$\begin{aligned} 0 &< |V(G) \cap ((D_2 \cup \Delta_1) \cap \Delta_1)| - |V(G) \cap (\Delta_1 \cap \Delta_2)| \\ &= |V(G) \cap (F_1 \cup F)| - |V(G) \cap (F_1 \cup F_2)| \\ &= |V(G) \cap F| - |V(G) \cap F_2|, \end{aligned}$$

as required.

$$(2) \quad |\Delta_1 \cap \Delta_2 \cap V(G)| = 4g - 4.$$

*Subproof.* Certainly  $|\Delta_1 \cap \Delta_2 \cap V(G)| \leq 4g - 4$ , since  $(G \cap \Delta_1, G \cap \Delta_2) \in \mathcal{T}$ . Suppose that  $|\Delta_1 \cap \Delta_2 \cap V(G)| \leq 4g - 5$ . Let  $r$  be a region of  $G$  with  $\Delta_1 \cap \Delta_2 \cap r \neq \emptyset$  (regions are open sets). If  $r$  is incident with some vertex  $v$  of  $G$  not in  $\Delta_1$ , then by (2.8) we can enlarge  $\Delta_1$  within  $r \cup \{v\}$  so that its boundary passes through  $v$  (and reduce  $\Delta_2$  correspondingly), contrary to the optimality of  $\Delta_1$ . Thus there is no such  $v$ . By a similar argument and (2.8), there is no edge of  $G \cap \Delta_2$  incident with  $r$  with both ends in  $\Delta_1 \cap \Delta_2$ . It follows that  $\Delta_2 \subseteq r \cup (\Delta_1 \cap \Delta_2)$ , and hence  $V(G \cap \Delta_1) = V(G)$ , contrary to the third axiom. This proves (2).

Let the vertices of  $G$  in  $\Delta_1 \cap \Delta_2$  be

$$a_1, a_2, \dots, a_g = b_1, b_2, \dots, b_g = c_1, c_2, \dots, c_g = d_1, d_2, \dots, d_g = a_1$$

in order. By (1) and (for example) [7, Theorem (3.6)], there are vertex-disjoint paths  $P_1, \dots, P_g$  of  $G \cap \Delta_2$  such that  $P_i$  has ends  $a_i, c_{g+1-i}$  for  $1 \leq i \leq g$ . Similarly, there are vertex-disjoint paths  $Q_1, \dots, Q_g$  of  $G \cap \Delta_2$  such that  $Q_i$  has ends  $b_i, d_{g+1-i}$  for  $1 \leq i \leq g$ . By a straightforward modification of the proof of [5, Theorem (4.1)], we may choose  $P_1, \dots, P_g$  and  $Q_1, \dots, Q_g$  so that  $P_i \cap Q_j$  is a path for all  $i, j$ . But then  $G$  has a  $g \times g$  grid minor, as required. ■

As a corollary, we have

(6.4). *If  $G$  is planar, with  $n$  vertices, then  $G$  has no tangle of order  $\geq 4\lfloor n^{1/2} \rfloor + 1$ .*

*Proof.* Let  $g = 1 + \lfloor n^{1/2} \rfloor$ . Since  $g^2 > n$ , it follows that  $G$  has no  $g \times g$  grid minor, and so by (6.3),  $G$  has no tangle of order  $4g - 3$ . The result follows. ■

Consequently:

(6.5). *If  $G$  is planar, with  $n$  vertices, there is a separation  $(A, B)$  of  $G$  of order  $\leq 4n^{1/2}$  with  $|V(A) - V(B)|, |V(B) - V(A)| \leq \frac{2}{3}|V(G)|$ .*

*Proof.* Let  $\mathcal{T}$  be the set of all separations  $(A, B)$  of  $G$  of order  $\leq 4n^{1/2}$  with  $|V(B) - V(A)| > \frac{2}{3}|V(G)|$ . By (6.4),  $\mathcal{T}$  is not a tangle, and yet the second and third axioms are satisfied (for if  $(A, B) \in \mathcal{T}$  then  $|V(A)| < \frac{1}{3}|V(G)|$ ). Hence the first is not satisfied, and the result follows. ■

(6.5) is a form of a theorem of Lipton and Tarjan [3], who in fact proved the same statement with  $4n^{1/2}$  replaced by  $2(2n)^{1/2}$ , using a completely different argument. The method given above was derived from [1] where the same method is used to show that (6.5) holds with  $4n^{1/2}$  replaced by  $3(2n)^{1/2}/2$ .

## REFERENCES

1. N. ALON, P. D. SEYMOUR, AND R. THOMAS, Planar separators, *SIAM J. Discrete Math.* **7** (1994), 184–193.
2. P. ERDŐS, Graph theory and probability, *Canad. J. Math.* **11** (1959), 34–38.
3. R. J. LIPTON AND R. E. TARJAN, A separator theorem for planar graphs, *SIAM J. Appl. Math.* **36** (1979), 177–189.
4. R. RADO, A theorem on independence relations, *Quart. J. Math. (Oxford)* **13** (1942), 83–89.
5. N. ROBERTSON AND P. D. SEYMOUR, Graph minors. III. Planar tree-width, *J. Combin. Theory Ser. B* **36** (1984), 49–64.
6. N. ROBERTSON AND P. D. SEYMOUR, Graph minors. V. Excluding a planar graph, *J. Combin. Theory Ser. B* **41** (1986), 92–114.
7. N. ROBERTSON AND P. D. SEYMOUR, Graph minors. VI. Disjoint paths across a disc, *J. Combin. Theory Ser. B* **41** (1986), 115–138.
8. N. ROBERTSON AND P. D. SEYMOUR, Graph minors. X. Obstructions to tree-decomposition, *J. Combin. Theory Ser. B* **52** (1991), 153–190.
9. H. WHITNEY, A theorem on graphs, *Ann. of Math.* **32** (1931), 378–390.