

# Infinite-state games with finitary conditions

CSL, September 5th, 2013

Krishnendu Chatterjee    Nathanaël Fijalkow

Institute of Informatics, Warsaw University – Poland

LIAFA, Université Paris 7 Denis Diderot – France

# In a nutshell

This paper is about  $\omega B$ -games, and the subclass of finitary games.

They are:

This paper is about  $\omega B$ -games, and the subclass of finitary games.

They are:

- two-player turn-based games played over **infinite** graphs,

This paper is about  $\omega B$ -games, and the subclass of finitary games.

They are:

- two-player turn-based games played over **infinite** graphs,
- the winning conditions involve counters,

This paper is about  $\omega B$ -games, and the subclass of finitary games.

They are:

- two-player turn-based games played over **infinite** graphs,
- the winning conditions involve counters,
- the first issue is to prove the existence of finite-memory strategies,

This paper is about  $\omega B$ -games, and the subclass of finitary games.

They are:

- two-player turn-based games played over **infinite** graphs,
- the winning conditions involve counters,
- the first issue is to prove the existence of finite-memory strategies,
- the second issue is to construct algorithms to decide the winner.

# Motivation: expressing boundedness properties




$\text{MSO} + \mathbb{U}$



cost MSO

A lot is known, and even more is not known about those two logics!

# Definition of $\omega B$ -games




controlled by Eve



controlled by Adam




### Definition of $\omega B$ -games




controlled by Eve

controlled by Adam

# Definition of $\omega B$ -games



# Definition of $\omega B$ -games




controlled by Eve



controlled by Adam

# Definition of $\omega B$ -games




controlled by Eve



controlled by Adam


# Definition of $\omega B$ -games



controlled by Eve




controlled by Adam




$\omega B$  winning condition:

parity  
and  
all counters  
are bounded



parity condition:  
the minimal priority  
seen infinitely often  
is even

# Definition of $\omega B$ -games



$$c_1 = 0$$

$$c_2 = 0$$


$\epsilon$  : nothing

$i$  : increment

$r$  : reset



# Definition of $\omega B$ -games



$$c_1 = 0$$

$$c_2 = 0$$


$\epsilon$  : nothing

$i$  : increment

$r$  : reset

# Definition of $\omega B$ -games

3



$$c_1 = 0$$

$$c_2 = 1$$


$\epsilon$  : nothing

$i$  : increment

$r$  : reset

# Definition of $\omega B$ -games

3



$$c_1 = 0$$


$$c_2 = 1$$

$\varepsilon$  : nothing

$i$  : increment

$r$  : reset

# Definition of $\omega B$ -games



$$c_1 = 1$$


$$c_2 = 0$$

$\epsilon$  : nothing

$i$  : increment

$r$  : reset

# Definition of $\omega B$ -games




$$c_1 = 1$$

$$c_2 = 0$$

$\epsilon$  : nothing

$i$  : increment

$r$  : reset



$\omega B$  winning condition:

parity  
and  
all counters  
are bounded

# Strategy (for Eve)

General form

$$\sigma : V^+ \rightarrow V$$

# Strategy (for Eve)

General form

$$\sigma : V^+ \rightarrow V$$

Positional or memoryless

$$\sigma : V \rightarrow V$$



# Strategy (for Eve)

General form

$$\sigma : V^+ \rightarrow V$$

Positional or memoryless

$$\sigma : V \rightarrow V$$

Theorem (Müller and Schupp)

*In parity games, both players have memoryless winning strategies.*

# Strategy (for Eve)

General form

$$\sigma : V^+ \rightarrow V$$

Positional or memoryless

$$\sigma : V \rightarrow V$$

Theorem (Müller and Schupp)

*In parity games, both players have memoryless winning strategies.*

What about  $\omega B$ -games?

# Strategy (for Eve)

General form

$$\sigma : V^+ \rightarrow V$$

Positional or memoryless

$$\sigma : V \rightarrow V$$

Theorem (Müller and Schupp)

*In parity games, both players have memoryless winning strategies.*

What about  $\omega B$ -games?

Finite-memory

$$\begin{cases} \sigma : V \times M \rightarrow V \\ \mu : M \times E \rightarrow M \end{cases}$$

# Quantification

Eve wins means:



$\exists \sigma$  (strategy for Eve),  
 $\forall \pi$  (paths),  
 $\exists N \in \mathbb{N}$ ,



$\exists \sigma$  (strategy for Eve),  
 $\exists N \in \mathbb{N}$ ,  
 $\forall \pi$  (paths),

$\pi$  satisfies parity and each counter is bounded by  $N$ .

# Quantification

Eve wins means:



$\exists \sigma$  (strategy for Eve),  
 $\forall \pi$  (paths),  
 $\exists N \in \mathbb{N}$ ,

$\pi$  satisfies parity and each counter is bounded by  $N$ .

non-uniform  
(MSO +  $\mathbb{U}$ )



$\exists \sigma$  (strategy for Eve),  
 $\exists N \in \mathbb{N}$ ,  
 $\forall \pi$  (paths),

uniform  
(cost MSO)

## Thomas Colcombet's habilitation:

le fait 2.22 et en deduire que la domination entre formules de la logique monadique de cost est décidable sur les arbres infinis. Ainsi, la conjecture 9.2 implique la conjecture 9.1.

En fait, il est possible de pointer avec encore plus de précision où se trouve la difficulté. Si l'on cherche à démontrer la conjecture 9.2, tout comme dans le cas des arbres finis, le point crucial est l'existence de stratégies gagnantes à mémoire finie. Il suffirait d'établir la conjecture suivante.

**Conjecture 9.3.** *Les objectifs  $\text{hB} \wedge \text{parité}$  et  $\neg \text{B} \wedge \text{parité}$  sont à  $\approx$ -mémoire finie, sur toutes les arènes/sur les arènes chronologiques/sur les arènes «arborescentes».*

## Existence of finite-memory strategies in (some) $\omega B$ -games

$\implies$  Decidability of cost MSO over infinite trees

$\implies$  Decidability of the index of the non-deterministic Mostowski's hierarchy (open for 40 years)!

Finitary conditions were introduced by Alur and Henzinger in 94, and are a subclass of  $\omega B$  conditions where the counters and parity are not independent.

Theorem (not in this talk)

*Over general graphs, Eve has finite-memory winning strategies in finitary games.*

Theorem (not in this talk)

*Solving pushdown finitary games with stack boundedness condition is EXPTIME-complete.*

## Theorem

*Over pushdown graphs, the uniform and non-uniform quantifications are **almost** equivalent.*


## Corollary

*Solving pushdown  $\omega B$ -games is decidable.*



- 1 Equivalence for pushdown  $\omega B$ -games
  - The case of finite graphs
  - The case of pushdown graphs

# A first example



Eve should maintain a low stack.

## Theorem

*For all pushdown games, the following are equivalent:*

- $\exists \sigma$  (strategy for Eve),  $\forall \pi$  (paths),  $\exists N \in \mathbb{N}$ ,  
 $\pi$  satisfies parity and each counter is bounded by  $N$ .
- $\exists \sigma$  (strategy for Eve),  $\exists N \in \mathbb{N}$ ,  $\forall \pi$  (paths),  
 $\pi$  satisfies parity and **eventually** each counter is bounded by  $N$ .


## Theorem

*For all pushdown games, the following are equivalent:*

- $\exists \sigma$  (strategy for Eve),  $\forall \pi$  (paths),  $\exists N \in \mathbb{N}$ ,  
 $\pi$  satisfies parity and each counter is bounded by  $N$ .
- $\exists \sigma$  (strategy for Eve),  $\exists N \in \mathbb{N}$ ,  $\forall \pi$  (paths),  
 $\pi$  satisfies parity and **eventually** each counter is bounded by  $N$ .



# Counter-example for the general case



Eve wins but she does not know the bound!

- 1 Equivalence for pushdown  $\omega B$ -games
  - The case of finite graphs
  - The case of pushdown graphs

Condition: parity and all counters are bounded.

Define:

- $\mathcal{W}_E(N)$  the set of vertices where Eve wins for the bound  $N$ .
- $\mathcal{W}_E$  the set of vertices where Eve wins for some (non-uniform) bound.

Condition: parity and all counters are bounded.

Define:

- $\mathcal{W}_E(N)$  the set of vertices where Eve wins for the bound  $N$ .
- $\mathcal{W}_E$  the set of vertices where Eve wins for some (non-uniform) bound.

**Lemma**

$$\textcircled{1} \quad \mathcal{W}_E(0) \subseteq \mathcal{W}_E(1) \subseteq \cdots \subseteq \mathcal{W}_E(N) \subseteq \mathcal{W}_E(N+1) \subseteq \cdots \subseteq \mathcal{W}_E.$$



Condition: parity and all counters are bounded.

Define:

- $\mathcal{W}_E(N)$  the set of vertices where Eve wins for the bound  $N$ .
- $\mathcal{W}_E$  the set of vertices where Eve wins for some (non-uniform) bound.

Lemma

- ①  $\mathcal{W}_E(0) \subseteq \mathcal{W}_E(1) \subseteq \dots \subseteq \mathcal{W}_E(N) \subseteq \mathcal{W}_E(N+1) \subseteq \dots \subseteq \mathcal{W}_E$ .
- ② *There exists  $N$  such that  $\mathcal{W}_E(N) = \mathcal{W}_E(N+1) = \dots$ .*

Condition: parity and all counters are bounded.

Define:


- $\mathcal{W}_E(N)$  the set of vertices where Eve wins for the bound  $N$ .
- $\mathcal{W}_E$  the set of vertices where Eve wins for some (non-uniform) bound.

## Lemma

- ①  $\mathcal{W}_E(0) \subseteq \mathcal{W}_E(1) \subseteq \dots \subseteq \mathcal{W}_E(N) \subseteq \mathcal{W}_E(N+1) \subseteq \dots \subseteq \mathcal{W}_E$ .
- ② *There exists  $N$  such that  $\mathcal{W}_E(N) = \mathcal{W}_E(N+1) = \dots$ .*
- ③ *For such  $N$ , Adam wins from  $V \setminus \mathcal{W}_E(N)$ , hence  $\mathcal{W}_E = \mathcal{W}_E(N)$ .*


- 1 Equivalence for pushdown  $\omega B$ -games
  - The case of finite graphs
  - The case of pushdown graphs

## Another example




Adam may use the stack as “credit”.

## Another example




Adam may use the stack as “credit”.

## Another example




Adam may use the stack as “credit”.

## Another example



Adam may use the stack as “credit”.


## Another example



Adam may use the stack as “credit”.




## Another example



Adam may use the stack as “credit”.

## Another example



Adam may use the stack as “credit”.

Condition: parity and all counters are bounded.

Define:

- $\mathcal{W}_E(N)$  the set of vertices where Eve wins for the bound  $N$  **in the limit**.
- $\mathcal{W}_E$  the set of vertices where Eve wins for some (non-uniform) bound.

Condition: parity and all counters are bounded.

Define:

- $\mathcal{W}_E(N)$  the set of vertices where Eve wins for the bound  $N$  **in the limit**.
- $\mathcal{W}_E$  the set of vertices where Eve wins for some (non-uniform) bound.

## Proposition

- 1  $\mathcal{W}_E(0) \subseteq \mathcal{W}_E(1) \subseteq \dots \subseteq \mathcal{W}_E(N) \subseteq \mathcal{W}_E(N+1) \subseteq \dots \subseteq \mathcal{W}_E$ .
- 2 *There exists  $N$  such that  $\mathcal{W}_E(N) = \mathcal{W}_E(N+1) = \dots$ .*
- 3 *For such  $N$ , Adam wins from  $V \setminus \mathcal{W}_E(N)$ , hence  $\mathcal{W}_E = \mathcal{W}_E(N)$ .*

Why is 2. true?



## Theorem (derived from Serre)

*For all  $N$ ,  $\mathcal{W}_E(N)$  is a regular set of configurations, recognized by an alternating automaton of size  $|Q|$  (**independent of  $N$** ).*

## Theorem

*For all pushdown games, the following are equivalent:*

- $\exists \sigma$  (strategy for Eve),  $\forall \pi$  (paths),  $\exists N \in \mathbb{N}$ ,  
 $\pi$  satisfies parity and each counter is bounded by  $N$ .
- $\exists \sigma$  (strategy for Eve),  $\exists N \in \mathbb{N}$ ,  $\forall \pi$  (paths),  
 $\pi$  satisfies parity and **eventually** each counter is bounded by  $N$ .

## Theorem

*For all pushdown games, the following are equivalent:*

- $\exists \sigma$  (strategy for Eve),  $\forall \pi$  (paths),  $\exists N \in \mathbb{N}$ ,  
 $\pi$  satisfies parity and each counter is bounded by  $N$ .
- $\exists \sigma$  (strategy for Eve),  $\exists N \in \mathbb{N}$ ,  $\forall \pi$  (paths),  
 $\pi$  satisfies parity and **eventually** each counter is bounded by  $N$ .

## Corollary

*Determining the winner in a pushdown  $\omega B$ -game is decidable.*

## Theorem

*For all pushdown games, the following are equivalent:*

- $\exists \sigma$  (strategy for Eve),  $\forall \pi$  (paths),  $\exists N \in \mathbb{N}$ ,  
 $\pi$  satisfies parity and each counter is bounded by  $N$ .
- $\exists \sigma$  (strategy for Eve),  $\exists N \in \mathbb{N}$ ,  $\forall \pi$  (paths),  
 $\pi$  satisfies parity and **eventually** each counter is bounded by  $N$ .

## Corollary

*Determining the winner in a pushdown  $\omega B$ -game is decidable.*

Remark: one can show that the collapse bound is doubly-exponential!