

# Nash equilibrium in Infinite extensive-form games


Stéphane Le Roux (TU Darmstadt)

HIGHLIGHTS 2013

September 17, 2013


# Nash equilibrium in **finite** extensive-form games

NE are traditionally obtained by **backward induction**:



# Nash equilibrium in **finite** extensive-form games


NE are traditionally obtained by **backward induction**:



Double lines represent strategic choices.

# Nash equilibrium in **finite** extensive-form games


NE are traditionally obtained by **backward induction**:



Double lines represent strategic choices.

# Nash equilibrium in **finite** extensive-form games

NE are traditionally obtained by **backward induction**:



Double lines represent strategic choices.

# Gale-Stewart games

Let  $C$  be a non-empty set.

Player <b>a</b>		$p_0 \in C$		$p_2 \in C$		$\dots$
Player <b>b</b>			$p_1 \in C$		$p_3 \in C$	$\dots$

# Gale-Stewart games

Let  $C$  be a non-empty set.

Player <b>a</b>		$p_0 \in C$		$p_2 \in C$		$\dots$
Player <b>b</b>			$p_1 \in C$		$p_3 \in C$	$\dots$

Let  $W \subseteq C^\omega$ , player **a** wins iff  $p_0 p_1 p_2 \dots \in W$ .


# Gale-Stewart games

Let  $C$  be a non-empty set.

Player <b>a</b>	$p_0 \in C$		$p_2 \in C$		$\dots$
Player <b>b</b>		$p_1 \in C$		$p_3 \in C$	$\dots$

Let  $W \subseteq C^\omega$ , player **a** wins iff  $p_0 p_1 p_2 \dots \in W$ .

Example with  $C = \{\text{left}, \text{right}\}$






# Gale-Stewart games

Let  $C$  be a non-empty set.

Player <b>a</b>	$p_0 \in C$	$p_2 \in C$	$\dots$
Player <b>b</b>	$p_1 \in C$	$p_3 \in C$	$\dots$

Let  $W \subseteq C^\omega$ , player **a** wins iff  $p_0 p_1 p_2 \dots \in W$ .


Example with  $C = \{\text{left}, \text{right}\}$



Leaf-free infinite extensive-form games: no backward induction!!!

# Quasi-Borel determinacy


Discrete topology on  $C$  and product topology on  $C^\omega$ .



$$\text{clopen balls} = \{uC^\omega \mid u \in C^*\}$$

# Quasi-Borel determinacy

Discrete topology on  $C$  and product topology on  $C^\omega$ .




$$\text{clopen balls} = \{uC^\omega \mid u \in C^*\}$$

Borel sets include the open sets of  $C^\omega$  and are close under:

1. complementation;
2. countable union;

# Quasi-Borel determinacy

Discrete topology on  $C$  and product topology on  $C^\omega$ .




$$\text{clopen balls} = \{uC^\omega \mid u \in C^*\}$$

**Quasi**-Borel sets include the open sets of  $C^\omega$  and are close under:

1. complementation;
2. countable union;
3. **open-separated union**. (For uncountable  $C$ .)

# Quasi-Borel determinacy

Discrete topology on  $C$  and product topology on  $C^\omega$ .



$$\text{clopen balls} = \{uC^\omega \mid u \in C^*\}$$

**Quasi**-Borel sets include the open sets of  $C^\omega$  and are close under:


1. complementation;
2. countable union;
3. **open-separated union**. (For uncountable  $C$ .)

**Theorem (Martin 1975, 1990)**


*If  $W$  is (quasi)-Borel, one player has a winning strategy.*

# Messy Gale-Stewart games

Gale-Stewart games,  
players play alternately:



**Messy** Gale-Stewart games,  
arbitrary order:



# First extension of Martin's theorem

## Lemma

*Let  $W$  be the winning set of a **messy** Gale-Stewart game.  
If  $W$  is quasi-Borel, one player has a winning strategy.*

Proof.




# First extension of Martin's theorem

## Lemma

*Let  $W$  be the winning set of a **messy** Gale-Stewart game. If  $W$  is quasi-Borel, one player has a winning strategy.*

Proof.






# First extension of Martin's theorem

## Lemma

Let  $W$  be the winning set of a **messy** Gale-Stewart game.  
If  $W$  is quasi-Borel, one player has a winning strategy.

## Proof.

Folklore transformation: insertion of dummy nodes.




# First extension of Martin's theorem

## Lemma


Let  $W$  be the winning set of a **messy** Gale-Stewart game.  
If  $W$  is quasi-Borel, one player has a winning strategy.

## Proof.

Folklore transformation: insertion of dummy nodes.



If winning set  $W$  is quasi-Borel,



modified  $W$  is quasi-Borel




# First extension of Martin's theorem

## Lemma

Let  $W$  be the winning set of a **messy** Gale-Stewart game.  
If  $W$  is quasi-Borel, one player has a winning strategy.

## Proof.

Folklore transformation: insertion of dummy nodes.




If winning set  $W$  is quasi-Borel,      modified  $W$  is quasi-Borel


Win. strat. after dummy insertion translate back to win. strat.



# Infinite game in extensive form




# Infinite game in extensive form




- $v : C^\omega \rightarrow O$  from plays to outcomes.

# Infinite game in extensive form



- ▶  $v : C^\omega \rightarrow O$  from plays to outcomes.
- ▶  $\prec_a$  is a binary relation over  $O$ , the preference of  $a$ .


# Infinite game in extensive form



- ▶  $v : C^\omega \rightarrow O$  from plays to outcomes.
- ▶  $\prec_a$  is a binary relation over  $O$ , the preference of  $a$ .

start  $\longrightarrow$  a  $\rightarrow$  a  $\rightarrow$  a  $\text{-----}$  a  $\rightarrow$  a  $\text{-----}$  0  
                   ↓      ↓      ↓                  ↓      ↓  
                   1      2      3                  n   n+1

# Infinite game in extensive form



- ▶  $v : C^\omega \rightarrow O$  from plays to outcomes.
- ▶  $\prec_a$  is a binary relation over  $O$ , the preference of  $a$ .


start  $\longrightarrow$  a  $\rightarrow$  a  $\rightarrow$  a  $\text{-----}$  a  $\rightarrow$  a  $\text{-----}$  0  
                   ↓      ↓      ↓                  ↓      ↓  
                   1      2      3                  n   n+1

## Theorem

The game has an NE, if the  $\prec_a^{-1}$  are strictly *well-founded*,



# Infinite game in extensive form




- ▶  $v : C^\omega \rightarrow O$  from plays to outcomes.
- ▶  $\prec_a$  is a binary relation over  $O$ , the preference of  $a$ .

$\text{start} \longrightarrow a \rightarrow a \rightarrow a \text{ ----- } a \rightarrow a \text{ ----- } 0$   
 $\qquad\qquad\qquad \downarrow \quad \downarrow \quad \downarrow \qquad\qquad\qquad \downarrow \quad \downarrow$   
 $\qquad\qquad\qquad 1 \quad 2 \quad 3 \qquad\qquad\qquad n \quad n+1$

## Theorem

The game has an NE, if the  $\prec_a^{-1}$  are strictly *well-founded*,  
 $v^{-1}(o)$  is *quasi-Borel* for all  $o$

# Infinite game in extensive form



- ▶  $v : C^\omega \rightarrow O$  from plays to outcomes.
- ▶  $\prec_a$  is a binary relation over  $O$ , the preference of  $a$ .


$\text{start} \longrightarrow a \rightarrow a \rightarrow a \text{ ----- } a \rightarrow a \text{ ----- } 0$   
                    $\downarrow$      $\downarrow$      $\downarrow$                      $\downarrow$      $\downarrow$   
                   1    2    3                    n   n+1

## Theorem

The game has an NE, if the  $\prec_a^{-1}$  are strictly *well-founded*,  
 $v^{-1}(o)$  is *quasi-Borel* for all  $o$  in *countable*  $O$ .

# Idea of the proof for finite games


Both players get the same payoffs.



- At the root,  $a$  seeks the best guaranteed payoff.

# Idea of the proof for finite games

Both players get the same payoffs.




- At the root,  $a$  seeks the best guaranteed payoff.

## 8

# Idea of the proof for finite games


Both players get the same payoffs.



- ▶ At the root, *a* seeks the best guaranteed payoff.
- ▶ In the subgame ignored by *a*, player *b* threatens *a* (old idea).
- ▶ Repeat this in the subgame chosen by *a*.

# Idea of the proof for finite games

Both players get the same payoffs.



- ▶ At the root,  $a$  seeks the best guaranteed payoff.
- ▶ In the subgame ignored by  $a$ , player  $b$  threatens  $a$  (old idea).
- ▶ Repeat this in the subgame chosen by  $a$ .

# Idea of the proof for infinite games

$$\begin{array}{ccccccccccc} a & \rightarrow & a & \rightarrow & a & \rightarrow & a & \rightarrow & a & \rightarrow & \dots & 0 \\ \downarrow & & \downarrow & & \downarrow & & \downarrow & & & & & \\ 1 & & 1 & & 1 & & 1 & & & & & \end{array}$$



# Idea of the proof for infinite games

$$\begin{array}{ccccccc} \mathbf{a} & \Rightarrow & a & \rightarrow & a & \rightarrow & a \text{ ----- } 0 \\ \downarrow & & \Downarrow & & \downarrow & & \downarrow \\ 1 & & 1 & & 1 & & 1 \end{array}$$

## Idea of the proof for infinite games

$$\begin{array}{ccccccc} a \Rightarrow \mathbf{a} \Rightarrow a \rightarrow a & \text{-----} & 0 \\ \downarrow & \downarrow & \Downarrow & \downarrow & & & \\ 1 & 1 & 1 & 1 & & & \end{array}$$

# Idea of the proof for infinite games

$$\begin{array}{ccccccc} a & \Rightarrow & a & \Rightarrow & \textcolor{red}{a} & \Rightarrow & a \text{ ----- } 0 \\ \downarrow & & \downarrow & & \downarrow & & \Downarrow \\ 1 & & 1 & & 1 & & 1 \end{array}$$

# Idea of the proof for infinite games

$$\begin{array}{ccccccc} a & \Rightarrow & a & \Rightarrow & \textcolor{red}{a} & \Rightarrow & a & = & = & = & = & = & 0 \\ \downarrow & & \downarrow & & \downarrow & & \downarrow & & & & & & \\ 1 & & 1 & & 1 & & 1 & & & & & & \end{array}$$

# Idea of the proof for infinite games

$$\begin{array}{ccccccc} a & \Rightarrow & a & \Rightarrow & \textcolor{red}{a} & \Rightarrow & a \\ \downarrow & & \downarrow & & \downarrow & & \downarrow \\ 1 & & 1 & & 1 & & 1 \end{array} \quad = = = = = 0$$

- To fix it, change subprofiles only if improving the guarantee.

# Idea of the proof for infinite games


$$\begin{array}{ccccccc} a & \Rightarrow & a & \Rightarrow & \mathbf{a} & \Rightarrow & a = = = = 0 \\ \downarrow & & \downarrow & & \downarrow & & \downarrow \\ 1 & & 1 & & 1 & & 1 \end{array}$$

- ▶ To fix it, change subprofiles only if improving the guarantee.
- ▶ No infinite ascending preference chain, so each player improves guarantee only finitely many times.

# Idea of the proof for infinite games

$$\begin{array}{ccccccc} a & \Rightarrow & a & \Rightarrow & \textcolor{red}{a} & \Rightarrow & a & = & = & = & = & 0 \\ \downarrow & & \downarrow & & \downarrow & & \downarrow & & & & & \\ 1 & & 1 & & 1 & & 1 & & & & & \end{array}$$


- ▶ To fix it, change subprofiles only if improving the guarantee.
- ▶ No infinite ascending preference chain, so each player improves guarantee only finitely many times.



# Idea of the proof for infinite games

$$\begin{array}{ccccccc} a & \Rightarrow & a & \Rightarrow & \textcolor{red}{a} & \Rightarrow & a & = & = & = & = & 0 \\ \downarrow & & \downarrow & & \downarrow & & \downarrow & & & & & \\ 1 & & 1 & & 1 & & 1 & & & & & \end{array}$$

- ▶ To fix it, change subprofiles only if improving the guarantee.
- ▶ No infinite ascending preference chain, so each player improves guarantee only finitely many times.






# Idea of the proof for infinite games

$$\begin{array}{ccccccc} a & \Rightarrow & a & \Rightarrow & \textcolor{red}{a} & \Rightarrow & a = = = = 0 \\ \downarrow & & \downarrow & & \downarrow & & \downarrow \\ 1 & & 1 & & 1 & & 1 \end{array}$$


- ▶ To fix it, change subprofiles only if improving the guarantee.
- ▶ No infinite ascending preference chain, so each player improves guarantee only finitely many times.



# Idea of the proof for infinite games

$$\begin{array}{ccccccc}
 a & \Rightarrow & a & \Rightarrow & \textcolor{red}{a} & \Rightarrow & a & = & = & = & = & 0 \\
 \downarrow & & \downarrow & & \downarrow & & \downarrow & & & & & \\
 1 & & 1 & & 1 & & 1 & & & & & 
 \end{array}$$


- ▶ To fix it, change subprofiles only if improving the guarantee.
- ▶ No infinite ascending preference chain, so each player improves guarantee only finitely many times.



# Idea of the proof for infinite games

$$\begin{array}{cccc}
 a & \Rightarrow & a & \Rightarrow & \textcolor{red}{a} & \Rightarrow & a & = & = & = & = & 0 \\
 \downarrow & & \downarrow & & \downarrow & & \downarrow & & & & \\
 1 & & 1 & & 1 & & 1 & & & & 
 \end{array}$$

- ▶ To fix it, change subprofiles only if improving the guarantee.
- ▶ No infinite ascending preference chain, so each player improves guarantee only finitely many times.




Transform  $g$  into a messy Gale-Stewart game:

$a$ 's opponents team up and win if  $a$  gets  $x' \leq_a x$ .

# Idea of the proof for infinite games

$$\begin{array}{cccc}
 a \Rightarrow a \Rightarrow \textcolor{red}{a} \Rightarrow a & = & = & = & = & 0 \\
 \downarrow & \downarrow & \downarrow & \downarrow & & \\
 1 & 1 & 1 & 1 & & 
 \end{array}$$

- ▶ To fix it, change subprofiles only if improving the guarantee.
- ▶ No infinite ascending preference chain, so each player improves guarantee only finitely many times.



Transform  $g$  into a messy Gale-Stewart game:

$a$ 's opponents team up and win if  $a$  gets  $x' \leq_a x$ .


Player  $a$  has no winning strategy, by definition of the guarantee.

The winning set is quasi-Borel, so  $a$ 's opponents have a win. strat.

# Idea of the proof for infinite games

$$\begin{array}{cccc}
 a \Rightarrow a \Rightarrow \textcolor{red}{a} \Rightarrow a & = & = & = & = & 0 \\
 \downarrow & \downarrow & \downarrow & \downarrow & & \\
 1 & 1 & 1 & 1 & & 
 \end{array}$$

- ▶ To fix it, change subprofiles only if improving the guarantee.
- ▶ No infinite ascending preference chain, so each player improves guarantee only finitely many times.



Transform  $g$  into a messy Gale-Stewart game:

$a$ 's opponents team up and win if  $a$  gets  $x' \leq_a x$ .

Player  $a$  has no winning strategy, by definition of the guarantee.

The winning set is quasi-Borel, so  $a$ 's opponents have a win. strat.

# Conclusion

The actual result works for more than just quasi-Borel sets:

- ▶ If a pointclass with reasonable closure properties yields determined games, it yields also games with NE.

# Conclusion

The actual result works for more than just quasi-Borel sets:

- ▶ If a pointclass with reasonable closure properties yields determined games, it yields also games with NE.
- ▶ Axiom of determinacy  $\Rightarrow$  every game has an NE (under preference condition).

# Conclusion

The actual result works for more than just quasi-Borel sets:

- ▶ If a pointclass with reasonable closure properties yields determined games, it yields also games with NE.
- ▶ Axiom of determinacy  $\Rightarrow$  every game has an NE (under preference condition).

Questions:

- ▶ Several proofs in logic invoke Borel determinacy, can the results be extended by invoking existence of NE?



# Conclusion

The actual result works for more than just quasi-Borel sets:

- ▶ If a pointclass with reasonable closure properties yields determined games, it yields also games with NE.
- ▶ Axiom of determinacy  $\Rightarrow$  every game has an NE (under preference condition).

Questions:

- ▶ Several proofs in logic invoke Borel determinacy, can the results be extended by invoking existence of NE?
- ▶ Is anyone familiar with the proof of Borel determinacy?