

Total Dual Integrality and Integer Polyhedra*

F. R. Giles†

Department of Operations Research
Cornell University
Ithaca, New York

and

W. R. Pulleyblank‡

CORE, de Croylaan 54
3030 Heverlee
Belgium

Submitted by Hans Schneider

ABSTRACT

A linear system $Ax \leq b$ (A, b rational) is said to be totally dual integral (TDI) if for any integer objective function c such that $\max \{cx : Ax \leq b\}$ exists, there is an integer optimum dual solution. We show that if P is a polytope all of whose vertices are integer valued, then it is the solution set of a TDI system $Ax \leq b$ where b is integer valued. This was shown by Edmonds and Giles [4] to be a sufficient condition for a polytope to have integer vertices.

1. INTRODUCTION

Let $Ax \leq b$ be a linear system with A and b rational. We say that this linear system is *totally dual integral* (or TDI) if for any integer valued c such that the linear program

$$\text{maximize } \{cx : Ax \leq b\}$$

*Research supported in part by the National Research Council of Canada.

†Current address: Department of Mathematics, University of Kentucky, Lexington, Kentucky.

‡Current address: Department of Computer Science, University of Calgary, Calgary, Alberta, Canada.

has an optimum solution, the corresponding dual linear program has an integer optimal solution. This concept was introduced by Edmonds and Giles [4], who showed:

THEOREM 1.1. *If a polyhedron is the solution set of a TDI system $Ax \leq b$, where b is integer valued, then every nonempty face of P contains an integer point—in particular, every vertex of P is integer valued.*

In Sec. 3 we observe that for any rational linear system $Ax \leq b$ there exists a rational α such that $(\alpha A)x \leq \alpha b$ is a TDI system. Of course, if the polyhedron defined by the system has some nonempty face which contains no integer point, then for any α which makes $(\alpha A)x \leq \alpha b$ a TDI system, we will have αb fractional. However, we also prove a converse to the Edmonds-Giles theorem, namely that if P is a polyhedron such that every nonempty face contains an integer point, then there exists a TDI system $Ax \leq b$ with b integer such that $P = \{x : Ax \leq b\}$. We give a proof of this result which is based on the fact that the set of objective functions maximized over a face of a polyhedron forms a convex cone. In Sec. 2 we present the basic definitions and results on cones which we require. In particular, we give a new short proof of a classical theorem of Hilbert which shows that a rational cone has a finite integer basis (see Theorem 2.1)).

2. RATIONAL CONES

Let D be a finite subset of \mathbb{R}^j . The cone $K(D)$ generated by D is the set of all vectors $x \in \mathbb{R}^j$ such that $x = \sum (\lambda_d d : d \in D)$, where for each $d \in D$, λ_d is a nonnegative real number. We call K a rational cone if $K = K(D)$ for some $D \subseteq \mathbb{R}^j$ such that every member of D is rational. We now prove a classical theorem of Hilbert [7] which shows that for any rational cone K there exists a finite set Z of integer members of K such that every integer $x \in K$ can be expressed as a nonnegative integer linear combination of members of Z .

This is not in general true for a cone with irrational generators. For example, consider the cone K in \mathbb{R}^2 generated by $(0, 1)$ and $(1, z)$, where z is some positive irrational. The line $\alpha \cdot (1, z)$ for $\alpha \geq 0$ contains no rational points, and hence for any finite subset Z of integer members of K , there must exist some rational $p \in K - K(Z)$. But then if we multiply p by a sufficiently large integer, we obtain an integer $\hat{p} \in K - K(Z)$, and so \hat{p} is certainly not a positive integer linear combination of members of Z .

THEOREM 2.1. *Let K be a rational cone. Then there exists a finite set Z of integer members of K such that every integer $x \in K$ is a nonnegative integer linear combination of members of Z .*

Proof. Let $K = K(D)$. We may assume that D is a set of integer vectors. Let

$$Z \equiv \{x \in \mathbb{R}^I : x \text{ is an integer vector,} \\ x = \sum (\lambda_d d : d \in D), \\ 0 \leq \lambda_d \leq 1 \text{ for all } d \in D\}.$$

Then Z is a finite subset of integer vectors in K . For any integer $x \in K$, there exists $(\lambda_d \geq 0 : d \in D)$ such that

$$x = \sum (\lambda_d d : d \in D) = \sum (\lfloor \lambda_d \rfloor d : d \in D) + \sum ((\lambda_d - \lfloor \lambda_d \rfloor) d : d \in D),$$

where for any real number t , $\lfloor t \rfloor$ denotes the greatest integer no greater than t . Because x and $\sum (\lfloor \lambda_d \rfloor d : d \in D)$ are integer vectors, $\sum ((\lambda_d - \lfloor \lambda_d \rfloor) d : d \in D)$ is an element of Z . Since $D \subseteq Z$, x is a nonnegative integer combination of members of Z . ■

3. INTEGER POLYHEDRA AND TOTAL DUAL INTEGRALITY

Let $P = \{x \in \mathbb{R}^n : Ax \leq b\}$ be a polyhedron. We say that P is an *integer polyhedron* if every nonempty face of P contains an integer point. If P is pointed, that is, has at least one vertex, this is equivalent to the assertion that every vertex of P is integer valued. For any nonempty face F of P , there exists a linear objective function c such that cx is maximized over P by precisely the members of F . Conversely, for any linear objective function c such that cx has a maximum over P , there exists a nonempty face F of P such that cx is maximized over P by the members of F . Thus it follows immediately that P is an integer polyhedron if and only if for every linear objective function c such that cx has a maximum over P , there is an integer member of P for which the maximum is attained. Edmonds and Giles [4] prove the following strengthening of this result.

THEOREM 3.1. *Let $P = \{x \in \mathbb{R}^n : Ax \leq b\}$ where A and b are rational. Then P is an integer polyhedron if and only if $\max\{cx : x \in P\}$ is integer valued for every integer objective function c having a maximum over P .*

The important fact about Theorem 3.1 is the assertion that if P is not an integer polyhedron, then there exists an *integer* objective function which when maximized over P takes on a *fractional* value.

If P has vertices, then this result can be proved using the Gomory cutting-plane algorithm [6], since if this algorithm finds a fractional vertex v of polyhedron P , it generates a cut $ax \leq \beta$ where a is integer valued, a is maximized over P at v , and av is not integer valued.

If a polyhedron P is the solution set of a TDI system with integer right hand sides, then for any integer objective function c which has a maximum over P , by linear programming duality this maximum value is an integer. Hence from Theorem 3.1 P is an integer polyhedron, and so Theorem 1.1 follows directly from Theorem 3.1. We now prove two converse results.

THEOREM 3.2. *For any rational linear system $Ax \leq b$ there is a positive rational number α such that $\alpha Ax \leq \alpha b$ is TDI.*

Proof. It is trivial to show that for a fixed integer valued c there exists an α such that the polyhedron $\{y \in \mathbb{R}^m : y(\alpha A) = c, y > 0\}$ is an integer polyhedron, since multiplying A by α has the effect of multiplying all extreme points by $1/\alpha$. However, we must show that it is possible to choose a single α which will work for every possible integer c .

We may assume that $A = (a_{ij} : i \in I, j \in J)$ is integer valued. Let N be the set of all $|I| \times |I|$ nonsingular submatrices of the concatenation of A with a $|I| \times |I|$ identity matrix. Let

$$\beta \equiv \left| \prod (\det(B) : B \in N) \right| \quad \text{and} \quad \alpha \equiv 1/\beta.$$

A simple application of Cramer's rule now shows that every component of every basic feasible solution y^* of $y(\alpha A) = c, y \geq 0$, is integer valued for any integer c . ■

This theorem makes clear the importance of the hypothesis in Theorem 1.1 that b should be integer valued. If $P = \{x : Ax \leq b\}$ is not an integer polyhedron, then for any rational α which makes $\alpha Ax \leq \alpha b$ a TDI system, we must have αb fractional. However, we show the following.

THEOREM 3.3. *Let $P = \{x : Ax \leq b\}$, where A and b are rational. If P is an integer polyhedron, then there exists a TDI linear system $A'x \leq b'$ with b' integer such that $P = \{x \in \mathbb{R}^J : A'x \leq b'\}$.*

Before proving Theorem 3.3 we give some definitions and a lemma. Let $A = (a_{ij} : i \in I, j \in J)$ and $b = (b_i : i \in I)$. For any $H \subseteq I$ we let $A[H] \equiv (a_{ij} : i \in H, j \in J)$ and $b[H] \equiv (b_i : i \in H)$. For any $i \in I$ we let $A[i]$ denote $(a_{ij} : j \in J)$. Let F be any nonempty face of the polyhedron $P = \{x \in \mathbb{R}^J : Ax \leq b\}$. Then there is a unique maximal subset H of I such that $F = \{x \in P : A[H]x = b[H]\}$. We call H the *equality set* of F (relative to the system $Ax \leq b$). Finally, we let $C(F)$ be the set of all $c \in \mathbb{R}^J$ such that every $x \in F$ maximizes cx over P .

LEMMA 3.4. *Let F be a nonempty face of $P = \{x : Ax \leq b\}$, and let H be the equality set of F . Then, if R is the set of rows of $A[H]$,*

(a) $C(F) = K(R)$ (that is, $C(F)$ is the cone generated by the rows of A indexed by the members of the equality set of F);

(b) for any $c \in C(F)$, if $c = \sum(\lambda_h A[h] : h \in H)$ where $\lambda_h \geq 0$ for all $h \in H$, then $\max \{cx : x \in P\} = \sum(\lambda_h b_h : h \in H)$.

Proof. If $c \in K(R)$, then $c = \sum(\lambda_h A[h] : h \in H)$, where $\lambda_h \geq 0$ for all $h \in H$. For any $x \in F$, $cx = \sum(\lambda_h A[h]x : h \in H) = \sum(\lambda_h b_h : h \in H)$. For any $x \in P$, $cx = \sum(\lambda_h A[h]x : h \in H) \leq \sum(\lambda_h b_h : h \in H)$. Hence $c \in C(F)$. Moreover (b) is established for every $c \in K(R)$.

Conversely, for any $c \in C(F)$ consider the dual linear program to $\max \{cx : x \in P\}$, namely $\min \{by : y \geq 0, yA = c\}$. There exists an optimal solution y^* , since $c \in C(F)$; and moreover, if $y_i^* > 0$, then by complementary slackness, $A[i]x = b_i$ for any $x \in P$ which maximizes cx over P . Hence $A[i]x = b_i$ for all $x \in F$, and so $i \in H$. Therefore y expresses c as a nonnegative linear combination of members of R , so $c \in K(R)$. Thus $C(F) = K(R)$, and the proof is complete. ■

Proof of Theorem 3.3. We may assume that A and b are integer. If c is an objective function that has a maximum value over P , then there is a minimal nonempty face F of P for whose members the maximum is attained, so it will be sufficient to show:

(c) For any minimal nonempty face F of P there is a linear system $D_F x \leq d_F$, with d_F integer, satisfied by every $x \in P$ and such that for any $c \in C(F)$ there exists an integer optimal dual solution to the linear program $\max \{cx : Ax \leq b, D_F x \leq d_F\}$.

Let F be a minimal nonempty face of P , with equality set H . By Lemma 3.4(a), $C(F)$ is the rational cone generated by the rows of $A[H]$. By Theorem 2.1 there is a finite "integer basis" Z of $C(F)$ such that every $z \in Z$ is an integer and every integer $c \in C(F)$ is a nonnegative integer linear combination of members of Z . For any $z \in Z$ let $\zeta_z = \max \{zx : x \in P\} = z\bar{x}$ for any $\bar{x} \in F$. Since F contains an integer point, ζ_z is integer valued. Moreover every $x \in P$ satisfies $zx \leq \zeta_z$. We let $D_F x \leq d_F$ be the linear system $(zx \leq \zeta_z : z \in Z)$. Now since every integer $c \in C(F)$ is a nonnegative integer linear combination of rows of D_F , (c) follows from (b). Hence a TDI linear system defining P is $(D_F x \leq d_F : F \text{ is a minimal nonempty face of } P)$, which is a finite system because P has a finite number of minimal nonempty faces. ■

For a given integer polyhedron P , a popular problem is that of finding a minimal linear system $Ax \leq b$ such that $P = \{x : Ax \leq b\}$. In view of Theorem 3.3 a second question that can be asked is the following. What is a minimal TDI system $A'x \leq b'$ with integer right-hand sides such that $P = \{x : A'x \leq b'\}$? For some classes of polyhedra these two linear systems can be "identical"; we can find a minimal linear system $Ax \leq b$ which defines P and which

is TDI with integer b . For example, if $G=(V, E)$ is a graph and $b=(b_i: i \in V)$ is a vector of positive integers, the matching polyhedron $P(G, b)$ is defined to be the convex hull of all nonnegative integer vectors $x=(x_j: j \in E)$ such that for each node i of G , the sum of the x_j over the edges of G incident with i is at most b_i . Let $\bar{1}$ be the unit vector indexed by V .

THEOREM 3.5 (Cunningham and Marsh [2]). *The minimal linear system $Ax \leq d$ such that $P(G, \bar{1}) = \{x: Ax \leq d\}$ (scaled in the "natural way" so that A is 0-1 valued) is a TDI system with d integer.*

However, this result does not generalize to matching problems with arbitrary b . In general a TDI linear system with integer right-hand side that defines $P(G, b)$ will be larger than a minimal linear system necessary to define $P(G, b)$.

Other examples of polyhedra for which the two linear systems above can be identical are the intersection of integral polymatroids (see Edmonds [3] and Giles [5]) and the convex hull of the incidence vectors of independent sets of nodes in a perfect graph (see Chvátal [1]).

Many of the ideas presented here arose during discussions with Michael Todd and Robert Jeroslow. In particular, conversations with Jeroslow concerning the subject matter of Sec. 2 were valuable.

REFERENCES

- 1 V. Chvátal, On certain polytopes associated with graphs, *J. Combinatorial Theory Ser. B* 18:138–154 (1975).
- 2 W. H. Cunningham and A. B. Marsh, A primal algorithm for optimum matching, Technical Report No. 262, Dept of Math. Sciences, Johns Hopkins Univ., 1976.
- 3 J. Edmonds, Submodular functions, matroids and certain polyhedra, in *Combinatorial Structures and Their Applications, Proceedings of the Calgary International Conference*, (Guy et al, Eds.) Gordon and Breach, New York, 1970, pp. 69–87.
- 4 J. Edmonds and F. R. Giles, A min-max relation for submodular functions on graphs, *Ann. Discrete Math.* 1:185–204 (1977).
- 5 F. R. Giles, Submodular functions, graphs and integer polyhedra, Ph. D. Thesis, Univ. of Waterloo, 1975.
- 6 R. E. Gomory, An algorithm for integer solutions to linear programs, in *Recent Advances in Mathematical Programming*, (Graves and Wolf, Eds.), McGraw-Hill, 1963, pp. 269–302.
- 7 D. Hilbert, Über die Theorie der algebraischen Formen, *Math. Ann.* 36:473–534 (1890).

Received 30 August 1977; revised 1 August 1978