

A PROOF OF KAMP’S THEOREM

ALEXANDER RABINOVICH

The Blavatnik School of Computer Science, Tel Aviv University
e-mail address: rabinoa@post.tau.ac.il

ABSTRACT. We provide a simple proof of Kamp’s theorem.

1. INTRODUCTION

Temporal Logic (*TL*) introduced to Computer Science by Pnueli in [10] is a convenient framework for reasoning about “reactive” systems. This has made temporal logics a popular subject in the Computer Science community, enjoying extensive research in the past 30 years. In *TL* we describe basic system properties by *atomic propositions* that hold at some points in time, but not at others. More complex properties are expressed by formulas built from the atoms using Boolean connectives and *Modalities* (temporal connectives): A k -place modality M transforms statements $\varphi_1, \dots, \varphi_k$ possibly on ‘past’ or ‘future’ points to a statement $M(\varphi_1, \dots, \varphi_k)$ on the ‘present’ point t_0 . The rule to determine the truth of a statement $M(\varphi_1, \dots, \varphi_k)$ at t_0 is called a *truth table* of M . The choice of particular modalities with their truth tables yields different temporal logics. A temporal logic with modalities M_1, \dots, M_k is denoted by $TL(M_1, \dots, M_k)$.

The simplest example is the one place modality $\Diamond P$ saying: “ P holds some time in the future.” Its truth table is formalized by $\varphi_\Diamond(x_0, P) := \exists x(x > x_0 \wedge P(x))$. This is a formula of the First-Order Monadic Logic of Order (*FOMLO*) - a fundamental formalism in Mathematical Logic where formulas are built using atomic propositions $P(x)$, atomic relations between elements $x_1 = x_2$, $x_1 < x_2$, Boolean connectives and first-order quantifiers $\exists x$ and $\forall x$. Two more natural modalities are the modalities Until (“*Until*”) and Since (“*Since*”). $X\text{Until}Y$ means that X will hold from now until a time in the future when Y will hold. $X\text{Since}Y$ means that Y was true at some point of time in the past and since that point X was true until (not necessarily including) now. Both modalities have truth tables in *FOMLO*. Most modalities used in the literature are defined by such *FOMLO* truth tables, and as a result, every temporal formula translates directly into an equivalent *FOMLO* formula. Thus, the different temporal logics may be considered as a convenient way to use fragments of *FOMLO*. *FOMLO* can also serve as a yardstick by which one is able to check the strength of temporal logics: A temporal logic is *expressively complete* for

2012 ACM CCS: [Theory of computation]: Logic—Modal and temporal logics.

Key words and phrases: Temporal Logic, Monadic Logic, Expressive Completeness.

a fragment L of $FOMLO$ if every formula of L with a single free variable x_0 is equivalent to a temporal formula.

Actually, the notion of expressive completeness refers to a temporal logic and to a model (or a class of models), since the question whether two formulas are equivalent depends on the domain over which they are evaluated. Any (partially) ordered set with monadic predicates is a model for TL and $FOMLO$, but the main, *canonical*, linear time intended models are the non-negative integers $\langle \mathbb{N}, < \rangle$ for discrete time and the reals $\langle \mathbb{R}, < \rangle$ for continuous time.

Kamp's theorem [8] states that the temporal logic with modalities **Until** and **Since** is expressively complete for $FOMLO$ over the above two linear time canonical¹ models.

This seminal theorem initiated the whole study of expressive completeness, and it remains one of the most interesting and distinctive results in temporal logic; very few, if any, similar 'modal' results exist. Several alternative proofs of it and stronger results have appeared; none of them are trivial (at least to most people) [7].

The objective of this paper is to provide a simple proof of Kamp's theorem.

The rest of the paper is organized as follows: In Section 2 we recall the definitions of the monadic logic, the temporal logics and state Kamp's theorem. Section 3 introduces formulas in a normal form and states their simple properties. In Section 4 we prove Kamp's theorem. The proof of one proposition is postponed to Section 5. Section 6 comments on the previous proofs of Kamp's theorem. Finally, in Section 7, we show that our proof can be easily modified to prove expressive completeness for the future fragment of $FOMLO$.

2. PRELIMINARIES

In this section we recall the definitions of the first-order monadic logic of order, the temporal logics and state Kamp's theorem.

Fix a set Σ of *atoms*. We use $P, Q, R, S \dots$ to denote members of Σ . The syntax and semantics of both logics are defined below with respect to such Σ .

2.1. First-Order Monadic Logic of Order. *Syntax*: In the context of $FOMLO$, the atoms of Σ are referred to (and used) as *unary predicate symbols*. Formulas are built using these symbols, plus two binary relation symbols: $<$ and $=$, and a set of first-order variables (denoted: x, y, z, \dots). Formulas are defined by the grammar:

$$\begin{aligned} \text{atomic} &::= x < y \mid x = y \mid P(x) \text{ (where } P \in \Sigma) \\ \varphi &::= \text{atomic} \mid \neg\varphi_1 \mid \varphi_1 \vee \varphi_2 \mid \varphi_1 \wedge \varphi_2 \mid \exists x\varphi_1 \mid \forall x\varphi_1 \end{aligned}$$

We also use the standard abbreviated notation for **bounded quantifiers**, e.g., $(\exists x)_{>z}(\dots)$ denotes $\exists x((x > z) \wedge (\dots))$, and $(\forall x)^{<z}(\dots)$ denotes $\forall x((x < z) \rightarrow (\dots))$, and $((\forall x)_{>z_1}^{<z_2}(\dots))$ denotes $\forall x((z_1 < x < z_2) \rightarrow (\dots))$, etc.

Semantics. Formulas are interpreted over *labeled linear orders* which are called *chains*. A Σ -*chain* is a triplet $\mathcal{M} = (T, <, \mathcal{I})$ where T is a set - the *domain* of the chain, $<$ is a linear order relation on T , and $\mathcal{I} : \Sigma \rightarrow \mathcal{P}(T)$ is the *interpretation* of Σ (where \mathcal{P} is the powerset notation). We use the standard notation $\mathcal{M}, t_1, t_2, \dots, t_n \models \varphi(x_1, x_2, \dots, x_n)$ to indicate that the formula φ with free variables among x_1, \dots, x_n is satisfiable in \mathcal{M} when

¹the technical notion which unifies $\langle \mathbb{N}, < \rangle$ and $\langle \mathbb{R}, < \rangle$ is Dedekind completeness.

x_i are interpreted as elements t_i of \mathcal{M} . For atomic $P(x)$ this is defined by: $\mathcal{M}, t \models P(x)$ iff $t \in \mathcal{I}(P)$; the semantics of $<, =, \neg, \wedge, \vee, \exists$ and \forall is defined in a standard way.

2.2. $TL(\text{Until}, \text{Since})$ Temporal Logic. In this section we recall the syntax and semantics of a temporal logic with *strict*-Until and *strict*-Since modalities, denoted by $TL(\text{Until}, \text{Since})$.

In the context of temporal logics, the atoms of Σ are used as *atomic propositions* (also called *propositional atoms*). Formulas of $TL(\text{Until}, \text{Since})$ are built using these atoms, Boolean connectives and *strict*-Until and *strict*-Since binary modalities. The formulas are defined by the grammar:

$$F ::= \text{True} \mid P \mid \neg F_1 \mid F_1 \vee F_2 \mid F_1 \wedge F_2 \mid F_1 \text{Until} F_2 \mid F_1 \text{Since} F_2,$$

where $P \in \Sigma$.

Semantics. Formulas are interpreted at *time-points* (or *moments*) in chains (elements of the domain). The semantics of $TL(\text{Until}, \text{Since})$ formulas is defined inductively: Given a chain $\mathcal{M} = (T, <, \mathcal{I})$ and $t \in T$, define when a formula F holds in \mathcal{M} at t - denoted $\mathcal{M}, t \models F$:

- $\mathcal{M}, t \models P$ iff $t \in \mathcal{I}(P)$, for any propositional atom P .
- $\mathcal{M}, t \models F_1 \vee F_2$ iff $\mathcal{M}, t \models F_1$ or $\mathcal{M}, t \models F_2$; similarly for \wedge and \neg .
- $\mathcal{M}, t \models F_1 \text{Until} F_2$ iff there is $t' > t$ such that $\mathcal{M}, t' \models F_2$ and $\mathcal{M}, t_1 \models F_1$ for all $t_1 \in (t, t')$.
- $\mathcal{M}, t \models F_1 \text{Since} F_2$ iff there is $t' < t$ such that $\mathcal{M}, t' \models F_2$ and $\mathcal{M}, t_1 \models F_1$ for all $t_1 \in (t', t)$.

We will use standard abbreviations. As usual $\Box F$ (respectively, $\Boxleftarrow F$) is an abbreviation for $\neg(\text{TrueUntil}(\neg F))$ (respectively, $\neg(\text{TrueSince}(\neg F))$), and $\mathbf{K}^+(F)$ (respectively, $\mathbf{K}^-(F)$) is an abbreviation for $\neg((\neg F) \text{Until} \text{True})$ (respectively, $\neg((\neg F) \text{Since} \text{True})$).

- (1) $\Box F$ (respectively, $\Boxleftarrow F$) holds at t iff F holds everywhere after (respectively, before) t .
- (2) $\mathbf{K}^-(F)$ holds at a moment t iff $t = \sup(\{t' \mid t' < t \text{ and } F \text{ holds at } t'\})$.
- (3) $\mathbf{K}^+(F)$ holds at a moment t iff $t = \inf(\{t' \mid t' > t \text{ and } F \text{ holds at } t'\})$.

Note that $\mathbf{K}^+(\text{True})$ (respectively, $\mathbf{K}^-(\text{True})$) holds at t in \mathcal{M} if t is a right limit (respectively, a left limit) point of the underlining order. In particular, both $\mathbf{K}^+(\text{True})$ and $\mathbf{K}^-(\text{True})$ are equivalent to False in the chains over $(\mathbb{N}, <)$,

2.3. Kamp's Theorem. Equivalence between temporal and monadic formulas is naturally defined: F is equivalent to $\varphi(x)$ over a class \mathcal{C} of structures iff for any $\mathcal{M} \in \mathcal{C}$ and $t \in \mathcal{M}$: $\mathcal{M}, t \models F \Leftrightarrow \mathcal{M}, t \models \varphi(x)$. If \mathcal{C} is the class of all chains, we will say that F is equivalent to φ .

A linear order $(T, <)$ is *Dedekind complete* if for every non-empty subset S of T , if S has a lower bound in T then it has a greatest lower bound, written $\inf(S)$, and if S has an upper bound in T then it has a least upper bound, written $\sup(S)$. The canonical linear time models $(\mathbb{N}, <)$ and $(\mathbb{R}, <)$ are Dedekind complete, while the order of the rationals is not Dedekind complete. A chain is Dedekind complete if its underlying linear order is Dedekind complete.

The fundamental theorem of Kamp's states that $TL(\text{Until}, \text{Since})$ is expressively equivalent to *FOMLO* over Dedekind complete chains.

Theorem 2.1 (Kamp [8]). (1) *Given any $TL(\text{Until}, \text{Since})$ formula A there is a FOMLO formula $\varphi_A(x)$ which is equivalent to A over all chains.*

- (2) *Given any FOMLO formula $\varphi(x)$ with one free variable, there is a $TL(\text{Until}, \text{Since})$ formula which is equivalent to φ over Dedekind complete chains.*

The meaning preserving translation from $TL(\text{Until}, \text{Since})$ to $FOMLO$ is easily obtained by structural induction. The contribution of our paper is a proof of Theorem 2.1 (2). The proof is constructive. An algorithm which for every $FOMLO$ formula $\varphi(x)$ constructs a $TL(\text{Until}, \text{Since})$ formula which is equivalent to φ over Dedekind complete chains is easily extracted from our proof. However, this algorithm is not efficient in the sense of complexity theory. This is unavoidable because there is a non-elementary succinctness gap between $FOMLO$ and $TL(\text{Until}, \text{Since})$ even over the class of finite chains, i.e., for every $m, n \in \mathbb{N}$ there is a $FOMLO$ formula $\varphi(x_0)$ of size $|\varphi| > n$ which is not equivalent (even over finite chains) to any $TL(\text{Until}, \text{Since})$ formula of size $\leq \exp(m, |\varphi|)$, where the m -iterated exponential function $\exp(m, n)$ is defined by induction on m so that $\exp(1, n) = 2^n$, and $\exp(m+1, n) = 2^{\exp(m, n)}$.

3. $\exists\forall$ FORMULAS

First, we introduce $\vec{\exists}\forall$ formulas which are instances of the Decomposition formulas of [3].

Definition 3.1 ($\vec{\exists}\forall$ -formulas). Let Σ be a set of monadic predicate names. An $\vec{\exists}\forall$ -formula over Σ is a formula of the form:

$$\begin{aligned} \psi(z_0, \dots, z_m) := & \exists x_n \dots \exists x_1 \exists x_0 \\ & \left(\bigwedge_{k=0}^m z_k = x_{i_k} \right) \wedge (x_n > x_{n-1} > \dots > x_1 > x_0) \text{ “ordering of } x_i \text{ and } z_j\text{”} \\ & \wedge \bigwedge_{j=0}^n \alpha_j(x_j) \text{ “Each } \alpha_j \text{ holds at } x_j\text{”} \\ & \wedge \bigwedge_{j=1}^n [(\forall y)_{>x_{j-1}}^{<x_j} \beta_j(y)] \text{ “Each } \beta_j \text{ holds along } (x_{j-1}, x_j)\text{”} \\ & \wedge (\forall y)_{>x_n} \beta_{n+1}(y) \text{ “}\beta_{n+1} \text{ holds everywhere after } x_n\text{”} \\ & \wedge (\forall y)^{<x_0} \beta_0(y) \text{ “}\beta_0 \text{ holds everywhere before } x_0\text{”} \end{aligned}$$

with a prefix of $n+1$ existential quantifiers and with all α_j, β_j quantifier free formulas with one variable over Σ , and $i_0, \dots, i_m \in \{0, \dots, n\}$. (ψ has $m+1$ free variables z_0, \dots, z_m and $n+1$ existential quantifiers, $m+1$ quantifiers are dummy and are introduced just in order to simplify notations.)

It is clear that

Lemma 3.2.

- (1) *Conjunction of $\vec{\exists}\forall$ -formulas is equivalent to a disjunction of $\vec{\exists}\forall$ -formulas.*
- (2) *Every $\vec{\exists}\forall$ -formula is equivalent to a conjunction of $\vec{\exists}\forall$ -formulas with at most two free variables.*
- (3) *For every $\vec{\exists}\forall$ -formula φ the formula $\exists x\varphi$ is equivalent to a $\vec{\exists}\forall$ -formula.*

Definition 3.3 ($\vee\vec{\exists}\forall$ -formulas). A formula is a $\vee\vec{\exists}\forall$ formula if it is equivalent to a disjunction of $\vec{\exists}\forall$ -formulas.

Lemma 3.4 (closure properties). *The set of $\vee \exists \forall$ formulas is closed under disjunction, conjunction, and existential quantification.*

Proof. By (1) and (3) of Lemma 3.2, and distributivity of \exists over \vee . \square

The set of $\vee \exists \forall$ formulas is not closed under negation². However, we show later (see Proposition 4.3) that the negation of a $\vee \exists \forall$ formula is equivalent to a $\vee \exists \forall$ formula in the expansion of the chains by all $TL(\text{Until}, \text{Since})$ definable predicates.

The $\vee \exists \forall$ formulas with one free variable can be easily translated to $TL(\text{Until}, \text{Since})$.

Proposition 3.5 (From $\vee \exists \forall$ -formulas to $TL(\text{Until}, \text{Since})$ formulas). *Every $\vee \exists \forall$ -formula with one free variable is equivalent to a $TL(\text{Until}, \text{Since})$ formula.*

Proof. By a simple formalization we show that every $\exists \forall$ -formula with one free variable is equivalent to a $TL(\text{Until}, \text{Since})$ formula. This immediately implies the proposition.

Let $\psi(z_0)$ be an $\exists \forall$ -formula

$$\begin{aligned} \exists x_n \dots \exists x_1 \exists x_0 \ z_0 = x_k \wedge (x_n > x_{n-1} > \dots > x_1 > x_0) \wedge \bigwedge_{j=0}^n \alpha_j(x_j) \\ \wedge \bigwedge_{j=1}^n (\forall y)_{>x_{j-1}}^{<x_j} \beta_j(y) \wedge (\forall y)_{>x_0}^{<x_0} \beta_0(y) \wedge (\forall y)_{>x_n} \beta_{n+1}(y) \end{aligned}$$

Let A_i and B_i be temporal formulas equivalent to α_i and β_i (A_i and B_i do not even use Until and Since modalities). It is easy to see that ψ is equivalent to the conjunction of

$$A_k \wedge (B_{k+1} \text{Until}(A_{k+1} \wedge (B_{k+2} \text{Until} \dots (A_{n-1} \wedge (B_n \text{Until}(A_n \wedge \Box B_{n+1}))) \dots)))$$

and

$$A_k \wedge (B_{k-1} \text{Since}(A_{k-1} \wedge (B_{k-2} \text{Since}(\dots A_1 \wedge (B_1 \text{Since}(A_0 \wedge \Box B_0)) \dots)))$$

4. PROOF OF KAMP'S THEOREM

The next definition plays a major role in the proof Kamp's theorem [3].

Definition 4.1. Let \mathcal{M} be a Σ chain. We denote by $\mathcal{E}[\Sigma]$ the set of unary predicate names $\Sigma \cup \{A \mid A \text{ is an } TL(\text{Until}, \text{Since})\text{-formula over } \Sigma\}$. The canonical $TL(\text{Until}, \text{Since})$ -expansion of \mathcal{M} is an expansion of \mathcal{M} to an $\mathcal{E}[\Sigma]$ -chain, where each predicate name $A \in \mathcal{E}[\Sigma]$ is interpreted as $\{a \in \mathcal{M} \mid \mathcal{M}, a \models A\}$ ³. We say that first-order formulas in the signature $\mathcal{E}[\Sigma] \cup \{<\}$ are equivalent over \mathcal{M} (respectively, over a class of Σ -chains \mathcal{C}) if they are equivalent in the canonical expansion of \mathcal{M} (in the canonical expansion of every $\mathcal{M} \in \mathcal{C}$).

²The truth table of $P\text{Until}Q$ is an $\exists \forall$ formula $(\exists x')_{>x}(Q(x') \wedge (\forall y)_{>x}^{<x'} P(y))$, yet we can prove that its negation is not equivalent to any $\vee \exists \forall$ formula.

³We often use “ $a \in \mathcal{M}$ ” instead of “ a is an element of the domain of \mathcal{M} ”

Note that if A is a $TL(\text{Until}, \text{Since})$ formula over $\mathcal{E}[\Sigma]$ predicates, then it is equivalent to a $TL(\text{Until}, \text{Since})$ formula over Σ , and hence to an atomic formula in the canonical $TL(\text{Until}, \text{Since})$ -expansions.

In this section and the next one we say that “formulas are equivalent in a chain \mathcal{M} ” instead of “formulas are equivalent in the canonical $TL(\text{Until}, \text{Since})$ -expansion of \mathcal{M} .” The $\vec{\exists}\forall$ and $\forall\vec{\exists}\forall$ formulas are defined as previously, but now they can use as atoms $TL(\text{Until}, \text{Since})$ definable predicates.

It is clear that all the results stated above hold for this modified notion of $\forall\vec{\exists}\forall$ formulas. In particular, every $\forall\vec{\exists}\forall$ formula with one free variable is equivalent to an $TL(\text{Until}, \text{Since})$ formula, and the set of $\forall\vec{\exists}\forall$ formulas is closed under conjunction, disjunction and existential quantification. However, now the set of $\forall\vec{\exists}\forall$ formulas is also closed under negation, due to the next proposition whose proof is postponed to Sect. 5.

Proposition 4.2. *(Closure under negation) The negation of $\vec{\exists}\forall$ -formulas with at most two free variables is equivalent over Dedekind complete chains to a disjunction of $\vec{\exists}\forall$ -formulas.*

As a consequence we obtain

Proposition 4.3. *Every first-order formula is equivalent over Dedekind complete chains to a disjunction of $\vec{\exists}\forall$ -formulas.*

Proof. We proceed by structural induction.

Atomic: It is clear that every atomic formula is equivalent to a disjunction of (even quantifier free) $\vec{\exists}\forall$ -formulas.

Disjunction: - immediate.

Negation: If φ is an $\vec{\exists}\forall$ -formula, then by Lemma 3.2(2) it is equivalent to a conjunction of $\vec{\exists}\forall$ formulas with at most two free variables. Hence, $\neg\varphi$ is equivalent to a disjunction of $\neg\psi_i$ where ψ_i are $\vec{\exists}\forall$ -formulas with at most two free variables. By Proposition 4.2, $\neg\psi_i$ is equivalent to a disjunction of $\vec{\exists}\forall$ formulas γ_i^j . Hence, $\neg\varphi$ is equivalent to a disjunction $\vee_i \vee_j \gamma_i^j$ of $\vec{\exists}\forall$ formulas.

If φ is a disjunction of $\vec{\exists}\forall$ formulas φ_i , then $\neg\varphi$ is equivalent to the conjunction of $\neg\varphi_i$. By the above, $\neg\varphi_i$ is equivalent to a $\forall\vec{\exists}\forall$ formula. Since, $\forall\vec{\exists}\forall$ formulas are closed under conjunction (Lemma 3.4), we obtain that $\neg\varphi$ is equivalent to a disjunction of $\vec{\exists}\forall$ formulas.

\exists -quantifier: For \exists -quantifier, the claim follows from Lemma 3.4. \square

Now, we are ready to prove Kamp’s Theorem:

Theorem 4.4. *For every FOMLO formula $\varphi(x)$ with one free variable, a $TL(\text{Until}, \text{Since})$ formula exists that is equivalent to φ over Dedekind complete chains.*

Proof. By Proposition 4.3, $\varphi(x)$ is equivalent over Dedekind complete chains to a disjunction of $\vec{\exists}\forall$ formulas $\varphi_i(x)$. By Proposition 3.5, $\varphi_i(x)$ is equivalent to a $TL(\text{Until}, \text{Since})$ formula. Hence, $\varphi(x)$ is equivalent over Dedekind complete chains to a $TL(\text{Until}, \text{Since})$ formula. \square

This completes our proof of Kamp’s theorem except Proposition 4.2 which is proved in the next section.

5. PROOF OF PROPOSITION 4.2

Let $\psi(z_0, z_1)$ be an $\vec{\exists}\forall$ -formula

$$\begin{aligned} \exists x_n \dots \exists x_1 \exists x_0 [z_0 = x_m \wedge z_1 = x_k \wedge (x_0 < x_1 < \dots < x_{n-1} < x_n) \wedge \bigwedge_{j=0}^n \alpha_j(x_j) \\ \wedge \bigwedge_{j=1}^n (\forall y)_{>x_{j-1}}^{<x_j} \beta_j(y) \wedge (\forall y)_{<x_0} \beta_0(y) \wedge (\forall y)_{>x_n} \beta_{n+1}(y)] \end{aligned}$$

We consider two cases. In the first case $k = m$, i.e., $z_0 = z_1$ and in the second $k \neq m$.

If $k = m$, then ψ is equivalent to $z_0 = z_1 \wedge \psi'(z_0)$, where ψ' is an $\vec{\exists}\forall$ -formula. By Proposition 3.5, ψ' is equivalent to an $TL(\text{Until}, \text{Since})$ formula A' . Therefore, ψ is equivalent to an $\vec{\exists}\forall$ -formula $\exists x_0 [z_0 = x_0 \wedge z_1 = x_0 \wedge A'(x_0)]$, and $\neg\psi$ is equivalent to a $\forall\vec{\exists}\forall$ formula $z_0 < z_1 \vee z_1 < z_0 \vee \exists x_0 [z_0 = x_0 \wedge z_1 = x_0 \wedge \neg A'(x_0)]$.

If $k \neq m$, w.l.o.g. we assume that $m < k$. Hence, ψ is equivalent to a conjunction of

(1) $\psi_0(z_0)$ defined as:

$$\begin{aligned} \exists x_0 \dots \exists x_{m-1} \exists x_m [z_0 = x_m \wedge (x_0 < x_1 < \dots < x_m) \wedge \bigwedge_{j=0}^m \alpha_j(x_j) \\ \wedge \bigwedge_{j=1}^m (\forall y)_{>x_{j-1}}^{<x_j} \beta_j(y) \wedge (\forall y)_{<x_0} \beta_0(y)] \end{aligned}$$

(2) $\psi_1(z_1)$ defined as:

$$\begin{aligned} \exists x_k \dots \exists x_{k+1} \exists x_n [z_1 = x_k \wedge (x_k < x_{k+1} < \dots < x_n) \wedge \bigwedge_{j=k}^n \alpha_j(x_j) \\ \wedge \bigwedge_{j=k+1}^n (\forall y)_{>x_{j-1}}^{<x_j} \beta_j(y) \wedge (\forall y)_{>x_n} \beta_{n+1}(y)] \end{aligned}$$

(3) $\varphi(z_0, z_1)$ defined as:

$$\begin{aligned} \exists x_m \dots \exists x_k [(z_0 = x_m < x_{m+1} < \dots < x_k = z_1) \wedge \bigwedge_{j=m}^k \alpha_j(x_j) \\ \wedge \bigwedge_{j=m+1}^k (\forall y)_{>x_{j-1}}^{<x_j} \beta_j(y)] \end{aligned}$$

The first two formulas are $\vec{\exists}\forall$ -formulas with one free variable. Therefore, (by Proposition 3.5) they are equivalent to $TL(\text{Until}, \text{Since})$ formulas (in the signature $\mathcal{E}[\Sigma]$). Hence, their negations are equivalent (over the canonical expansions) to atomic (and hence to $\vec{\exists}\forall$) formulas.

Therefore, it is sufficient to show that the negation of the third formula is equivalent over Dedekind complete chains to a disjunction of $\vec{\exists}\forall$ -formulas. This is stated in the following lemma:

Lemma 5.1. *The negation of any formula of the form*

$$\exists x_0 \dots \exists x_n [(z_0 = x_0 < \dots < x_n = z_1) \wedge \bigwedge_{j=0}^n \alpha_j(x_j) \wedge \bigwedge_{j=1}^n (\forall y)_{>x_{j-1}}^{<x_j} \beta_j(y)] \quad (5.1)$$

where α_i, β_i are quantifier free, is equivalent (over Dedekind complete chains) to a disjunction of $\vec{\exists}\forall$ -formulas.

In the rest of this section we prove Lemma 5.1. Our proof is organized as follows. In Lemma 5.3 we prove an instance of Lemma 5.1 where α_0, α_n and all β_i are equivalent to True. Then we derive a more general instance (Corollary 5.4) where β_n is equivalent to true. Finally we prove the full version of Lemma 5.1.

First, we introduce some helpful notations.

Notation 5.2. We use the abbreviated notation $[\alpha_0, \beta_1, \dots, \alpha_{n-1}, \beta_n, \alpha_n](z_0, z_1)$ for the $\vec{\exists}\forall$ -formula as in (5.1).

In this notation Lemma 5.1 can be rephrased as $\neg[\alpha_0, \beta_1, \dots, \alpha_{n-1}, \beta_n, \alpha_n](z_0, z_1)$ is equivalent (over Dedekind complete chains) to a $\vec{\exists}\forall$ formula.

We start with the instance of Lemma 5.1 where all β_i are True.

Lemma 5.3. $\neg\exists x_1 \dots \exists x_n (z_0 < x_1 < \dots < x_n < z_1) \wedge \bigwedge_{i=1}^n P_i(x_i)$ is equivalent over Dedekind complete chains to a $\vec{\exists}\forall$ formula $O_n(P_1, \dots, P_n, z_0, z_1)$.

Proof. We proceed by induction on n .

Basis: $\neg(\exists x_1)_{>z_0}^{<z_1} P_1(x_1)$ is equivalent to $(\forall y)_{>z_0}^{<z_1} \neg P_1(y)$.

Inductive step: $n \mapsto n+1$. We assume that a $\vec{\exists}\forall$ formula O_n has already defined and construct a $\vec{\exists}\forall$ formula O_{n+1} .

Observe that if the interval (z_0, z_1) is non-empty, then one of the following cases holds:

Case 1: P_1 does not occur in (z_0, z_1) , i.e. $(\forall y)_{>z_0}^{<z_1} \neg P_1(y)$. Then $O_{n+1}(P_1, \dots, P_{n+1}, z_0, z_1)$ should be equivalent to True.

Case 2: If case 1 does not hold then let $r_0 = \inf\{z \in (z_0, z_1) \mid P_1(z)\}$ (such r_0 exists by Dedekind completeness. Note that $r_0 = z_0$ iff $\mathbf{K}^+(P_1)(z_0)$. If $r_0 > z_0$ then $r_0 \in (z_0, z_1)$ and r_0 is definable by the following $\vec{\exists}\forall$ formula:

$$\begin{aligned} INF(z_0, r_0, z_1, P_1) := & z_0 < r_0 < z_1 \wedge (\forall y)_{>z_0}^{<r_0} \neg P_1(y) \wedge \\ & \wedge (P_1(r_0) \vee \mathbf{K}^+(P_1)(r_0)) \end{aligned} \quad (5.2)$$

Subcase $r_0 = z_0$: In this subcase $O_n(P_2, \dots, P_n, z_0, z_1)$ and $O_{n+1}(P_1, \dots, P_{n+1}, z_0, z_1)$ should be equivalent.

Subcase $r_0 \in (z_0, z_1)$: Now $O_n(P_2, \dots, P_n, r_0, z_1)$ and $O_{n+1}(P_1, \dots, P_{n+1}, z_0, z_1)$ should be equivalent.

Hence, $O_{n+1}(P_1, \dots, P_{n+1}, z_0, z_1)$ can be defined as the disjunction of “ (z_0, z_1) is empty” and the following formulas:

- (1) $(\forall y)_{>z_0}^{<z_1} \neg P_1(y)$
- (2) $\mathbf{K}^+(P_1)(z_0) \wedge O_n(P_2, \dots, P_n, z_0, z_1)$
- (3) $(\exists r_0)_{>z_0}^{<z_1} (INF(z_0, r_0, z_1, P_1) \wedge O_n(P_2, \dots, P_n, r_0, z_1))$

The first formula is a $\vec{\exists}\forall$ formula. By the inductive assumptions O_n is a $\vec{\exists}\forall$ formula. $\mathbf{K}^+(P_1)(z_0)$ is an atomic (and hence a $\vec{\exists}\forall$) formula in the canonical expansion, and $INF(z_0, r_0, z_1, P_1)$ is a $\vec{\exists}\forall$ formula. Since $\vec{\exists}\forall$ formulas are closed under conjunction, disjunction and the existential quantification, we conclude that O_{n+1} is a $\vec{\exists}\forall$ formula. \square

As a consequence we obtain

Corollary 5.4.

- (1) $\neg(\exists z)_{>z_0}^{<z_1}[\alpha_0, \beta_1, \alpha_1, \beta_2, \dots, \alpha_{n-1}, \beta_n, \alpha_n](z_0, z)$ over Dedekind complete chains is equivalent to a $\bigvee \overrightarrow{\exists} \forall$ formula.
- (2) $\neg(\exists z)_{>z_0}^{<z_1}[\alpha_0, \beta_1, \alpha_1, \beta_2, \dots, \alpha_{n-1}, \beta_n, \alpha_n](z, z_1)$ over Dedekind complete chains is equivalent to a $\bigvee \overrightarrow{\exists} \forall$ formula.

Proof. (1) Define

$$F_n := \alpha_n$$

$$F_{i-1} := \alpha_{i-1} \wedge (\beta_i \text{Until} F_i) \quad \text{for } i = 1, \dots, n$$

Observe that there is $z \in (z_0, z_1)$ such that $[\alpha_0, \beta_1, \alpha_1, \beta_2, \dots, \alpha_{n-1}, \beta_n, \alpha_n](z_0, z)$ iff $F_0(z_0)$ and there is an increasing sequence $x_1 < \dots < x_n$ in an open interval (z_0, z_1) such that $F_i(x_i)$ for $i = 1, \dots, n$. Indeed, the direction \Rightarrow is trivial. The direction \Leftarrow is easily proved by induction.

The *basis* is trivial.

Inductive step: $n \mapsto n+1$. Assume $F_0(z_0)$ holds and that (z_0, z_1) contains an increasing sequence $x_1 < \dots < x_{n+1}$ such that $F_i(x_i)$ for $i = 1, \dots, n+1$. By the inductive assumption there is $y_1 \in (z_0, x_{n+1})$ such that

$$[\alpha_0, \beta_1, \alpha_1, \beta_2, \dots, \beta_{n-1} \alpha_{n-1}, \beta_n, (\alpha_n \wedge \beta_{n+1} \text{Until} \alpha_{n+1})](z_0, y_1).$$

In particular, y_1 satisfies $(\alpha_n \wedge \beta_{n+1} \text{Until} \alpha_{n+1})$. Hence, there is $y_2 > y_1$ such that y_2 satisfies α_{n+1} and β_{n+1} holds along (y_1, y_2) .

If $y_2 \leq x_{n+1}$ then the required $z \in (z_0, z_1)$ equals to y_2 , and we are done. Otherwise, $x_{n+1} < y_2$. Therefore, $x_{n+1} \in (y_1, y_2)$ and β_{n+1} holds along (y_1, x_{n+1}) . Hence, the required z equals to x_{n+1} .

The above observation and Lemma 5.3 imply that $\neg F_0(z_0) \vee O_n(F_1, \dots, F_n, z_0, z_1)$ is a $\bigvee \overrightarrow{\exists} \forall$ formula that is equivalent to $\neg(\exists z)_{>z_0}^{<z_1}[\alpha_0, \beta_1, \alpha_1, \beta_2, \dots, \alpha_{n-1}, \beta_n, \alpha_n](z_0, z)$.

(2) is the mirror image of (1) and is proved similarly. \square

Now we are ready to prove Lemma 5.1, i.e.,

$$\neg[\alpha_0, \beta_1 \dots, \beta_{n-1}, \alpha_{n-1}, \beta_n, \alpha_n](z_0, z_1) \text{ is equivalent}$$

$$\text{over Dedekind complete chains to a } \bigvee \overrightarrow{\exists} \forall \text{ formula.}$$

Proof. (of Lemma 5.1) If the interval (z_0, z_1) is empty then the assertion is immediate. We assume that (z_0, z_1) is non-empty. Hence, at least one of the following cases holds:

Case 1: $\neg\alpha_0(z_0)$ or $\mathbf{K}^+(\neg\beta_1)(z_0)$.

Case 2: $\alpha_0(z_0)$, and β_1 holds along (z_0, z_1) .

Case 3: (1) $\alpha_0(z_0) \wedge \neg\mathbf{K}^+(\neg\beta_1)(z_0)$, and

(2) there is $x \in (z_0, z_1)$ such that $\neg\beta_1(x)$.

For each of these cases we construct a $\bigvee \overrightarrow{\exists} \forall$ formula $Cond_i$ that describes it (i.e., Case i holds iff $Cond_i$ holds) and show that if $Cond_i$ holds, then $\neg[\alpha_0, \beta_1 \dots, \beta_{n-1}, \alpha_{n-1}, \beta_n, \alpha_n](z_0, z_1)$ is equivalent to a $\bigvee \overrightarrow{\exists} \forall$ formula $Form_i$. Hence, $\neg[\alpha_0, \beta_1 \dots, \beta_{n-1}, \alpha_{n-1}, \beta_n, \alpha_n](z_0, z_1)$ is equivalent to $\bigvee_i [Cond_i \wedge Form_i]$ which is a $\bigvee \overrightarrow{\exists} \forall$ formula.

Case 1 This case is already explicitly described by the $\bigvee \overrightarrow{\exists} \forall$ formula (in the canonical expansion). In this case $\neg[\alpha_0, \beta_1 \dots, \beta_{n-1}, \alpha_{n-1}, \beta_n, \alpha_n](z_0, z_1)$ is equivalent to True.




FIGURE 1. $B_2(z_0, z, z_1) := [\alpha_0, \beta_1, \alpha_1, \beta_2, \beta_2](z_0, z) \wedge [\beta_2, \beta_2, \alpha_2, \beta_3, \alpha_3](z, z_1)$

Case 2 This case is described by a $\bigvee \overrightarrow{\exists} \forall$ formula $\alpha_0(z_0) \wedge (\forall z)_{>z_0}^{<z_1} \beta_1$. In this case $\neg[\alpha_0, \beta_1, \dots, \beta_{n-1}, \alpha_{n-1}, \beta_n, \alpha_n](z_0, z_1)$ is equivalent to “there is no $z \in (z_0, z_1)$ such that $[\alpha_1, \beta_2, \dots, \beta_n, \alpha_n](z, z_1)$.” By Corollary 5.4(2) this is expressible by a $\bigvee \overrightarrow{\exists} \forall$ formula.

Case 3 The first condition of Case 3 is already explicitly described by a $\bigvee \overrightarrow{\exists} \forall$ formula. When the first condition holds, then the second condition is equivalent to “there is (a unique) $r_0 \in (z_0, z_1)$ such that $r_0 = \inf\{z \in (z_0, z_1) \mid \neg\beta_1(z)\}$ ” (If $\neg\mathbf{K}^+(\neg\beta_1)$ holds at z_0 and there is $x \in (z_0, z_1)$ such that $\neg\beta_1(x)$, then such r_0 exists because we deal with Dedekind complete chains.) This r_0 is definable by the following $\bigvee \overrightarrow{\exists} \forall$ formula, i.e., it is a unique z which satisfies it⁴:

$$INF^{\neg\beta_1}(z_0, z, z_1) := z_0 < z < z_1 \wedge (\forall y)_{>z_0}^{<z} \beta_1(y) \wedge (\neg\beta_1(z) \vee \mathbf{K}^+(\neg\beta_1)(z)) \quad (5.3)$$

Hence, Case 3 is described by $\alpha_0(z_0) \wedge \neg\mathbf{K}^+(\neg\beta_1)(z_0) \wedge (\exists z)_{>z_0}^{<z_1} INF^{\neg\beta_1}(z_0, z, z_1)$ which is equivalent to an $\overrightarrow{\exists} \forall$ formula.

It is sufficient to show that $(\exists z)_{>z_0}^{<z_1} INF^{\neg\beta_1}(z) \wedge \neg[\alpha_0, \beta_1, \alpha_1, \dots, \beta_{n+1}, \alpha_{n+1}](z_0, z_1)$ is equivalent to a $\bigvee \overrightarrow{\exists} \forall$ formula.

We prove this by induction on n .

The *basis* is trivial.

Inductive step $n \mapsto n + 1$.

Define:

$$\begin{aligned} A_i^-(z_0, z) &:= [\alpha_0, \beta_1, \dots, \beta_i, \alpha_i](z_0, z) & i = 1, \dots, n \\ A_i^+(z, z_1) &:= [\alpha_i, \beta_{i+1}, \dots, \beta_{n+1}, \alpha_{n+1}](z, z_1) & i = 1, \dots, n \\ A_i(z_0, z, z_1) &:= A_i^-(z_0, z) \wedge A_i^+(z, z_1) & i = 1, \dots, n \\ B_i^-(z_0, z) &:= [\alpha_0, \beta_1, \dots, \beta_{i-1}, \alpha_{i-1}, \beta_i, \beta_i](z_0, z) & i = 1, \dots, n + 1 \\ B_i^+(z, z_1) &:= [\beta_i, \beta_i, \alpha_i \beta_{i+1} \alpha_{i+1}, \dots, \beta_{n+1}, \alpha_{n+1}](z, z_1) & i = 1, \dots, n + 1 \\ B_i(z_0, z, z_1) &:= B_i^-(z_0, z) \wedge B_i^+(z, z_1) & i = 1, \dots, n + 1 \end{aligned}$$

⁴We will use only existence and will not use uniqueness.

If the interval (z_0, z_1) is non-empty, these definitions imply

$$\begin{aligned} [\alpha_0, \beta_1, \alpha_1, \dots, \beta_{n+1}, \alpha_{n+1}](z_0, z_1) &\Leftrightarrow (\forall z)_{>z_0}^{\leq z_1} \left(\bigvee_{i=1}^n A_i \vee \bigvee_{i=1}^{n+1} B_i \right) \\ [\alpha_0, \beta_1, \alpha_1, \dots, \beta_{n+1}, \alpha_{n+1}](z_0, z_1) &\Leftrightarrow (\exists z)_{>z_0}^{\leq z_1} \left(\bigvee_{i=1}^n A_i \vee \bigvee_{i=1}^{n+1} B_i \right) \end{aligned}$$

Hence, for every φ

$$(\exists z)_{>z_0}^{\leq z_1} \varphi(z) \wedge \neg[\alpha_0, \beta_1, \alpha_1, \dots, \beta_{n+1}, \alpha_{n+1}](z_0, z_1)$$

is equivalent to

$$(\exists z)_{>z_0}^{\leq z_1} \left(\varphi(z) \wedge \bigwedge_{i=1}^n \neg A_i \wedge \bigwedge_{i=1}^{n+1} \neg B_i \right)$$

In particular,

$$(\exists z)_{>z_0}^{\leq z_1} INF^{\neg\beta_1}(z) \wedge \neg[\alpha_0, \beta_1, \alpha_1, \dots, \beta_{n+1}, \alpha_{n+1}](z_0, z_1)$$

is equivalent to

$$(\exists z)_{>z_0}^{\leq z_1} \left(INF^{\neg\beta_1}(z) \wedge \bigwedge_{i=1}^n \neg A_i \wedge \bigwedge_{i=1}^{n+1} \neg B_i \right),$$

where $INF^{\neg\beta_1}(z)$ was defined in equation (5.3).

By the inductive assumption

(a): $\neg A_i$ is equivalent to a $\bigvee \vec{\exists} \forall$ formula for $i = 1, \dots, n$.

(b): $\neg B_i$ is equivalent to a $\bigvee \vec{\exists} \forall$ formula for $i = 2, \dots, n$.

Recall $B_1 := B_1^- \wedge B_1^+$ and $B_{n+1} := B_{n+1}^- \wedge B_{n+1}^+$.

(c): $\neg B_1^-$ and $\neg B_{n+1}^+$ are equivalent to $\bigvee \vec{\exists} \forall$ formulas, by the induction basis.

(d): $INF^{\neg\beta_1}(z) \wedge \neg B_1^+(z, z_1)$ is equivalent to $INF^{\neg\beta_1}(z)$, because if $INF^{\neg\beta_1}(z)$, then for no $x > z$, β_1 holds along $[z, x]$.

(e): $INF^{\neg\beta_1}(z) \wedge \neg B_{n+1}^-(z_0, z)$ is equivalent to $INF^{\neg\beta_1}(z) \wedge (\text{"}\beta_1 \text{ holds on } (z_0, z)\text{"} \wedge \neg B_{n+1}^-(z_0, z))$. Since, by case 2, $\text{"}\beta_1 \text{ holds on } (z_0, z)\text{"} \wedge \neg B_{n+1}^-(z_0, z)$ is equivalent to a $\bigvee \vec{\exists} \forall$ formula, and $INF^{\neg\beta_1}(z)$ is a $\bigvee \vec{\exists} \forall$ formula, we conclude that $INF^{\neg\beta_1}(z) \wedge \neg B_{n+1}^-(z_0, z)$ is equivalent to a $\bigvee \vec{\exists} \forall$ formula.

Since the set of $\bigvee \vec{\exists} \forall$ formulas is closed under conjunction, disjunction and \exists , by (a)-(e) we obtain that $(\exists z)_{>z_0}^{\leq z_1} (INF^{\neg\beta_1}(z) \wedge \bigwedge_{i=1}^n \neg A_i \wedge \bigwedge_{i=1}^{n+1} \neg B_i)$ is equivalent to a $\bigvee \vec{\exists} \forall$ formula.

Therefore, $(\exists z)_{>z_0}^{\leq z_1} INF^{\neg\beta_1}(z) \wedge \neg[\alpha_0, \beta_1, \alpha_1, \dots, \beta_{n+1}, \alpha_{n+1}](z_0, z_1)$ is also a $\bigvee \vec{\exists} \forall$ formula.

This completes our proof of Lemma 5.1 and of Proposition 4.2. \square

6. RELATED WORKS

Kamp's theorem was proved in

- (1) Kamp's thesis [8] (proof > 100pages).
- (2) Outlined by Gabbay, Pnueli, Shelah and Stavi [3] (Sect. 2) for \mathbb{N} and stated that it can be extended to Dedekind complete orders using game arguments.
- (3) Was proved by Gabbay [1] by separation arguments for \mathbb{N} , and extended to Dedekind complete order in [2].
- (4) Was proved by Hodkinson [5] by game arguments and simplified in [6] (unpublished).

A temporal logic has the *separation* property if its formulas can be equivalently rewritten as a boolean combination of formulas, each of which depends only on the past, present or future. The separation property was introduced by Gabbay [1], and surprisingly, a temporal logic which can express \Box and \Box has the separation property (over a class \mathcal{C} of structures) iff it is expressively complete for *FOMLO* over \mathcal{C} .

The separation proof for $TL(\text{Until}, \text{Since})$ over \mathbb{N} is manageable; however, over the real (and over Dedekind complete) chains it contains many rules and transformations and is not easy to follow. Hodkinson and Reynolds [7] write:

The proofs of theorems 18 and 19 [Kamp's theorem over naturals and over reals, respectively] are direct, showing that each formula can be separated. They are tough and tougher, respectively. Nonetheless, they are effective, and so, whilst not quite providing an algorithm to determine if a set of connectives is expressively complete, they do suggest a potential way of telling in practice whether a given set of connectives is expressively complete – in Gabbay's words, *try to separate and see where you get stuck!*

The game arguments are easier to grasp, but they use complicated inductive assertions. The proof in [6] proceeds roughly as follows. Let \mathcal{L}_r be the set of $TL(\text{Until}, \text{Since})$ formulas of nesting depth at most r . A formula of the form: $\exists \bar{x} \forall y \chi(\bar{x}, y, \bar{z})$ where \bar{x} is an n -tuple of variables and χ is a quantifier free formula over $\{<, =\}$ and \mathcal{L}_r -definable monadic predicates is called $\langle n, r \rangle$ -decomposition formula. The main inductive assertion is proved by “unusual back-and-forth games” and can be rephrased in logical terms as there is a function $f : \mathbb{N} \rightarrow \mathbb{N}$ such that for every $n, r \in \mathbb{N}$, the negation of positive Boolean combinations $\langle n, r \rangle$ -decomposition formula is equivalent to a positive Boolean combination of $\langle f(n), (n + r) \rangle$ -decomposition formulas.

Our proof is inspired by [3] and [6]; however, it avoids games, and it separates general logical equivalences and temporal arguments.

The temporal logic with the modalities *Until* and *Since* is not expressively complete for *FOMLO* over the rationals. Stavi introduced two additional modalities *Until^s* and *Since^s* and proved that $TL(\text{Until}, \text{Since}, \text{Until}^s, \text{Since}^s)$ is expressively complete for *FOMLO* over all linear orders [2]. In the forthcoming paper we prove Stavi's theorem. The proof is similar to our proof of Kamp's theorem; however, it treats some additional cases related to gaps in orders, and replaces $\exists \forall$ -formulas by slightly more general formulas.

7. FUTURE FRAGMENT OF *FOMLO*

Many temporal formalisms studied in computer science deal only with future formulas, whose truth value at any moment is determined by what happens from a current moment on.

A formula (temporal, or monadic with a single free first-order variable) F is (*semantically*) *future* if for every chain \mathcal{M} and moment $t_0 \in \mathcal{M}$:

$$\mathcal{M}, t_0 \models F \text{ iff } \mathcal{M}|_{\geq t_0}, t_0 \models F,$$

where $\mathcal{M}|_{\geq t_0}$ is the subchain of \mathcal{M} over the interval $[t_0, \infty)$. For example, $P\text{Until}Q$ and $\mathbf{K}^+(P)$ are future formulas, while $P\text{Since}Q$ and $\mathbf{K}^-(P)$ are not future ones.

For a set B of modalities we denote by $TL(B)$ the temporal logic which uses only modalities from B . In particular, $TL(\text{Until})$ is the temporal logic which uses the modality *Until* and $TL(\text{Until}, \mathbf{K}^-)$ is the temporal logic with modalities *Until* and \mathbf{K}^- .

It was shown in [3] that Kamp's theorem holds also for *future formulas* of *FOMLO* over $\omega = \langle \mathbb{N}, < \rangle$:

Theorem 7.1 (Gabbay, Pnueli, Shelah, Stavi [3]). *Every future FOMLO formula is equivalent over ω -chains to a $TL(\text{Until})$ formula.*

The situation is radically different for the continuous time $\langle \mathbb{R}, < \rangle$. In [4] it was shown that $TL(\text{Until})$ is not expressively complete for the future fragment of *FOMLO* and there is no easy way to remedy it. In fact, it was shown in [4] that there is no temporal logic with a finite set of modalities which is expressively equivalent to the future fragment of *FOMLO* over the Reals.

From the separation proof of Kamp's theorem in [2] it follows that every future *FOMLO* formula is equivalent over Dedekind complete chains to a $TL(\text{Until}, \mathbf{K}^-)$ formula.

This future-past mixture of *Until* and \mathbf{K}^- is somewhat better than the standard *Until* - *Since* basis in the following sense: although \mathbf{K}^- is (like *Since*) a past modality, it does not depend on much of the past. The formula $\mathbf{K}^-(P)$ depends just on an arbitrarily short 'near past', and is actually independent of most of the past. In this sense, we may say that it is an "almost" future formula.

Definition 7.2 (Syntactically future $TL(\text{Until}, \mathbf{K}^-)$ formulas). A $TL(\text{Until}, \mathbf{K}^-)$ formula is syntactically future if it is a boolean combination of atomic formulas and formulas of the form $\varphi_1 \text{Until} \varphi_2$, where φ_1 and φ_2 are arbitrary $TL(\text{Until}, \mathbf{K}^-)$ formulas.

The following lemma immediately follows from the definition and the observation that $\mathcal{M}|_{\geq t_0}, t_0 \models \neg \mathbf{K}^-(\varphi)$.

Lemma 7.3. *A syntactically future $TL(\text{Until}, \mathbf{K}^-)$ formula is future. A $TL(\text{Until}, \mathbf{K}^-)$ formula is future iff it is equivalent to a syntactically future $TL(\text{Until}, \mathbf{K}^-)$ formula.*

The next theorem (implicitly) appears in [2] (Chapter 8).

Theorem 7.4. *Every future FOMLO formula is equivalent over Dedekind complete chains to a syntactically future $TL(\text{Until}, \mathbf{K}^-)$ formula.*

Since $\mathbf{K}^-\varphi$ is equivalent to *False* over discrete orders, we obtain that Theorems 7.1 is an instance of Theorem 7.4.

Theorem 7.4 is easily obtained by a slight refinement of our proof of Kamp's theorem. We outline its proof in the rest of this section.

Definition 7.5 ((z_0, z_1) - $\vec{\exists}\forall$ formula). Let z_0 and z_1 be two variables. A formula $z_0 > z_1$, $z_0 = z_1$ or of the form $[\alpha_0, \beta_1 \dots, \beta_{n-1}, \alpha_{n-1}, \beta_n, \alpha_n](z_0, z_1)$ is called a (z_0, z_1) - $\vec{\exists}\forall$ formula. A formula is (z_0, z_1) - $\vee \vec{\exists}\forall$ formula if it is equivalent to a disjunction of (z_0, z_1) - $\vec{\exists}\forall$ formulas.

Lemma 7.6 (closure properties). *The set of $(z_0, z_1)\text{-}\bigvee \overrightarrow{\exists} \forall$ formulas is closed under disjunction and conjunction. If φ_1 is a $(z_0, z_1)\text{-}\bigvee \overrightarrow{\exists} \forall$ formula and φ_2 is a $(z_1, z_2)\text{-}\bigvee \overrightarrow{\exists} \forall$ formula, then $(\exists z_1)_{> z_0}^{< z_2} (\varphi_1 \wedge \varphi_2)$ is a $(z_0, z_2)\text{-}\bigvee \overrightarrow{\exists} \forall$ formula.*

The set of $(z_0, z_1)\text{-}\bigvee \overrightarrow{\exists} \forall$ formulas is not closed under negation. However, we show that the negation of a $\bigvee \overrightarrow{\exists} \forall$ formula is equivalent to a $(z_0, z_1)\text{-}\bigvee \overrightarrow{\exists} \forall$ formula in the expansion of the chains by all $TL(Until, \mathbf{K}^-)$ definable predicates.

Definition 7.7 (The canonical $TL(Until, \mathbf{K}^-)$ and $TL(Since, \mathbf{K}^+)$ expansions). Let \mathcal{M} be a Σ chain. We denote by $\mathcal{E}[\Sigma, TL(Until, \mathbf{K}^-)]$ the set of unary predicate names $\Sigma \cup \{A \mid A \text{ is an } TL(Until, \mathbf{K}^-) \text{ formula over } \Sigma\}$. The canonical $TL(Until, \mathbf{K}^-)$ -expansion of \mathcal{M} is an expansion of \mathcal{M} to an $\mathcal{E}[\Sigma, TL(Until, \mathbf{K}^-)]$ -chain, where each predicate's name $A \in \mathcal{E}[\Sigma, TL(Until, \mathbf{K}^-)]$ is interpreted as $\{a \in \mathcal{M} \mid \mathcal{M}, a \models A\}$. We say that first-order formulas in the signature $\mathcal{E}[\Sigma, TL(Until, \mathbf{K}^-)] \cup \{<\}$ are equivalent over \mathcal{M} (respectively, over a class of Σ -chains \mathcal{C}) if they are equivalent in the canonical $TL(Until, \mathbf{K}^-)$ -expansion of \mathcal{M} (in the canonical $TL(Until, \mathbf{K}^-)$ -expansion of every $\mathcal{M} \in \mathcal{C}$). The canonical $TL(Since, \mathbf{K}^+)$ -expansion of a chain \mathcal{M} is defined similarly.

The next lemma implies that Lemma 5.1 holds over the canonical $TL(Since, \mathbf{K}^+)$ -expansions and over canonical $TL(Until, \mathbf{K}^-)$ -expansions.

Lemma 7.8.

- (1) $\neg[\alpha_0, \beta_1 \dots, \beta_{n-1}, \alpha_{n-1}, \beta_n, \alpha_n](z_0, z_1)$ is equivalent over the canonical $TL(Since, \mathbf{K}^+)$ -expansions of Dedekind complete chains to a $(z_0, z_1)\text{-}\bigvee \overrightarrow{\exists} \forall$ -formula.
- (2) Dually, $\neg[\alpha_0, \beta_1 \dots, \beta_{n-1}, \alpha_{n-1}, \beta_n, \alpha_n](z_0, z_1)$ is equivalent over the canonical $TL(Until, \mathbf{K}^-)$ -expansions of Dedekind complete chains to a $(z_0, z_1)\text{-}\bigvee \overrightarrow{\exists} \forall$ -formula.

Proof. Actually, our proof of Lemma 5.1, as it is, works for the canonical $TL(Since, \mathbf{K}^+)$ -expansions of Dedekind complete chains, when “ $\overrightarrow{\exists} \forall$ formulas” are replaced by “ $(z_0, z_1)\text{-}\overrightarrow{\exists} \forall$ formulas.”

Indeed, Lemma 5.3 uses only modality \mathbf{K}^+ . Thus, exactly the same proof works for $TL(Since, \mathbf{K}^+)$ -expansions and for $TL(Until, \mathbf{K}^-)$ -expansions (because \mathbf{K}^+ is equivalent to a $TL(Until)$ formula).

In the proof of Corollary 5.4(1) we used Lemma 5.3 and *Until* modality. Hence, it holds for $TL(Until, \mathbf{K}^-)$ -expansions. Corollary 5.4(2) is dual and it holds for $TL(Since, \mathbf{K}^+)$ -expansions.

In proof of Lemma 5.1 we use standard logical equivalences and Corollary 5.4(2). Hence, it works as it is for the canonical $TL(Since, \mathbf{K}^+)$ -expansions of Dedekind complete chains. This proves (1). Item (2) is the mirror image of (1). \square

Notation 7.9. We use the abbreviated notation $[\alpha_0, \beta_1 \dots, \alpha_{n-1}, \beta_n, \alpha_n, \beta_{n+1}](z_0, \infty)$ for

$$\begin{aligned} & \exists x_n \dots \exists x_1 \exists x_0 z_0 = x_0 \wedge (x_n > \dots > x_1 > x_0) \\ & \wedge \bigwedge_{j=0}^n \alpha_j(x_j) \wedge \bigwedge_{j=1}^n (\forall y)_{> x_{j-1}}^{< x_j} \beta_j(y) \wedge (\forall y)_{> x_n} \beta_{n+1}(y); \end{aligned}$$

such formulas will be called (z_0, ∞) -formulas; we use the similarly abbreviated notation $[\beta_0, \alpha_0, \beta_1 \dots, \alpha_{n-1}, \beta_n, \alpha_n](-\infty, z_0)$ for the $\vec{\exists}\forall$ -formula

$$\begin{aligned} \exists x_n \dots \exists x_1 \exists x_0 z_0 = x_n \wedge (x_n > \dots > x_1 > x_0) \\ \wedge \bigwedge_{j=0}^n \alpha_j(x_j) \wedge \bigwedge_{j=1}^n (\forall y)_{>x_{j-1}}^{<x_j} \beta_j(y) \wedge (\forall y)^{<x_0} \beta_0(y). \end{aligned}$$

Lemma 7.10. $[\alpha_0, \beta_1 \dots, \alpha_{n-1}, \beta_n, \alpha_n, \beta_{n+1}](z_0, \infty)$ over canonical $TL(\text{Until}, \mathbf{K}^-)$ -expansions is equivalent to a $TL(\text{Until}, \mathbf{K}^-)$ formula.

Proof. By a straightforward formalization as in the proof of Proposition 3.5. \square

Definition 7.11 (Syntactically future FOMLO formulas). A FOMLO formula $\varphi(z_0)$ is syntactically future if all its quantifiers are bounded quantifiers of the form $(\forall y)_{>z_0}$ and $(\exists y)_{>z_0}$.

The following lemma immediately follows from the definition.

Lemma 7.12. A syntactically future FOMLO formula is future. A FOMLO formula $\varphi(z_0)$ is future iff it is equivalent to a syntactically future FOMLO formula.

Definition 7.13. Let (z_0, z_1, \dots, z_k) be a sequence of distinct variables. A formula is $(z_0, z_1, \dots, z_k, \infty)$ - $\vec{\exists}\forall$ formula if it is a conjunction $\bigwedge_{i \leq k} \varphi_i$, where φ_k is (z_k, ∞) - $\vec{\exists}\forall$ formula and φ_i is (z_i, z_{i+1}) - $\vec{\exists}\forall$ formulas for $i < k$. A formula is a $(z_0, z_1, \dots, z_k, \infty)$ - $\vee \vec{\exists}\forall$ formula if it is equivalent to a disjunction of $(z_0, z_1, \dots, z_k, \infty)$ - $\vec{\exists}\forall$ formulas.

Lemma 7.14. Let $\varphi(z_0, z_1, \dots, z_k)$ be a FOMLO formula with free variables in $\{z_i \mid i \leq k\}$ and all its quantifiers are bounded quantifiers of the form $(\forall y)_{>z_0}$ and $(\exists y)_{>z_0}$. Then, there is $(z_0, z_1, \dots, z_k, \infty)$ - $\vee \vec{\exists}\forall$ formula ψ such that $z_0 < z_1 < \dots < z_k \wedge \varphi$ is equivalent over the canonical $TL(\text{Until}, \mathbf{K}^-)$ -expansions of Dedekind complete chains to $z_0 < z_1 < \dots < z_k \wedge \psi$.

Proof. By Lemmas 7.6, 7.8(2), 7.10 and a straightforward structural induction. \square

Now we are ready to prove Theorem 7.4.

Proof. (of Theorem 7.4) Assume that $\varphi(z_0)$ is a future FOMLO formula. By Lemma 7.12 w.l.o.g we can assume that all its quantifiers are bounded quantifiers of the form $(\forall y)_{>z_0}$ and $(\exists y)_{>z_0}$. By 7.14, it is equivalent to (z_0, ∞) - $\vee \vec{\exists}\forall$ formula. Hence, by Lemma 7.10, it is equivalent to a $TL(\text{Until}, \mathbf{K}^-)$ formula. Therefore, by Lemma 7.3 it is equivalent to a syntactically future $TL(\text{Until}, \mathbf{K}^-)$ formula. \square

In [9] we erroneously stated that the analog of Proposition 4.3 holds for $TL(\text{Until}, \mathbf{K}^-)$ -expansions. However, “ P is unbounded from below” is expressible by a FOMLO sentence $\forall x \exists y (y < x \wedge P(y))$; yet there is no $\vee \vec{\exists}\forall$ formula which expresses “ P is unbounded from below” over the canonical $TL(\text{Until}, \mathbf{K}^-)$ -expansions of integer-chains. We state the following Proposition for the sake of completeness.

Proposition 7.15. Every FOMLO formula is equivalent over the canonical $TL(\text{Until}, \mathbf{K}^-)$ -expansions of Dedekind complete chains to a positive boolean combination of $\vec{\exists}\forall$ formulas and sentences of the form “ P is unbounded from below.”

The additional step needed for the proof of Proposition 7.15 is an observation that $\neg[\beta_0, \alpha_0, \beta_1 \dots, \alpha_{n-1}, \beta_n, \alpha_n](-\infty, z_0)$ is equivalent over the canonical $TL(\text{Until}, \mathbf{K}^-)$ -expansions of Dedekind complete chains to a positive boolean combinations of $(-\infty, z_0) \rightarrow \exists \forall$ formulas and sentences of the form “ P is unbounded from below.” This is proved almost in the same way as Lemma 5.1.

Acknowledgments I am very grateful to Yoram Hirshfeld for numerous insightful discussions, and to the anonymous referees for their helpful suggestions.

REFERENCES

- [1] D. Gabbay. Expressive functional completeness in tense logic (preliminary report). In U. Monnich, editor, *Aspects of Philosophical Logic*, pages 91-117. Reidel, Dordrecht, 1981.
- [2] D. Gabbay, I. Hodkinson, and M. Reynolds. *Temporal logic: Mathematical Foundations and Computational Aspects*. Oxford University Press, 1994.
- [3] D. Gabbay, A. Pnueli, S. Shelah and J. Stavi. On the Temporal Analysis of Fairness. In *POPL 1980*, pp. 163-173, 1980.
- [4] Y. Hirshfeld, A. Rabinovich. Future temporal logic needs infinitely many modalities. *Information and Computation*, 187(2003), 196-208.
- [5] I. Hodkinson. Expressive completeness of Until and Since over dedekind complete linear time. *Modal logic and process algebra*, ed. A. Ponse, M. de Rijke, Y. Venema, *CSLI Lecture Notes 53*, 1995.
- [6] I. Hodkinson. Notes on games in temporal logic. *Lecture notes for LUATCS meeting, Johannesburg, Dec 1999*. <http://www.doc.ic.ac.uk/~imh/index.html>
- [7] I. Hodkinson and M. Reynolds. Temporal Logic Chapter 11 (pp. 655-720) in *Handbook of Modal Logic*, Patrick Blackburn, Johan van Benthem, and Frank Wolter, eds., Elsevier Science, 2006.
- [8] H. W. Kamp. Tense logic and the theory of linear order. *Phd thesis, University of California, Los Angeles*, 1968.
- [9] D. Pardo and A. Rabinovich: A Finite Basis for ‘Almost Future’ Temporal Logic over the Reals. *MFCS 2012*: 740-751
- [10] A. Pnueli (1977). The temporal logic of programs. In *Proc. IEEE 18th Annu. Symp. on Found. Comput. Sci.*, pages 46-57, New York, 1977.