

Estimating Riparian ET through Remote sensing

By

Zohrab Samani, Salim Bawazir

Max Bleiweiss, Rhonda Skaggs


Thomas Schmugge

New Mexico State University

Rio Grande River Spring, 2003

Water is scarce in New Mexico






Legend

- Road, Primary
- Road, Secondary
- Road, Unimproved
- ~ Canal, River, etc.
-  Wx Tower
- ▲ Research Tower

- 1.... North Saltcedar Research Tower
- 2.... South Saltcedar Research Tower
- 3.... South Cottonwood Research Tower
- 4.... South Bosque Weather Station

106° 54' 38"
33° 48' 36"

106° 50' 47"
33° 48' 39"



106° 54' 35"
33° 46' 11"


106° 50' 44"
33° 46' 14"



Rio Grande Riparian Vegetation




ColorIR Satellite Image of Bosque del Apache NWR





Energy Balance for ET

ET is calculated as a "residual" of the energy balance



Ground ET Measurements



Eddy Covariance

$$H = \rho c_p COV[wT]$$

$$LE = \lambda COV[wq]$$



Why Satellites?

$$ET = K_c \cdot ET_o$$




- Traditional method for ET:
 - ET_o is calculated from weather data (weather station)
 - K_c assume “well-watered” situation (*impacts of stress are difficult to quantify*)
 - K_c for most riparian vegetation is **not available**

Net Radiation

Net Surface Radiation = Gains – Losses

$$R_n = (1-\alpha)R_{S\downarrow} + R_{L\downarrow} - R_{L\uparrow} - (1-\varepsilon_o)R_{L\downarrow}$$

Albedo, NDVI, LST



How It Works

Parameters from Satellite:


- Albedo
- Vegetation indices (NDVI)
- Surface temperature
- Wind speed (*from ground station*)

Satellite


- Satellite Images are available from:
 - NASA-Landsat (30 m, every 16 days)
- since 1982. Landsat 7 went dead in May 2003.
 - NOAA-AVHRR (*advanced very high resolution radiometer*)
(1 km, daily) - since 1980's
 - NASA-MODIS (*moderate resolution imaging spectroradiometer*), *daily*, (250 m, for NDVI & ALBEDO, but 1 Km for Temp. - since 1999
 - NASA-ASTER (*Terra, Advanced Spaceborne Thermal Emission and Reflection Radiometer*)- since 1999

Predicting Net Radiation


Rn, SaltCedar, Bosque 2003



Predicting Net Radiation



Estimating Ground Flux, G_n




Sensible Heat is calculate at anchor points

$$ET = R_n - G - H$$

$$H = R_n - ET - G$$

$$H = \frac{\rho \cdot C_p \cdot dT}{rah} + \text{Monin-Obukhov Similarity}$$


Temperature Gradient Versus Surface Temperature




Calculating Evaporative fraction, E_f

$$E_f = \frac{R_n - G - H}{R_n - G}$$


North Saltcedar Tower, Bosque NWR




South Saltcedar Tower, Bosque NWR



South Cottonwood Tower, Bosque, 2003



ET NSCT-2003
Bosque del Apache



Regional ET, Bosque, 2003

