```
0.00
Created by Ye Cheng (Lewis) Chen and Jesko Sirker
The interacting disordered SSH model
Calculation for Loschmidt echo using SPH and formula |det(1-C+C*exp(-iHt))|^2
import numpy as np
import argparse
from bisect import bisect left as findIndex
# parameters for entanglement calculations
parser = argparse.ArgumentParser(description=
'Calculation of Loschmidt echo for disordered fermions')
parser.add argument('-L', type=int, default=4,
                   help='# system length L')
parser.add_argument('-V', type=float, default=1.0,
                   help='V interaction term')
parser.add argument('-W', type=float, default=0.0,
                   help='width box potential disorder')
parser.add argument('-delta', type=float, default=-0.5,
                   help='# quench parameter delta')
parser.add_argument('-tint', type=float, default=0,
                   help='# initial time')
parser.add_argument('-tmax', type=float, default=20,
                   help='# maximum time range')
parser.add_argument('-dt', type=float, default=0.01,
                   help='# discrete time interval')
parser.add argument('-sample', type=int, default=1,
                   help='# samples')
args=parser.parse args()
# State function configurations in spinless fermions
def manyPsi(particle, site):
   a=np.arange(2**site)
   bitcount=np.array([bin(x).count("1") for x in a])
   b=a.compress(bitcount==particle).tolist()[::-1]
   aList=[]
    for item in b:
       a=bin(int(item))[2:].zfill(site)
       aList.append(a)
    return aList
# C+C fermionic operators on Neel configurations
def cpc(l,j,Psi):
   Psi2=Psi.copy()
    for item in Psi2:
       k = Psi2.index(item)
       if item[j-1]=='0':
           Psi2[k]=list(item)
           Psi2[k][j-1]='1'
           Psi2[k]="".join(Psi2[k])
```

if item[j-1]=='1':

Psi2[k]=list(item) Psi2[k][j-1]='0'