

L9.slides

2020年4月2日 星期四 下午5:38



Lecture9
print

MATH 661

Lecture 9
Chapter 5, 6

Instructor: Padma Natarajan

Interpreting confidence interval

- In practice, we obtain only one random sample and calculate one confidence interval.
- We then say that the observed interval brackets the true value of μ with confidence $100(1-\alpha)\%$.

3/29/2020

Common Levels of Confidence

If the level of confidence is 0.90, this means that we are 90% confident that the interval contains the population mean, μ .

$z_{\alpha/2} = 1.645$, $z_{0.05} = 1.96$, $z_{0.025} = 2.05$

The corresponding t values are 1.645, 1.98, 2.075.

Example

A random sample of 32 textbook prices is taken from a local college bookstore. The sample mean \bar{x} is \$20.75. Use a 95% confidence level and find the margin of error for the mean price of all textbooks in the bookstore.

$\bar{x} = 20.75$, $n = 32$, $\sigma = 2.00$

$z_{\alpha/2} = 1.96$, $t_{0.025} = 2.05$

$M = z_{\alpha/2} \frac{\sigma}{\sqrt{n}} = 1.96 \cdot \frac{2.00}{\sqrt{32}} = 1.20$

20.75 ± 1.20 include true value.

5

Common Levels of Confidence

If the level of confidence is 0.95, this means that we are 95% confident that the interval contains the population mean, μ .

$z_{\alpha/2} = 1.96$, $t_{0.025} = 2.05$

$t = 0.95$, $t = 1 - 0.95 = 0.05$, $\sigma^2 = 0.052 = 0.025$

$z_{\alpha/2} (t_{0.025}) = 1.96$

6

Example

A random sample of 32 textbook prices is taken from a local college bookstore. The sample mean \bar{x} is \$20.75. Use a 95% confidence level and find the margin of error for the mean price of all textbooks in the bookstore.

$\bar{x} = 20.75$, $n = 32$, $\sigma = 2.00$

$t = 0.95$, $t = 1 - 0.95 = 0.05$

$M = z_{\alpha/2} \frac{\sigma}{\sqrt{n}} = 1.96 \cdot \frac{2.00}{\sqrt{32}} = 1.20$

20.75 ± 1.20

1

6x Z_{0.05} for 95% and 99% CI. (Formative)

$z_{0.05} = 1.645$
 $\alpha = 0.05$
 $\frac{1}{2} = 0.025$
 $z_{0.025} = 2(1 - 0.025) = 2(0.975) = 1.96$

$z_{0.01} = 2.326$
 $\alpha = 0.01$
 $\frac{1}{2} = 0.005$

$z_{0.005} = 2(1 - 0.005) = 2(0.995) = 2.57$
 2.57

2

5x Z_{0.025} for 95% and 99% CI. (Formative)

$z_{0.025} = 1.96$
 $\alpha = 0.025$
 $\frac{1}{2} = 0.0125$
 $z_{0.0125} = 2(1 - 0.0125) = 2(0.9875) = 2.326$
 2.326

3

5x Z_{0.005} for 95% and 99% CI. (Formative)

$z_{0.005} = 2.57$
 $\alpha = 0.005$
 $\frac{1}{2} = 0.0025$
 $z_{0.0025} = 2(1 - 0.0025) = 2(0.9975) = 2.673$
 2.673

4

5x Z_{0.01} for 95% and 99% CI. (Formative)

$z_{0.01} = 2.326$
 $\alpha = 0.01$
 $\frac{1}{2} = 0.005$
 $z_{0.005} = 2(1 - 0.005) = 2(0.995) = 2.57$
 2.57

5

5x Z_{0.001} for 95% and 99% CI. (Formative)

$z_{0.001} = 2.673$
 $\alpha = 0.001$
 $\frac{1}{2} = 0.0005$
 $z_{0.0005} = 2(1 - 0.0005) = 2(0.9995) = 2.673$
 2.673

6

upwind work week quiz

Example 1

The age at diagnosis of hypertension in a random sample of patients with hypertension is listed below:

32.8	40	41	42	45.5	47	48.5	50	51
52	54	59.2	61.2	62.5	64.5	65.5	67.5	69

Assume that age at diagnosis is approximately normally distributed with a standard deviation of 7.2 years.

Proportion

Obtain 95% CI for the true age, μ , at diagnosis.

$\bar{x} = 51.96$, $n = 9$

$M = z_{\alpha/2} \frac{\sigma}{\sqrt{n}} = 1.96 \cdot \frac{7.2}{\sqrt{9}} = 1.96 \cdot 2.4 = 4.704$

51.96 ± 4.704

Confidence level and Precision of estimation

The length of a CI = $2z_{\alpha/2} \frac{\sigma}{\sqrt{n}}$

$\frac{1}{2} = 0.05$

• It is desirable to obtain a confidence interval that is short enough for decision-making purposes and that also has an adequate confidence.

• One way to achieve this is by choosing a sample size n to be large enough to give a CI of specified length.

9

Solution

$z(2z_{\alpha/2}) = 1.96$

$\bar{x} = \frac{\sum x_i}{n} = \frac{563}{12} = 46.91$

$46.91 \pm 1.96 \cdot \frac{7.2}{\sqrt{12}} = 46.91 \pm 4.1$

We are 95% confident that the mean age at diagnosis of hypertension is between 42.8 and 51.0 years. The point estimate is 46.9 years, with a margin of error of 4.1 years.

8

choose Sample size.

• The appropriate sample size can be found by choosing n such that $z_{\alpha/2} \frac{\sigma}{\sqrt{n}} = m$

• Solving this equation gives the formula for n

$n = \frac{(z_{\alpha/2})^2}{m^2}$

If the right hand side of the equation is not an integer, it must be rounded up. This will ensure that the level of confidence does not fall below 95% (i.e. the length of the interval).

10

Example

$z(\frac{1}{2}) = 0.86$

What sample size should be selected to estimate the mean age of the tails of the scorpion to within 0.5 years at the 95% confidence level if the standard deviation for the tails is 1.0?

Solution

We are given that $(1-\alpha) = 0.95$, $\bar{x}_0 = 1.96$, $\sigma = 3.5$, and $m = 1$.

Substituting into the formula, we get

$n = \frac{(z_{\alpha/2})^2}{m^2} = \frac{(0.86)^2}{1^2} = 47.596 \approx 48$

11

Example

You want to estimate the mean price of all the textbooks in the college bookstore. How many books should be included in your sample to be 95% confident that the sample mean is within 10% of the true mean?

$\sigma = 2.575$

$n = \frac{(z_{\alpha/2})^2}{m^2} = \frac{(1.96)^2}{0.1^2} = 384.144 \approx 385$

You should include at least 385 books in your sample.

12

Solution

$z(\frac{1}{2}) = 0.86$

$z(\frac{1}{2}) = 0.86 = (\frac{1.96 \cdot 3.5}{m})^2$

$= (1.96 \cdot 3.5)^2 / 0.86^2 = 1.96^2 \cdot 3.5^2 / 0.86^2 = 67.0546 \approx 67$

2

Instructor: Padma Natarajan

Example

Suppose we want to estimate the proportion of patients in a particular physician's practice with osteoarthritis. In a random sample of 200 patients, 38 are diagnosed with osteoarthritis.

• Find a 95% confidence interval on mean tire life.

$\hat{p} = 0.19$, $n = 200$

$\hat{p} \pm z_{\alpha/2} \sqrt{\hat{p}(1-\hat{p})/n} = 0.19 \pm 1.96 \sqrt{0.19(1-0.19)/200} = 0.19 \pm 0.038 = [0.152, 0.228]$

Using Minitab to find t-value

• In Minitab, choose Graph > Probability Plot.

• Select View Probability, then click OK.

• From Distribution, select t.

• In Degrees of freedom, enter 19.

• Click Shaded Area. Both Tails.

• Input α .

3

Solution

The appropriate confidence interval for μ is given by

$\bar{x} \pm z_{\alpha/2} \frac{\sigma}{\sqrt{n}}$

From Table D for 95% confidence $t = 2.131$

$106 + 2(1.96) \cdot \frac{12.4}{\sqrt{106}} = 106 + 2(1.96) \cdot 1.24 = 116.33$

We are 95% confident that the mean IQ score for all 22 year olds is between 99.4 and 112.6.

13

Example

Determine t that is required to construct each of the following confidence intervals:

- confidence level=95%, degrees of freedom=12
- confidence level=95%, sample size=25
- confidence level=95%, degrees of freedom=13
- confidence level=99%, sample size=16

$t_{0.05/2,11} = 2.179$, $t_{0.05/2,24} = 2.064$, $t_{0.05/2,12} = 3.012$, $t_{0.05/2,15} = 4.073$

Example

Suppose we want to estimate the mean systolic blood pressure of members of a health maintenance organization (HMO) who are 25 years of age.

Solution

A random sample of 100 HMO members is chosen and the mean systolic blood pressure of the population is found to be 120 with a standard deviation of 15.

$t = 2.227$

$120 \pm 2.227 \cdot \frac{15}{\sqrt{100}} = 120 \pm 3.34 = [116.66, 133.34]$

3/29/2020

Example

Suppose we wish to estimate the proportion of patients in a particular physician's practice with osteoarthritis.

• In a random sample of 200 patients, 38 are diagnosed with osteoarthritis.

• Compute a 95% confidence interval for the proportion of all patients in this physician's practice with osteoarthritis.

$\hat{p} = 0.19$, $n = 200$

$\hat{p} \pm z_{\alpha/2} \sqrt{\hat{p}(1-\hat{p})/n} = 0.19 \pm 1.96 \sqrt{0.19(1-0.19)/200} = 0.19 \pm 0.038 = [0.152, 0.228]$

Choice of sample size

If we set $m = z_{\alpha/2} \sqrt{\hat{p}(1-\hat{p})/n}$ and solve for n , we can obtain an approx. sample size.

$n = \frac{z_{\alpha/2}^2 \cdot \hat{p}(1-\hat{p})}{m^2}$

$n = 0.19^2 \cdot 0.19(1-0.19) / 0.038^2 = 106$

$n = 0.19^2 \cdot 0.19(1-0.19) / 0.038^2 = 106$

We are 95% confident that the mean IQ score among 22 year olds is between 99.4 and 112.6.

31

Example

Suppose we want to estimate the proportion of patients in a particular physician's practice with osteoarthritis.

• In a random sample of 200 patients, 38 are diagnosed with osteoarthritis.

• Compute a 95% confidence interval for the proportion of all patients in this physician's practice with osteoarthritis.

$\hat{p} = 0.19$, $n = 200$

$\hat{p} \pm z_{\alpha/2} \sqrt{\hat{p}(1-\hat{p})/n} = 0.19 \pm 1.96 \sqrt{0.19(1-0.19)/200} = 0.19 \pm 0.038 = [0.152, 0.228]$

Solution

The appropriate confidence interval for μ is given by

$\bar{x} \pm z_{\alpha/2} \frac{\sigma}{\sqrt{n}}$

From Table D for 95% confidence $t = 2.131$

$106 + 2(1.96) \cdot \frac{12.4}{\sqrt{106}} = 106 + 2(1.96) \cdot 1.24 = 116.33$

We are 95% confident that the mean IQ score for all 22 year olds is between 99.4 and 112.6.

32

Example

Suppose we want to estimate the true proportion of high school seniors who smoke, how many objects would be required to ensure that a margin of error of 0.05 with 95% confidence?

Solution

Based on a pilot study involving a random sample of 1000 high school seniors, they reported that they were smokers.

$\hat{p} = 0.19$, $n = 1000$

$n = \frac{z_{\alpha/2}^2 \cdot \hat{p}(1-\hat{p})}{m^2} = \frac{1.96^2 \cdot 0.19(1-0.19)}{0.05^2} = 384.144 \approx 385$

33

Example

Suppose we want to estimate the true proportion of high school seniors who smoke, how many objects would be required to ensure that a margin of error of 0.05 with 95% confidence?

Solution

Based on a pilot study involving a random sample of 1000 high school seniors, they reported that they were smokers.

$\hat{p} = 0.19$, $n = 1000$

$n = \frac{z_{\alpha/2}^2 \cdot \hat{p}(1-\hat{p})}{m^2} = \frac{1.96^2 \cdot 0.19(1-0.19)}{0.05^2} = 384.144 \approx 385$

34

Example

Suppose we want to estimate the true proportion of high school seniors who smoke, how many objects would be required to ensure that a margin of error of 0.05 with 95% confidence?

Solution

Based on a pilot study involving a random sample of 1000 high school seniors, they reported that they were smokers.

$\hat{p} = 0.19$, $n = 1000$

$n = \frac{z_{\alpha/2}^2 \cdot \hat{p}(1-\hat{p})}{m^2} = \frac{1.96^2 \cdot 0.19(1-0.19)}{0.05^2} = 384.144 \approx 385$

35

Example

An investigator wishes to estimate the proportion of patients in a new physician's office diagnosed with osteoarthritis. He plans to use a new policy and wants the estimate to be within 0.05 of the true proportion of all patients with osteoarthritis with 95% confidence.

$\hat{p} = 0.19$, $n = 100$

$n = \frac{z_{\alpha/2}^2 \cdot \hat{p}(1-\hat{p})}{m^2} = \frac{1.96^2 \cdot 0.19(1-0.19)}{0.05^2} = 384.144 \approx 385$

36

Example

Suppose we want to estimate the true proportion of high school seniors who smoke, how many objects would be required to ensure that a margin of error of 0.05 with 95% confidence?

Solution

Based on a pilot study involving a random sample of 1000 high school seniors, they reported that they were smokers.

$\hat{p} = 0.19$, $n = 1000$

$n = \frac{z_{\alpha/2}^2 \cdot \hat{p}(1-\hat{p})}{m^2} = \frac{1.96^2 \cdot 0.19(1-0.19)}{0.05^2} = 384.144 \approx 385$

37

Example

An investigator wishes to estimate the proportion of patients in a new physician's office diagnosed with osteoarthritis. He plans to use a new policy and wants the estimate to be within 0.05 of the true proportion of all patients with osteoarthritis with 95% confidence.

$\hat{p} = 0.19$, $n = 100$

$n = \frac{z_{\alpha/2}^2 \cdot \hat{p}(1-\hat{p})}{m^2} = \frac{1.96^2 \cdot 0.19(1-0.19)}{0.05^2} = 384.144 \approx 385$

38

Example

Suppose we want to estimate the true proportion of high school seniors who smoke, how many objects would be required to ensure that a margin of error of 0.05 with 95% confidence?

Solution

Based on a pilot study involving a random sample of 1000 high school seniors, they reported that they were smokers.

$\hat{p} = 0.19$, $n = 1000$

$n = \frac{z_{\alpha/2}^2 \cdot \hat{p}(1-\hat{p})}{m^2} = \frac{1.96^2 \cdot 0.19(1-0.19)}{0.05^2} = 384.144 \approx 385$

39

Example

An estimate for p is available from pilot study.

$\hat{p} = 0.19$, $n = 100$, $m = 1.96$, $t = 2.131$

$n = \hat{p}(1-\hat{p}) \left(\frac{z_{\alpha/2}}{m} \right)^2$

$n = 0.19(1-0.19) \left(\frac{2.131}{1.96} \right)^2 = 106$

In order to estimate the true proportion of high school seniors who smoke with a margin of error of 0.05, we need at least 106 high school seniors.

40

Example

Suppose we want to estimate the proportion of patients in a particular physician's practice with osteoarthritis.

• In a random sample of 200 patients, 38 are diagnosed with osteoarthritis.

• Compute a 95% confidence interval for the proportion of all patients in this physician's practice with osteoarthritis.

$\hat{p} = 0.19$, $n = 200$

$\hat{p} \pm z_{\alpha/2} \sqrt{\hat{p}(1-\hat{p})/n} = 0.19 \pm 1.96 \sqrt{0.19(1-0.19)/200} = 0.19 \pm 0.038 = [0.152, 0.228]$

Choice of sample size

If we set $m = z_{\alpha/2} \sqrt{\hat{p}(1-\hat{p})/n}$ and solve for n , we can obtain an approx. sample size.

$n = \frac{z_{\alpha/2}^2 \cdot \hat{p}(1-\hat{p})}{m^2}$

$n = 0.19^2 \cdot 0.19(1-0.19) / 0.038^2 = 106$

$n = 0.19^2 \cdot 0.19(1-0.19) / 0.038^2 = 106$

We are 95% confident that the mean IQ score among 22 year olds is between 99.4 and 112.6.

41

Example

Suppose we want to estimate the true proportion of patients in a particular physician's practice with osteoarthritis.

• In a random sample of 200 patients, 38 are diagnosed with osteoarthritis.

• Compute a 95% confidence interval for the proportion of all patients in this physician's practice with osteoarthritis.

$\hat{p} = 0.19$, $n = 200$

$\hat{p} \pm z_{\alpha/2} \sqrt{\hat{p}(1-\hat{p})/n} = 0.19 \pm 1.96 \sqrt{0.19(1-0.19)/200} = 0.19 \pm 0.038 = [0.152, 0.228]$

Solution

The appropriate confidence interval for μ is given by

$\bar{x} \pm z_{\alpha/2} \frac{\sigma}{\sqrt{n}}$

From Table D for 95% confidence $t = 2.131$

$106 + 2($