

Notes for Differential Galois Theory by P. Jossen

Texed by Lin-Da Xiao

October 27, 2017

Contents

1	Differential rings and modules	2
2	Picard-Vessiot extension	5
3	The Differential Galois Groups	10
4	Algebraic Group: A Detour	11
4.1	Application of the fact that differential Galois group is an algebraic group	14
5	The Galois-correspondence of differential equations	16
6	local theory of differential equations	17
7	Personal Reads	18
7.1	Monodromy, Riemann-Hilbert Problem	18

About the Course

Classical Galois theory:

- Fields \mathcal{K}
- Fields extensions \mathcal{K}'/\mathcal{K} .
- $Gal(\mathcal{K}'/\mathcal{K}) = Aut_{\mathcal{K}}(\mathcal{K}')$.
- Solution of polynomials, $\mathcal{K}' = \text{"}\mathcal{K}(\text{solutions of polynomials})\text{"}$, which is what we call **splitting field**.

In the Differential setting,

- Fields \mathcal{K} with derivation ∂ , e.g. $\mathbb{Q}(t)$, with usual derivative:
- Differential field extension \mathcal{K}'/\mathcal{K}
- $Gal(\mathcal{K}'/\mathcal{K}) = Aut_{(\mathcal{K}, \partial)}(\mathcal{K}', \partial)$
- Solution of differential equations. $\mathcal{K}' = \text{"}\mathcal{K}(\text{solution of differential equation})\text{"}$, which is called **Picard-Vessiot Field**

1 Differential rings and modules

Convention: Ring= commutative ring with unit.

Definition 1.1. Let \mathcal{R} be a ring. A **derivation** on \mathcal{R} is a map $\partial : \mathcal{R} \rightarrow \mathcal{R}$ s.t.

1. $\partial(a + b) = \partial a + \partial b$,
2. $\partial(ab) = a\partial b + b\partial a$.

then we call (\mathcal{R}, ∂) a differential ring. A morphism of diff-rings $\varphi : (\mathcal{R}_1, \partial_1) \rightarrow (\mathcal{R}_2, \partial_2)$ is a ring morphism s.t. $\partial_2 \varphi = \varphi \partial_1$

Example 1.2. $\mathbb{Q}(t), \mathcal{K}(t)$ with usual derivations and $\mathbb{Q}[t], \mathcal{K}[t], C^\infty([0, 1])$

Definition 1.3. Let (\mathcal{R}, ∂) be a diff-ring, Call $\mathcal{C} \subset \mathcal{R}$ **constant** if $\partial c = 0, \forall c \in \mathcal{C}$.

Proposition 1.4. Let $\mathcal{C} \subset \mathcal{R}$ be the set of constant elements.

1. \mathcal{C} is a subring,

2. If \mathcal{R} is a field, then so is \mathcal{C}

Proof. $1 \in \mathcal{C}$, $\partial 1 = \partial(1 \cdot 1) = 2\partial 1 \implies \partial 1 = 0$

$a, b \in \mathcal{C} \implies a + b \in \mathcal{C}$, and $ab \in \mathcal{C}$ by Leibnitz rule.

Suppose \mathcal{R} is a field, $c \in \mathcal{C}$, $c \neq 0$, $0 = \partial 1 = \partial(c \cdot c^{-1}) = c\partial(c^{-1}) \implies \partial c^{-1} = 0 \quad \square$

Example 1.5. *Caution:* $\mathcal{R} = \mathbb{F}_p[t]$, ∂ is the usual derivative. Here, constant $\mathcal{C} = \mathbb{F}_p[t^p] \supsetneq \mathbb{F}_p$, because

$$\partial(a_0 + a_1 t^p + a_2 t^{2p} + \dots + a_n t^{np}) = 0.$$

Exercise 1.6. Show that $\mathcal{C} \subseteq \mathcal{R}$ is a algebraically closed in \mathcal{R} . i.e. $x \in \mathcal{R}$ algebraic over $\mathcal{C} \implies x \in \mathcal{C}$. (Notice it does not mean $\mathcal{C} = \overline{\mathcal{C}}$ in general)

Exercise 1.7. Show that for differential field \mathcal{K} with constants \mathcal{C} , consider a field extension \mathcal{R}/\mathcal{K} , an element $x \in \mathcal{R}$ satisfying $x' = 0 \in \mathcal{K}$ is algebraic over $\mathcal{K} \implies x$ is algebraic over \mathcal{C} .

Solution:

x is algebraic over \mathcal{K} , consider the minimal monic polynomial $p(X) = X^n + \dots + a_0$ with coefficients in \mathcal{K} . Then $p(x) = 0 \implies p(x)' = (a'_{n-1})x^{n-1} + \dots + (a'_0) = 0$ by the minimality of $p(X)$, we conclude that each $a'_i = 0$, thus finished the proof.

Definition 1.8. A **differential (\mathcal{R}, ∂) -module** (M, ∂) is a \mathcal{R} -module M , together with $\partial : M \rightarrow M$ satisfying:

$$1. \partial(m + n) = \partial m + \partial n$$

$$2. \partial_M(am) = \partial_{\mathcal{R}} a \cdot m + a \cdot \partial_M m.$$

Think of (M, ∂) as a differential equation, with solutions $\ker(\partial : M \rightarrow M)$.

Suppose $\mathcal{R} = \mathcal{K}$ is a field (over that M is free), M has finite dimension. Choose a \mathcal{K} -basis, (e_1, \dots, e_n) of M . Set

$$\partial e_i = - \sum_{j=1}^n a_{ij} e_j,$$

where $A = (a_{ij}) \in M_{n \times n}(\mathcal{K})$. The matrix A characterizes $\partial : M \rightarrow M$, uniquely, by additivity and Leibnitz:

$$m \in M, m = \sum_{i=1}^n \lambda_i e_i,$$

$$\begin{aligned}\partial m &= \sum \partial(\lambda_i e_i) = \sum (\partial \lambda_i) e_i + \sum \lambda_i \partial e_i \\ &= \sum (\partial \lambda_i) e_i - \sum \sum \lambda_i a_{ij} e_j.\end{aligned}$$

The differential equation corresponding to (M, ∂) is the equation

$$u' = Au.$$

Remark 1.9. *the matrix A depends on the choice of the \mathcal{K} -basis of M . Choosing a different basis yields an equation $u' = \tilde{A}u$ with $\tilde{A} = S^{-1}AS - S^{-1}S'$, where $S \in GL_n(\mathcal{K})$ is the base change matrix. we called A, \tilde{A} equivalent*

Remark 1.10. *Let M be a differential \mathcal{K} -module, $\mathcal{C} \subset \mathcal{K}$ be the set of constants, then we have $\partial : M \rightarrow M$. is \mathcal{C} -linear, follows from Leibnitz. In particular $\ker \partial \subseteq M$ is a \mathcal{C} -module (vector space)*

Lemma 1.11. *Let $u' = Au$ be a differential equation with $A \in M_{n \times n}(\mathcal{K})$. Let $v_1, \dots, v_r \in \mathcal{K}^n$ be solutions, i.e. $v'_i = Av_i$. If v_1, \dots, v_r are linear dependent over \mathcal{K} , then they are linear dependent over \mathcal{C} . In particular,*

$$\dim_{\mathcal{C}}(\ker \partial) \leq n.$$

Proof. Induction on r . For $r = 1$, trivial. Fix $r \geq 2$, suppose lemma holds for $< r$ solutions. Suppose w.l.o.g that no proper subset of $\{v_1, \dots, v_r\}$ is linear dependent over \mathcal{K} . We find that there is a unique linear dependence relation

$$v_1 = \sum_{i=2}^r b_i v_i, \quad b_i \in \mathcal{K}$$

$$\begin{aligned}0 &= v'_1 - Av_1 = \sum b'_i v_i + b_i v'_i - \sum_{i=2}^r b_i Av_i \\ &= \sum_{i=2}^r b'_i v_i + \sum_{i=2}^r b_i (v'_i - Av_i) = \sum_{i=2}^r b'_i v_i\end{aligned}$$

so $b'_i = 0$ for $i = 2, \dots, r \implies b_i \in \mathcal{C}$, and v_1, \dots, v_r linear dependent over \mathcal{C} . \square

Compactify the notation, v_1, \dots, v_r columns of a matrix $V \in M_{n \times r}(\mathcal{K})$, then $v'_i = Av_i \implies V' = AV$. Know that $\text{rank}_{\mathcal{C}} V \leq n$. What we usually seek is a $V \in GL_n(\mathcal{K})$ with $V' = AV$. The columns of such V provide a basis of the solution space of the differential equation and thus also a basis of the differential module itself.

Definition 1.12. \mathcal{K} is a diff-field, $A \in M_{n \times n}(\mathcal{K})$, let \mathcal{R} be a differential \mathcal{K} -algebra, which means we have a diff-ring morphism from $(\mathcal{R}, \partial_{\mathcal{R}})$ to $(\mathcal{K}, \partial_{\mathcal{K}})$. We also suppose \mathcal{R} has same constants as \mathcal{K} (every constant of \mathcal{R} lies in \mathcal{K}). A matrix $V \in GL_n(\mathcal{R})$ is said to be a **fundamental matrix** of solutions of the differential equation $u' = Au$, if $V' = AV$.

Remark 1.13. Let V, \tilde{V} be fundamental matrices of solutions of $u' = Au$, $V, \tilde{V} \in GL_n(\mathcal{R})$, $\tilde{V} = V \cdot S$ $S = V^{-1}\tilde{V}$

$$A\tilde{V} = \tilde{V}' = (VS)' = V'S + VS' = AVS + VS' = A\tilde{V} + VS'$$

$$\implies VS' = 0, V \text{ is invertible} \implies S' = 0 \implies S \in GL_n(C).$$

Definition 1.14. Let $v_1, \dots, v_n \in \mathcal{K}$, The **Wronski matrix** of $\underline{v} \in \mathcal{K}^n$ is

$$Wr(\underline{v}) = \begin{pmatrix} v_1 & v_2 & \cdots & v_n \\ v_1^{(1)} & v_2^{(1)} & \cdots & v_n^{(1)} \\ \vdots & \vdots & \ddots & \vdots \\ v_1^{(n-1)} & v_2^{(n-1)} & \cdots & v_n^{(n-1)} \end{pmatrix}$$

the **Wronskian** is the determinant of the Wronski matrix, i.e. $\det(Wr(\underline{v})) =: wr(\underline{v})$.

2 Picard-Vessiot extension

Through out this section, we will assume \mathcal{K} is a differential field with $\text{char}(\mathcal{K}) = 0$ (It contains \mathbb{Q} as subfield). The set of constants $\mathcal{C} \subset \mathcal{K}$ is a field and we assume it to be algebraic closed. For example, think of $\mathcal{K} = \mathbb{C}(t)$, $\partial = d/dt$ and $\mathcal{C} = \mathbb{C}$.

Definition 2.1. Let \mathcal{R} be a diff. \mathcal{K} -algebra. An ideal $I \subset \mathcal{R}$ is a **differential ideal** if $\partial I \subseteq I$. Say that \mathcal{R} is **simple** if $\{0\}$ and \mathcal{R} are the only differential ideal in \mathcal{R}

Remark 2.2. $I \subseteq \mathcal{R}$ is differential ideal, then the derivation on \mathcal{R} induces a derivation on \mathcal{R}/I . Given any morphism of diff. rings $\varphi: \mathcal{R} \rightarrow \mathcal{R}'$, then $\text{Ker}(\varphi)$ is a differential ideal.

Definition 2.3. Let $A \in M_n(\mathcal{K})$, consider the matrix differential equation

$$u' = Au$$

A differential \mathcal{K} -algebra \mathcal{R} is said to be a **Picard-Vessiot extension** for $u' = Au$ if

1. \mathcal{R} is simple
2. The equation $u' = Au$ admits a fundamental matrix of solution in \mathcal{R} , i.e. $\exists V \in M_n(\mathcal{R})$, invertible such that

$$V' = AV.$$

3. As \mathcal{K} -algebra, \mathcal{R} is generated by the coefficients v_{ij} of V and $\det(V)^{-1}$.

Some references require in addition the constants of \mathcal{R} are \mathcal{C} . We will see this additional requirement can be derived by 1-3 in our setting.

A **Picard-Vessiot extension of a differential module** M is a Picard-Vessiot extension for any of the corresponding matrix differential equation. Exercise: check this.

Alternatively, we can define the Picard-Vessiot extension of a differential module M directly. Given a diff. module (M, ∂_M) , a Picard-Vessiot extension for (M, ∂_M) is a diff. \mathcal{K} -algebra \mathcal{R} s.t.

1. \mathcal{R} is simple
2. $\dim(\partial_{\mathcal{R} \otimes M}) = \dim_{\mathcal{K}} M$, where $\partial_{\mathcal{R} \otimes M} : \mathcal{R} \otimes M \longrightarrow \mathcal{R} \otimes M$, $\partial_{\mathcal{R} \otimes M}(r \otimes m) = r' \otimes m + r \otimes \partial_M m$.
3. \mathcal{R} is minimal with these properties.

Exercise: Check that the two definitions of PV extension for diff. module coincide.

Example 2.4. $\mathcal{K} = \mathbb{C}(t), \mathcal{C} = \mathbb{C}$, Consider the 2nd order homogeneous linear differential equation

$$t \cdot u'' + u' = 0.$$

We can set $v := u', v' = u''$, then we have a new 1st order

$$\begin{pmatrix} u' \\ v' \end{pmatrix} = \begin{pmatrix} 0 & 1 \\ 0 & -1/t \end{pmatrix} \begin{pmatrix} u \\ v \end{pmatrix},$$

where the matrix is called **companion matrix**. What we want is $\mathcal{R} \supseteq \mathbb{C}(t), V \in GL_2(\mathcal{R})$ s.t. $V' = AV$. The general solution of the 2nd order equation is

$$a + b \log(t).$$

Solution to $(u', v')^T = A \cdot (u, v)^T$ are

$$\begin{pmatrix} 1 \\ 0 \end{pmatrix}, \begin{pmatrix} \log(t) \\ 1/t \end{pmatrix},$$

The corresponding fundamental matrix

$$V = \begin{pmatrix} 1 & \log(t) \\ 0 & 1/t \end{pmatrix}.$$

A candidate of the Picard-Vessiot ring is

$$\mathcal{R} = \mathbb{C}(t)[X]$$

the Differential on \mathcal{R} , we just need to set $\partial X = X' = 1/t$, in this case, we don't have to adjoint $\det(V)^{-1}$. The only thing left to check is whether \mathcal{R} is simple.

Yes, $I \subseteq \mathcal{R}$ diff. ideal. \mathcal{R} principal $I = f(X)\mathcal{R}$, $\partial I \subset I$, where $\partial I = (\partial f)\mathcal{R}$, derive it sufficiently many times. until $\partial^n f \in \mathbb{C}(t) \implies \mathcal{R} = \partial^n f \mathcal{R} \subseteq I$.

Lemma 2.5. *Let \mathcal{R} be simple diff. \mathcal{K} -algebra. Then*

1. \mathcal{R} is integral domain
2. Suppose \mathcal{R} is finitely generated as \mathcal{K} -algebra, then $\text{Frac}(\mathcal{R}) = \mathcal{L}$ is a differential field with constants equal to \mathcal{C} .

Proof. The proof of the second part relies on the assumption that \mathcal{C} is algebraic closed, we will postpone it a little.

Proof of (1) Pick $a \in \mathcal{R}$ and $a \neq 0$. Consider the ideal $I = \{b \in \mathcal{R} | a^n \cdot b = 0, \text{ for some } n \geq 1\}$. This is a differential ideal. (Check it by derive it and multiply it with a) If a is not nilpotent, then $1 \notin I$ then I has to be 0, so a is not a zero divisor. It follows that if a is a zero divisor, then a has to be nilpotent.

Let $I \subset \mathcal{R}$ be the nil radical of \mathcal{R} . Again, I is a diff. ideal. But $1 \notin I$, I has to be $\{0\}$. \square

Lemma 2.6. *(Criterion for Algebraicity) Let \mathcal{K} be a field of char 0. \mathcal{R} is a finitely generated \mathcal{K} -algebra, integral domain, and suppose that $x \in \mathcal{R}$ is such that the set $S := \{c \in \mathcal{K} | x - c \in \mathcal{R}^\times\}$ is infinite. Then x is algebraic over \mathcal{K} .*

For example $\mathcal{K} = \bar{\mathbb{Q}}$, $\mathcal{R} = \bar{\mathbb{Q}}[x, y, 1/x, 1/(x^2 + 2)]$. For $x - c$ to be unit, then only possibility is $c = 0$, which means x is not algebraic over $\bar{\mathbb{Q}}$. While in the case $\mathcal{K} = \mathbb{Q}$, $\mathcal{R} = \mathbb{Q}[\sqrt{2}]$, $\sqrt{2} - c$ is always unit in \mathcal{R}^\times , which means $\sqrt{2}$ is algebraic over \mathbb{Q} .

Proof. Say $\mathcal{R} = \mathcal{K}[x_1, \dots, x_n]$ and w.l.o.g. $x_1 = x$. Set $\mathcal{L} = \text{Frac}(\mathcal{R})$ (\mathcal{R} is integral domain), suppose x is **not** algebraic over \mathcal{K} , so x is transcendental. Suppose w.l.o.g. that Reorder x_2, \dots, x_n such that x_1, \dots, x_r are a transcendence base for \mathcal{L}/\mathcal{K} , i.e. x_1, \dots, x_r are algebraic independent and $\mathcal{L}/\mathcal{K}(x_1, \dots, x_r)$ is a finite algebraic extension. Recall the [Lemma of primitive element](#) (every finite field extension of char 0, can be generated by one element). Pick $y \in \mathcal{R}$ such that $\mathcal{L} = \mathcal{K}(x_1, \dots, x_r)[y]$, look at the minimal polynomial of y over $\mathcal{K}[x_1, \dots, x_r]$

$$a_N(x_1, \dots, x_r)T^N + a_{N-1}(x_1, \dots, x_r)T^{N-1} + \dots,$$

where $a_i \in \mathcal{K}[x_1, \dots, x_r]$

Pick $G \in \mathcal{K}[x_1, \dots, x_r]$ s.t.

- 1) $a_N | G$ and
- 2) $x_1, \dots, x_n \in \mathcal{K}[x_1, \dots, x_r, y, G^{-1}]$

For $s > r$, $x_s \in \mathcal{R} \subseteq \mathcal{L} = \text{Frac}(\mathcal{K}[x_1, \dots, x_r])[y]$

$$x_s = \frac{P_s(x_1, \dots, x_r, y)}{Q_s(x_1, \dots, x_r)} = \frac{\tilde{P}_s(x_1, \dots, x_r, y)}{G}$$

G has to be a multiple of all those denominators Q_s , it is always possible to pick such a G .

Since the set $S \subseteq \mathcal{K}$ is infinite, we can find $s_1, \dots, s_r \in S$ with $G(s_1, \dots, s_r) \neq 0$. Fix such elements $s_1, \dots, s_r \in S$, we can define a ring homomorphism $\mathcal{K}[x_1, \dots, x_r, y, G^{-1}] \xrightarrow{\varphi} \bar{\mathcal{K}}$ where $x_i \mapsto s_i$, $y \mapsto$ any root of the minimal polynomial evaluated in s_i . $a_N(s_1, \dots, s_r)T^N + \dots$ and $G^{-1} \mapsto G(s_1, \dots, s_r)^{-1}$. since $G(s_1, \dots, s_r) \neq 0$ also we have $a_N(s_1, \dots, s_r) \neq 0$ (The minimal polynomial of y with coefficients evaluated in s_i indeed has nontrivial roots in $\bar{\mathcal{K}}$). The ring homomorphism is well-defined and $\mathcal{R} \subseteq \mathcal{K}[x_1, \dots, x_r, y, G^{-1}]$ $\varphi(x_1 - s_s) = 0$, where $(x_1 - s_s)$ is invertible in \mathcal{R} , which makes the contradiction. \square

Lemma 2.7. (Second half of Lemma 2.5) \mathcal{K} is a differential field and \mathcal{C} is the field of constant, $\mathcal{C} = \bar{\mathcal{C}}$ and $\text{char} \mathcal{C} = 0$. \mathcal{R}/\mathcal{K} simple differential ring which is finitely generated as \mathcal{K} -algebra. \implies the field of constants of \mathcal{R} is \mathcal{C} .

Proof. We already know \mathcal{R} is an integral domain. Let $\mathcal{L} = \text{Frac}(\mathcal{R})$, fix $a \in \mathcal{L}, a \neq 0, a' = 0$. Suppose $a \notin \mathcal{C}$, consider the ideal $I := \{b \in \mathcal{R} | a \cdot b \in \mathcal{R}\} \subseteq \mathcal{R}$. This is a differential ideal because $b \in I \implies ab' = a'b + ab' = (ab)' \in \mathcal{R}$. By the assumption \mathcal{R} is simple differential ring $\implies I = \mathcal{R}$. Then $1 \in I \implies a \cdot 1 \in \mathcal{R}$. a has an inverse in \mathcal{L} , denote it by c . Then $e \neq 0, e' = 0$ we can proceed the similar construction $J := \{b \in \mathcal{R} | e \cdot b \in \mathcal{R}\} \subseteq \mathcal{R}$ it also indicates that $e \in \mathcal{R}$, hence we get the conclusion that $a \in \mathcal{R}^\times$

Same argument for $a + c$ for any $c \in \mathcal{C}$ shows $(a + c) \in \mathcal{R}^\times, \forall c \in \mathcal{C} \implies a$ is algebraic over $\mathcal{K} \xrightarrow{\text{Exercise 1.7}} a$ is algebraic over $\mathcal{C} = \overline{\mathcal{C}} \implies a \in \mathcal{C}$ \square

Proposition 2.8. \mathcal{K} is a differential field with constants $\mathcal{C} = \overline{\mathcal{C}}$ Let $u' = Au$ be a matrix differential equation over \mathcal{K} .

- (1) A Picard-Vessiot extension for $u' = Au$ exists.
- (2) Any two P-V extension for $u' = Au$ are isomorphic.
- (3) The field of constant of any P-V extension is $\mathcal{C} = \overline{\mathcal{C}}$

Proof. The previous Lemma \implies (3)

For (1) consider the ring $\mathcal{R}_0 = \mathcal{K}[(X_{ij})_{1 \leq i, j \leq n}, \det(X)^{-1}]$. Define a differentiation on \mathcal{R}_0 by

$$X' = AX$$

$$X'_{ij} = (AX)_{ij} \text{ a polynomial in } \mathcal{K}[X_{ij}, \dots, X_{nn}]$$

and together with the Leibnitz rule it is a well-defined differentiation on \mathcal{R}_0 .

Pick any maximal differential ideal $I \subseteq \mathcal{R}_0$ and set $\mathcal{R} = \mathcal{R}_0/I$. \mathcal{R} is a P-V ring:

Simple because I is maximal.

Fundamental matrix of solutions is X (the classes of X in \mathcal{R}_0/I)

\mathcal{R} is generated by X_{ij} and $\det(X)^{-1}$.

For (2) Let $\mathcal{R}_1, \mathcal{R}_2$ be P-V rings. Consider $\mathcal{R} = \mathcal{R}_1 \otimes \mathcal{R}_2$ with differential $(a \otimes b)' = a' \otimes b + a \otimes b'$. Choose $I \subseteq \mathcal{R}$ maximal differential ideal. Consider $\varphi_1 : \mathcal{R}_1 \longrightarrow \mathcal{R}/I | \varphi_1(a) = a \otimes 1$ and $\varphi_2 : \mathcal{R}_2 \longrightarrow \mathcal{R}/I | \varphi_2(1) = 1 \otimes b$. φ_1 and φ_2 are morphism of differential rings and since $\mathcal{R}_1, \mathcal{R}_2$ are simple, φ_1, φ_2 are injective. Let $V_1 \in M_n(\mathcal{R}_1), V_2 \in M_n(\mathcal{R}_2)$ be fundamental matrices of solution of $u' = Au$. $\varphi_1(V_1)$ and $\varphi_2(V_2)$ are fundamental matrices of solution in \mathcal{R}/I . \mathcal{R}/I is simple finitely generated \implies constants in \mathcal{R}/I are \mathcal{C} . $\exists S \in GL_n(\mathcal{C})$ with $\varphi_1(V_1) = \varphi_2(V_2)S$

$\varphi_1(\mathcal{R}_1)$ is isomorphic to the algebra in \mathcal{R}/I generated by $\varphi_1(V_{1,ij})$ and $\varphi_1(\det(V_1))^{-1}$ = the algebra in \mathcal{R}/I generated by $\varphi_2(V_{2,ij})$ and $\varphi_2(\det(V_2))^{-1}$ $\cong \varphi_2(\mathcal{R}_2)$

Then $\mathcal{R}_1 \cong \varphi_1(\mathcal{R}_1) = \varphi_2(\mathcal{R}_2) \cong \mathcal{R}_2$ \square

3 The Differential Galois Groups

Assumption: \mathcal{K} - differential field with $\text{char}0$, \mathcal{C} is the set of constants in \mathcal{K} and $\mathcal{C} = \overline{\mathcal{C}}$.

Definition 3.1. Let \mathcal{R} be a Picard-Vessiot ring of a differential equation $u' = Au$ or of a differential module (M, ∂) over \mathcal{K} . We call **Galois group of the equation/ module** the group $\text{Aut}^\partial(\mathcal{R}/\mathcal{K}) = \{\mathcal{K}\text{-algebra isomorphism } \varphi : \mathcal{R} \rightarrow \mathcal{R} \text{ compatible with the differentiations}\}$. Usually we denote it with $\text{Gal}^\partial(\mathcal{R}/\mathcal{K})$

Exercise: Let \mathcal{L}/\mathcal{K} be a finite Galois extension.

- (1) There is unique differentiation on \mathcal{L} extending that of \mathcal{K} .
- (2) Look at \mathcal{L} as a \mathcal{K} -module (differential module), Then a Picard-Vessiot extension for \mathcal{L} is \mathcal{L} as a \mathcal{K} -algebra.
- (3) $\text{Gal}^\partial(\mathcal{L}/\mathcal{K}) = \text{Gal}(\mathcal{L}/\mathcal{K})$

$\text{Gal}^\partial(\mathcal{R}/\mathcal{K})$ can be seen as a subgroup of $GL_n(\mathcal{C})$ Let $V \in GL_n(\mathcal{R})$ be a fundamental matrix of solutions. Pick $g \in G = \text{Gal}^\partial(\mathcal{R}/\mathcal{K})$. then $gV = g(v_{ij})$ is again a fundamental matrix of solutions.

$$(gV)' = gV' = gAV = A(gV)$$

$$g(V) = V \cdot \gamma(g)$$

$\gamma \in GL_n(\mathcal{C})$ is unique, because two fundamental matrices are linked with a unique matrix in $GL_n(\mathcal{C})$ (Remark 1.13). Then we get a group homomorphism:

$$\begin{aligned} \gamma : G &\hookrightarrow GL_n(\mathcal{C}) \\ g &\longmapsto \gamma(g) \end{aligned}$$

It is injective: $\gamma(g) = \mathbb{1} \implies gV = V$, but \mathcal{R} is generated by entries of $V \implies g = \text{id}_{\mathcal{R}} = \mathbb{1}_G$.

What makes differential Galois groups a powerful tool is that they are linear algebraic groups and, moreover, establish a Galois correspondence, analogous to the classical Galois correspondence. Torsors will explain the connection between the Picard-Vessiot ring and the differential Galois group. The Tannakian approach to linear differential equations provides new insight and useful methods. Some of this is rather technical in nature. We will try to explain theorems and proofs on various levels of abstraction.

Example 3.2. $\mathcal{K} = \mathbb{C}(t), \mathcal{C} = \mathbb{C} \ u' = Au$

$$A = \begin{pmatrix} 0 & 1 \\ 0 & -1/t \end{pmatrix}$$

$X := \log(t)$. The Picard-Vessiot ring $\mathcal{R} = \mathbb{C}(t)[X]$, with differential defined by $X' = \frac{1}{t} + \text{Leibnitz}$. $\text{Aut}^\partial(\mathcal{R}/\mathcal{K}) \ni g$, the action of g is $\mathbb{C}(t)$ -linear.

$$\begin{aligned} g : \mathbb{C}(t)[X] &\Longrightarrow \mathbb{C}(t)[X] \\ X &\longmapsto g(X) \end{aligned}$$

It is compatible with the differentiation

$$\begin{aligned} g(X)' &= g(X') = g(1/t) = 1/t \\ g(X) &= X + a, \quad a \in \mathbb{C} \end{aligned}$$

For a fundamental matrix

$$V = \begin{pmatrix} 1 & X \\ 0 & 1/t \end{pmatrix}$$

$$g(V) = \begin{pmatrix} 1 & X+a \\ 0 & 1/t \end{pmatrix} = \begin{pmatrix} 1 & X \\ 0 & 1/t \end{pmatrix} \cdot \begin{pmatrix} 1 & a \\ 0 & 1 \end{pmatrix}$$

Then

$$\text{Gal}^\partial(\mathcal{R}/\mathcal{K}) = \left\{ \begin{pmatrix} 1 & a \\ 0 & 1 \end{pmatrix} \middle| a \in \mathbb{C} \right\} \cong (\mathbb{C}, +)$$

4 Algebraic Group: A Detour

Last week, we introduced differential Galois group, where \mathcal{K} is a differential field with algebraic closed constants $\mathcal{C} = \overline{\mathcal{C}}$. And the Picard-Vessiot extension (for equation or module). $\text{Gal}(\mathcal{R}/\mathcal{K}) = \text{Aut}_{\mathcal{K}}^\partial(\mathcal{R})$. If $F \in \text{Cl}_n(\mathcal{R})$ is a fundamental matrix, $g \in \text{Gal}(\mathcal{R}/\mathcal{K})$, then gF is again a fundamental matrix,

hence of the form $gF = F\gamma(g)$, where $\gamma(g) \in GL_n(\mathcal{C})$. Then we obtain the embedding of $Gal(\mathcal{R}/\mathcal{K}) \xrightarrow{\gamma} GL_n(\mathcal{C})$.

Intrinsic variant:

if \mathcal{R} is the Picard-Vessiot extension associated to a differential module M , then the $Gal(\mathcal{R}/\mathcal{K})$ acts on $\mathcal{R} \otimes M$ and leaves

$$V = Ker(\mathcal{R} \otimes M \xrightarrow{\partial} \mathcal{R} \otimes M)$$

invariant. Which says $Gal(\mathcal{R}/\mathcal{K})$ acts on V , get

$$\gamma : Gal \hookrightarrow GL(V)$$

Exercise 4.1. \mathcal{R} is the Picard-Vessiot extension of \mathcal{K} $\mathcal{L} = \text{frac}(\mathcal{R})$, show that

$$Aut_{\mathcal{K}}^{\partial}(\mathcal{L}) = Aut_{\mathcal{K}}^{\partial}(\mathcal{R}) = Gal$$

Definition 4.2. A subgroup $G \subseteq GL_n(\mathcal{C})$ is **algebraic** if there exists polynomials $f_1, \dots, f_N \in \mathcal{C}[x_{11}, \dots, x_{nn}, \det(X)^{-1}]$ such that $G = \{g \in GL_n(\mathcal{C}) | f_i(g) = 0 \forall i = 1, \dots, N\}$

Alternatively, because $\mathcal{C}[X, \dots, X]$ is Noetherian:

Definition 4.3. A subgroup $G \subseteq GL_n(\mathcal{C})$ is **algebraic** if there exists an ideal $\mathcal{I} \subset \mathcal{C}[x_{11}, \dots, x_{nn}, \det(X)^{-1}]$ s.t. $G = \{g \in GL_n(\mathcal{C}) | f(g) = 0, \forall f \in \mathcal{I}\}$. One can suppose that $\mathcal{I} = \sqrt{\mathcal{I}} = \{f | f^N \in \mathcal{I}\}$

Example 4.4. • $G = GL_2(\mathcal{C}), \mathcal{I} = \{0\}$

• $SL_2(\mathcal{C}), \mathcal{I} = \langle X_{11}X_{22} - X_{12}X_{21} - 1 \rangle$

•

$$\begin{pmatrix} 1 & * \\ 0 & 1 \end{pmatrix}$$

$$\mathcal{I} = \langle X_{21}, X_{11} - 1, X_{22} - 1 \rangle$$

FACT:

$G = GL_n(\mathcal{C})$ algebraic, then G is the zero set of the ideal $\{f \in \mathcal{C}[X_{11}, \dots, X_{nn}, \det(X)^{-1}] | f(g) = 0, \forall g \in G\}$, i.e $G = \{g \in GL_n(\mathcal{C}) | f(g) = 0 \forall f \in \mathcal{I}\}$. This is what we call **Hilbert Nullstaltensatz** (\mathcal{C} is algebraically closed.)

Consider the algebra $\mathcal{O}_G = \mathcal{A} = \mathcal{C}[X_{11}, \dots, X_{nn}, \det(X)^{-1}]/\mathcal{I}$, where we assume $\mathcal{I} = \sqrt{\mathcal{I}}$ is called **algebra of regular functions** on G . Given $a \in \mathcal{A}, a = [f], f \in \mathcal{C}[X_{11}, \dots]$ s.t.

$$\begin{aligned} G &\subseteq GL_n(\mathcal{C}) \longrightarrow \mathcal{C} \\ g &\mapsto G \mapsto f(g) \end{aligned}$$

only depends on a .

Theorem 4.5. *Let \mathcal{R}/\mathcal{K} be a Picard-Vessiot extension with Galois group G . The group G , as a subgroup of $GL_n(\mathcal{C})$, is an algebraic group.*

Proof. W.L.O.G, suppose \mathcal{R} is the P-V ring of an equation $u' = Au$, $A \in M_n(\mathcal{K})$. $\mathcal{R} = \mathcal{R}_0/\mathcal{I}$, where $\mathcal{R}_0 = \mathcal{K}[X_{11}, \dots, X_{nn}, \det(X)^{-1}]$, $X' = AX$ $\mathcal{I} \subseteq \mathcal{R}_0$ maximal differential ideal. Choose generators h_1, \dots, h_N of $\mathcal{I} \subseteq \mathcal{R}_0$. Choose $(e_s)_{s \in S}$ a \mathcal{C} -basis of \mathcal{R} . Can identify the $G \subseteq GL_n(\mathcal{C})$ with the group of matrices M for which

$$\begin{aligned} \sigma_M : \mathcal{R}_0 &\longrightarrow \mathcal{R}_0 \\ X_{ij} &\longmapsto (X \cdot M)_{ij} \end{aligned}$$

sends \mathcal{I} to \mathcal{I} . Write $\sigma_M(h_i) \bmod \mathcal{I} = \sum_{s \in S} c_{i,s}(M)e_s$, where $c_{i,s}(M)$ is uniquely determined.

Claim: $c_{is}(M)$ is a polynomial expression in the coefficients $M_{ij}, \det(M)^{-1}$. i.e. $\exists c_{is}(M) \in \mathcal{C}[Y_{11}, \dots, Y_{nn}, \det(Y)^{-1}]$. Knowing this $G = \{M \in GL_n(\mathcal{C}) | c_{is}(M) = 0\}$. $G = \{M \in GL_n(\mathcal{C}) | c_{is}(M) = 0 \forall i, s\}$ = zero set of the ideal generated by $c_{is}(Y) \subseteq \mathcal{C}[Y_{11}, \dots, \det(Y)^{-1}]$.

Verification of claim, For a \mathcal{C} -algebra \mathcal{B} , set $G(\mathcal{B}) = \{\mathcal{B} \otimes \mathcal{K}$ -linear automorphism of $\mathcal{B} \otimes \mathcal{R}$ replacing t by $t + \sigma_M(t)$ for $M \in GL_n(\mathcal{B})$. Alternatively, $G(\mathcal{B})$ is the group of these matrices $M \in GL_n(\mathcal{B})$ s.t.

$$\begin{aligned} \sigma_M : \mathcal{B} \otimes \mathcal{R}_0 &\longrightarrow \mathcal{B} \otimes \mathcal{R}_0 \\ X &\longmapsto XM \end{aligned}$$

sends $\mathcal{B} \otimes \mathcal{I}$ to $\mathcal{B} \otimes \mathcal{I}$

Given a morphism $\mathcal{B}_1 \xrightarrow{\varphi} \mathcal{B}_2$, we get a group homomorphism

$$\begin{aligned} G(\mathcal{B}) &\longrightarrow G(\mathcal{B}_2) \\ M &\longmapsto \varphi(M) \end{aligned}$$

Notice:

$$\begin{array}{ccc} \mathcal{B}_1 \otimes \mathcal{R}_0 & \xrightarrow{\sigma_M} & \mathcal{B}_1 \otimes \mathcal{R}_0 \\ \downarrow \varphi & & \downarrow \varphi \\ \mathcal{B}_2 \otimes \mathcal{R}_0 & \xrightarrow{\sigma_{\varphi(M)}} & \mathcal{B}_2 \otimes \mathcal{R}_0 \end{array}$$

For $M \in GL_n(\mathcal{B})$ have $\sigma_M(1 \otimes h_i) \bmod \mathcal{B} \otimes \mathcal{I} = \sum_{s \in S} c_{is}(M) e_s$ $c_{is} \in \mathcal{B}$ unique. Given a morphism $\varphi : \mathcal{B} = \mathcal{B}_1 \rightarrow \mathcal{B}_2$, get: Apply φ to relation, or consider relation for $\varphi(M)$, Get $\varphi(c_{is}(M)) = c_{is}(\varphi(M))$. Consider the special case $\mathcal{B} = \mathcal{C}[Y_{11}, \dots, Y_{nn}, \det(Y)^{-1}] \xrightarrow{\varphi} \mathcal{C} : Y \mapsto M$. By the previous result we have $c_{is}(M) = c_{is}(\varphi(Y)) = \varphi_{is}(c_{is}(Y))$ \square

Back to the algebraic groups. Let $G \subseteq GL_n(\mathcal{C})$ be an algebraic group ideal $\mathcal{I} = \sqrt{\mathcal{I}}$, algebra of functions \mathcal{A} .

$$G = Hom_{\mathcal{C}}(\mathcal{A}, \mathcal{C})$$

$$g \in G \mapsto [ev_g : \mathcal{A} \rightarrow \mathcal{C} | f \mapsto f(g)]$$

$$\varphi(x) \leftarrow [\varphi : \mathcal{A} \rightarrow \mathcal{C}]$$

For a \mathcal{C} -algebra \mathcal{B} , consider $G(\mathcal{B}) = Hom_{\mathcal{C}}(\mathcal{A}, \mathcal{B}) \hookrightarrow GL_n(\mathcal{B}) : \varphi \mapsto \varphi(X)$. FACT: $G(\mathcal{B}) \subseteq GL_n(\mathcal{B})$ is a subgroup.

Explanation: Consider the following algebra morphisms:

$$\mathcal{C}[X_{11}, \dots, \det^{-1}] \rightarrow \mathcal{C}[Y_{ij}, \det(Y)^{-1}] \otimes \mathcal{C}[Z_{ij}, \det(Z)^{-1}] = \mathcal{C}[Y_{ij}, Z_{ij}, \det(Y)^{-1}, \det(Z)^{-1}]$$

~~~~~1

Get from  $\mu$  a composition law on  $Hom_{\mathcal{C}}(\mathcal{A}, \mathcal{B})$ , For  $\varphi, \psi \in Hom_{\mathcal{C}}(\mathcal{A}, \mathcal{B})$ , define  $\varphi \cdot \psi$  as

$$\mathcal{A} \xrightarrow{\mu} \mathcal{A} \otimes \mathcal{A} \xrightarrow{\varphi \otimes \psi} \mathcal{B} \otimes \mathcal{B} \xrightarrow{\text{multiplication}} \mathcal{B}$$

This is compatible with the group law in  $GL_n(\mathcal{B}) \supseteq G(\mathcal{B}) = Hom_{\mathcal{C}}(\mathcal{A}, \mathcal{B})$ .  $\mathcal{A}$  is a Hopf Algebra

Then the algebraic group  $G$  defines a functor  $G\{\mathcal{C}\text{-algebra}\} \rightarrow \{\text{Groups}\}$ ,  $\mathcal{B} \mapsto G(\mathcal{B}) = Hom_{\mathcal{C}}(\mathcal{A}, \mathcal{B})$ .

**Definition 4.6.** A representable functor  $G\{\mathcal{C}\text{-algebra}\} \rightarrow \{\text{Groups}\}$ ,  $\mathcal{B} \mapsto G(\mathcal{B}) = Hom_{\mathcal{C}}(\mathcal{A}, \mathcal{B})$  for some unique Hopf algebra  $\mathcal{A}$ . is called an **Affine group scheme**.  $G = Spec(\mathcal{A})$ . Moreover, if  $\mathcal{A}$  is finitely generated, then we call  $G$  a **linear algebraic group over  $\mathcal{C}$** .

#### 4.1 Application of the fact that differential Galois group is a algebraic group

laast week:  $\mathcal{R}/\mathcal{K}$  Picard-Vessiot extension with Galois group  $G = GL_n(\mathcal{C})$ . The main message of last week is that  $G$  is a algebraic group. you can see it in different ways.

$$G = Spec(\mathcal{A}),$$

where  $\mathcal{A}$  is a Hopf Algebra. Or equivalently, we can see  $G$  as a functor:

$$\{\mathcal{C}\text{-algebra}\} \longrightarrow \{\text{Groups}\}$$

$$\mathcal{B} \longmapsto G(\mathcal{B}) = \text{Hom}_{\mathcal{C}\text{-Alg}}(\mathcal{A}, \mathcal{B}).$$

Call **dimension** of  $G$  the transcendence degree of  $\mathcal{A}$  over  $\mathcal{C}$ .

$$\dim(G) \leq n^2$$

This notion of dimension coincides with the intuitive ideas, and with actual dimension as a complex variety if  $\mathcal{C} = \mathbb{C}$ .

denote by  $G_{\mathcal{K}}$  the base-change of  $G$  to  $\mathcal{K}$ , i.e.

$$G_{\mathcal{K}} = \text{Spec}(\mathcal{K} \otimes \mathcal{A})$$

$$\{\mathcal{K}\text{-alg}\} \longrightarrow \{\text{Groups}\}$$

$$\mathcal{L} \longmapsto \text{Hom}_{\mathcal{K}}(\mathcal{K} \otimes \mathcal{A}, \mathcal{L}) = \text{Hom}_{\mathcal{C}}(\mathcal{A}, \mathcal{L}).$$

Set  $X = \text{Spec}(\mathcal{R})$ , and consider a functor from  $\mathcal{K}$ -algebra to sets:

$$\{\mathcal{K}\text{-alg}\} \longrightarrow \{\text{Sets}\}$$

$$\mathcal{L} \longmapsto \text{Hom}_{\mathcal{K}}(\mathcal{R}, \mathcal{L})$$

and in fact, for every  $\mathcal{K}$ -algebra  $\mathcal{L}$ , the group  $G_{\mathcal{K}}(\mathcal{L})$  acts on the set  $X(\mathcal{L})$

$$\begin{array}{ccc} G_{\mathcal{K}}(\mathcal{L}) \times X(\mathcal{L}) & \longrightarrow & X(\mathcal{L}) \\ \parallel & & \parallel \\ \text{Hom}_{\mathcal{C}}(\mathcal{A}, \mathcal{L}) \times \text{Hom}_{\mathcal{K}}(\mathcal{R}, \mathcal{L}) & \longrightarrow & \text{Hom}_{\mathcal{K}}(\mathcal{R}, \mathcal{L}) \\ \parallel & & \parallel \\ G(\mathcal{L}) \times \text{Hom}_{\mathcal{L}}(\mathcal{R} \otimes \mathcal{L}, \mathcal{L}) & \longrightarrow & \text{Hom}_{\mathcal{L}}(\mathcal{R} \otimes \mathcal{L}, \mathcal{L}) \end{array}$$

where  $G(\mathcal{L}) = \text{Aut}(\mathcal{R} \otimes \mathcal{L}|_{\mathcal{L}})$

$$(g, x) \longmapsto [\mathcal{R} \otimes \mathcal{L} \xrightarrow{g} \mathcal{R} \otimes \mathcal{L} \xrightarrow{x} \mathcal{L}]$$

This action is natural in  $\mathcal{L}$ , An action of the algebraic group  $G$  on the algebraic variety  $X$ .

**Theorem 4.7.** For every  $\mathcal{K}$ -algebra  $\mathcal{L}$  either  $X(\mathcal{L}) = \emptyset$  or  $G(\mathcal{L})$  acts simply transitively on  $X(\mathcal{L})$ . In other words:

$$G \times X \longrightarrow G \times X$$

$$(g, x) \longmapsto (g, gx)$$

is a bijection for all  $\mathcal{L}$ , i.e. an isomorphism of functors or an isomorphism of affine algebra varieties.  $X$  is a  $G$ -torsor.

**Corollary 4.8.**  $\dim_{\mathcal{C}}(G) = \dim_{\mathcal{K}}(G_{\mathcal{K}}) = \dim_{\mathcal{K}}(X) = \text{tr.deg}(\mathcal{R}/\mathcal{K})$

**Example 4.9.** As an illustration:  $\mathcal{K} = \mathbb{C}(t), \mathcal{R} = \mathbb{C}(1, \log(t))$ . remember that

$$G = \begin{pmatrix} 1 & * \\ 0 & 1 \end{pmatrix}$$

and  $\dim(G) = 1$ , then  $\text{tr.deg}(\mathbb{C}(t, \log(t))) = 1 \implies \log(t)$  is not algebraic over  $\mathbb{C}(t)$

**Exercise 4.10.** Consider similarly,  $\mathcal{R} = (t, \exp(t))$ ,  $G = ?$

## 5 The Galois-correspondence of differential equations

Recall our setting:  $\mathcal{K}$  differential field,  $\mathcal{C}$  is constants,  $\mathcal{C} = \overline{\mathcal{C}}$  with  $\text{char} = 0$

**Theorem 5.1.** Let  $\mathcal{R}/\mathcal{K}$  be a Picard-Vessiot extension, with Galois group  $G$ . Set  $\mathcal{L} = \text{frac}(\mathcal{R})$ .

$$\left\{ \begin{array}{l} \text{Subfields } M \subseteq \mathcal{L} \\ \text{containing } \mathcal{K}, \text{ stable} \\ \text{under differentiation} \end{array} \right\} \longleftrightarrow \{ \text{algebraic subgroups } H \subseteq G \}$$

$$M \longmapsto \text{Gal}(\mathcal{L}/M) \subseteq \text{Gal}(\mathcal{L}/\mathcal{K}) = G$$

$$\mathcal{L}^H \longleftarrow H$$

where  $\mathcal{L}^H$  is the subfield of  $\mathcal{L}$  which is fixed under every element of  $H$ . There maps are bijection inverse of each other.

If  $N \subseteq G$  is normal subgroup, then the map  $G \longrightarrow \text{Gal}(\mathcal{L}^N/\mathcal{K})$  is surjective, and  $\mathcal{L}^N$  is a Picard-Vessiot extension with Galois group  $G/N$ .



**Proposition 5.2.** *In the setup of the theorem*

$$L^G = \mathcal{K}$$

*Proof.* The inclusion  $\supseteq$  is evident. For the converse inclusion. Pick  $x \in \mathcal{L}, x \notin \mathcal{K}$  and we have to produce  $g \in G$  with  $gx \neq x$ . Set  $x = \frac{p}{q}, p, q \in \mathcal{R}, q \neq 0$ . Consider  $r = p \otimes q - q \otimes p \in \mathcal{R} \otimes \mathcal{R}$ .

Two easy facts:  $r \neq 0$  (because  $\frac{p}{q} \notin \mathcal{K}$ ) and  $\mathcal{R} \otimes \mathcal{R}$  has no nilpotents (Char=0)

Choose a maximal differential ideal  $J$  in  $\mathcal{R} \otimes \mathcal{R}[1/r]$ . Have two canonical morphisms of differential algebras

$$\begin{aligned} \mathcal{R} &\xrightarrow{\varphi_1} \frac{(\mathcal{R} \otimes \mathcal{R})[1/r]}{J} \xleftarrow{\varphi_2} \mathcal{R} \\ a &\longmapsto (a \otimes 1), (1 \otimes b) \longleftarrow b \end{aligned}$$

$\varphi_1, \varphi_2$  are injective ( $\mathcal{R}$  simple) and images of  $\varphi_1, \varphi_2$  are equal (c.f. proof of unicity of P-V-extension) say  $S$

$$\begin{array}{ccccc} \mathcal{R} & \xrightarrow{\varphi_1} & S & \xleftarrow{\varphi_2} & \mathcal{R} \\ & \searrow & & \nearrow & \\ & & g = \varphi_2^{-1} \circ \varphi_1 & & \end{array}$$

$g \in \text{Gal}(\mathcal{R}/\mathcal{K}) = G$ . The class of  $r$  in  $\mathcal{R} \otimes \mathcal{R}(1/r)/J$  is the unit

$$\begin{aligned} 0 \neq [p \otimes p - q \otimes p] &= \varphi_1(p)\varphi_2(q) - \varphi_1(q) - \varphi_2(p) \\ &= \varphi_1(p \cdot (gq)) - \varphi_1(q \cdot (gp)) \\ &\implies p \cdot gq \neq q \cdot gp \\ &x \neq gx \end{aligned}$$

□

## 6 local theory of differential equations

Setups:  $\mathcal{C} = \overline{\mathcal{C}}$  field of  $\text{char} = -$ .  $\mathbb{C} = \mathcal{A}$ . choose  $\xi_m \in \mathcal{C}$ -primitive  $m$ -th roots of unity.  $\xi_m^n = \xi_{m/n}$  if  $n|m$ . e.a.  $xi_m = e^{2\pi i/m}$ .  $\mathcal{K} = \mathcal{C}((t)) = \text{frac}(\mathcal{C}[[t]])$ ??? with usual derivative. Goal: classify differential equation of differential modules over  $\mathcal{K}$ .

Ingredients to classification:

1. Set  $\mathfrak{D} = \mathcal{K}[\partial]$  the noncommutative ring of polynomial expressions differential operators  $\mathcal{L} = a_n \partial^n + a_{n-1} \partial^{n-1} + \dots + a_0$ .  $\mathfrak{D}$  = the ring of differential operators. multiplication:  $\partial \cdot a = a' + a\partial, a \in \mathcal{K}$ . Differential modules over *calk* are the same as left  $\mathfrak{D}$ -modules. Given  $\epsilon \in \mathfrak{D}$ , get a  $\mathfrak{D}$ -module

$$\mathcal{K}[\partial]/\mathcal{K}[\partial] \cdot \mathcal{L} = \mathfrak{D}/\mathfrak{D}L$$

This  $\mathfrak{D}$ -modules is the differential module of the equation  $L(u) = 0$ .

2. Finite field extension of  $\mathcal{K}$ . Let  $m \geq 1$  integer. Set  $\mathcal{K}_m = \mathcal{C}((t^{1/m})) = \mathcal{C}((t)[s])/\langle s^m - t \rangle$ . The field extension  $\mathcal{K}_m/\mathcal{K}$  is Galois with group  $\mathbb{Z}/m\mathbb{Z}$ . For  $[a] \in \mathbb{Z}/m\mathbb{Z}$  consider the automorphism

$$\begin{aligned} \mathcal{K}_m &\xrightarrow{\epsilon(a)} \mathcal{K}_m \\ t^{1/m} &\longmapsto \xi_m^a \cdot t^{1/m}, \end{aligned}$$

where  $\epsilon : \mathbb{Z}/m\mathbb{Z} \xrightarrow{\cong} \text{Gal}(\mathcal{K}_m/\mathcal{K})$  isomorphism.

If  $n|m$ , then  $\mathcal{K}_n \subseteq \mathcal{K}_m$ . Can consider  $\overline{\mathcal{K}} = \cup_{m \geq 1} \mathcal{K}_m$ .

**Theorem 6.1.**  $\overline{\mathcal{K}}$  is algebraically closed!

$$\text{Gal}(\overline{\mathcal{K}}|\mathcal{K}) = \lim_m \mathbb{Z}/m\mathbb{Z} = \widehat{\mathbb{Z}}$$

## 7 Personal Reads

### 7.1 Monodromy, Riemann-Hilbert Problem

$U$  is an open connected subset of the complex sphere  $\mathcal{P}^1$  and let  $Y' = AY$  be a differential equation on  $U$ , with  $A$   $n \times n$  matrix with coefficients are meromorphic functions on  $U$ . We assume that the equation is regular at every point  $p \in U$  (might be regular singular point (which means the growth of solutions is bounded)). Let  $F_p$  be a matrix whose columns are the  $n$  independent solutions, then  $F_p$  is a fundamental matrix with entries in  $\mathbb{C}(\{z - p\})$

**Does there exist on all of  $U$ , a solution space for the equation having dimension  $n$**

**Definition 7.1.** Two differential operator  $\frac{d}{dz} - A$  and  $\frac{d}{dz} - B$  are called equivalent if there exists a  $F \in GL_n(\mathcal{K})$  s.t.,  $F^{-1}(\frac{d}{dz} - A)F = \frac{d}{dz} - B$ . A matrix differential operator  $\frac{d}{dz} - A$  is called **regular singular** if the equation is equivalent to  $\frac{d}{dz} - B$  such that the entries of  $B$  have poles at  $z = 0$  of order at most 1.